

Factors Affecting Tuberculosis Patients' Quality of Life in Surabaya, Indonesia

This article was published in the following Dove Press journal:
Journal of Multidisciplinary Healthcare

Ni Njoman Juliasih¹
Ni Made Mertaniasih²
Cholichul Hadi³
Soedarsono⁴
Reny Mareta Sari¹
Ilham Nur Alfian⁵

¹Laboratory of Tuberculosis, Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia; ²Department of Medical Microbiology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia; ³Department of Industrial and Organizational Psychology, Faculty of Psychology, Universitas Airlangga, Surabaya, Indonesia; ⁴Department of Pulmonology and Respiratory Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia; ⁵Department of Personality and Social Psychology, Faculty of Psychology, Universitas Airlangga, Surabaya, Indonesia

Purpose: Patients with tuberculosis need to religiously take medication daily. However, they experience several side effects from these medications. The main reason for measuring the quality of life is to explain closely related factors that affect the patient's daily life that have been compromised with illness, while considering a patients' well-being that has associations with individual characteristics.

Patients and Methods: This study included 157 patients with tuberculosis at 5 primary health-care centers and 2 hospitals in Surabaya. Quality of life is determined based on eight domains: general health, pain, social functioning, physical functioning, role limitation due to physical health, role limitation due to emotional problems, energy, and emotional well-being. The research instrument used to measure the quality of life is the RAND-36 Item Health Survey, whereas that used to measure mental distress is the Self-Reporting Questionnaire.

Results: Our study results show that, of the eight domains measuring the quality of life, only age exhibited a significant effect on general health ($P = 0.018$); sex did not significantly affect the quality of life in all domains. The level of education exhibited a significant effect only on role limitation due to emotional problems ($P = 0.014$). Mental distress demonstrated a significant effect on the quality of life in all domains.

Conclusion: There are several factors affecting TB patients' quality of life. The study found that age, level of education, and comorbidity affect quality of life in several domains. However, mental distress affects quality of life in all domains.

Keywords: demographic variables, mental distress, quality of life, tuberculosis

Introduction

Tuberculosis (TB) remains a major challenge in the health sector.¹ In Indonesia, its prevalence in 2018 was 321 per 100,000 people.² The Ministry of Health has set targets for the National TB Control Program to reduce morbidity by 80% in 2030 and 90% in 2035 and to reduce mortality by 90% in 2030 and 95% in 2035. The duration of TB treatment and the side effects that arise influence the daily life of TB patients, which then affects their quality of life.³⁻⁶

The World Health Organization defined quality of life as an individual's perception of his/her position in life within the cultural context and value system in how they live. Quality of life is also related to one's goals, hopes, standards, and concerns.⁷ In addition, it refers to an individual's assessment of his/her satisfaction and meaningfulness in living life.⁸

There are several factors that affect the quality of life of TB patients, including social support, medical factors, psychological factors, demographic factors, and educational and counseling programs.⁹ TB patients tend to have poor quality of life and

Correspondence: Ni Njoman Juliasih
Institute of Tropical Disease,
Universitas Airlangga, Surabaya 60115,
Indonesia
Tel +628113642237
Fax +62315992445
Email njomanjuliasih@staf.unair.ac.id

Ni Made Mertaniasih
Department of Medical Microbiology,
Faculty of Medicine, Universitas Airlangga,
Surabaya 60115, Indonesia
Email nmademertaniasih@gmail.com

a high risk of experiencing depression.¹⁰ Quality of life can also affect a TB patient's adherence to treatment.^{11,12} Social support, age, and education contribute to the quality of life of TB patients.¹³

The quality of life of TB patients is related to psychological domain, environmental conditions, social relationships, and physical conditions (physical domain). Quality of life is also closely connected to psychological and environmental conditions. Age, sex, income, and duration of treatment variables affect the psychological condition aspects of quality of life. Social relations and environmental conditions are only affected by income and treatment duration.¹⁴ The results of the research conducted by Dawar et al on 100 TB patients in 2016 revealed that 32 of the patients were diagnosed with major depressive disorder, including 7 with mild depression, 20 with moderate depression, and 5 with severe depression. The variables that affect the quality of life in the physical condition aspects are income, treatment duration, and employment status.^{15–17}

There are several factors, such as sex, age, marital status, educational level, employment status, monthly income, chronic illness, and body mass index, that significantly affect a person's quality of life.¹⁸ The research conducted by Salodia, Sethi, and Khokhar in 2019 demonstrated that 23.6% of TB patients experience depression.¹⁹ The main reason for measuring the quality of life is to explain closely related factors that affect a patient's daily life that have been compromised with illness, while considering a patients' well-being that has associations with individual characteristics.

Patients and Methods

The current study was conducted as observational research using a cross-sectional approach. A total of 157 TB patients from 5 primary health-care centers and 2 hospitals were included in the study. The sample were all TB patients chosen randomly based on data given by primary health-care centers and hospitals excluding pregnant women and children. The study was approved by the Ethics Committee in Health Research of Dr. Soetomo Hospital with ethical clearance number 1636/KEPK/XI/2019 and was approved on 9 September 2019. The study was non-intervention research, data were obtained by interviewing TB patients as respondents. Before the interview, the respondents were given an explanation of the research and publications to be carried out. All respondents information is kept confidential and only used for

research purposes. After getting an explanation, the respondent is allowed to refuse the interview or resign in the middle of the interview. The respondents gave their written consent and permission for publication of the letters and to participate in the research. We confirm that all the research meets the ethical guidelines and has been submitted to the ethics committee.

Quality of life is determined based on eight domains: general health, pain, social functioning, physical functioning, role limitation due to physical health, role limitation due to emotional problems, energy, and emotional well-being. Quality of life was measured using the RAND 36-item health survey. Table 1 contains an explanation of the domains from the RAND 36-Item Health Survey.

Every domain was scored from 0 to 100. The higher score equating to the better quality of life in every domain. In general health, the higher score showed the better health status. In domain pain the higher score showed that there is less pain and how pain did not affect daily activities. The higher score in social functioning and physical functioning showed that there is no limitation in doing social activities and daily activities. In role limitation due to physical health and role limitation due to emotional problem, the higher score showed that there is no limitation in doing daily activities. In domain energy, the higher score means the more energy or less fatigue they have. In domain emotional well-being, the higher score means the absence of feeling sad or nervousness.

The independent variables that were considered were sex, age, level of education, comorbidities, mental distress,

Table 1 The Explanation of the Domains from RAND 36-Item Health Survey

Domain	Explanation
General health	The evaluation of general health status
Pain	Presence of pain and its limitation due to pain
Social functioning	Limitation in social activities
Physical functioning	Limitation in daily activities
Role limitation due to physical health	Difficulties in doing daily activities due to physical health problem
Role limitation due to emotional problems	Difficulties in doing daily activities due to emotional problem
Energy	Loss energy or presence of fatigue
Emotional well-being	Presence of sad feeling or nervousness

working status, monthly income, and body mass index (BMI). Age variables were classified based on the recommendations of the Ministry of Health in Indonesia: teenagers (15–25 y), young adults (26–35 y), late adults (36–45 y), early elderly (46–55 y), middle elderly (56–65 y), and late elderly (>65 y).² Comorbidities were considered when TB patients were diagnosed with other illnesses in addition to TB. Mental distress variables were measured using the Self-Reporting Questionnaire. Scores were then categorized into a probable non-case (score ≤ 7) and/or probable case (score ≥ 8).²⁰ Working status was categorized as working and not working. Respondents who classified as not working are house wife, student, and people who do not have a job. Body mass index (BMI) was categorized, based on the Indonesian Ministry of Health, into severely underweight (BMI < 17.0); light underweight (BMI = 17.0–18.4); normal weight (BMI = 18.5–25.0); light overweight (BMI = 25.1–27.0); and severely overweight (BMI > 27.0). Statistical analysis was conducted using the General Linear Model with α set at 0.05 to analyze the determinants of quality of life.

Results

Table 2 shows the characteristics of the respondents, including sex, age, educational level, comorbidities, and mental distress.

Table 2 indicates that some respondents were female. The age groups of teenagers (23.6%) and late adults (22.3%) had the highest numbers of respondents, and the highest level of education was senior high school (47.1%). As many as 63.7% of TB patients did not exhibit comorbidity, and 36.9% were suspected of having mental distress. Most of the respondents had a monthly income less than IDR 500,000. The large number of respondents who had low monthly income can be caused by the high number of respondents who do not work (46.5%). Body mass index of TB patients ranged from severely underweight to severely overweight. Most of the respondents had a normal BMI. However, there are 22.3% who were severely underweight and 12.7% who were underweight. It can be seen that TB patients tend to be severely underweight to normal BMI.

Quality of life consists of eight domains: general health, pain, social functioning, physical functioning, role limitation due to physical health, role limitation due to emotional problems, energy, and emotional well-being. In every domain, the higher score means the better quality of life the respondent have.

Table 2 Respondent Characteristics

Characteristics	Category	n	%
Sex	Male	75	47.8
	Female	82	52.2
	Total	157	100
Age	Teenager (15–25 y)	37	23.6
	Young adult (26–35 y)	30	19.1
	Late adult (36–45 y)	35	22.3
	Early elderly (46–55 y)	25	15.9
	Middle elderly (56–65 y)	27	17.2
	Late elderly (>65 y)	3	1.9
	Total	157	100
Level of education	Elementary school	32	20.4
	Junior high school	26	16.6
	Senior high school	74	47.1
	Diploma and above	25	15.9
	Total	157	100
Comorbidity	Yes	57	36.3
	No	100	63.7
	Total	157	100
Mental distress	Probable case (score ≥ 8)	58	36.9
	Probable non-case (score ≤ 7)	99	63.1
	Total	157	100
Working Status	Yes	84	53.5
	No	73	46.5
	Total	157	100
Monthly income Indonesian Rupiah (IDR)	< 500,000	55	35.0
	500,000–1,000,000	21	13.4
	1,000,000–2,000,000	25	15.9
	2,000,000–3,000,000	34	21.7
	>3,000,000	22	14.0
	Total	157	100
Body mass index	Severe underweight	35	22.3
	Underweight	20	12.7
	Normal	87	55.4
	Overweight	6	3.8
	Severe Overweight	9	5.7
	Total	157	100

Table 3 indicates that the physical functioning domain has the highest average value (70.97) and it means that generally the respondents had good physical function and there is no limitation in doing daily activities. Role limitation due to physical health domain has the lowest average

Table 3 Quality of Life of Patients with Tuberculosis

Dependent Variable	Domain	Minimum	Maximum	Mean	Standard Deviation
Quality of life	General health	20	95	63.67	17.184
	Pain	0	100	55.22	29.019
	Social functioning	0	100	70.70	26.468
	Physical functioning	0	100	70.97	24.004
	Role limitation due to physical health	0	100	36.78	41.753
	Role limitation due to emotional problems	0	100	48.30	43.379
	Energy	10	90	59.60	18.322
	Emotional well-being	16	88	65.77	16.450

value (36.78). It means that generally the respondents' physical health tended to affect their daily activities.

Table 4 indicates that sex did not significantly affect quality of life, and only age significantly affected the general health domain ($P = 0.018$). The level of education significantly affected the physical functioning domain ($P = 0.016$) and role limitation due to physical health ($P = 0.020$). Comorbidity had a significant effect on general health ($P = 0.029$), pain ($P = 0.026$), physical functioning ($P = 0.012$), role limitation due to physical health ($P = 0.001$), and role limitation due to emotional problems ($P = 0.024$). Moreover, mental distress significantly affected the quality of life of TB patients in all domains. Working status, monthly income, and BMI did not affect quality of life in every domain.

Discussion

Patients with TB had a lower quality of life in all domains compared with healthy people.²¹ Some research results revealed that sex does not affect the quality of life of TB

patients.^{13,22} Quality of life is influenced by various factors, including education, employment status, and income.¹⁸ In Indonesia, both men and women have the same rights in obtaining education and employment.²³ The number of health facilities in Indonesia continues to increase, and TB patients in larger cities have easy access to facilities.²⁴ No one should be discriminated against, especially when obtaining health services, social facilities, and equal opportunities.²⁴ As no differences were observed in obtaining education, employment, or access to health facilities, both men and women can achieve an equal quality of life.

Age influences general health because, as people age, their health conditions decline in general. Older people are less likely to maintain their diet and health conditions.^{25,26} Age did not influence any of the other domains, because older people can overcome and adjust to these conditions and thus feel that they do not experience obstacles in running their daily lives.^{27,28}

The results revealed that the level of education affected the quality of life in role limitation due to emotional

Table 4 P-Value of Factors Affecting the Quality of Life of TB Patients

Factors	Quality of Life (P-value)							
	General Health	Pain	Social Functioning	Physical Functioning	Role Limitation due to Physical Health	Role Limitation due to Emotional Problems	Energy	Emotional Well-Being
Sex	0.406	0.248	0.426	0.181	0.346	0.297	0.427	0.097
Age	0.018*	0.551	0.262	0.093	0.227	0.306	0.403	0.238
Educational level	0.300	0.139	0.163	0.016*	0.120	0.020*	0.273	0.408
Comorbidity	0.029*	0.026*	0.161	0.012*	0.001*	0.024*	0.360	0.964
Mental distress	0.001*	0.001*	0.001*	0.001*	0.001*	0.004*	0.001*	0.001*
Working status	0.183	0.441	0.560	0.201	0.763	0.408	0.302	0.838
Monthly income	0.841	0.806	0.363	0.212	0.789	0.097	0.343	0.260
BMI	0.523	0.802	0.537	0.965	0.900	0.713	0.848	0.470

Note: *Significantly affects quality of life.

problems. Educational level was a strong predictor of quality of life.²⁹ An individual who has a higher education more easily receives new information.^{30,31} The ability to receive information has an impact on one's way of thinking, including the ability to overcome problems, such as emotional issues. Individuals with a higher education tend to have more knowledge and, thus, find ways to overcome problems as they have fundamental knowledge, reasoning abilities, emotional self-regulation, and interactional abilities. Thus, the level of education can improve one's ability to overcome limitations due to emotional problems.^{31,32} One study has reported that emotions are closely related to ways of thinking and solving problems.³³

The presence of comorbidities results in TB patients experiencing more symptoms and pain, aside from TB itself. The number of symptoms experienced by patients affects their quality of life.³⁴ The decline in physical condition and the increase in the number of chronic symptoms are closely related to mental distress, which can have an impact on quality of life.^{35,36}

The prevalence of depression in TB patients is higher than in the general population.³⁷ Mental distress among TB patients can affect their health conditions and quality of life,³⁸ a finding which is similar to our research result. Poor quality of life is often experienced by people with mental distress.³⁹ This is in line with the research by Uddin et al who stated that an individual without mental distress has a better quality of life.⁴⁰

Conclusion

Sex does not significantly influence the quality of life of TB patients, and age only affects the general health domain. Moreover, the level of education affects physical functioning and role limitation due to emotional problems. Comorbidity affects several domains, namely, general health, pain, physical functioning, role limitation due to physical health, and role limitation due to emotional problems. Mental distress has a significant effect on quality of life in all domains.

Future study is warranted to investigate the implementation of practices to improve quality of life factors for effective TB patient care with optimal clinical outcomes. Management of TB patients should include the implementation of care to support the quality of life factors in addition to increased patient immunity and appropriate anti-TB drug therapy.

Acknowledgments

The authors would like to thank Manukan Kulon Primary Health Care (PHC), Medokan Ayu PHC, Gunung Anyar PHC, Dupak PHC, and Rangkah PHC; and the Directors of Dr. Soetomo General Hospital and Sidotopo Lung Hospital. We also thank Mr. Lantip and Mr. Danny for their technical support and all the students who helped to collect the data used in this study.

Disclosure

The authors declare no conflicts of interest.

References

1. Sandhu G. Tuberculosis: current situation, challenges and overview of its control programs in India. *J Glob Infect Dis.* 2011;3(2):143–150. doi:10.4103/0974-777X.81691
2. Ministry of Health. *Hasil Utama RISKESDAS 2018.* 2018.
3. Bauer M, Ahmed S, Benedetti A, et al. Health-related quality of life and tuberculosis: a longitudinal cohort study. *Health Qual Life Outcomes.* 2015;13(1):1–13.
4. Malik M, Nasir R, Hussain A. Health related quality of life among TB patients: question mark on performance of TB DOTS in Pakistan. *J Trop Med.* 2018;2018:1–7.
5. Salehitali S, Noorian K, Hafizi M, Dehkordi AH. Quality of life and its effective factors in tuberculosis patients receiving directly observed treatment short-course (DOTS). *J Clin Tuberc Mycobact Dis.* 2019;15:100093. doi:10.1016/j.jctube.2019.100093
6. Sartika I, Insani W, Abdulah R. Assessment of health-related quality of life among tuberculosis patients in a public primary care facility in Indonesia. *J Glob Infect Dis.* 2019;11(3):102–106. doi:10.4103/jgid.jgid_136_18
7. WHO. WHOQOL user manual. L [Internet]; 2012. Available from: http://apps.who.int/iris/bitstream/handle/10665/77932/WHO_HIS_HSI_Rev.2012.03protect_LY1extunderscore_eng.pdf;jsessionid=6BC7AC984CA0F8801C86C8296D9D4B2A?sequence=1%0Ahttp://www.springerreference.com/index/doi/10.1007/SpringerReference_28001.
8. Sarafino EP, Smith TW. *Health Psychology Biopsychosocial Interaction.* New York: John Wiley & Sons, Inc; 2011.
9. Kakhki AD, Masjedi MR. Factors associated with health-related quality of life in tuberculosis patients referred to the national research institute of tuberculosis and lung disease in Tehran. *Tuberc Respir Dis (Seoul).* 2015;78(4):309–314. doi:10.4046/trd.2015.78.4.309
10. Jaber AAS, Khan AH, Sulaiman SAS, Ahmad N, Anaam MS, Hasnain SE. Evaluation of health-related quality of life among tuberculosis patients in two cities in Yemen. *PLoS One.* 2016;11(6):1–19. doi:10.1371/journal.pone.0156258
11. Mohammed S, Nagla S, Morten S, Asma E, Arja A. Illness perceptions and quality of life among tuberculosis patients in gezira, sudan. *Afr Health Sci.* 2015;15(2):385–393. doi:10.4314/ahs.v15i2.11
12. Kastien-Hilka T, Abulfathi A, Rosenkranz B, Bennett B, Schwenkglens M, Sinanovic E. Health-related quality of life and its association with medication adherence in active pulmonary tuberculosis- a systematic review of global literature with focus on South Africa. *Health Qual Life Outcomes.* 2016;14.
13. Magdalena L. Factors affecting life quality of pulmonary tuberculosis sufferers at sanglah hospital denpasar. *IOSR J Nurs Health Sci.* 2013;2(5):30–34. doi:10.9790/1959-0253034

14. Farias SN, Medeiros CR, Paz EP, Lobo AD, Ghelman LG. Completeness in caring: study of quality of life in clients with tuberculosis. *Esc Anna Nery Rev Enferm*. 2013;17:749–754.
15. Adeyeye OO, Ogunleye OO, Coker A, et al. Factors influencing quality of life and predictors of low quality of life scores in patients on treatment for pulmonary tuberculosis: a cross sectional study. *J Public Health Africa*. 2014;5(2):88–92. doi:10.4081/jphia.2014.366
16. Dawar U, Garg A, Oswal R, Patel K. A study of prevalence of depression in patients with multi drug resistant tuberculosis. In: 102 tuberculosis [Internet]. European Respiratory Society; 2016. Available from: https://erj.ersjournals.com/content/48/suppl_60/PA2690. Accessed October 10, 2020.
17. Rajkumar LC, Karibeeran S. Quality of life of multi drug resistant tuberculosis patients in India. *Int Multidiscip Res J*. 2015;5 (2 August):1–8.
18. Karyani AK, Rashidian A, Sefiddashti SE, Sari AA. Self-reported health-related quality of life (HRQoL) and factors affecting HRQoL among individuals with health insurance in Iran. *Epidemiol Health*. 2016;38:e2016046. doi:10.4178/epih.e2016046
19. Salodia UP, Sethi S, Khokhar A. Depression among tuberculosis patients attending a DOTS centre in a rural area of Delhi: a cross-sectional study umang. *Indian J Public Health*. 2018;63:39–43.
20. Stratton KJ, Richardson LK, Tran TL, et al. Using the SRQ-20 factor structure to examine changes in mental distress following typhoon exposure. *Psychol Assess*. 2014;26(2):528–538. doi:10.1037/a0035871
21. Mamani M, Majzoobi MM, Ghahfarokhi SM, Esna-Ashari F, Keramat F. Assessment of health-related quality of life among patients with tuberculosis in Hamadan, Western Iran. *Oman Med J*. 2014;29(2):102–105. doi:10.5001/omj.2014.25
22. Jankowska-Polanska BK, Kamińska M, Uchmanowicz I, Rycombel A. Quality of life and health behaviours of patients with tuberculosis — sex differences. *Pneumonol Alergol Pol*. 2015;83 (4):256–265. doi:10.5603/PiAP.2015.0046
23. Australia Indonesia Partnership for Economic Governance MU. Women's economic participation in Indonesia: a study of gender inequality in employment, entrepreneurship, and key enablers for change; 2017. Available from: <https://www.monash.edu/business/cdes/research/publications/publications2/Womens-economic-participation-in-Indonesia-June-2017.pdf>. Accessed October 10, 2020.
24. Mahendradhata Y, Trisnantoro L, Listiyadewi S, et al. *The Republic of Indonesia Health System Review*. Vol. 7. 2017:1.
25. Dionigi RA. Stereotypes of aging: their effects on the health of older adults. *J Geriatr*. 2015;2015:1–9. doi:10.1155/2015/954027
26. Irwan AM, Kato M, Kitaoka K, Kido T, Taniguchi Y, Shogenji M. Self-care practices and health-seeking behavior among older persons in a developing country: theories-based research. *Int J Nurs Sci*. 2016;3:11–23.
27. Abedi H, Mostafavidarani F, Riji HM. The elderly perception and views on their health - facilitating and inhibiting factors in elderly health care in Iran: a qualitative study. *Procedia Soc Behav Sci*. 2010;5:2222–2226. doi:10.1016/j.sbspro.2010.07.439
28. Miguel L, Ram JM. The impact of quality of life on the health of older. *J Aging Res*. 2018;2018:1–9.
29. Javed S, Javed S, Khan A. Effect of education on quality of life and well being. *Int J Indian Psychol*. 2016;3:1–10.
30. Hamid HG. *Breast Cancer Among Women Attending Primary Health Center in Kufa City. The Impact of Educational Level on Knowledge, Attitude and Practices Toward Breast Cancer Among Women Attending Primary Health Center in Kufa City*. 2019.
31. Diaz-Quijano FA, Martínez-Vega RA, Rodríguez-Morales AJ, Rojas-Calero RA, Luna-González ML, Díaz-Quijano RG. Association between the level of education and knowledge, attitudes and practices regarding dengue in the Caribbean region of Colombia. *BMC Public Health*. 2018;18(1):1–10. doi:10.1186/s12889-018-5055-z
32. Hahn R, Truman B. Education improves health and promotes health equality. *Int J Health Serv*. 2015;45(4):657–678. doi:10.1177/0020731415585986
33. Jung N, Wranke C, Hamburger K, Knauff M. How emotions affect logical reasoning: evidence from experiments with mood-manipulated participants, spider phobics, and people with exam anxiety. *Front Psychol*. 2014;5:1–12.
34. Megari K. Quality of life in chronic disease patients. *Health Psychol Res*. 2013;1:17–20.
35. Peltzer K. Tuberculosis non-communicable disease comorbidity and multimorbidity in public primary care patients in South Africa. *Afr J Prim Health Care Fam Med*. 2018;10:1–6.
36. Somrongthong R, Hongthong D, Wongchalee S, Wongtongkam N. The influence of chronic illness and lifestyle behaviors on quality of life among older thais. *Biomed Res Int*. 2016;2016.
37. Molla A, Mekuriaw B, Kerebih H. Depression and associated factors among patients with tuberculosis in Ethiopia: a cross-sectional study. *Neuropsychiatr Dis Treat*. 2019;15:1887–1893. doi:10.2147/NDT.S208361
38. Louw JS, Mabaso M, Peltzer K, Shankar EM. Change in health-related quality of life among pulmonary tuberculosis patients at primary health care settings in South Africa: a prospective cohort study. *PLoS One*. 2016;11(5):1–13. doi:10.1371/journal.pone.0151892
39. Connell J, Brazier J, O'Cathain A, Lloyd-Jones M, Paisley S. Quality of life of people with mental health problems: a synthesis of qualitative research. *Health Qual Life Outcomes*. 2012;10(1):1. doi:10.1186/1477-7525-10-138
40. Uddin MN, Bhar S, Al Mahmud A, Islam FMA. Psychological distress and quality of life: rationale and protocol of a prospective cohort study in a rural district in Bangladesh. *BMJ Open*. 2017;7.

Journal of Multidisciplinary Healthcare

Publish your work in this journal

The Journal of Multidisciplinary Healthcare is an international, peer-reviewed open-access journal that aims to represent and publish research in healthcare areas delivered by practitioners of different disciplines. This includes studies and reviews conducted by multidisciplinary teams as well as research which evaluates the results or conduct of such teams or healthcare processes in general. The journal

covers a very wide range of areas and welcomes submissions from practitioners at all levels, from all over the world. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-inflammation-research-journal>

Dovepress