Toll-Like Receptor 4 and Inflammatory Micro-Environment of Pancreatic Islets in Type-2 Diabetes Mellitus: A Therapeutic Perspective

This article was published in the following Dove Press journal:
Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Zhaoping Wang¹
Xiaolin Ni^{1,2}
Li Zhang¹
Liang Sun¹
Xiaoquan Zhu¹
Qi Zhou¹
Ze Yang¹
Huiping Yuan¹

¹The MOH Key Laboratory of Geriatrics, Beijing Hospital, National Center of Gerontology, Beijing, People's Republic of China; ²Graduate School of Peking Union Medical College, Chinese Academy of Medical Sciences, Beijing, People's Republic of China

Abstract: Patients with type-2 diabetes mellitus (T2DM) display chronic low-grade inflammation induced by activation of the innate immune system. Toll-like receptor (TLR)4 is a pattern recognition receptor that plays a vital part in activation of the innate immune system. Results from animal and computer-simulation studies have demonstrated that targeting TLR4 to block the TLR4-nuclear factor-kappa B (NF- κ B) pathway reduces the inflammatory response and complications associated with T2DM. Therefore, TLR4-targeted therapy has broad prospects. Here, we reviewed the role of TLR4 in inflammation during chronic hyperglycemia in T2DM and its therapeutic prospects.

Keywords: T2DM, TLRs, TLR4, inflammation, TLR4 treatment

Introduction

The International Diabetes Federation showed that the number of people aged 20–79 years with diabetes mellitus (DM) worldwide was estimated to be 463 million in 2019. Almost 90% of these people will have type-2 diabetes mellitus (T2DM), and this number may increase to 700 million by 2045. However, 1 in 2 (~232 million) patients do not know that they are suffering from DM, and the number of people at risk for T2DM is ~374 million.¹ T2DM has become a major public-health problem worldwide.

T2DM is characterized by impaired insulin resistance and a dysregulated immune response. T2DM patients display chronic initiation of the innate immune system in pancreatic islets, which occurs in insulin-sensitive tissues and at the sites of diabetic complications. Several studies have shown that T2DM is associated significantly with activation of the innate immune system and that, in many aspects, T2DM is an inflammatory disease. Therefore, the proteins associated with components of the innate immune system could be potential targets for studying T2DM and its complications, including diabetic kidney disease (DKD), atherosclerosis and diabetic retinopathy.

Toll-like receptors (TLRs) are receptors that recognize pathogen-associated molecular patterns (PAMPs) and danger-associated molecular patterns (DAMPs).² TLR4 has important roles in modulating innate immunity. TLR4 is expressed in macrophages, adipose tissue, pancreatic β -cells, vascular endothelial cells, skeletal muscle, airway epithelia and smooth muscle cells. TLR4 interacts with other endogenous and exogenous substances, such as heat-shock proteins, fibronectin,

Correspondence: Huiping Yuan
The MOH Key Laboratory of Geriatrics,
Beijing Hospital, National Center of
Gerontology, Dongdan DaHua Road 1#,
Beijing 100730, People's Republic of China
Tel +86-10-58115043
Fax +86-10-65237929
Email yuanhuiping@126.com

fibrinogen, free fatty acids (FFAs), and saturated fatty acids. This protein is also a crucial receptor for the lipopolysaccharide (LPS) component of Gram-negative bacteria. Most of the proinflammatory cytokines induced by activated TLR4 are involved in insulin resistance, which is associated significantly with T2DM risk.³ Given this information, the relationship between T2DM and TLR4 will be highlighted in this review.

TLRs Family

The first identification of a Toll receptor was in the embryonic development of *Drosophila*. Results from subsequent studies showed that the *Drosophila* Toll receptor had a major role in the antifungal responses of the adult fly.⁴ Since then, researchers have focused intensively on the role of Toll receptors in the innate immune response of insects and mammals. Consequently, the Toll receptor was identified in mammals, and revealed the important roles of Toll receptors in recognizing microorganisms.

Medzhitov and colleagues first identified that the mammalian homolog of the *Drosophila* Toll receptor was hToll (now termed TLR4) in 1997.⁵ Consequently, many studies have suggested that several proteins that are structurally associated with TLR4 are called TLRs. To date, 10 TLRs in humans (TLR1–TLR10) and 12 TLRs in mice (TLR1 to TLR9 and TLR11 to TLR13) have been identified. It has been demonstrated that TLR-mediated recognition of microorganisms is complex. Different TLRs can recognize and bind with the specific patterns of microbial components.⁶

Structure of TLRs

TLRs are evolutionarily conserved proteins that can detect PAMPs. TLRs are important for the innate immune response and are vital regulators of innate and adaptive immune responses.^{2,7} As type-I transmembrane glycoproteins, TLRs share three common domains: an intracellular Toll-interleukin 1 receptor (TIR) homology signaling domain; a single transmembrane domain; an extracellular recognition domain.⁸ Since the first reported crystal structures of unliganded hTLR3 provided atomic detail,^{9,10} the hybrid long-range radar method has been established to crystallize TLRs.¹¹ Subsequently, eight structures of TLR complexes activated by ligand induction (TLR1–TLR2–triacyl lipopeptide,¹² TLR2–TLR6–diacylated lipopeptide,¹³ TLR3–dsRNA,^{10,14} TLR4–MD-2–Eritoran (–LPS),^{11,15} TLR5–flagellin,¹⁶ TLR8–imidazoquinoline derivatives,¹⁷

TLR9–ssDNA,¹⁸ and TLR13–ssRNA)¹⁹ have been confirmed based on the hybrid long-range radar method.

Structures of the Intracellular TIR Domains and Signaling Adapters

The TIR domain is a pivotal module that adapts the pathways of the innate immune response. The TIR domain is mediated by TLRs and is contained in the C-terminal regions of all mammalian TLRs. Initiation of downstream signaling requires homo- or heterotypic dimerization of TIR domains. The latter and a five-stranded parallel β -sheet (β A– β E) in the center have a common fold, and there are five α -helices (α A– α E) on both sides.^{20,21} Including three highly conserved residues (arginine BB3, aspartic acid BB4 and glycine BB8), the BB loop connects helix α B and strand β B, and has a vital role in the dimerization of TIR and/or recruitment of adaptors.^{21,22} The BB loop can use different conformations to form TIR dimers.²²

Structures of Extracellular Domains

The extracellular recognition domains of TLRs belong to the leucine-rich repeat (LRR) family. The domains contain 16–28 LRR modules in the middle of the N-terminal and C-terminal modules.²³ Each of the LRR modules includes 20–30 amino acids that contain a mutable part and a conserved “LxxLxLxxN” motif.^{24,25} Evidence from crystal structures has shown that all LRR modules are horseshoe-like. The concave face and convex surface of the horseshoe shape consist of parallel β strands and short 3_{10} helices and parallel loops, respectively.²⁶ There are two structural transitions in the central β -sheets of TLR4 so the LRR domains of TLR4 can be separated into three subdomains: C-terminal, central, and N-terminal.¹¹

TLR4 Signaling Pathways

The main pathways activated by TLR4 are the myeloid differentiation primary response protein (MyD)88-dependent and -independent pathways. The initiating step of each pathway is TLR4 identification of ligands. The complex combination of LPS and LPS binding protein (LBP) attaches to another protein known as cluster of differentiation (CD)14.²⁷ As a membrane-binding protein on innate immune cells, CD14 can also cycle in the cytoplasm in a soluble form. It has been suggested that the major role of CD14 is to strengthen TLR4 signaling by accelerating the transport of TLR4 to “lipid rafts” on the cellular membrane.

Subsequently, the coreceptor MD-2 is recruited to facilitate TLR4 translocation to the cellular membrane.²⁸ Endocytosis of the TLR4/MD-2 complex occurs after translocation. The heterotrimer CD14/TLR4/MD-2 recognizes LPS and then induces the initiation of MyD88-dependent and MyD88-independent pathways. The general pattern of these pathways is shown in Figure 1.

MyD88-Dependent Pathway

The MyD88-dependent pathway was the first to be discovered, and it is the most vital pathway. Evidence from several studies has suggested that MyD88 knockout mice do not respond to TLR ligands.²⁹ Through this mechanism, after

stimulation, the binding of TLR4 and MyD88 at the cytoplasmic TIR domain³⁰ results in recruitment of the interleukin-1 receptor-associated kinase (IRAK)4, which is associated with the IL-1 receptor.³¹ After binding to and activating tumor necrosis factor receptor-associated factor (TRAF)6, activated IRAK4 dissociates from MyD88, resulting in the formation of Complex-1. Subsequently, after dissociating from TLR4, this triple association stimulates the recruitment of numerous proteins, including TGF α -associated kinase (TAK)1 and TAK1 binding protein (TAB) 1–3, which form the core components of Complex-2. These processes result in the activation and initiation of two sub-pathways. In the first sub-pathway, TAK-1 phosphorylation

Figure 1 TLR4 signaling pathway (schematic). After LBP- and CD14-mediated LPS binding or FetA-mediated FFA binding, the TLR4/MD-2 complex recruits MyD88 or TRIF via TIRAP and TRAM.²⁷ This action, in turn, induces (1) the MyD88-dependent pathway.²⁹ MyD88 binds with the IRAK/TRAF6 complex (Complex I) and TAK1/TAB complex (Complex II), subsequently triggering the activation of IKK. IκBα is phosphorylated by activation of IKK and stimulates the translocation of NF-κB into the nucleus, where NF-κB leads to the production of ROS and proinflammatory cytokines. Complex II can also trigger the MAPK pathway and lead to the activation of p38, JNK and ERK1/2. This activation stimulates the translocation of AP-1 into the nucleus, and proinflammatory cytokines are subsequently transcribed. (2) In the MyD88-independent pathway,³⁴ TLR4 is stimulated by LPS and associates with translocating chain-associated membrane protein (TRAM) and TIR-domain-containing adaptor-inducing interferon-β (TRIF). The dimerization of TRAM and TRIF activates the TRAF6-TAK1-IKK pathway. Alternatively, TBK1 kinase can activate interferon regulatory factor (IRF)3 by phosphorylating it. Activated IRF3 can migrate to the nucleus and act as a transcription factor. Interferon (IFN)-β binds to activated IRF3 and subsequently activates the production proinflammatory cytokines. Furthermore, two potential targeting points blocking TLR4 signal-pathway have been found according to the literature: (a) FP7 blocks the TLR4 signaling pathway by targeting MD2 and CD14; (b) dioscin inhibits the TLR4 signaling pathway by decreasing NF-κB expression.

activates the I κ B kinase (IKK) complex, which in turn induces the phosphorylation of inhibitor of kappa B (I κ B) protein. This action results in proteasomal degradation of I κ B and stimulates the translocation of nuclear factor-kappa B (NF- κ B) to the nucleus.³² NF- κ B then induces the activation of proinflammatory mediators, which are extremely effective in activating inducible nitric oxide synthase (iNOS) and inducible cyclo-oxygenase (COX)-2, and stimulates the release of many major inflammatory cytokines, such as tumor necrosis factor (TNF)- α , IL-1 β , and IL-6.³³ In the second sub-pathway, TRAF6 induces the activation of the mitogen-activated protein kinases (MAPKs) c-Jun N-terminal kinase (JNK), extracellular signal-regulated kinase (ERK)1/2, and p38 via phosphorylation, subsequently modulating the activation of some transcription factors, including activator protein (AP)-1.³¹

MyD88-Independent Pathway

Kawai and colleagues revealed that, following stimulation of TLRs, JNK and NF- κ B were produced in MyD88-knockout mice. This result suggests that there are other MyD88-independent pathways.³⁴ After stimulation of TLR4 with its ligand, TLR4 recruits translocating chain-associated membrane protein (TRAM) and TIR-domain-containing adaptor-inducing interferon- β (TRIF). The dimerization of TRAM and TRIF activates the TRAF6-TAK1-IKK pathway. This interaction activates NF- κ B and destroys I κ Bs.³⁵ Alternatively, TBK1 kinase activates interferon regulatory factor (IRF)3 by phosphorylating it. Activated IRF3 can translocate to the nucleus and act as a transcription factor. After binding to activated IRF3, interferon 1 β (IFN-1 β) subsequently activates the production of reactive oxygen species (ROS) and proinflammatory cytokines. Several endogenous inhibitors can regulate TLR4 overactivation, such as the ubiquitin ligase TRIAD3A, which strengthens ubiquitination and then degrades TLR4.³⁶ In addition, a homolog of the TLR4 protein RP105 can also antagonize TLR4 signaling on the surface of the cell membrane.³⁷ It has been shown that posttranslational modification of TLR4 signaling can be altered by the acetylation of lysine residues.³⁸

Role of TLR4 in T2DM Pathogenesis Inflammation and T2DM

T2DM has become a growing issue as people become older, especially for those who have genetic and epigenetic predispositions. T2DM is associated mainly with obesity and

physical inactivity. In people with predispositions, a defect in insulin secretion can be detected concomitantly with a reduced response to glucose uptake stimulated by insulin in the liver and adipose tissues, and this condition is called insulin resistance.^{39–42} In individuals, insulin resistance remains relatively constant over time and increases only slightly with age. However, an initial increase in insulin production was followed by a continuous deterioration in the insulin-secretory capacity of pancreatic β -cells, which causes the initiation and development of T2DM. Finally, the increased demand for peripheral insulin cannot be compensated for by insulin secretion. In T2DM, mechanisms associated with defective insulin secretion and responses include glucotoxicity, lipotoxicity, oxidative stress, endoplasmic reticulum (ER) stress, endocannabinoids, alterations in the gut microbiota and the formation of amyloid deposits in pancreatic islets.^{43–47} Interestingly, these factors are all associated with the inflammatory response.^{48–56} For example, increased glucose concentrations and LPS activate TLR4, which upregulates expression of caspase-3 and poly ADP ribose polymerase (PARP), and modifies the ratio of expression of B-cell lymphoma-2 (Bcl-2)/Bcl-2-associated X protein, which results in enhanced oxidative stress, increased release of ROS and apoptosis.^{57–59} Armann et al showed that ROS generation in pancreatic islets is linked to the percentage of cells undergoing apoptosis and the functional potency of pancreatic islets in vivo.⁶⁰ Duprez et al revealed that oxidative stress led to the apoptosis of pancreatic-islet β -cells and dysfunction of insulin secretion in rats.⁶¹ ROS and the byproducts induced by ROS (degraded hyaluronic acid, oxidize phospholipids and activated high-mobility group protein B1) are recognized as DAMPs, which can activate the TLR4 pathway, thereby leading to increased production of ROS and inflammatory mediators. Consequently, there is a “vicious cycle” between TLR activation and ROS generation that contributes to the maintenance of chronic inflammation in pancreatic islets.⁶² Furthermore, the increased ROS leads to activation of the NOD-, LRR- and pyrin domain-containing (NLRP)3 inflammasome and caspase-1, which promotes the production of mature interleukin-1 β (IL-1 β).

Initially, this inflammation can promote β -cell proliferation and the production of insulin to compensate for insulin resistance.⁶³ Various cytokines and chemokines are induced by IL-1 β , such as monocyte chemoattractant protein (MCP) 1; also called (CCL2) and TNF, which results in the infiltration of macrophages and other immune cells.^{64–66} Such infiltration of immune cells is increased due to the vicious

cycle of IL-1 β autostimulation.⁶⁷ Therefore, macrophage infiltration into resident pancreatic islets induces their dysfunction. In adipocytes, lipid overload can result in adipocyte death, further triggering an inflammatory response, which forms a vicious cycle. In addition, a review by Kitamura indicated the contribution of Forkhead box protein O1 (FOXO)1 (a multifunctional protein that regulates proliferation, apoptosis, senescence, differentiation, stress resistance, autophagy and metabolism) signaling to the initiation and development of loss of pancreatic β -cells in T2DM.⁶⁸ The inflammatory microenvironment in pancreatic islets stimulated by FFAs and high glucose includes macrophage polarization, and the conversion of M2 macrophages to M1 macrophages. Moreover, activation of TLR4 in M1 macrophages and β -cells mediates the inflammatory pathway, which leads to β -cell dysfunction (reduced production and secretion of insulin) and apoptosis/death. Eventually, these processes contribute to T2DM development (Figure 2). Insulin resistance and β -cell dysfunction are hallmarks of T2DM.

Expression of TLR4 in T2DM

T2DM has become the most prevalent metabolic disease worldwide. The effects of T2DM have become the main cause of mortality in humans.⁶⁹ Moreover, the results

obtained from several studies have suggested that proinflammatory cytokines in low-grade inflammation are involved in the pathological processes that lead to T2DM.^{70,71} TLR4 is expressed not only in various immune-related cells (eg, neurons, oligodendrocytes, astrocytes, microglia and the cerebral endothelium) but also functions in several types of nonimmune cells. Garay-Malpartida et al observed variable levels of TLR4 expression in pancreatic β -cells isolated from adult nondiabetic brain-dead donors. Their results showed that LPS could increase TLR4 expression and impair the viability of pancreatic β -cells, which led to a decrease in the synthesis and secretion of insulin.⁷² Moreover, activation of TLRs induced by any stimuli in injured tissue plays a key part in the human inflammatory response: TLRs are important mediators of the human inflammatory response because they facilitate the production of inflammatory cytokines. Given the findings mentioned above, Gupta et al measured TLR1-10 expression in neutrophils and monocytes derived from patients with T2DM and healthy participants. They showed that in neutrophils, mRNA expression of most TLRs was enhanced in glycemic control (GC) patients compared with that in normal glucose tolerance (NGT) cases. Similarly, mRNA expression of TLRs was increased in monocytes derived from GC cases.⁷³ The data collected

Figure 2 Implications of TLR4 in inflammation in pancreatic islets. TLR4 on the surface of β -cells and macrophages is activated by FFA and high glucose, which induces oxidative stress and ROS generation by upregulating expression of caspase-3 and PARP. The cytokine IL-1 β is released via the TLR4-NF- κ B signaling pathway in pancreatic β -cells and macrophages. Eventually, the generation of IL-1 β and ROS induces the pancreatic β -cell dysfunction (decreased production and secretion of insulin) and the apoptosis/cell death of β -cells. Moreover, M2 macrophages from bone marrow or adipose tissue enter pancreatic-islet tissue and translate into M1 macrophages. Activated M1 macrophages increase the level of ROS and IL-1 β to aggravate inflammation and further deteriorate dysfunction of β -cell in pancreatic islets.⁶²

from T2DM patients with good GC were consistent, and studies have suggested that TLR4 expression was enhanced in monocytes isolated from patients with T2DM; the mean duration of such expression was 29 months.⁷⁴ Moreover, data from other studies indicated that mRNA expression of TLR4 was enhanced in peripheral blood mononuclear cells (PBMCs) derived from patients with T2DM⁷⁵ (though the mean duration of disease was not stated).⁷⁶ Mohan et al showed that high glucose could stimulate the generation of nicotinamide adenine dinucleotide phosphate (NADPH) oxidase in human monocytes, and that TLR4 expression was induced by NADPH oxidase via protein kinase C (PKC)- δ .⁷⁷ Recently, Sindhu et al reported that expression of TLR10 (an immune marker of metabolic inflammation) was induced by oxidative stress, and that ROS-mediated oxidative stress induced TLR10 expression in monocytic cells, an ER stress response, as well as NF- κ B and MAPK signaling pathways.⁷⁸

Besides macrophages⁷⁹ or immunocompetent cells (such as dendritic cells, DCs),⁸⁰ TLR4 is also expressed in the cardiovascular system, including endothelial cells,⁷⁹ vascular smooth muscle cells (VSMCs)⁸¹ and adventitial fibroblasts.⁸² Several studies have identified the role of TLR4 in the development and progression of atherosclerotic disease. For instance, TLR4 on adventitial fibroblasts engaged in neointima formation in atherosclerosis and the TLR4 pathway lead to expression of apoptotic molecules in the Fas death pathway.⁸³ Moreover, we should also pay close attention to the expression and role of TLR4 in adipose tissue because it can communicate with multiple organs or tissues through many adipocytokines, which affects many physiologic and pathophysiologic processes. Several studies have shown that adipose tissue in obese people is infiltrated with macrophages; they may participate in the inflammatory process activated in adipose tissue.^{84–86} Notably, the infiltration of macrophages and expression of inflammation-associated genes in adipose tissue precede insulin resistance.^{84,85} Hence, infiltrating macrophages play an essential part in inflammation initiation in adipose tissue. Accumulated evidences have suggested that TLR4 has a vital role in inflammation of adipose tissue induced by obesity and systemic glucose and lipid metabolism.^{87–90} TLR4 is expressed in macrophages more abundantly than in adipocytes, so the chronic inflammatory responses induced by the interaction between macrophages and adipocytes could be mediated via TLR4 in macrophages.⁹¹

Role of TLR4 as an Inflammatory Target in T2DM

The innate immune system is activated by TLRs, particularly TLR4, which plays a vital part in chronic, low-grade inflammation. A high-fat diet can lead to an increase in plasma LPS concentrations in T2DM cases and healthy individuals,^{92,93} whereas LPS induces insulin resistance rapidly in healthy people.⁹⁴ LPS is derived from a group of Gram-negative bacteria⁹⁵ called Enterobacteriaceae (ie, *Escherichia coli*),⁹⁶ as well as fungal and viral products⁹⁷ that are well-known extracellular pathogens and can induce TLR4 activation. Similarly, an excessive amount of glucose and FFAs leads to increased expression of TLR4 mRNA and protein in patients with untreated T2DM. FFAs induce activation of the inflammatory response and metabolic signaling in insulin resistance, primarily via the TLR4-NF- κ B signaling pathway in pancreatic β -cells and macrophages.⁹⁸ However, TLR4 cannot bind to FFAs directly; thus, the interaction between TLR4 and FFAs is mediated by fetuin A (FetA), which may be regarded as a new therapeutic target.^{99,100}

Therapeutic Prospects of TLR4 in T2DM

Application of TLR4 Antagonists in T2DM

Having shown the effects of TLR4 in T2DM development, one could hypothesize that blockade of TLR4 signaling would inhibit the progression of inflammation in pancreatic islets. Consistent with this hypothesis, Shi et al observed that a lack of TLR4 could improve obesity-induced insulin resistance.⁹⁸ Moreover, Wang et al established that pretreatment with TLR4-short hairpin (sh)RNA and a TLR4 antibody led to significantly reduced expression of TLR4 compared with that observed upon treatment with LPS. After pretreatment with a TLR4 antibody and TLR4-shRNA, insulin secretion was increased significantly in isolated pancreatic islets in rats.¹⁰¹ This result showed that inhibition of TLR4 expression may protect pancreatic islet cells from the damage induced by ligands.

One study suggested, for the first time, that a TLR4 antagonist improved atherogenesis and vascular inflammation in diabetic apolipoprotein E (ApoE)^{-/-} mice and lowered serum levels of triglyceride and cholesterol in nondiabetic ApoE^{-/-} mice.¹⁰² Furthermore, Perrin-Cocon et al examined the activity of FP7 (a small-molecule synthetic TLR4 antagonist) on human monocytes and monocyte-derived DCs in vitro and on mice with

influenza-virus infection in vivo. They showed that FP7 inhibited activation of TLR4-mediated monocytes and DCs, thereby avoiding excessive TLR4 signaling in inflammatory disorders.¹⁰³ FP7 binds to MD-2 and CD14 and inhibits LPS-induced TLR4 signaling.¹⁰⁴ Inhibition was TLR4-specific, suggesting a more precise molecular interaction with MD-2.¹⁰³ Therefore, TLR4 antagonists such as FP7 could be developed as new drugs that target TLR4-related inflammatory diseases, such as T2DM.

Blockade of the TLR4 Signaling Pathway in T2DM Complications

Recently, several studies have revealed that blocking TLR4 expression is a strategy for treating T2DM complications. Diabetic retinopathy (DR), atherosclerosis and DKD are three of the most common complications of DM. Hu et al treated retinal ganglion cells (RGCs) isolated from retinas of Sprague–Dawley rats with a selective inhibitor of TLR4, TAK-242, under a high-glucose environment for 48 h. Subsequently, they showed that treatment with TAK-242 decreased expression of TLR4 downstream signaling molecules and significantly reduced levels of inflammatory cytokines, accompanied by a reduced rate of RGC apoptosis. Hu et al identified the potential effect of TAK-242 for treating the degeneration of primary RGCs in DR.¹⁰⁵

Moreover, low-density lipoprotein receptor-deficient (LDLR^{-/-}) mice (which are atherosclerotic and can be induced to develop T2DM with a high-fat diet) were treated with *Rhodobacter sphaeroides* LPS (Rs-LPS), an established TLR4 antagonist, in a study by Lu et al. They found that the number of monocytes and macrophages in atherosclerotic lesions was reduced, and that expression of IL-6 and matrix metalloproteinase-9 in atherosclerotic plaques was inhibited; they also showed that Rs-LPS inhibited atherosclerosis in diabetic LDLR^{-/-} mice and reduced collagen levels in diabetic mice.¹⁰⁶

Emerging evidence has identified that the TLR4-NF-κB pathway in the kidney plays a vital part in DKD pathogenesis.^{107,108} Wu et al suggested that high glucose would increase expression of microRNA (miR)-199a-5p to decrease expression of Klotho, leading to inflammation and fibrosis by activating the TLR4/NF-κB p65/NGAL signaling pathway. In contrast, they showed that initiation of the TLR4/NF-κB p65/NGAL signaling pathway might be attenuated under high glucose conditions by inhibiting miR-199a-5p or adding Klotho exogenously, thereby preventing the

progression of inflammation and fibrosis in the kidney.¹⁰⁹ Furthermore, Liu et al found that 1,25-dihydroxyvitamin D3 (1,25(OH)2D3) treatment reduced TLR4 expression, inhibited expression of NF-κB and MyD88 in cultured renal tubular epithelial cells, and reduced expression of collagen-associated proteins, resulting in a significant decrease in the inflammatory response in the renal tissues of rats with DKD. Immunohistochemistry results showed that 1,25(OH)2D3 treatment could suppress the immune response by reducing the infiltration of macrophages significantly. Therefore, 1,25(OH)2D3 could downregulate the innate immune TLR4-NF-κB pathway to protect against DKD.¹¹⁰ Recently, Cai et al reported the potential role of dioscin in treating DKD in patients suffering from DM.¹¹¹

Conclusions and Future Perspectives

TLR4, which is expressed in pancreatic β-cells, adipocytes and macrophages, plays a key part in the activation of chronic low-grade inflammation. This action induces the initiation and progression of T2DM under high glucose concentrations in peripheral blood in obese people. Hence, TLR4 could be an ideal therapeutic target to inhibit the inflammatory pathway and improve T2DM and its complications. Moreover, recent studies have shown that treatments targeting TLR4 could inhibit the initiation and progression of inflammation effectively, thereby eliciting a potential hypoglycemic effect. Ullah et al used a computer-based study and found that curcumin analogs are natural antagonists of TLR4 that could inhibit the TLR4 pathway.¹¹² Lakhia et al reported that miR-214 could attenuate inflammation by inhibiting TLR4 expression.¹¹³ Shah et al found that MAL/MyD88-inhibitory peptide 2, an inhibitor, has broad specificity for TLRs and attenuates the activation of TLRs (including TLR4) that is induced by microbial or environmental factors.¹¹⁴ Yang et al demonstrated that *D. officinale* polysaccharide could inhibit TLR4 expression to achieve potent anti-inflammatory effects.¹¹⁵ Therefore, TLR4 represents a promising therapeutic target in patients with T2DM, and researchers prefer to identify natural anti-inflammatory drugs. However, another important feature is the consequence of blocking the TLR4 pathway, which has a substantial role in initiating the immune response. Zheng et al suggested that TLR4 knockout influenced the growth, development, and reproduction of mice.¹¹⁶ Furthermore,

Coenen et al showed no differences in body composition or plasma lipids between M θ TLR4^{+/+} mice (recipient mice with TLR4 expression in bone marrow) and M θ TLR4^{-/-} mice (recipient mice with TLR4 knockdown in bone marrow) on any of the diets. Moreover, similar insulin sensitivity between M θ TLR4^{+/+} mice and M θ TLR4^{-/-} mice fed a high-fat diet with saturated fatty acids was found.¹¹⁷ Similarly, Eritoran (E5564, TLR4 antagonist) has shown its safety in several studies.^{118,119} Therefore, further research and understanding of related mechanisms are necessary to explore the potential future clinical importance of these treatments.

Abbreviations

T2DM, type-2 diabetes mellitus; TLR4, Toll-like receptor 4; NF- κ B, nuclear factor κ B; TLRs, toll-like receptors; PAMPs, pathogen-associated molecular patterns; DAMPs, danger-associated molecular patterns; LPS, lipopolysaccharide; TIR, Toll-interleukin 1 receptor; MD2, myeloid differentiation protein 2; LRR, leucine-rich repeat; MyD88, myeloid differentiation primary response protein 88; LBP, LPS-binding protein; CD14, cluster of differentiation 14; IRAK4, interleukin-1 receptor-associated kinase 4; TRAF6, TNF receptor-associated factor 6; TAK-1, TGF α -associated kinase-1; TAB, TAK1 binding protein; IKK, I κ B kinase; I κ B, inhibitor of kappa B; iNOS, inducible nitric oxide synthase; AP-1, activator protein-1; TRAM, translocating chain-associated membrane protein; TRIF, TIR-domain-containing adaptor-inducing interferon- β ; IRF3, interferon regulatory factor 3; IFN-1 β , interferon 1 β ; ROS, reactive oxygen species; ER, endoplasmic reticulum; NLRP3, NOD-, LRR- and pyrin domain-containing 3; IL-1 β , interleukin-1 β ; MCP1, monocyte chemoattractant protein 1; TNF, tumor necrosis factor; GC, glycemic control; NGT, normal glucose tolerance; PBMCs, peripheral blood mononuclear cells; MAPK, mitogen-activated protein kinase; FFAs, free fatty acids; DCs, dendritic cells; DR, diabetic retinopathy; DKD, diabetic kidney disease; RGCs, retinal ganglion cells; Rs-LPS, *Rhodobacter sphaeroides* LPS; NGAL, Neutrophil Gelatinase-Associated Lipocalin; MIP2, MAL/MyD88-inhibitory peptide 2; DOP, *D. officinale* polysaccharide; M θ TLR4^{+/+}, recipient mice with TLR4 expression in bone marrow; M θ TLR4^{-/-}, recipient mice with TLR4 knockdown in bone marrow; PARP, poly ADP ribose polymerase.

Acknowledgments

All authors made substantial contributions to conception and design, acquisition of data, or analyses and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed to submit to this journal; gave approval of the final version to be published; and agree to be accountable for all aspects of the work.

This work was supported by the Natural Science Foundation of China (81870552, 81400790, 81872096, 81571385, 91849118, and 91849132), National Key R&D Program of China (2018YFC2000400), Beijing Hospital Doctoral Scientific Research Foundation (BJ-2018-024), Beijing Hospital Nova Project (BJ-2018-139) and Non-profit Central Research Institute Fund of Chinese Academy of Medical Sciences (2018RC330003).

Disclosure

The authors report no conflicts of interest in this work.

References

1. IDF Diabetes atlas, 9th edn. International Diabetes Federation. Available from: <https://diabetesatlas.org/en/>. Accessed June 4, 2020.
2. West AP, Kobiansky AA, Ghosh S. Recognition and signaling by toll-like receptors. *Annu Rev Cell Dev Biol*. 2006;22:409–437. doi:10.1146/annurev.cellbio.21.122303.115827
3. Garibotto G, Carta A, Picciotto D, Viazzi F, Verzola D. Toll-like receptor-4 signaling mediates inflammation and tissue injury in diabetic nephropathy. *J Nephrol*. 2017;30(6):719–727. doi:10.1007/s40620-017-0432-8
4. Imler JL, Hoffmann JA. Toll receptors in innate immunity. *Trends Cell Biol*. 2001;11(7):304–311. doi:10.1016/s0962-8924(01)02004-9
5. Medzhitov R, Preston-Hurlburt P, Janeway CA. A human homologue of the *Drosophila* Toll protein signals activation of adaptive immunity. *Nature*. 1997;388(6640):394–397. doi:10.1038/411131
6. Takeda K, Akira S. Toll-like receptors. *Curr Protoc Immunol*. 2015;109:1–10. doi:10.1002/0471142735.im1412s109
7. Akira S, Takeda K. Toll-like receptor signalling. *Nat Rev Immunol*. 2004;4(7):499–511. doi:10.1038/nri1391
8. Brodsky I, Medzhitov R. Two modes of ligand recognition by TLRs. *Cell*. 2007;130(6):979–981. doi:10.1016/j.cell.2007.09.009
9. Bell JK, Botos I, Hall PR, et al. The molecular structure of the Toll-like receptor 3 ligand-binding domain. *Proc Natl Acad Sci U S A*. 2005;102(31):10976–10980. doi:10.1073/pnas.0505077102
10. Choe J, Kelker MS, Wilson IA. Crystal structure of human toll-like receptor 3 (TLR3) ectodomain. *Science*. 2005;309(5734):581–585. doi:10.1126/science.1115253
11. Kim HM, Park BS, Kim JI, et al. Crystal structure of the TLR4-MD-2 complex with bound endotoxin antagonist Eritoran. *Cell*. 2007;130(5):906–917. doi:10.1016/j.cell.2007.08.002
12. Jin MS, Kim SE, Heo JY, et al. Crystal structure of the TLR1-TLR2 heterodimer induced by binding of a tri-acylated lipopeptide. *Cell*. 2007;130(6):1071–1082. doi:10.1016/j.cell.2007.09.008

13. Kang JY, Nan X, Jin MS, et al. Recognition of lipopeptide patterns by Toll-like receptor 2-Toll-like receptor 6 heterodimer. *Immunity*. 2009;31(6):873–884. doi:10.1016/j.immuni.2009.09.018
14. Liu L, Botos I, Wang Y, et al. Structural basis of toll-like receptor 3 signaling with double-stranded RNA. *Science*. 2008;320(5874):379–381. doi:10.1126/science.1155406
15. Park BS, Song DH, Kim HM, Choi BS, Lee H, Lee JO. The structural basis of lipopolysaccharide recognition by the TLR4-MD-2 complex. *Nature*. 2009;458(7242):1191–1195. doi:10.1038/nature07830
16. Yoon SI, Kurnasov O, Natarajan V, et al. Structural basis of TLR5-flagellin recognition and signaling. *Science*. 2012;335(6070):859–864. doi:10.1126/science.1215584
17. Tanji H, Ohto U, Shibata T, Miyake K, Shimizu T. Structural reorganization of the Toll-like receptor 8 dimer induced by agonistic ligands. *Science*. 2013;339(6126):1426–1429. doi:10.1126/science.1229159
18. Ohto U, Shibata T, Tanji H, et al. Structural basis of CpG and inhibitory DNA recognition by Toll-like receptor 9. *Nature*. 2015;520(7549):702–705. doi:10.1038/nature14138
19. Song W, Wang J, Han Z, et al. Structural basis for specific recognition of single-stranded RNA by Toll-like receptor 13. *Nat Struct Mol Biol*. 2015;22(10):782–787. doi:10.1038/nsmb.3080
20. Nyman T, Stenmark P, Flodin S, Johansson I, Hammarström M, Nordlund P. The crystal structure of the human toll-like receptor 10 cytoplasmic domain reveals a putative signaling dimer. *J Biol Chem*. 2008;283(18):11861–11865. doi:10.1074/jbc.C800001200
21. Xu Y, Tao X, Shen B, et al. Structural basis for signal transduction by the Toll/interleukin-1 receptor domains. *Nature*. 2000;408(6808):111–115. doi:10.1038/35040600
22. Tao X, Xu Y, Zheng Y, Beg AA, Tong L. An extensively associated dimer in the structure of the C713S mutant of the TIR domain of human TLR2. *Biochem Biophys Res Commun*. 2002;299(2):216–221. doi:10.1016/S0006-291X(02)02581-0
23. Matsushima N, Tanaka T, Enkhbayar P, et al. Comparative sequence analysis of leucine-rich repeats (LRRs) within vertebrate toll-like receptors. *BMC Genom*. 2007;8:124. doi:10.1186/1471-2164-8-124
24. Kajava AV. Structural diversity of leucine-rich repeat proteins. *J Mol Biol*. 1998;277(3):519–527. doi:10.1006/jmbi.1998.1643
25. Kobe B, Kajava AV. The leucine-rich repeat as a protein recognition motif. *Curr Opin Struct Biol*. 2001;11(6):725–732. doi:10.1016/S0959-440X(01)00266-4
26. Gay NJ, Gangloff M. Structure and function of Toll receptors and their ligands. *Annu Rev Biochem*. 2007;76:141–165. doi:10.1146/annurev.biochem.76.060305.151318
27. Poltorak A, He X, Smirnova I, et al. Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene. *Science*. 1998;282(5396):2085–2088. doi:10.1126/science.282.5396.2085
28. Shimazu R, Akashi S, Ogata H, et al. MD-2, a molecule that confers lipopolysaccharide responsiveness on Toll-like receptor 4. *J Exp Med*. 1999;189(11):1777–1782. doi:10.1084/jem.189.11.1777
29. Kawai T, Adachi O, Ogawa T, Takeda K, Akira S. Unresponsiveness of MyD88-deficient mice to endotoxin. *Immunity*. 1999;11(1):115–122. doi:10.1016/S1074-7613(00)80086-2
30. Laird MH, Rhee SH, Perkins DJ, et al. TLR4/MyD88/PI3K interactions regulate TLR4 signaling. *J Leukoc Biol*. 2009;85(6):966–977. doi:10.1189/jlb.1208763
31. Takeda K, Akira S. TLR signaling pathways. *Semin Immunol*. 2004;16(1):3–9. doi:10.1016/j.smim.2003.10.003
32. Delaney JR, Mlodzik M. TGF-beta activated kinase-1: new insights into the diverse roles of TAK1 in development and immunity. *Cell Cycle (Georgetown, Tex)*. 2006;5(24):2852–2855. doi:10.4161/cc.5.24.3558
33. Saha RN, Pahan K. Regulation of inducible nitric oxide synthase gene in glial cells. *Antioxid Redox Signal*. 2006;8(5–6):929–947. doi:10.1089/ars.2006.8.929
34. Kawai T, Takeuchi O, Fujita T, et al. Lipopolysaccharide stimulates the MyD88-independent pathway and results in activation of IFN-regulatory factor 3 and the expression of a subset of lipopolysaccharide-inducible genes. *J Immunol*. 2001;167(10):5887–5894. doi:10.4049/jimmunol.167.10.5887
35. Okun E, Griffioen KJ, Lathia JD, Tang SC, Mattson MP, Arumugam TV. Toll-like receptors in neurodegeneration. *Brain Res Rev*. 2009;59(2):278–292. doi:10.1016/j.brainresrev.2008.09.001
36. Chuang TH, Ulevitch RJ. Triad3A, an E3 ubiquitin-protein ligase regulating Toll-like receptors. *Nat Immunol*. 2004;5(5):495–502. doi:10.1038/ni1066
37. Divanovic S, Trompette A, Atabani SF, et al. Inhibition of TLR4/MD-2 signaling by RP105/MD-1. *J Endotoxin Res*. 2005;11(6):363–368. doi:10.1179/096805105X67300
38. Hu X, Yu Y, Eugene Chin Y, Xia Q. The role of acetylation in TLR4-mediated innate immune responses. *Immunol Cell Biol*. 2013;91(10):611–614. doi:10.1038/icb.2013.56
39. Cerasi E. Insulin deficiency and insulin resistance in the pathogenesis of NIDDM: is a divorce possible? *Diabetologia*. 1995;38(8):992–997. doi:10.1007/BF00400591
40. Kahn SE, Hull RL, Utzschneider KM. Mechanisms linking obesity to insulin resistance and type 2 diabetes. *Nature*. 2006;444(7121):840–846. doi:10.1038/nature05482
41. Bonner-Weir S. Islet growth and development in the adult. *J Mol Endocrinol*. 2000;24(3):297–302. doi:10.1677/jme.0.0240297
42. Kahn BB. Type 2 diabetes: when insulin secretion fails to compensate for insulin resistance. *Cell*. 1998;92(5):593–596. doi:10.1016/S0092-8674(00)81125-3
43. Robertson RP, Harmon J, Tran POT, Poitout V. Beta-cell glucose toxicity, lipotoxicity, and chronic oxidative stress in type 2 diabetes. *Diabetes*. 2004;53(Suppl 1):S119–S124. doi:10.2337/diabetes.53.2007.S119
44. Weir GC, Bonner-Weir S. Five stages of evolving beta-cell dysfunction during progression to diabetes. *Diabetes*. 2004;53(Suppl 3):S16–S21. doi:10.2337/diabetes.53.suppl_3.S16
45. Prentki M, Nolan CJ. Islet beta cell failure in type 2 diabetes. *J Clin Invest*. 2006;116(7):1802–1812. doi:10.1172/JCI29103
46. Hull RL, Westermark GT, Westermark P, Kahn SE. Islet amyloid: a critical entity in the pathogenesis of type 2 diabetes. *J Clin Endocrinol Metab*. 2004;89(8):3629–3643. doi:10.1210/jc.2004-0405
47. Harding HP, Ron D. Endoplasmic reticulum stress and the development of diabetes: a review. *Diabetes*. 2002;51(Suppl 3):S455–S461. doi:10.2337/diabetes.51.2007.S455
48. Hotamisligil GS, Erbay E. Nutrient sensing and inflammation in metabolic diseases. *Nat Rev Immunol*. 2008;8(12):923–934. doi:10.1038/nri2449
49. Donath MY, Størling J, Maedler K, Mandrup-Poulsen T. Inflammatory mediators and islet beta-cell failure: a link between type 1 and type 2 diabetes. *J Mol Med*. 2003;81(8):455–470. doi:10.1007/s00109-003-0450-y
50. Ehres JA, Ellingsgaard H, Böni-Schnetzler M, Donath MY. Pancreatic islet inflammation in type 2 diabetes: from alpha and beta cell compensation to dysfunction. *Arch Physiol Biochem*. 2009;115(4):240–247. doi:10.1080/13813450903025879
51. Westwell-Roper CY, Ehres JA, Verchere CB. Resident macrophages mediate islet amyloid polypeptide-induced islet IL-1 β production and β -cell dysfunction. *Diabetes*. 2014;63(5):1698–1711. doi:10.2337/db13-0863

52. Westwell-Roper C, Dai DL, Soukhatcheva G, et al. IL-1 blockade attenuates islet amyloid polypeptide-induced proinflammatory cytokine release and pancreatic islet graft dysfunction. *J Immunol*. 2011;187(5):2755–2765. doi:10.4049/jimmunol.1002854
53. Masters SL, Dunne A, Subramanian SL, et al. Activation of the NLRP3 inflammasome by islet amyloid polypeptide provides a mechanism for enhanced IL-1 β in type 2 diabetes. *Nat Immunol*. 2010;11(10):897–904. doi:10.1038/ni.1935
54. Jourdan T, Godlewski G, Cinar R, et al. Activation of the Nlrp3 inflammasome in infiltrating macrophages by endocannabinoids mediates beta cell loss in type 2 diabetes. *Nat Med*. 2013;19(9):1132–1140. doi:10.1038/nm.3265
55. Oslowski CM, Hara T, O'Sullivan-Murphy B, et al. Thioredoxin-interacting protein mediates ER stress-induced β cell death through initiation of the inflammasome. *Cell Metab*. 2012;16(2):265–273. doi:10.1016/j.cmet.2012.07.005
56. Lerner AG, Upton J-P, Praveen PVK, et al. IRE1 α induces thioredoxin-interacting protein to activate the NLRP3 inflammasome and promote programmed cell death under irremediable ER stress. *Cell Metab*. 2012;16(2):250–264. doi:10.1016/j.cmet.2012.07.007
57. Ling PR, Smith RJ, Bistrian BR. Acute effects of hyperglycemia and hyperinsulinemia on hepatic oxidative stress and the systemic inflammatory response in rats. *Crit Care Med*. 2007;35(2):555–560. doi:10.1097/01.Ccm.0000253310.02180.C2
58. Apostolova N, Garcia-Bou R, Hernandez-Mijares A, Herance R, Rocha M, Victor VM. Mitochondrial antioxidants alleviate oxidative and nitrosative stress in a cellular model of sepsis. *Pharm Res*. 2011;28(11):2910–2919. doi:10.1007/s11095-011-0528-0
59. Lu H, Koshkin V, Allister EM, Gyulhandanyan AV, Wheeler MB. Molecular and metabolic evidence for mitochondrial defects associated with beta-cell dysfunction in a mouse model of type 2 diabetes. *Diabetes*. 2010;59(2):448–459. doi:10.2337/db09-0129
60. Armann B, Hanson MS, Hatch E, Steffen A, Fernandez LA. Quantification of basal and stimulated ROS levels as predictors of islet potency and function. *Am J Transpl*. 2007;7(1):38–47. doi:10.1111/j.1600-6143.2006.01577.x
61. Duprez J, Roma LP, Close AF, Jonas JC, Nadal A. Protective antioxidant and antiapoptotic effects of ZnCl₂ in rat pancreatic islets cultured in low and high glucose concentrations. *PLoS One*. 2012;7(10):e46831. doi:10.1371/journal.pone.0046831
62. Lucas K, Maes M. Role of the Toll Like receptor (TLR) radical cycle in chronic inflammation: possible treatments targeting the TLR4 pathway. *Mol Neurobiol*. 2013;48(1):190–204. doi:10.1007/s12035-013-8425-7
63. Maedler K, Fontana A, Ris F. FLIP switches Fas-mediated glucose signaling in human pancreatic β cells from apoptosis to cell replication. *Proc Natl Acad Sci USA*. 2002;99:8236–8241. doi:10.1073/pnas.122686299
64. Ehses JA, Perren A, Eppler E, et al. Increased number of islet-associated macrophages in type 2 diabetes. *Diabetes*. 2007;56(9):2356–2370. doi:10.2337/db06-1650
65. Richardson SJ, Willcox A, Bone AJ, Foulis AK, Morgan NG. Islet-associated macrophages in type 2 diabetes. *Diabetologia*. 2009;52(8):1686–1688. doi:10.1007/s00125-009-1410-z
66. Butcher MJ, Hallinger D, Garcia E, et al. Association of proinflammatory cytokines and islet resident leucocytes with islet dysfunction in type 2 diabetes. *Diabetologia*. 2014;57(3):491–501. doi:10.1007/s00125-013-3116-5
67. Böni-Schnetzler M, Thorne J, Parnaud G, et al. Increased interleukin (IL)-1 β messenger ribonucleic acid expression in β cells of individuals with type 2 diabetes and regulation of IL-1 β in human islets by glucose and autostimulation. *J Clin Endocrinol Metab*. 2008;93(10):4065–4074. doi:10.1210/jc.2008-0396
68. Kitamura T. The role of FOXO1 in beta-cell failure and type 2 diabetes mellitus. *Nat Rev Endocrinol*. 2013;9(10):615–623. doi:10.1038/nrendo.2013.157
69. Guest CB, Park MJ, Johnson DR, Freund GG. The implication of proinflammatory cytokines in type 2 diabetes. *Front Biosci*. 2008;13:5187–5194. doi:10.2741/3074
70. Festa A, D'Agostino R, Howard G, Mykkanen L, Tracy RP, Haffner SM. Chronic subclinical inflammation as part of the insulin resistance syndrome: the Insulin Resistance Atherosclerosis Study (IRAS). *Circulation*. 2000;102(1):42–47. doi:10.1161/01.cir.102.1.42
71. Navarro JF, Mora C. Diabetes, inflammation, proinflammatory cytokines, and diabetic nephropathy. *ScientificWorldJournal*. 2006;6:908–917. doi:10.1100/tsw.2006.179
72. Garay-Malpartida HM, Mourao RF, Mantovani M, Santos IA, Sogayar MC, Goldberg AC. Toll-like receptor 4 (TLR4) expression in human and murine pancreatic beta-cells affects cell viability and insulin homeostasis. *BMC Immunol*. 2011;12:18. doi:10.1186/1471-2172-12-18
73. Gupta S, Maratha A, Siednienko J, et al. Analysis of inflammatory cytokine and TLR expression levels in Type 2 Diabetes with complications. *Sci Rep*. 2017;7(1):7633. doi:10.1038/s41598-017-07230-8
74. Dasu MR, Devaraj S, Park S, Jialal I. Increased toll-like receptor (TLR) activation and TLR ligands in recently diagnosed type 2 diabetic subjects. *Diabetes Care*. 2010;33(4):861–868. doi:10.2337/dc09-1799
75. Sepehri Z, Kiani Z, Nasiri AA, et al. Human Toll like receptor 4 gene expression of PBMCs in diabetes mellitus type 2 patients. *Cell Mol Biol (Noisy-Le-Grand)*. 2015;61(3):92–95.
76. Ahmad R, Al-Mass A, Atizado V, et al. Elevated expression of the toll like receptors 2 and 4 in obese individuals: its significance for obesity-induced inflammation. *J Inflamm (Lond)*. 2012;9(1):48. doi:10.1186/1476-9255-9-48
77. Dasu MR, Devaraj S, Zhao L, Hwang DH, Jialal I. High glucose induces toll-like receptor expression in human monocytes: mechanism of activation. *Diabetes*. 2008;57(11):3090–3098. doi:10.2337/db08-0564
78. Sindhu S, Akhter N, Kochumon S, et al. Increased expression of the innate immune receptor TLR10 in obesity and type-2 diabetes: association with ROS-mediated oxidative stress. *Cell Physiol Biochem*. 2018;45(2):572–590. doi:10.1159/000487034
79. Edfeldt K, Swedenborg J, Hansson GK, Yan ZQ. Expression of toll-like receptors in human atherosclerotic lesions: a possible pathway for plaque activation. *Circulation*. 2002;105(10):1158–1161. doi:10.1161/circ.105.10.1158
80. Bobryshev YV, Lord RS. S-100 positive cells in human arterial intima and in atherosclerotic lesions. *Cardiovasc Res*. 1995;29(5):689–696. doi:10.1016/S0008-6363(96)88642-1
81. Stoll LL, Denning GM, Li WG, et al. Regulation of endotoxin-induced proinflammatory activation in human coronary artery cells: expression of functional membrane-bound CD14 by human coronary artery smooth muscle cells. *J Immunol*. 2004;173(2):1336–1343. doi:10.4049/jimmunol.173.2.1336
82. Vink A, Schoneveld AH, van der Meer JJ, et al. In vivo evidence for a role of toll-like receptor 4 in the development of intimal lesions. *Circulation*. 2002;106(15):1985–1990. doi:10.1161/01.cir.0000032146.75113.ee
83. Li H, Sun B. Toll-like receptor 4 in atherosclerosis. *J Cell Mol Med*. 2007;11(1):88–95. doi:10.1111/j.1582-4934.2007.00111.x
84. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW. Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest*. 2003;112(12):1796–1808. doi:10.1172/jci19246

85. Xu H, Barnes GT, Yang Q, et al. Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J Clin Invest*. 2003;112(12):1821–1830. doi:10.1172/jci19451
86. Clément K, Viguerie N, Poitou C, et al. Weight loss regulates inflammation-related genes in white adipose tissue of obese subjects. *FASEB J*. 2004;18(14):1657–1669. doi:10.1096/fj.04-2204com
87. Suganami T, Mieda T, Itoh M, Shimoda Y, Kamei Y, Ogawa Y. Attenuation of obesity-induced adipose tissue inflammation in C3H/HeJ mice carrying a Toll-like receptor 4 mutation. *Biochem Biophys Res Commun*. 2007;354(1):45–49. doi:10.1016/j.bbrc.2006.12.190
88. Shi L, Kishore R, McMullen MR, Nagy LE. Lipopolysaccharide stimulation of ERK1/2 increases TNF-alpha production via Egr-1. *Am J Physiol Cell Physiol*. 2002;282(6):C1205–C1211. doi:10.1152/ajpcell.00511.2001
89. Poggi M, Bastelica D, Gual P, et al. C3H/HeJ mice carrying a toll-like receptor 4 mutation are protected against the development of insulin resistance in white adipose tissue in response to a high-fat diet. *Diabetologia*. 2007;50(6):1267–1276. doi:10.1007/s00125-007-0654-8
90. Tsukumo DM, Carvalho-Filho MA, Carnevali JB, et al. Loss-of-function mutation in Toll-like receptor 4 prevents diet-induced obesity and insulin resistance. *Diabetes*. 2007;56(8):1986–1998. doi:10.2337/db06-1595
91. Saberi M, Woods NB, de Luca C, et al. Hematopoietic cell-specific deletion of toll-like receptor 4 ameliorates hepatic and adipose tissue insulin resistance in high-fat-fed mice. *Cell Metab*. 2009;10(5):419–429. doi:10.1016/j.cmet.2009.09.006
92. Ghanim H, Abuaysheh S, Sia CL, et al. Increase in plasma endotoxin concentrations and the expression of Toll-like receptors and suppressor of cytokine signaling-3 in mononuclear cells after a high-fat, high-carbohydrate meal: implications for insulin resistance. *Diabetes Care*. 2009;32(12):2281–2287. doi:10.2337/dc09-0979
93. Harte AL, Varma MC, Tripathi G, et al. High fat intake leads to acute postprandial exposure to circulating endotoxin in type 2 diabetic subjects. *Diabetes Care*. 2012;35(2):375–382. doi:10.2337/dc11-1593
94. van der Crabben SN, Blümler RME, Stegenga ME, et al. Early endotoxemia increases peripheral and hepatic insulin sensitivity in healthy humans. *J Clin Endocrinol Metab*. 2009;94(2):463–468. doi:10.1210/jc.2008-0761
95. Takeuchi O, Akira S. Toll-like receptors; their physiological role and signal transduction system. *Int Immunopharmacol*. 2001;1(4):625–635. doi:10.1016/S1567-5769(01)00010-8
96. Garate I, Garcia-Bueno B, Madrigal JL, et al. Stress-induced neuroinflammation: role of the Toll-like receptor-4 pathway. *Biol Psychiatry*. 2013;73(1):32–43. doi:10.1016/j.biopsych.2012.07.005
97. Schaefer TM, Desouza K, Fahey JV, Beagley KW, Wira CR. Toll-like receptor (TLR) expression and TLR-mediated cytokine/chemokine production by human uterine epithelial cells. *Immunology*. 2004;112(3):428–436. doi:10.1111/j.1365-2567.2004.01898.x
98. Shi H, Kokoeva MV, Inouye K, Tzameli I, Yin H, Flier JS. TLR4 links innate immunity and fatty acid-induced insulin resistance. *J Clin Invest*. 2006;116(11):3015–3025. doi:10.1172/JCI28898
99. Shen X, Yang L, Yan S, et al. Fetuin A promotes lipotoxicity in beta cells through the TLR4 signaling pathway and the role of pioglitazone in anti-lipotoxicity. *Mol Cell Endocrinol*. 2015;412:1–11. doi:10.1016/j.mce.2015.05.014
100. Pal D, Dasgupta S, Kundu R, et al. Fetuin-A acts as an endogenous ligand of TLR4 to promote lipid-induced insulin resistance. *Nat Med*. 2012;18(8):1279–1285. doi:10.1038/nm.2851
101. Wang X, Ge QM, Bian F, Dong Y, Huang CM. Inhibition of TLR4 protects rat islets against lipopolysaccharide-induced dysfunction. *Mol Med Rep*. 2017;15(2):805–812. doi:10.3892/mmr.2016.6097
102. Lu Z, Zhang X, Li Y, Jin J, Huang Y. TLR4 antagonist reduces early-stage atherosclerosis in diabetic apolipoprotein E-deficient mice. *J Endocrinol*. 2013;216(1):61–71. doi:10.1530/JOE-12-0338
103. Perrin-Cocon L, Aublin-Gex A, Sestito SE, et al. TLR4 antagonist FP7 inhibits LPS-induced cytokine production and glycolytic reprogramming in dendritic cells, and protects mice from lethal influenza infection. *Sci Rep*. 2017;7:40791. doi:10.1038/srep40791
104. Cighetti R, Ciaramelli C, Sestito SE, et al. Modulation of CD14 and TLR4/MD-2 activities by a synthetic lipid A mimetic. *Chembiochem*. 2014;15(2):250–258. doi:10.1002/cbic.201300588
105. Hu L, Yang H, Ai M, Jiang S. Inhibition of TLR4 alleviates the inflammation and apoptosis of retinal ganglion cells in high glucose. *Graefes Arch Clin Exp Ophthalmol*. 2017;255(11):2199–2210. doi:10.1007/s00417-017-3772-0
106. Lu Z, Zhang X, Li Y, Lopes-Virella MF, Huang Y. TLR4 antagonist attenuates atherogenesis in LDL receptor-deficient mice with diet-induced type 2 diabetes. *Immunobiology*. 2015;220(11):1246–1254. doi:10.1016/j.imbio.2015.06.016
107. Foresto-Neto O, Albino AH, Arias SCA, et al. NF-kappaB system is chronically activated and promotes glomerular injury in experimental type 1 diabetic kidney disease. *Front Physiol*. 2020;11:84. doi:10.3389/fphys.2020.00084
108. Yuan S, Liu X, Zhu X, et al. The role of TLR4 on PGC-1alpha-mediated oxidative stress in tubular cell in diabetic kidney disease. *Oxid Med Cell Longev*. 2018;2018:6296802. doi:10.1155/2018/6296802
109. Wu C, Lv C, Chen F, Ma X, Shao Y, Wang Q. The function of miR-199a-5p/Klotho regulating TLR4/NF-kappaB p65/NGAL pathways in rat mesangial cells cultured with high glucose and the mechanism. *Mol Cell Endocrinol*. 2015;417:84–93. doi:10.1016/j.mce.2015.09.024
110. Liu P, Li F, Xu X, et al. 1,25(OH)2D3 provides protection against diabetic kidney disease by downregulating the TLR4-MyD88-NF-kappaB pathway. *Exp Mol Pathol*. 2020;114:104434. doi:10.1016/j.yexmp.2020.104434
111. Cai S, Chen J, Li Y. Dioscin protects against diabetic nephropathy by inhibiting renal inflammation through TLR4/NF-kB pathway in mice. *Immunobiology*. 2020;225(3):151941. doi:10.1016/j.imbio.2020.151941
112. Ullah MA, Johora FT, Sarkar B, Araf Y, Rahman MH. Curcumin analogs as the inhibitors of TLR4 pathway in inflammation and their drug like potentialities: a computer-based study. *J Recept Signal Transduct Res*. 2020;1–15. doi:10.1080/10799893.2020.1742741
113. Lakhia R, Yheskel M, Flaten A, et al. Interstitial microRNA miR-214 attenuates inflammation and polycystic kidney disease progression. *JCI Insight*. 2020;5(7). doi:10.1172/jci.insight.133785
114. Shah M, Kim GY, Achek A, et al. The alphaC helix of TIRAP holds therapeutic potential in TLR-mediated autoimmune diseases. *Biomaterials*. 2020;245:119974. doi:10.1016/j.biomaterials.2020.119974
115. Yang K, Zhan L, Lu T, et al. Dendrobium officinale polysaccharides protected against ethanol-induced acute liver injury in vivo and in vitro via the TLR4/NF-kappaB signaling pathway. *Cytokine*. 2020;130:155058. doi:10.1016/j.cyt.2020.155058
116. Zheng Y, Wang JH, Liu LL, Li GY, Peng Y, Zhao TJ. Experimental study on growth and reproduction of Tlr4 knockout mice. *Chin J Compar Med*. 2018;28(12):41–43+50. doi:10.3969/j.issn.1671-7856.2018.12.007

117. Coenen KR, Gruen ML, Lee-Young RS, Puglisi MJ, Wasserman DH, Hasty AH. Impact of macrophage toll-like receptor 4 deficiency on macrophage infiltration into adipose tissue and the artery wall in mice. *Diabetologia*. 2009;52(2):318–328. doi:10.1007/s00125-008-1221-7
118. Lynn M, Rossignol DP, Wheeler JL, et al. Blocking of responses to endotoxin by E5564 in healthy volunteers with experimental endotoxemia. *J Infect Dis*. 2003;187(4):631–639. doi:10.1086/367990
119. Opal SM, Laterre PF, Francois B, et al. Effect of Eritoran, an antagonist of MD2-TLR4, on mortality in patients with severe sepsis the ACCESS randomized trial article. *JAMA*. 2013;309(11):1154–1162. doi:10.1001/jama.2013.2194

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Dovepress

Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert opinion

and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-targets-and-therapy-journal>