


Food Hygiene Practice and Its Determinants Among Food Handlers at University of Gondar, Northwest Ethiopia, 2019

This article was published in the following Dove Press journal:
International Journal of General Medicine

Ketseladingle Lema
Negasi Abuhay
Walegn Kindie
Henok Dagne 
Tadesse Guadu

Department of Environmental and Occupational Health and Safety, Institute of Public Health, College of Medicine and Health Sciences, University of Gondar, Gondar, Ethiopia

Introduction: Food hygiene is an essential matter of public health for protecting or preventing diseases caused by unsafe food due to lack of good quality from production to consumption.

Objective: The current study aimed at assessing the food hygiene practice and determinant factors among food handlers working at the University of Gondar.

Methods: Univariate and multivariable binary logistic regression analyses were used to test the association of covariates with the food safety practice. Variables with p -value <0.2 were candidates for multivariable analysis. The adjusted odds ratio with 95% confidence interval and p -values less than 0.05 were used to report associations in the final model.

Results: A total of 184(46.7%) of the study subjects had good self-reported food hygiene practice. Being male [Adjusted odds ratio (AOR): 2.37, 95% confidence interval (CI) (1.34, 4.19)], educational status (primary [AOR: 2.54, 95% CI (1.16, 5.58)] and secondary [AOR: 2.20, 95% CI (1.11, 4.37)]), workers with greater than 2 years work experience [AOR: 1.86, 95% CI (1.06, 3.25)], monthly income of 2044–4867ETB/month [AOR: 2.05, 95% CI (1.01, 4.16)] were independent predictors of food safety practice of food handlers.

Conclusion and Recommendations: Below half of the study subjects had good self-reported food hygiene practice. Sex, educational status, and income were factors associated with the food hygiene practice. There should be continuous supportive supervision to raise the skills of food handlers to comply to better food hygiene practice. Food hygiene training should be given especially to female food handlers. Frequent audits are also required to ensure the permanence of effective and continuous training. Regular medical check up and strict hygiene follow-up should be encouraged to prevent foodborne disease outbreaks at universities.

Keywords: food hygiene, knowledge, attitude and practice

Introduction

Food hygiene is an essential matter of public health for protecting or preventing diseases caused by unsafe food due to lack of good quality from production to consumption.¹

Foodborne disease (FBD) is of public health significance both in developed and developing nations. About 600 million individuals become ill every year due to consumption of contaminated food and an approximately 420,000 of these victims die per annum.² The World Health Organization (WHO) disclosed that 1 in 10 individuals worldwide are sick from foodborne illnesses.³ Foodborne infectious

Correspondence: Henok Dagne
Email enoch2313@gmail.com

diseases have been estimated to affect 550 million persons and cause 230,000 deaths globally in 2010 although it is difficult to determine the exact mortality associated with foodborne diseases.⁴ It is difficult to quantify the burden of foodborne diseases since most of the hazards that cause foodborne diseases are not transmitted exclusively by food.⁵

The consumption of contaminated food is correlated with an estimated 70% of diarrheal diseases in developing countries. The occurrence of foodborne illnesses is more common in developing nations because of poor hygiene, absence of drinking water, contaminated and inappropriate food storage equipment and absence of food safety education.⁶ In addition in low-income countries, in particular, FBDs outbreak is more serious due to inadequate sanitation, insufficient food safety regulations, weak regulatory structures, unsafe raw food, abused temperature, poor storage infrastructures, inadequate cooking, poor personal hygiene, improper handling methods, and cross-contamination of cooked food with uncooked raw food.⁷⁻⁹

Potential FBDs in institutions with a high number of people is a public health concern, since outbreaks in those places may affect a large number of consumers at once. Food handlers are expected to have excellent hygiene practice to reduce cross contamination and protect the consumers from foodborne diseases.^{10,11} Poor personal hygiene frequently contributes to foodborne illness which indicates that food handlers' knowledge and handling practices needs to be improved. Studies on the conditions of food and drink establishments have been scanty in Ethiopia.^{12,13}

In most developing countries hygiene is important since hygiene preventable diseases are prevalent. In Ethiopia these diseases account for 80% of the illnesses together with other infectious diseases and malnutrition.¹⁴

Food handlers with poor personal hygiene and lack of awareness of important issues in preventing foodborne diseases, working in food establishments could be potential sources of infections of many intestinal helminthes of protozoa and estrogenic pathogens.¹⁵

Institutional foodservice is an important sector of the food industry. Foods consumed at such institutions have been identified as important sources of foodborne disease outbreaks and often feature prominently in many national statistics on outbreaks of foodborne illness.¹⁶ Food poisoning could result in the institutional food service suffering huge financial losses and public confidence.¹⁰ This study, therefore was aimed at assessing the food hygiene practice

and determinant factors among food handlers at the University of Gondar.

Method

Study Design, Area and Period

An institution-based cross-sectional study design was used to assess food hygiene practice and determinant factors among food handlers at the University of Gondar (UoG) from April 1 to April 15, 2019. The UoG currently has five campuses; Atse Fasil campus (Technology College), College of Medicine and Health sciences (CMHS), Tseda Campus (Agriculture College), Atse Tewodros Campus (College of Natural and Computational Sciences), and Maraki Campus (College of Social Sciences and Humanities). A total of 645 food handlers (492 females and 153 males) are serving the five campuses.¹⁷

Study Population and Unit

All food handlers in the UoG students' cafeteria, 645 in total, were the source of our study population. The study units were all randomly selected food handlers from the study population. Workers who were absent during data collection time due to different reasons (maternity leave, sick and absent from work for any reason) were excluded from the study.

Sample Size Determination

Sample size (n) was determined by using a single population proportion formula,¹⁸ based on the following assumptions. The proportion of good food hygiene practice (p) was 47.7% from a study conducted among Addis Ababa university food handlers,¹⁹ standard normal distribution confidence interval ($z_{\alpha/2}$) (1.96), and margin of error (d) = 0.05.

$$n = \frac{(z_{\alpha/2})^2 \times p(1-p)}{d^2} \quad n = \frac{(1.96)^2 \times 0.477(1-0.477)}{0.05^2} \\ = 384$$

Taking 5% none response, the total sample size became 403.

Sampling Technique and Procedures

A simple random sampling technique was employed for selecting the study units and self-reported food hygiene practice was assessed. The total sample size was proportionally distributed in the five campuses based on the number of food handlers at each campus.

Operational Definitions

Food Hygiene Practice

The food handlers were asked 19 questions regarding their food hygiene practice. Study subjects who scored less than the mean value of the score of the practice questions were considered as having “poor food hygiene practices” and those who scored mean and above the mean value of the practice questions were considered as having “good food hygiene practice.”^{20–22}

Food Hygiene Knowledge

Food hygiene knowledge was assessed by asking food handlers 16 knowledge items (yes or no questions). Study subjects who scored below the mean score were considered as having “poor knowledge” and those who had a score of mean and above the mean were classified as having “good knowledge” about food hygiene.^{21,23,24}

Food Hygiene Attitude

Food hygiene attitude was measured by asking respondents 14 questions about their attitude. Study subjects with mean and above score of the attitude questions were considered as holding desirable (good) attitude and otherwise “poor attitude” towards food hygiene practice.^{21,22}

Data Collection Tool and Procedure

Data were collected using a self-administered structured questionnaire which included: sociodemographic characteristics, 19 practice questions with a three-point Likert scale (1=always, 2=sometimes and 3=never), 16 knowledge items (yes/no) regarding food-borne disease transmission, knowledge of personal hygiene, knowledge of cross contamination and knowledge of temperature control and 14 attitudinal questions with a four-point Likert scale (1-strongly agree, 2-agree, 3-disagree strongly and 4-disagree). The data collection tool was prepared after a careful literature survey.^{21–27} Content validity was done by doing a pretest. The pretest was done among 5% (ie, 21 food handlers out of the study population) and editorial and language adjustments were done based on the pretest results.

Data Processing and Analysis

Complete items were coded and entered onto Epi Info version 7 and transported to Statistical Package for the Social Science (SPSS) version 21 software for analysis. Univariate and multivariable binary logistic regression were employed to test the association of covariates with the outcome variable. Association was declared at p -value < 0.05.

Results

Overall, 394 study participants (98.2% response rate) were included in the current study. About 263 (66.8%) were female. The mean age of the study participants was 28 years. The majority of participants (62.4%) were married. About half of the participants (53.6%) had over 2 years' experience (Table 1).

Above three-fifths (61.9%) of the study subjects had good knowledge. Above half (54.8%) of the respondents had a desirable/good attitude but only 46.7% of the participants had good practice regarding food hygiene (Figure 1).

Factors Associated with Food Hygiene Practice

Sex, age, educational status, work experience, monthly income, family size, knowledge, attitude, job category and orientation about food hygiene had a p -value < 0.2 in the univariate binary logistic regression and hence were candidates to be used in the multivariable logistic analysis. Only monthly income, work experience, educational status, age and sex were associated with the food hygiene practice in the final model (p -value < 0.05).

Male food handlers were 2.37 times more likely to have better food hygiene practice than females [AOR: 2.37, 95% CI (1.34, 4.19)].

Study subjects with primary educational status were 2.54 times more likely to have better food hygiene practice [AOR: 2.54, 95% CI (1.16, 5.58)] and respondents with secondary educational status were 2.20 times more likely to have better food hygiene practice [AOR: 2.20, 95% CI (1.11, 4.37)] than those with college and above educational status.

Study subjects who had more than 2 years work experience were 1.86 times more likely to have better food hygiene practice than those with 2 years and less than 2 years' experience [AOR: 1.86, 95% CI (1.06, 3.25)].


Study subjects with 2044–4867 ETB average monthly income were 2.05 times more likely to report good food hygiene practice than those with an average income of 1100–1743 ETB [AOR: 2.05, 95% CI (1.01, 4.16)] (Table 2).

Discussion

This study was aimed at assessing the food safety practice and associated factors among food handlers in the University of Gondar. Accordingly, 46.7% with 95% CI (42.4%, 51.5%) of UoG food handlers had good food hygiene practice. Sex, work experience, educational status,

Table I Sociodemographic Characteristics of University of Gondar Food Handlers, 2019 (n=394)

Variables	Category	Frequency (n)	Percent (%)
Sex	Female	263	66.8
	Male	131	33.2
Age in years	19–26	108	27.4
	27–28	92	23.4
	29–32	78	19.8
	33–58	116	29.4
Marital status	Single	118	29.9
	Married	246	62.4
	Divorced	30	7.6
Educational level	Primary	123	31.2
	Secondary	208	52.8
	College and above	63	16.0
Religion	Orthodox	376	95.4
	Muslim	5	1.3
	Protestant	13	3.3
Experience in years	<2	211	53.6
	≥2	183	46.4
Income per month in ETB	1100–1743	136	34.5
	1743–2000	146	37.1
	2000–2043	16	4.1
	2043–4867	96	24.4
Family size	1–2	113	28.7
	2–3	102	25.9
	3–4	117	29.7
	4–9	62	15.7
Job category	Ticker/shift leader/manager	61	15.5
	Waiter	111	28.2
	Cooker/baker/chef	153	38.8
	Laborer	69	17.5
Food hygiene orientation	Monthly	109	27.7
	Biannually	114	28.9
	Annually	171	43.4

**Figure 1** Knowledge, attitude and practice of food hygiene among University of Gondar food handlers 2019 (n=394).

and monthly income were factors associated with food hygiene practice.

The proportion of food handlers with good food handling practice in the current survey is consistent with studies conducted in Ghana¹⁶ and Addis Ababa, Ethiopia.²⁸ However, the current food hygiene practice is lower than the practice levels reported in Malaysia,²⁹ and Nigeria.³⁰ The discrepancy in food hygiene practice level might be due to variations in the study tool used, time of the study and variation in sociodemographic and the socioeconomic status. Above half of the study subjects had poor food hygiene practice which could have a huge impact in the disease pathogenesis. According to the Codex Alimentarius Commission (2003), poor food handling is a main reason for the occurrence of foodborne diseases.³¹ A study done in Gondar town among food handlers earlier in 2008 indicated that 29.1% and 3.1% of the study subjects were positive for stool parasites and enteropathogenic bacterial species respectively.³² A study among Arba Minch University food handlers also revealed that among stool cultures 6.9% of *Salmonella* and 3% *Shigella* isolates were reported.³³ Several other studies conducted among food handlers at universities in Ethiopia also revealed that food handlers were vehicles of disease causing microorganisms such as *Shigella* and *Salmonella*.^{34–38} Food safety culture is required to create proper food handling and establishing regulations concerning food hygiene and safety.^{39,40}

Male study participants had better food hygiene practice than females in the current study. This is in line with a study conducted regarding food hygiene practice in Iran.⁴¹ But in other studies^{29,42} females had better food hygiene practice than males. There was no significant difference regarding food safety and hygiene

Table 2 Factors Associated with Food Hygiene Practice Among University of Gondar Food Handlers, 2019 (n=394)

Variables	Categories	Practice		COR (95% CI)	AOR (95% CI)
		Good (%)	Poor (%)		
Sex	Male	89(67.9%)	42(32.1%)	3.75(2.40,5.85)	2.37(1.34,4.19)***
	Female	95(36.1%)	168(63.9%)	I	I
Age	19–26	41(38.0%)	67(62.0%)	I	I
	27–28	44(47.8%)	48(52.2%)	1.50(0.85,2.63)	1.14(0.60,2.15)
	29–32	65(56.0%)	51(44.0%)	2.08(1.22,3.56)	1.64(0.83,3.25)
	33–58	34(43.6%)	44(56.4%)	1.26(0.70,2.28)	0.96(0.44,2.13)
Educational status	Primary	65(52.8%)	58(47.2%)	2.41(1.27,4.56)	2.54(1.16,5.58)*
	Secondary	99(47.6%)	109(52.4%)	1.95(1.08,3.55)	2.20(1.11,4.37)*
	College and above	20(31.7%)	43(68.3%)	I	I
Experience	≤2	89(42.2%)	122(57.8%)	I	I
	>2	95(51.9%)	88(48.1%)	1.48(0.99,2.20)	1.86(1.06,3.25)*
Monthly income ETB ^a	1100–1743	49(36.0%)	87(64.0%)	I	I
	1744–2000	74(50.7%)	72(49.3%)	1.82(1.13,2.94)	1.69(0.97,2.94)
	2001–2043	10(62.5%)	6(37.5%)	2.96(1.01,8.64)	2.11(0.60,7.39)
	2044–4867	51(53.1%)	45(46.9%)	2.01(1.18,3.43)	2.05(1.01,4.16)*
Family size	1–2	54(47.8%)	59(52.2%)	1.55(0.82,2.92)	2.03(0.94,4.39)
	2–3	45(44.1%)	57(55.9%)	1.34(0.70,2.56)	1.28(0.61,2.71)
	3–4	62(53.0%)	55(47.0%)	1.91(1.02,3.60)	1.63(0.80,3.33)
	4–9	23(37.1%)	39(62.9%)	I	I
Knowledge	Good	102(55.4%)	142(67.6%)	0.60(0.40,0.90)	0.78(0.49,1.26)
	Poor	82(44.6%)	68(32.4%)	I	I
Attitude	Good	112(51.9%)	104(48.1%)	1.58(1.06,2.37)	1.36(0.84,2.20)
	Poor	72(40.4%)	106(59.6%)	I	I
Job category	Ticker/shift leader/manager	30(49.2%)	31(%)	I	I
	Waiter	52(46.8%)	59(53.2%)	0.91(0.49,1.70)	0.73(0.33,1.63)
	Cooker/baker/chef	50(32.7%)	103(67.3%)	0.50(0.27,0.92)	0.56(0.26,1.21)
Food hygiene orientation	Monthly	59(54.1%)	50(45.9%)	1.79(1.10,2.90)	1.24(0.70,2.21)
	Biannually	57(50%)	57(50%)	1.52(0.94,2.44)	1.19(0.68,2.09)
	Annually	68(39.8%)	103(60.2%)	I	I

Notes: * $p < 0.05$, *** $p < 0.001$ Hosmer and Lemeshow goodness-of-fit 0.364, at approximately the rate of 1 USD = 27 ETB.

practice based on difference in gender in some previous studies.^{6,43–47} This might be due to other factors such as educational status and work role which could predict the food safety practice better than gender.

Educational status was inversely associated with food hygiene practice in the current study. However, in other studies educational status was positively associated with food hygiene practice.^{13,20,29,41,48} Other studies reported

no association between educational status and food hygiene practice.^{6,11,49,50} The food handlers who attended college and above education had poor food hygiene practice. This might be due to the fact that these workers are not frequently engaged in food handling and preparation.

Work experience was associated with food hygiene practice of respondents. Experienced food handlers had reported better food hygiene practice. This association is consistent with earlier studies conducted regarding food hygiene practice and determinant factors.^{20,28,29,41,42,51–54} However other studies reported that work experience has no association with level of food safety practice.⁴⁴ Work experience is important to develop better food hygiene practice as it enables workers better opportunity to undergo food hygiene training and orientation.

Monthly income was another factor associated with food hygiene practice of food handlers. Study participants with better income had better food hygiene practice in the current study. This is consistent with earlier studies.^{55–58} But in another study income was not significantly associated.²²

Knowledge, attitude and practice of food safety play a basic role in preventing and controlling food poisoning outbreaks.⁵⁹ Knowledge and attitude were not associated with food hygiene practice of study subjects in the current study. Numerous other previous studies^{11,60–64} also indicated no association between knowledge and food hygiene practice. In other studies,^{23,29,65} however, it was significantly associated with the participants' food hygiene practice. Earlier studies^{21,26,51,63,66–68} revealed that knowledge has a positive correlation with food hygiene practice. Whereas other literature^{11,69} showed that knowledge of food hygiene may not be translated to food hygiene practice. In another study it was negatively associated with food hygiene practice.⁷⁰ The difference may be due to the study participants having surface level knowledge that cannot bring about behavioral change. Attitude, also was not associated with the respondents of food hygiene practice in the current study. This is consistent with previous studies.^{71–78} However in other studies attitude was significantly associated with food hygiene practice.^{25,54,63,79–81}

Limitations of the Study

This study is susceptible to the participants being biased and does not represent the actual practice because the self-reported practices and behaviors were assessed. Self-reported practice is usually over estimated.⁸² Observational data provide the most reliable information

denoting actual food safety behavior.⁸² The other limitation is the lack of generalizability to the large population as this study was conducted among food handlers at university colleges. Finally, cross-sectional designs lack the capacity to definitely demonstrate cause-effect relationships because of the inherent limitation of the design. Also, because self-reported hygiene practice was assessed it may be affected by social desirability bias.

Conclusion and Recommendation

Only less than half of the study participants had good food hygiene practice. Sex, educational status, work experience and monthly income were factors significantly associated with food handlers' food hygiene practice. Strategies targeted at improving the hygienic practice of food handlers are imperative. There should be continuous supportive supervision to raise the skills of food handlers to comply to better food hygiene practice. Targeted food hygiene improvement training should be given especially to female food handlers. Additionally, frequent audits are also required to ensure the permanence of effective and continuous training. Regular medical check ups and strict hygiene follow-up should be encouraged to prevent foodborne disease outbreaks at universities.

Abbreviations

AOR, Adjusted odds ratio; COR, Crude odds ratio; CI, Confidence interval; SPSS, Statistical package for social sciences.

Data Sharing Statement

Data will be available upon request from the corresponding author.

Ethics Approval and Consent to Participate

Ethical clearance was obtained from the Environmental and Occupational Health and Safety ethical review committee at the University of Gondar with ethical approval number of EOHS/841/2011. The purpose of the study was clearly explained to the study subjects. Written consent was obtained. Confidentiality of the information was maintained at all levels of the study. Health education was given for study subjects about good food hygiene practice after the data collection was over. This research was carried out in accordance with the principles of the Declaration of Helsinki.

Consent for Publication

This paper does not contain any individual person's data.

Acknowledgments

The authors are pleased to acknowledge study participants for their unreserved contributions to the success of this study. The authors are also pleased to acknowledge the University of Gondar for covering the questionnaire duplication fee.

Author Contributions

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed on the journal to which the article was submitted; gave final approval of the version to be published; and agreed to be accountable for all aspects of the work.

Funding

There was no funding for this research but the University of Gondar has covered the questionnaire duplication fee.

Disclosure

The authors report no conflicts of interest for this work.

References

- Ethiopian Ministry of Health. Hygiene And Environment. 2015.
- Zanin LM, da Cunha DT, de Rosso VV, Capriles VD, Stedefeldt E. Knowledge, attitudes and practices of food handlers in food safety: an integrative review. *Food Res Int*. 2017;100(Pt1):53–62. doi:10.1016/j.foodres.2017.07.042
- World Health Organization. *WHO Estimates of the Global Burden of Foodborne Diseases: Foodborne Disease Burden Epidemiology Reference Group 2007–2015*. World Health Organization; 2015.
- Pal M, Ayele Y. Emerging role of foodborne viruses in public health. *Biomed Res*. 2020;5:01–4.
- Hald T, Aspinall W, Devleeschauwer B, et al. World Health Organization estimates of the relative contributions of food to the burden of disease due to selected foodborne hazards: a structured expert elicitation. *PLoS One*. 2016;11(1):e0145839. doi:10.1371/journal.pone.0145839
- Stratev D, Odeyemi OA, Pavlov A, Kyuchukova R, Fatehi F, Bamidele FA. Food safety knowledge and hygiene practices among veterinary medicine students at Trakia University, Bulgaria. *J Infect Public Health*. 2017;10(6):778–782. doi:10.1016/j.jiph.2016.12.001
- Odeyemi OA, Bamidele FA. Harnessing the potentials of predictive microbiology in microbial food safety and quality research in Nigeria. *Future Sci*. 2016;2. doi:10.4155/fso.15.91
- Odeyemi OA, Sani NA. Antibiotic resistance and burden of foodborne diseases in developing countries. *Future Sci*. 2016;2:FSO139. doi:10.4155/fsoa-2016-0023
- Lamuka P. Public health measures: challenges of developing countries in management of food safety. In: Motarjemi Y, editor. *Encyclopedia of Food Safety*, Volume 4, pp. 20–26. Waltham, MA: Academic Press; 2014.
- Abdul-Mutalib N-A, Abdul-Rashid M-F, Mustafa S, Amin-Nordin S, Hamat RA, Osman M. Knowledge, attitude and practices regarding food hygiene and sanitation of food handlers in Kuala Pilah, Malaysia. *Food Control*. 2012;27(2):289–293.
- Akabanda F, Hlortsi EH, Owusu-Kwarteng J. Food safety knowledge, attitudes and practices of institutional food-handlers in Ghana. *BMC Public Health*. 2017;17(1):40. doi:10.1186/s12889-016-3986-9
- Gemeda T, Asayehu T, Abdisa M, Fekadu H. Assessment of knowledge, attitude and practices of food handlers in Nekemte Referral Hospital, Wollega, Ethiopia. *J Nutr Health Food Eng*. 2018;8(1):00262.
- Lestantyo D, Husodo AH, Iravati S, Shaluhiah Z. Safe food handling knowledge, attitude and practice of food handlers in hospital kitchen. *IJPHS*. 2017;6(4):324. doi:10.11591/ijphs.v6i4.10778
- Gebreeyessus GD, Adem DB. Knowledge, attitude, and practice on hygiene and morbidity status among tertiary students: the case of Kotebe Metropolitan University, Addis Ababa, Ethiopia. *J Environ Public Health*. 2018;2018:2094621. doi:10.1155/2018/2094621
- Kubde S, Pattankar J, Kokiwar P. Knowledge and food hygiene practices among food handlers in food establishments. *Int J Commun Med Public Health*. 2016;251–256. doi:10.18203/2394-6040.ijcmph20151572
- Parry-Hanson Kunadu A, Ofosu DB, Aboagye E, Tano-Debrah K. Food safety knowledge, attitudes and self-reported practices of food handlers in institutional foodservice in Accra, Ghana. *Food Control*. 2016;69:324–330. doi:10.1016/j.foodcont.2016.05.011
- University of Gondar. University of Gondar official site; 2019. Available from: <http://www.uog.edu.et>.
- Arifin WN. Introduction to sample size calculation. *Educ Med J*. 2013;5(2). doi:10.5959/eimj.v5i2.130
- Meleko A, Henok A, Tefera W, Lamaro T. Assessment of the sanitary conditions of catering establishments and food safety knowledge and practices of food handlers in Addis Ababa University Students' Cafeteria. *Science*. 2015;3(5):733–743.
- Legesse DTM, Agedew E, Haftu D. Food handling practices and associated factors among food handlers in Arba Minch Town public food establishments in Gamo Gofa Zone, Southern Ethiopia. *Epidemiology (Open Access)*. 2017;07(02).
- Dagne H, Raju R, Andualem Z, Hagos T, Addis K. Food safety practice and its associated factors among mothers in Debarq Town, Northwest Ethiopia: community-based cross-sectional study. *Biomed Res Int*. 2019;2019.
- Tessema AG, Gelaye KA, Chercos DH. Factors affecting food handling Practices among food handlers of Dangila town food and drink establishments, North West Ethiopia. *BMC Public Health*. 2014;14(1):571. doi:10.1186/1471-2458-14-571
- Kibret M, Abera B. The sanitary conditions of food service establishments and food safety knowledge and practices of food handlers in Bahir Dar town. *Ethiop J Health Sci*. 2012;22(1):27–35.
- Legesse D, Tilahun M, Agedew E, Haftu D. Food handling practices and associated factors among food handlers in Arba Minch Town public food establishments in Gamo Gofa Zone, Southern Ethiopia. *Epidemiology (Sunnyvale)*. 2017;7(02):2161. doi:10.4172/2161-1165.1000302
- Chekol FA, Melak MF, Belew AK, Zeleke EG. Food handling practice and associated factors among food handlers in public food establishments, Northwest Ethiopia. *BMC Res Notes*. 2019;12(1):20. doi:10.1186/s13104-019-4047-0
- Baraki N, Egata G. *Food Safety Practice and Associated Factors Among Food Handlers in Public Food Establishments of Godey Town*. Somali Region, Eastern Ethiopia: Haramaya University; 2018.
- Demssie A, Daniel D, Tefera A, Kindu H, Abebe S, Sanbata H. Knowledge, attitude and practice (KAP) of hand washing among mothers of under five children in Gotu Kebele Wondogenet Woreda Oromia, Ethiopia. *Int J Environ Sci Technol*. 2017;6(4):146–153.

28. Meleko A. Assessment of the sanitary conditions of catering establishments and food safety knowledge and practices of food handlers in Addis Ababa University students' cafeteria. *Sci J Public Health*. 2015;3(5):733. doi:10.11648/j.sjph.20150305.30
29. Sharif L, Obaidat MM, Al-Dalalah M-R. Food hygiene knowledge, attitudes and practices of the food handlers in the military hospitals. *Food Nutr Sci*. 2013;04(03):245–251. doi:10.4236/fns.2013.43033
30. Faremi FA, Olatubi MI, Nnabuike GC. Food safety and hygiene practices among food vendors in a Tertiary Educational Institution in South Western Nigeria. *Eur J Nutr Food Saf*. 2018;8:59–70. doi:10.9734/EJNFS/2018/39368
31. Codex Alimentarius Commission. *Recommended International Code of Practice General Principles of Food Hygiene*. Rome, Italy: Codex Alimentarius Commission; 2003.
32. Andargie G, Kassu A, Moges F, Tiruneh M, Huruy K. Prevalence of bacteria and intestinal parasites among food-handlers in Gondar town, northwest Ethiopia. *J Health Popul Nutr*. 2008;26(4):451.
33. Mama M, Alemu G. Prevalence, antimicrobial susceptibility patterns and associated risk factors of Shigella and Salmonella among food handlers in Arba Minch University, South Ethiopia. *BMC Infect Dis*. 2016;16(1):686. doi:10.1186/s12879-016-2035-8
34. Dagnew M, Tiruneh M, Moges F, Gizachew M. Bacterial profile and antimicrobial susceptibility pattern among food handlers at Gondar University Cafeteria, Northwest Ethiopia. *J Infect Dis Ther*. 2013;1–6.
35. Mengist A, Mengistu G, Reta A. Prevalence and antimicrobial susceptibility pattern of Salmonella and Shigella among food handlers in catering establishments at Debre Markos University, Northwest Ethiopia. *Int J Infect Dis*. 2018;75:74–79. doi:10.1016/j.ijid.2018.08.008
36. Marami D, Hailu K, Tolera M. Prevalence and antimicrobial susceptibility pattern of Salmonella and Shigella species among asymptomatic food handlers working in Haramaya University cafeterias, Eastern Ethiopia. *BMC Res Notes*. 2018;11(1):74. doi:10.1186/s13104-018-3189-9
37. Gebreyesus A, Adane K, Negash L, et al. Prevalence of Salmonella typhi and intestinal parasites among food handlers in Mekelle University student cafeteria, Mekelle, Ethiopia. *Food Control*. 2014;44:45–48. doi:10.1016/j.foodcont.2014.03.040
38. Diriba K, Awulachew E, Ashuro Z. Prevalence and antimicrobial resistance pattern of Salmonella, Shigella, and intestinal parasites and associated factor among food handlers in Dilla University Student Cafeteria, Dilla, Ethiopia. *Int J Microbiol*. 2020;2020:1–10. doi:10.1155/2020/3150539
39. de Andrade ML, Stedefeldt E, Zanin LM, da Cunha DT. Food safety culture in food services with different degrees of risk for foodborne diseases in Brazil. *Food Control*. 2020;112:107152. doi:10.1016/j.foodcont.2020.107152
40. Nyarugwe SP, Linnemann A, Nyanga LK, Fogliano V, Luning PA. Food safety culture assessment using a comprehensive mixed-methods approach: a comparative study in dairy processing organisations in an emerging economy. *Food Control*. 2018;84:186–196. doi:10.1016/j.foodcont.2017.07.038
41. Mehrdad Askarian M, Gholamhosein Kabir P, Maria A, et al. Knowledge, attitudes, and practices of food service staff regarding food hygiene in Shiraz, Iran. *Infect Control Hosp Epidemiol*. 2004;25(1):16–20.
42. Nee SO, Sani NA. Assessment of knowledge, attitudes and practices (KAP) among food handlers at residential colleges and canteen regarding food safety. *Sains Malaysiana*. 2011;40(4):403–410.
43. Alqurashi NA, Priyadarshini A, Jaiswal AK. Evaluating food safety knowledge and practices among foodservice staff in Al Madinah Hospitals, Saudi Arabia. *Safety*. 2019;5(1):9. doi:10.3390/safety5010009
44. Baluka SA, Miller R, Kaneene JB. Hygiene practices and food contamination in managed food service facilities in Uganda. *Afri J Food Sci*. 2015;9(1):31–42. doi:10.5897/AJFS2014.1170
45. Suryani D, Sutomo AH, Aman AT. The factors associated with food safety practices on food handlers in primary school canteens. *Unnes J Public Health*. 2019;8(1):1–9. doi:10.15294/ujph.v8i1.22830
46. Admasu M, Kelbessa W. Food safety knowledge, handling practice and associated factors among food handlers of hotels/restaurants in Asosa Town, North Western Ethiopia. *SM J Public Health Epidemiol*. 2018;4(1):1051.
47. Woh PY, Thong KL, Behnke JM, Lewis JW, Zain SNM. Evaluation of basic knowledge on food safety and food handling practices amongst migrant food handlers in Peninsular Malaysia. *Food Control*. 2016;70:64–73. doi:10.1016/j.foodcont.2016.05.033
48. Onyeneho S, Hedberg C. An assessment of food safety needs of restaurants in Owerri, Imo State, Nigeria. *Int J Environ Res Public Health*. 2013;10(8):3296–3309. doi:10.3390/ijerph10083296
49. Acikel CH, Ogur R, Yaren H, Gogeldi E, Ucar M, Kir T. The hygiene training of food handlers at a teaching hospital. *Food Control*. 2008;19(2):186–190. doi:10.1016/j.foodcont.2007.03.008
50. Al-Mohaithef M. *Food Hygiene in Hospitals: Evaluating Food Safety Knowledge, Attitudes and Practices of Foodservice Staff and Prerequisite Programs in Riyadh's Hospitals*. Saudi Arabia: University of Birmingham; 2014.
51. Çakıroğlu FP, Uçar A. Employees' perception of hygiene in the catering industry in Ankara (Turkey). *Food Control*. 2008;19(1):9–15. doi:10.1016/j.foodcont.2007.01.001
52. Thelwell-Reid MA. Food safety knowledge and self-reported practices of food handlers in Jamaica. Dissertation. Walden University; 2014.
53. Saad M, See TP, Adil MAM. Hygiene practices of food handlers at Malaysian government institutions training centers. *Procedia Social Behav Sci*. 2013;85:118–127. doi:10.1016/j.sbspro.2013.08.344
54. Lee H, Abdul Halim H, Thong K, Chai L. Assessment of food safety knowledge, attitude, self-reported practices, and microbiological hand hygiene of food handlers. *Int J Environ Res Public Health*. 2017;14(1):55. doi:10.3390/ijerph14010055
55. Gizaw Z, Gebrehiwot M, Tekla Z. Food safety knowledge, attitude and associated factors of food handlers working in substandard food establishments in Gondar Town, Northwest Ethiopia, 2013. *Int J Med Health Sci Res*. 2014;1:37–49.
56. Olumakaiye MF, Bakare KO. Training of food providers for improved environmental conditions of food service outlets in urban area Nigeria. *Food Nutr Sci*. 2013;4(07):99. doi:10.4236/fns.2013.47A012
57. Adebukola OC, Opeyemi AO, Ayodeji AI. Knowledge of food borne infection and food safety practices among local food handlers in Ijebu-Ode Local Government Area of Ogun State. *J Public Health Epidemiol*. 2015;7(9):268–273.
58. Gizaw Z, Gebrehiwot M, Tekla Z. Food safety practice and associated factors of food handlers working in substandard food establishments in Gondar Town, Northwest Ethiopia, 2013/14. *Int J Food Sci Nutr Diet*. 2014;3(7):138–146.
59. Sharif L, Al-Malki T. Knowledge, attitude and practice of Taif University students on food poisoning. *Food Control*. 2010;21(1):55–60. doi:10.1016/j.foodcont.2009.03.015
60. Grema HA, Kwaga J, Bello M, Onimisi HU. Assessment of food hygiene knowledge, attitudes and practices of fish handlers in Kaduna State, Nigeria. *Adv Anim Veter Sci*. 2018;7(3):131–137.
61. Garayoa R, Vitas AI, Diez-Leturia M, García-Jalón I. Food safety and the contract catering companies: food handlers, facilities and HACCP evaluation. *Food Control*. 2011;22(12):2006–2012. doi:10.1016/j.foodcont.2011.05.021
62. Bassyouni RH, El-Sherbiny N, Hefzy EH, Wegdan -A-A. Perception of food safety and prevalence of Staphylococcus aureus and Salmonella species carriers among Fayoum University food handlers. *Life Sci J*. 2012;9(4):2934–2940.
63. Rahman MM, Arif MT, Bakar K, Talib Z. Food safety knowledge, attitude and hygiene practices among the street food vendors in Northern Kuching City, Sarawak. *Borneo Sci*. 2016;31.

64. Vo TH, Le NH, Le ATN, Minh NNT, Nuorti JP. Knowledge, attitudes, practices and training needs of food-handlers in large canteens in Southern Vietnam. *Food Control*. 2015;57:190–194. doi:10.1016/j.foodcont.2015.03.042
65. Bamidele JO, Adebimpe WO, Oladele EA, Adeoye OA. Hygiene practices among workers in local eateries of Orolu community in South Western Nigeria. *Ann Med Health Sci Res*. 2015;5(4):235–240.
66. Iwu AC, Uwakwe KA, Duru CB, et al. Knowledge, attitude and practices of food hygiene among food vendors in Owerri, Imo State, Nigeria. *Occup Dis Environ Med*. 2017;5(1):11–25. doi:10.4236/odem.2017.51002
67. Bamidele J, Adebimpe W, Oladele E, Adeoye O. Hygiene practices among workers in local eateries of Orolu community in South Western Nigeria. *Ann Med Health Sci Res*. 2015;5(4):235–240.
68. Al-Shabib NA, Mosilhey SH, Husain FM. Cross-sectional study on food safety knowledge, attitude and practices of male food handlers employed in restaurants of King Saud University, Saudi Arabia. *Food Control*. 2016;59:212–217. doi:10.1016/j.foodcont.2015.05.002
69. Ansari-Lari M, Soodbakhsh S, Lakzadeh L. Knowledge, attitudes and practices of workers on food hygienic practices in meat processing plants in Fars, Iran. *Food Control*. 2010;21(3):260–263. doi:10.1016/j.foodcont.2009.06.003
70. Rossi MSC, Stedefeldt E, da Cunha DT, de Rosso VV. Food safety knowledge, optimistic bias and risk perception among food handlers in institutional food services. *Food Control*. 2017;73:681–688. doi:10.1016/j.foodcont.2016.09.016
71. Al-Kandari D, Al-abdeen J, Sidhu J. Food safety knowledge, attitudes and practices of food handlers in restaurants in Kuwait. *Food Control*. 2019;103:103–110. doi:10.1016/j.foodcont.2019.03.040
72. Angelillo IF, Viggiani NM, Rizzo L, Bianco A. Food handlers and foodborne diseases: knowledge, attitudes, and reported behavior in Italy. *J Food Prot*. 2000;63(3):381–385. doi:10.4315/0362-028X-63.3.381
73. Zain MM, Naing NN. Sociodemographic characteristics of food handlers and their knowledge, attitude and practice towards food sanitation: a preliminary report. *Southeast Asian J Trop Med Public Health*. 2002;33(2):410–417.
74. Askarian M, Kabir G, Aminbaig M, Memish ZA, Jafari P. Knowledge, attitudes, and practices of food service staff regarding food hygiene in Shiraz, Iran. *Infect Control Hosp Epidemiol*. 2004;25(1):16–20.
75. Tokuç B, Ekuklu G, Berberoğlu U, Bilge E, Dedeler H. Knowledge, attitudes and self-reported practices of food service staff regarding food hygiene in Edirne, Turkey. *Food Control*. 2009;20(6):565–568. doi:10.1016/j.foodcont.2008.08.013
76. Faour-Klingbeil D, Kuri V, Todd E. Investigating a link of two different types of food business management to the food safety knowledge, attitudes and practices of food handlers in Beirut, Lebanon. *Food Control*. 2015;55:166–175. doi:10.1016/j.foodcont.2015.02.045
77. Samapundo S, Thanh TC, Khaferi R, Devlieghere F. Food safety knowledge, attitudes and practices of street food vendors and consumers in Ho Chi Minh city, Vietnam. *Food Control*. 2016;70:79–89. doi:10.1016/j.foodcont.2016.05.037
78. Ko W-H. The relationship among food safety knowledge, attitudes and self-reported HACCP practices in restaurant employees. *Food Control*. 2013;29(1):192–197. doi:10.1016/j.foodcont.2012.05.076
79. Ismail FH, Chik CT, Muhammad R, Yusoff NM. Food safety knowledge and personal hygiene practices amongst mobile food handlers in Shah Alam, Selangor. *Procedia Social Behav Sci*. 2016;222:290–298. doi:10.1016/j.sbspro.2016.05.162
80. Lim T-P, Chye FY, Sulaiman MR, Suki NM, Lee J-S. A structural modeling on food safety knowledge, attitude, and behaviour among Bum Bum Island community of Semporna, Sabah. *Food Control*. 2016;60:241–246. doi:10.1016/j.foodcont.2015.07.042
81. da Cunha DT, de Rosso VV, Pereira MB, Stedefeldt E. The differences between observed and self-reported food safety practices: a study with food handlers using structural equation modeling. *Food Res Int*. 2019;125:108637. doi:10.1016/j.foodres.2019.108637
82. Evans EW, Redmond EC. Behavioral observation and microbiological analysis of older adult consumers' cross-contamination practices in a model domestic kitchen. *J Food Prot*. 2018;81(4):569–581. doi:10.4315/0362-028X.JFP-17-378

International Journal of General Medicine

Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-general-medicine-journal>

Dovepress

across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.