

Intravenous Administration of Triptonide Attenuates CFA-Induced Pain Hypersensitivity by Inhibiting DRG AKT Signaling Pathway in Mice

This article was published in the following Dove Press journal:
Journal of Pain Research

Yue-Juan Ling^{1,2}
Ting-Yu Ding^{1,2}
Fu-Lu Dong³
Yong-Jing Gao^{1,2}
Bao-Chun Jiang^{1,2}

¹Institute of Pain Medicine, Nantong University, Nantong, Jiangsu 226019, People's Republic of China; ²Institute of Special Environmental Medicine, Nantong University, Nantong, Jiangsu 226019, People's Republic of China; ³School of Medicine, Nantong University, Nantong, Jiangsu, People's Republic of China

Background: Currently, medical treatment of inflammatory pain is limited by a lack of safe and effective therapies. Triptonide (TPN), a major component of *Tripterygium wilfordii Hook.f.* with low toxicity, has been shown to have good anti-inflammatory and neuroprotective effects. The present study aims to investigate the effects of TPN on chronic inflammatory pain.

Materials and Methods: Inflammatory pain was induced by intraplantar injection of complete Freund's adjuvant (CFA). TPN's three different doses were intravenously administered to compare the analgesic efficacy: 0.1 mg/kg, 0.5 mg/kg, and 2.0 mg/kg. The foot swelling was quantitated by measuring paw volume. Mechanical allodynia and thermal hyperalgesia were assessed with von Frey filament testing and Hargreaves' test, respectively. Western blots, qRT-PCR and immunofluorescence tests were used to analyze the expression of pAKT, tumor necrosis factor- α (TNF- α), interleukin 1 beta (IL-1 β), and interleukin 6 (IL-6). Two AKT inhibitors, AKT inhibitor IV and MK-2206, were used to examine AKT's effects on pain behavior and cytokines expression.

Results: Intravenous treatment with TPN attenuated CFA-induced paw edema, mechanical allodynia, and thermal hyperalgesia. Western blotting and immunofluorescence results showed that CFA induced AKT activation in the dorsal root ganglion (DRG) neurons. However, these effects were suppressed by treatment with TPN. Furthermore, TPN treatment inhibited CFA-induced increase of pro-inflammatory cytokines, including TNF- α , IL-1 β , and IL-6. Consistent with the in vivo data, TPN inhibited LPS-induced Akt phosphorylation and inflammatory mediator production in ND7/23 cells. Finally, intrathecal treatment with AKT inhibitor IV or MK-2206, attenuated CFA-induced mechanical allodynia and thermal hyperalgesia, and simultaneously decreased the mRNA expression of TNF- α , IL-1 β , and IL-6 in DRG.

Conclusion: These data indicate that TPN attenuates CFA-induced pain potentially via inhibiting AKT-mediated pro-inflammatory cytokines production in DRG. TPN may be used for the treatment of chronic inflammatory pain.

Keywords: triptonide, AKT, DRG, inflammatory pain, pro-inflammatory factors

Correspondence: Bao-Chun Jiang
Institute of Pain Medicine, Nantong University, 9 Seyuan Road, Nantong, Jiangsu 226019, People's Republic of China
Tel +86-513-55003373
Email jiangbaochun@ntu.edu.cn

Introduction

Tripterygium wilfordii Hook. f., a member of the Celastraceae family of plants, is one of the fundamental herbs in traditional Chinese herbal medicine. The diterpenoid triepoxide chemicals are the main biologically active ingredients in the root of *Tripterygium wilfordii*. They have been used as a part of systematic medication to treat a wide variety

of diseases, including cancer, lupus, rheumatoid arthritis, Alzheimer's disease, Parkinson's disease, and rheumatoid arthritis.^{1–3}

Triptolide (TPL) was the first monomer isolated from *Tripterygium wilfordii* and structurally characterized in 1972.⁴ TPL was shown to be able to relieve many different types of pain. It can effectively relieve neuropathic pain by inhibiting microglial activation and astrocytes in the spinal dorsal horn.^{5,6} TPL also attenuates cancer pain via suppressing the up-regulation of *Chemokine (C-C motif) ligand 5* and histone deacetylases in the spinal glial cells.^{7,8} In addition, TPL inhibits the activation of extracellular signal-regulated kinase (ERK) pathway and the production of inflammatory cytokines in the spinal cord dorsal horn induced by inflammation.⁹ Recently, TPL was reported to have a potent anti-depressive function by its influences on hippocampal neuroinflammation in a rat model of depression comorbidity of chronic pain.² However, clinical reports showed that TPL exposure resulted in the injury of some organs, including liver, kidney, heart, testes, and ovary in humans.^{10–12} Severe hepatotoxicity was also shown after TPL exposure in animals.^{10,13,14}

Triptonide (TPN) is another bioactive component of *Tripterygium wilfordii*. It was reported that TPN does not induce severe liver toxicity in animals.^{1,3,15} TPN acts as a novel potent anticancer drugs with low toxicity.^{16–19} The differences between the chemical structures of TPL and TPN are the substituent groups at C-14 position in which TPL is with C-14-hydroxyl and TPN is with C-14-carbonyl (Figure 1A and B). The metabolomics study shows that the hydroxyl group at C-14 in the molecular structure of TPL plays an important role in its hepatotoxicity.^{15,20}

Sensitization of peripheral nociceptors by pro-inflammatory factors is a major factor of neuropathic pain and inflammation pain.^{21,22} The *tumor necrosis factor-alpha* (TNF- α), *interleukin-1 beta* (IL-1 β) and *interleukin-6* (IL-6)

in the dorsal root ganglia (DRG) contributes to the pathogenesis of chronic pain.^{23–25} Phosphoinositide 3-kinase/protein kinase B (PI3K/AKT) pathway regulate the TNF- α , IL-1 β and IL-6 expression in different cells.^{26,27} PI3K/AKT pathway could be activated in DRG neurons and spinal glial cells in different pain models.^{28–30} Recent studies showed that TPN can diminish AKT signaling pathways in cervical cancer cells or lymphoma cells.^{1,31} Although the analgesic function of TPL has been studied, whether TPN has an analgesic effect and the underlying mechanisms remain unknown.

The intraplantar injection of complete Freund's adjuvant (CFA) is frequently used to establish a chronic inflammatory pain model since it can trigger inflammatory responses, including the formation of an inflammatory infiltrate, swelling and cytokine release. In this study, we investigated whether systemic treatment with TPN can attenuate nociceptive behaviors in the CFA-induced inflammatory pain model. We also explored the possible analgesic mechanisms of TPN by assaying the activation of AKT pathway and the production of inflammatory cytokines.

Materials and Methods

Animals

Adult ICR male mice (6 to 8 weeks) were obtained from the Animal Care and Use Committee of Nantong University. Animals were kept under a 12/12 h light/dark cycle at a temperature of $23 \pm 2^\circ\text{C}$, humidity (50–60%) with free-feeding. All behavioral experiments were performed between 9 am and 6 pm. All in vivo studies were performed in accordance with the UK Animals Scientific Procedures Act (1986) and were approved by the Animal Care and Use Committee of Nantong University.

CFA Pain Model and Drug Administration

Inflammatory pain was induced by intraplantar injection of 20 μL CFA (100%; Sigma-Aldrich) in the left hind paws




Figure 1 Chemical structures of triptolide (A) and triptonide (B).

under brief anesthesia with isoflurane (2%; RWD Life Science, Shenzhen, China). TPN was purchased from Chengdu Mansite Biotechnology (Sichuan, China). To evaluate if TPN has analgesic effects, We used four experimental treatments: control group (saline injection), light-dose TPN treatment group (0.1 mg/kg), medium-dose TPN treatment group (0.5 mg/kg), and high-dose TPN treatment group (2.0 mg/kg). Each treatment started 1 hour before CFA injection by intravenous injection of a single daily bolus (0.25 mL) for 5 consecutive days (Figure 2A). To analyze CFA's effect on AKT's phosphorylation and cytokines' expression in vivo, we had set a naive group that only received an intraplantar injection of saline (0.9% NaCl, 20 μ L/mouse) in the left hind paw. To investigate the involvement of AKT activation in CFA-induced inflammatory pain, intrathecal injection (i.t.) of the AKT inhibitors: AKT

inhibitor IV (Sigma-Aldrich, Saint Louis, MO) or MK-2206 (MCE, NJ, USA) was performed in mice.

Paw Edema Measurement

Paw volumes were measured by plethysmometry (Ugo Basile Plethysmometer, Comerio, Italy) before CFA injections and 1, 2, 3, 4, and 5 days after TPN administration. Edema was expressed by paw volume (mL).

Behavioral Analysis

For the von Frey test, the animals were put in boxes on an elevated metal mesh floor daily for at least 2 d before baseline testing and allowed 30 min for habituation before the examination. The plantar surface of the left hind paw was stimulated with a series of von Frey hairs with logarithmically incrementing stiffness (0.02–2.56 g, Stoelting).




Figure 2 TPN treatment attenuates CFA-induced paw swelling, mechanical allodynia, and thermal hyperalgesia. **(A)** Schematic diagram of the timeline for CFA injection, drug treatment, and behavioral testing. **(B)** Change in paw volume, **(C)** Mechanical allodynia and **(D)** Thermal hyperalgesia of CFA-injected mice treated with various doses of TPN. Two-way ANOVA followed by Bonferroni's test was applied for statistical comparison between groups, $n = 7-8$ mice/group, $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, CFA vs CFA + TPN (0.5mg/kg), $^{\#}P < 0.05$, $^{\#\#}P < 0.01$, $^{\#\#\#}P < 0.001$, CFA vs CFA + TPN (2 mg/kg).

The 50% paw withdrawal threshold was determined using Dixon's up-down method. For the Hargreaves test, the animals were put in a plastic box placed on a glass plate, and the plantar surface was exposed to a beam of radiant heat through a transparent glass surface (Life Science). The baseline latencies were adjusted to 10–14 s with a maximum of 20 s as a cutoff to prevent potential injury. All the behavioral experiments were done by individuals who were blinded to the treatment of the mice.

Cell Cultures

Mouse neuroblastoma/rat DRG neuron hybrid ND7/23 (RRID: CVCL_4259) cells were grown under standard cell culture conditions (5% CO₂, 37°C) in High Glucose 'Dulbecco's Modified Eagle Medium (DMEM, Gibco) supplemented with 10% fetal bovine serum and 100 U/mL Penicillin-Streptomycin. ND7/23 cells were incubated with 1 µg/mL Lipopolysaccharides (LPS, Sigma-Aldrich, St. Louis, MO, USA) for 6 h. The treatment of the 1 µg/mL TPN or the treatment of 1 µg/mL AKT inhibitor IV was started 1 h prior to LPS treatment.

Western Blotting

Western blotting analysis of pAKT levels were performed using samples collected from four experimental groups (Naive n = 3, CFA n = 3, CFA + TPN (0.5 mg/kg), n = 3, CFA + TPN (2 mg/kg), n = 3). The tissues or cells were homogenized in a lysis buffer containing protease and phosphatase inhibitors (Sigma-Aldrich) and 30 µg of proteins were loaded for each lane and separated on SDS-PAGE gel (10%). After the transfer, the blots were incubated overnight at 4°C with the pAKT primary antibody (RRID: AB_2716452; rabbit, 1:1000; Cell Signaling Technology, USA). For loading control, the blots were probed with AKT primary antibody (RRID: AB_2225340; rabbit, 1:1000; Cell Signaling Technology, USA). Then the membranes were incubated with secondary antibodies (Dylight 800, Donkey Anti-Rabbit IgG, 1:10,000, Millipore). Protein bands were detected using the Odyssey System (Li-COR, NE, USA), and the intensity of each band was quantified using Image J (NIH, MD, USA).

Immunohistochemistry

Animals were deeply anesthetized with isoflurane and perfused through the ascending aorta with saline followed by 4% paraformaldehyde with 0.1 M phosphate buffer. After the perfusion, the L4-L5 DRG were removed and postfixed in the same fixative overnight. DRG (8–15 µm) were cut in a cryostat and processed for immunofluorescence. DRG was

first blocked with 1% bovine serum albumin for 2 h at room temperature, then incubated overnight at 4°C with the pAKT primary antibody (RRID: AB_2716452; rabbit, 1:200; Cell Signaling Technology, USA). The sections were then incubated for 2 h at room temperature with Cy3-conjugated secondary antibodies (1:1000; Jackson ImmunoResearch). The stained sections were examined with Leica SP8 Gated STED confocal microscope (Leica Microsystems, Wetzlar, Germany). The pAKT fluorescence intensity were measured by ImageJ.

Real-Time qPCR

Total RNA was extracted from L4-6 DRG with the Trizol reagent (Invitrogen, Carlsbad, CA, USA). One microgram of total RNA was converted into cDNA using PrimeScript RT reagent kit (Takara, Shiga, Japan). qPCR analysis was performed with the Real-Time Detection System by SYBR green I dye detection (Takara). The cDNA was amplified using the following primers: GAPDH forward, 5'-AAA TGG TGA AGG TCG GTG TGA AC-3'; GAPDH reverse, 5'-CAA CAA TCT CCA CTT TGC CAC TG-3'; TNF-α forward, 5'-GTT CTA TGG CCC AGA CCC TCA C-3'; reverse, 5'-GGC ACC ACT AGT TGG TTG TCT TTG-3'; IL-1β forward, 5'-TCC AGG ATG AGG ACA TGA GCA C-3'; reverse, 5'-GAA CGT CAC ACA CCA GCA GGT TA-3'; IL-6 forward, 5'-TAG TGG ATG CTA CCA AAC TGG A-3'; reverse, 5'-TGT GAC TCC AGC TTA TCT CTT G G-3'.

Quantification and Statistics

All data were analyzed by researchers blinded to the reagents used. All data were analyzed by GraphPad Prism (version 5.01) and presented as mean ± SEM. *P* < 0.05 was considered statistically significant. Behavioral data were analyzed using two-way ANOVA. Western blotting, immunofluorescence density and qPCR data were compared using one-way ANOVA. Student's *t*-test (2-tailed) was used to analyze qPCR data if only 2 groups were applied.

Results

Triptonide Attenuates CFA-Induced Inflammatory Pain

To test the anti-nociceptive effect of TPN in CFA-induced inflammatory pain, different doses of TPN or vehicle were intravenously injected daily for 5 consecutive days, with the first injection of TPN given at 1 h before CFA (Figure 2A). On Day 1, the left hind paw volume of the CFA group was

increased by approximately 2-fold compared to the baseline (Figure 2B). TPN at 0.5 mg/kg or 2.0 mg/kg caused a statistically significant decrease of paw swelling from day 1 to day 5 after CFA injection compared with untreated CFA mice [Figure 2B, (Treatment $F_{(3, 150)} = 30.08$, $P < 0.0001$; Time $F_{(5, 150)} = 61.09$, $P < 0.0001$; treatment * time interaction $F_{(15, 150)} = 1.201$, $P = 0.2770$; two-way ANOVA followed by Bonferroni's multiple comparisons test (compare treatment means), *** $P < 0.001$, CFA vs CFA+TPN (0.5mg/kg), #### $P < 0.001$, CFA vs CFA+TPN (2.0mg/kg)]. Bonferroni's multiple comparisons test (within each time point, compare treatments) was further used to assess statistical significance between controls and the treatments at the same time point. At almost each time point, both 0.5 mg/kg or 2.0 mg/kg TPN treatments can significantly attenuate CFA-induced footpad edema [Figure 2B, ** $P < 0.01$, *** $P < 0.001$, CFA vs CFA+TPN (0.5mg/kg), # $P < 0.05$, ## $P < 0.01$, ### $P < 0.001$, CFA vs CFA+TPN (2.0mg/kg)]. This revealed that TPN alleviated CFA-induced paw edema and had a strong anti-inflammatory effect.

In order to evaluate the effect of TPN on inflammatory pain, behavioral tests were performed in all groups of animals. CFA induced significant mechanical allodynia by decreasing the mechanical threshold induced by mechanical stimuli from day 1 to day 5 after CFA (Figure 2C). The intravenous administration of TPN significantly attenuated mechanical allodynia at 0.5 mg/kg or 2.0 mg/kg, but not at 0.1 mg/kg [(Treatment $F_{(3, 150)} = 7.196$, $P = 0.0002$; Time $F_{(5, 150)} = 72.60$, $P < 0.0001$; treatment * time interaction $F_{(15, 150)} = 1.334$, $P = 0.1885$; two-way ANOVA followed by Bonferroni's multiple comparisons test (compare treatment means), ** $P < 0.01$, CFA vs CFA+TPN (0.5mg/kg), ## $P < 0.01$, CFA vs CFA+TPN (2.0mg/kg)]. The Bonferroni's multiple comparisons test (within each time point, compare treatments) showed that TPN at 2 mg/kg or at 0.5 mg/kg had no effect on CFA-induced mechanical allodynia during the first 3 days following CFA injection, but started to show a reversal effect at 4 days and maintained till 5 days [Figure 2C, * $P < 0.05$, ** $P < 0.01$, CFA vs CFA+TPN (0.5mg/kg), # $P < 0.05$, ### $P < 0.001$, CFA vs CFA+TPN (2.0mg/kg)].

Meanwhile, TPN substantially attenuated CFA-induced thermal hyperalgesia at 2 mg/kg or at 0.5 mg/kg [Figure 2D, (Treatment $F_{(3, 150)} = 21.73$, $P < 0.0001$; Time $F_{(5, 150)} = 42.25$, $P < 0.0001$; treatment * time interaction $F_{(15, 150)} = 2.563$, $P = 0.002$; Two-way ANOVA followed by Bonferroni's multiple comparisons test (compare treatment means), ** $P < 0.01$, CFA vs CFA+TPN (0.5mg/kg), #### $P < 0.001$, CFA vs CFA+TPN (2.0mg/kg)]. TPN attenuated

thermal hyperalgesia at days 3, 4 and 5 after CFA injection [Figure 2D, Bonferroni's multiple comparisons test (within each time point, compare columns), * $P < 0.05$, CFA vs CFA+TPN (0.5mg/kg), #### $P < 0.001$, CFA vs CFA+TPN (2.0mg/kg)]. These data suggest that repeated TPN administration attenuates CFA-induced pain hypersensitivity, both mechanical allodynia and thermal hyperalgesia.

TPN Inhibits CFA-Induced AKT Activation in DRG

To determine whether the analgesic effects of TPN were associated with inhibition of the AKT signaling pathways, we evaluated the expression level of phosphorylation of AKT (pAKT) in the DRG. Western blotting showed that pAKT was significantly increased at 5 days after CFA injection. Treatment with TPN (2 mg/kg) significantly decreased CFA-induced pAKT upregulation (Figure 3A and B, $F_{(3, 8)} = 9.506$, $P = 0.0051$, one-way ANOVA followed by 'Bonferroni's multiple comparisons test, # $P < 0.05$, Naive vs CFA, ** $P < 0.01$, CFA vs CFA+TPN). Immunofluorescence analysis also showed upregulation of pAKT expression in DRG from animals injected with CFA. Treatment with TPN (2 mg/kg) significantly attenuated the CFA-induced upregulation of pAKT (Figure 3C and D, Area $F_{(2, 10)} = 56.24$, $P < 0.001$, Intensity $F_{(2, 10)} = 41.81$, $P < 0.001$, one-way ANOVA followed by 'Bonferroni's multiple comparisons test, ### $P < 0.001$, Naive vs CFA, *** $P < 0.001$, CFA vs CFA+TPN), which was consistent with the Western blotting results. Double immunostaining further showed that pAKT was expressed in the DRG neurons (Figure 3E).

TPN Inhibits CFA-Induced Increase of TNF- α , IL-1 β , and IL-6 in the DRG

Because TPN is involved in the regulation of *GFAP* and *IBA1* expression, which are markers for astrocytes and microglia, respectively, we examined their mRNA expression in the DRG after repeated TPN administration. CFA did not induce *GFAP* and *IBA1* expression, and TPN treatments also did not affect expression levels of them in DRG (Figure 4A and B, *GFAP*, $F_{(3, 20)} = 2.086$, $P = 0.1343$; *IBA1*, $F_{(3, 20)} = 0.5977$, $P = 0.6239$, one-way ANOVA). *TNF- α* , *IL-1 β* and *IL-6* are important pro-inflammatory cytokines in mediating peripheral sensitization and neuropathic pain.^{32–34} To check whether the anti-nociceptive effect of TPN is associated with the downregulation of pro-inflammatory cytokines, we checked *TNF- α* , *IL-1 β* , and *IL-6* expression. qPCR results showed that, compared with control animals, *TNF- α* , *IL-1 β* ,




Figure 3 TPN treatment decreases pAKT expression in the DRG neurons. **(A and B)** pAKT expression of CFA-injected mice treated or untreated with TPN. pAKT expression was increased in DRG at 5 days after CFA injection, and recovered in CFA + TPN (2 mg/kg) animals. $n = 3$, one-way ANOVA followed by Bonferroni's multiple comparisons test, $^{\#}P < 0.05$, Naive vs CFA, $^{**}P < 0.01$, CFA vs CFA+TPN; **(C)** Immunostaining of pAKT in the DRG; **(D)** Statistical data show the number of pAKT positive neurons and pAKT staining intensity in the DRG. $n = 5$ mice/group, one-way ANOVA followed by Bonferroni's multiple comparisons test, $^{###}P < 0.001$, Naive vs CFA, $^{***}P < 0.001$, CFA vs CFA+TPN. **(E)** Double staining of pAKT with a neuronal marker, β -tubulin (TUJ1) in the DRG at 5 days after CFA. Scale bars: 250 μ m.

and *IL-6* mRNAs were significantly increased in animals after 5 days of CFA injection. Both the dose of 0.5 mg/kg and 2.0 mg/kg TPN treatments significantly reduced the CFA-induced mRNA increases of *TNF- α* , *IL-1 β* , and *IL-6* (Figure 4C–E, *TNF- α* , $F_{(3, 15)} = 5.568$, $P = 0.009$; *IL-1 β* , $F_{(3, 18)} = 9.987$, $P = 0.0004$; *IL-6*, $F_{(3, 15)} = 5.944$, $P = 0.07$; one-way ANOVA followed by Bonferroni's multiple comparisons test, $^{\#}P < 0.05$, $^{###}P < 0.01$, Naive vs CFA, $^{*}P < 0.05$, $^{**}P < 0.01$, $^{***}P < 0.001$, CFA vs CFA+TPN). These data indicate TPN's inhibitory effect on CFA-induced *TNF- α* , *IL-1 β* , and *IL-6* expression in the DRG.

TPN Attenuates LPS-Induced pAKT Activation, and *TNF- α* , *IL-1 β* , and *IL-6* Expression in the ND7/23 Cells

To further confirm that TPN can attenuate pAKT activation in DRG neurons under inflammatory conditions, we used the

classical exogenous *toll-like receptor 4* ligand LPS to stimulate ND7/23 cells, a DRG neuron cell line, to mimic inflammatory conditions in vitro. We first checked pAKT expression after incubation with LPS (1 μ g/mL) for 6 hours. Western blotting showed that LPS induced a rapid and dramatic increase of pAKT expression. Pre-incubation with TPN, 1 hour before LPS application, markedly inhibited LPS-induced AKT phosphorylation by 43.2% (Figure 5A and B, $F_{(2, 6)} = 22.85$, $P = 0.0016$, one-way ANOVA followed by Bonferroni's multiple comparisons test, $^{###}P < 0.01$, Vehicle vs LPS, $^{**}P < 0.01$, LPS vs LPS+TPN), which was consistent with the in vivo results (Figure 3A and B).

To examine whether TPN inhibit *TNF- α* , *IL-1 β* , and *IL-6* expression via pAKT, we pretreated ND7/23 cells with AKT inhibitor, AKT inhibitor IV, 1 h before LPS application. As shown in Figure 5C–E, the qRT-PCR results showed that TPN (1 μ g/mL) treatment blocked CFA-induced *TNF- α* , *IL-1 β* , and *IL-6* increase (*TNF- α* , $F_{(2, 13)} = 20.70$, $P < 0.0001$; *IL-1 β* ,




Figure 4 TPN decreases CFA-induced upregulation of *TNF-α*, *IL-1β*, and *IL-6* increases. (**A** and **B**) Effect of TPN treatment on expression of *GFAP* (**A**) and *IBA1* (**B**) in DRG from CFA-treated mice. (**C–E**) Effects of TPN on the mRNA expression levels of *TNF-α* (**C**), *IL-1β* (**D**), and *IL-6* (**E**). TPN treatment with both medium dose (0.5 mg/kg) and high dose (2 mg/kg) decreased *TNF-α*, *IL-1β*, and *IL-6* upregulation after CFA injection. $n = 5$ mice/group, one-way ANOVA followed by Bonferroni's multiple comparisons test, # $P < 0.05$, ## $P < 0.01$, Naive vs CFA, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, CFA vs CFA+TPN.

$F_{(2, 11)} = 8.133$, $P = 0.0068$, *IL-6*, $F_{(2, 15)} = 61.09$, $P < 0.0001$; one-way ANOVA followed by Bonferroni's multiple comparisons test, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, LPS vs LPS+TPN). In consistent with TPN's effect, pretreatment with AKT inhibitor IV (1 $\mu\text{g/mL}$) also significantly decreased LPS-induced *TNF-α*, *IL-1β*, and *IL-6* upregulation (Figure 5C–E, ## $P < 0.01$ and ### $P < 0.001$, one-way ANOVA). These data suggest that TPN is involved in pAKT-mediated upregulation of *TNF-α*, *IL-1β*, and *IL-6* in activated ND7/23 cells by LPS.

AKT Inhibitor Attenuates CFA-Induced Pain Hypersensitivity and Upregulation of *TNF-α*, *IL-1β*, and *IL-6* in DRG

To define whether AKT is associated with CFA-induced pain hypersensitivity and the upregulation of inflammatory cytokines, mice were injected intravenously with the AKT inhibitors AKT inhibitor IV (1 $\mu\text{g}/10 \mu\text{L}$) or MK-2206 (1 $\mu\text{g}/10 \mu\text{L}$) 3 days after CFA injection. The behavioral results showed that the administration of AKT inhibitor IV slightly reduced mechanical allodynia without significant difference (Figure 6A). But, it can attenuate thermal

hyperalgesia at 6 h (Figure 6B, Treatment, $F_{(1, 55)} = 7.773$ and $P = 0.0073$, Time, $F_{(4, 55)} = 8.480$ and $P < 0.0001$; treatment * time interaction, $F_{(4, 55)} = 4.208$ and $P = 0.0048$, two-way ANOVA followed by Bonferroni's multiple comparisons test, *** $P < 0.001$). Intravenous administration of MK-2206 attenuated both mechanical allodynia (Figure 6C, Treatment, $F_{(1, 60)} = 5.872$, $P = 0.0184$, Time, $F_{(4, 60)} = 202.1$, $P < 0.0001$; treatment * time interaction, $F_{(4, 60)} = 5.713$, $P = 0.0006$, Two-way ANOVA followed by Bonferroni's multiple comparisons test, *** $P < 0.001$) and thermal hyperalgesia (Figure 6D, Treatment, $F_{(1, 60)} = 26.76$, $P < 0.0001$, Time, $F_{(4, 60)} = 20.52$, $P < 0.0001$; treatment * time interaction, $F_{(4, 60)} = 4.355$, $P = 0.0037$, two-way ANOVA followed by Bonferroni's multiple comparisons test, ** $P < 0.01$, *** $P < 0.001$). These results revealed that inhibition of AKT could relieve CFA-induced pain behavior.

As shown in Figure 6E and F, AKT inhibitor IV or MK-2206 treatment also significantly decreased the mRNA expression of *TNF-α*, *IL-1β* and *IL-6* (* $P < 0.05$, ** $P < 0.01$, Student's *t*-test). Taken together, these data suggest that




Figure 5 TPN decreases LPS-induced pAKT expression and the production of inflammatory cytokines in ND7/23 cells. (A and B) pAKT expression of LPS-induced ND7/23 cells treated or untreated with TPN (1 μ g/mL). pAKT expression was increased after LPS application and recovered in the LPS + TPN group. $n = 3$, one-way ANOVA followed by Bonferroni's multiple comparisons test, $^{###}P < 0.01$, Vehicle vs LPS, $^{**}P < 0.01$, LPS vs LPS+TPN. (C–E) Effects of TPN, and AKT inhibitor IV on the mRNA expression levels of *TNF- α* (C), *IL-1 β* (D), and *IL-6* (E). Both TPN treatment (1 μ g/mL) and AKT inhibitor IV treatment (1 μ g/mL) decreased *TNF- α* , *IL-1 β* , and *IL-6* upregulation after LPS incubation. $n = 5$, one-way ANOVA followed by Bonferroni's multiple comparisons test, $^{*}P < 0.05$, $^{**}P < 0.01$, $^{***}P < 0.001$, LPS vs LPS+TPN, $^{#}P < 0.05$ and $^{####}P < 0.001$, LPS vs LPS+ AKT inhibitor IV.

the AKT signaling pathway is involved in the development of CFA-induced inflammatory pain, and inhibition of AKT phosphorylation reduces the upregulation of inflammatory cytokines in the DRG.

Discussion

In this study, we investigated the anti-nociceptive effect of systemic injection of TPN on CFA-induced pain hypersensitivity and explored possible mechanisms. Our results demonstrated that TPN administration attenuated CFA-induced edema and pain hypersensitivity. Meanwhile, TPN treatment inhibited the AKT signaling pathway and the expression of *TNF- α* , *IL-1 β* , and *IL-6* in the DRG after CFA injection. AKT pathway acts as an upstream signaling pathway in CFA-induced upregulation of *TNF- α* , *IL-1 β* , and *IL-6*. TPN may exert an analgesic effect via inhibition of the CFA-induced DRG AKT-*TNF- α* /*IL-1 β* /*IL-6* signaling pathway (Figure 7).

Previous studies showed that TPL could prevent and attenuate neuropathic pain, cancer pain, and inflammatory

pain in mice or rats via inhibiting central immune response,³⁵ inhibiting the upregulation of the expression of histone deacetylases in spinal glial cells,⁸ or inhibiting spinal glial activation.⁹ However, the role of TPN in chronic pain has not yet been studied. We first demonstrated that repeated intravenous injection of TPN before CFA injection persistently attenuated CFA-induced thermal hyperalgesia and mechanical allodynia. TPN dose-dependently inhibited CFA-induced thermal hyperalgesia. However, there is no significant difference in mechanical allodynia between the 0.5 mg/kg and 2.0 mg/kg TPN groups. This may result from the target molecules or signaling pathways of TPN play more important roles in thermal pain sensitivity processing. Besides, the injection of TPN could also effectively reduce paw edema caused by CFA. Overall, these findings indicate that TPN has analgesic properties.

Although the pAKT signaling pathway typically regulates cell growth and survival, increasing evidence indicates the involvement of this pathway in the development and maintenance of chronic pain.^{36,37} Our data further




Figure 6 AKT inhibitor attenuates CFA-induced pain hypersensitivity and production of *TNF-α*, *IL-1β*, and *IL-6* in the DRG. (A–D) Effects of AKT inhibitors administration, AKT inhibitor IV or MK-2206, on mechanical allodynia and thermal hyperalgesia in CFA-induced inflammatory pain mice. Two-way ANOVA followed by Bonferroni's multiple comparisons test, ** $p < 0.01$, *** $p < 0.001$. (E and F) Intrathecal injection of AKT inhibitor IV or MK-2206 decreased expression of *TNF-α*, *IL-1β*, and *IL-6* in the DRG. $n = 6$. * $p < 0.05$, ** $p < 0.01$, AKT inhibitor IV vs CFA group, or MK-2206 vs CFA, Student's *t*-test.




Figure 7 Schematic representation of the mechanism of action of TPN in analgesia. Black arrows depict promotion. Red T-bar represents inhibition. Intraplantar injection of CFA can activate the AKT signaling pathway in DRG neurons, promoting the expression of pro-inflammatory genes, including *TNF-α*, *IL-1β*, and *IL-6*. Intravenous TPN could inhibit Akt phosphorylation and downstream gene expression, thereby relieving pain. The illustration was created with BioRender (BioRender.com).

showed that pAKT was predominantly expressed in DRG neurons after CFA injection. In agreement with our results, pAKT expression in neurons was found in the DRG following paclitaxel treatment.³⁸ Here, intraplantar injection of CFA induced pain hypersensitivity and increased pAKT expression in the DRG, which was inhibited by TPN administration, suggesting that the anti-nociceptive effect of TPN may be mediated by pAKT-mediated peripheral sensitization in the DRG. In accordance with our results, emerging evidence indicates that TPN suppressed the expression of pAKT on in vitro or in vivo models of different kinds of cancers.^{1,3,31} One report has indicated that TPN inhibits pAKT expression by suppressing a proto-oncogene *Lyn* in lymphoma.¹ Specifically, how the TPN inhibited pAKT signaling pathways in DRG neurons requires further exploration.

TNF- α , *IL-1 β* and *IL-6* are well-known pro-inflammatory cytokines that have been implicated in inflammatory pain.³⁹ Our data showed that the expression of *TNF- α* , *IL-1 β* , and *IL-6* was increased in DRG neurons after CFA injection and was inhibited by TPN treatment. *TNF- α* was expressed in the majority of the voltage-gated sodium channel (Nav) 1.3-positive or Nav1.8-positive neurons and up-regulated the expression of Nav1.3 and Nav1.8 in DRG neurons following peripheral nerve injury.³² Cleaved *IL-1 β* expression was significantly increased in small-sized DRG neurons after CFA injection into the hind paw.⁴⁰ *IL-6* was up-regulated in the ipsilateral L4 and L5 DRG neurons and in the bilateral lumbar spinal cord following L5-ventral root transection and contributed to the development of neuropathic pain.³⁴ Therefore, TPN may attenuate pain hypersensitivity via the inhibition of *TNF- α* , *IL-1 β* , and *IL-6*-mediated neuroinflammation.

Pharmacological inhibition of pAKT inhibited *TNF- α* , *IL-1 β* , and *IL-6* production in vitro. Previous studies have demonstrated that inhibition of pAKT pathway prevented the LPS-induced expression of *TNF- α* in human bronchial epithelial cells.⁴¹ Diesel exhaust particles exposure can activate the AKT signaling pathway and further up-regulate *IL-1 β* protein expression in primary human bronchial epithelial cells.⁴¹ Studies from Caco-2 cells and fibroblast-like synoviocytes showed that *IL-17*-mediated induction of *IL-6* was transduced via activation of AKT and NF- κ B,⁴² while mitogen-activated protein kinase (MAPK) are not likely to participate in the process.³⁸ Therefore, it is highly likely that the p-AKT pathway acts as an upstream signaling pathway, up-regulated the expression of cytokines *TNF- α* , *IL-1 β* , and *IL-6* in the

different cells. Both AKT Inhibitor IV and MK-2206 can relieve CFA-induced chronic inflammatory pain and the mRNA expression of *TNF- α* , *IL-1 β* , and *IL-6* in vivo. However, AKT Inhibitor IV showed a slightly weaker inhibition effect. Possible reasons are as follows; 1. The two drugs may have different absorption rates and degradation rates in DRG; 2. AKT inhibitor IV targets the ATP-binding site of a kinase upstream of AKT downstream of PI3K. In contrast, MK-2206 inhibits the autophosphorylation of AKT at threonine 308 and serine 473 directly; 3. The different action mechanisms of them may lead to a different degree of compensatory effects. Collectively, these findings suggest that the TPN may exert its analgesic effect through the inhibition of DRG AKT- *TNF- α* /*IL-1 β* /*IL-6* signaling.

Conclusion

Our present study demonstrated that intravenous treatment with TPN significantly attenuated CFA-induced pain hypersensitivity, associated with decreased *TNF- α* , *IL-1 β* , and *IL-6* expression, and decreased pAKT activation in DRG neurons. These data suggest that TPN could attenuate inflammatory pain via the inhibition of pAKT/*TNF- α* -*IL-6*-*IL-1 β* signaling pathway axis in DRG neurons. Our work provides valuable information for the preclinical and clinical study of TPN to treat chronic inflammatory pain.

Abbreviations

TPN, triptonide; PI3K, phosphoinositide 3-kinase; AKT, protein kinase B; mTOR, the mammalian target of rapamycin; CFA, complete Freund's adjuvant; DRG, dorsal root ganglion; *TNF- α* , tumor necrosis factor alpha; *IL-1 β* , interleukin 1 beta; *IL-6*, interleukin 6; TPL, triptolide; ERK, extracellular signal-regulated kinase; LPS, lipopolysaccharides; *GAPDH*, glyceraldehyde 3-phosphate dehydrogenase; Nav, voltage-gated sodium channel.

Data Sharing Statement

The datasets used and/or analyzed during the current study available from the corresponding author on reasonable request.

Ethics Approval

All in vivo studies were performed following the UK Animals Scientific Procedures Act (1986) and were approved by the Animal Care and Use Committee of Nantong University.

Funding

This work was funded by grants from the National Natural Science Foundation of China (NSFC 81771197, 81971054, 81300954 and 81400915), the Natural Science Foundation of Jiangsu Province (BK20171255), the Jiangsu University Qing Lan Project, and the Six talent peaks project in Jiangsu Province (SWYY-070).

Disclosure

The authors report no conflicts of interest for this work.

References

- Yang P, Dong FL, Zhou QS. Triptonide acts as a novel potent anti-lymphoma agent with low toxicity mainly through inhibition of proto-oncogene Lyn transcription and suppression of Lyn signal pathway. *Toxicol Lett.* 2017;278:9–17. doi:10.1016/j.toxlet.2017.06.010
- Hu X, Dong Y, Jin X, et al. The novel and potent anti-depressive action of triptolide and its influences on hippocampal neuroinflammation in a rat model of depression comorbidity of chronic pain. *Brain Behav Immun.* 2017;64:180–194. doi:10.1016/j.bbi.2017.03.005
- Dong F, Yang P, Wang R, et al. Triptonide acts as a novel antiprostata cancer agent mainly through inhibition of mTOR signaling pathway. *Prostate.* 2019;79(11):1284–1293. doi:10.1002/pros.23834
- Kupchan SM, Court WA, Dailey RG, Gilmore CJ, Bryan RF. Triptolide and triptidiolide, novel antileukemic diterpenoid triepoxides from *Tripterygium wilfordii*. *J Am Chem Soc.* 1972;94(20):7194–7195. doi:10.1021/ja00775a078
- Tang J, Li Z-H, Ge S-N, et al. The inhibition of spinal astrocytic JAK2-STAT3 pathway activation correlates with the analgesic effects of triptolide in the rat neuropathic pain model. *Evid Based Complement Alternat Med.* 2012;2012:1–13. doi:10.1155/2012/185167
- Huang Y, Zhu N, Chen T, et al. Triptolide Suppressed the Microglia Activation to Improve Spinal Cord Injury Through miR-96/IKK beta/NF-kappa B Pathway. *Spine.* 2019;44(12):E707–E714. doi:10.1097/BRS.0000000000002989
- Hang LH, Li SN, Shao DH, Chen Z, Chen YF, Shu WW. Evidence for involvement of spinal RANTES in the anti-nociceptive effects of triptolide, a diterpene triepoxide, in a rat model of bone cancer pain. *Basic Clin Pharmacol Toxicol.* 2014;115(6):477–480. doi:10.1111/bcpt.12265
- Hu X-F, He X-T, Zhou K-X, et al. The analgesic effects of triptolide in the bone cancer pain rats via inhibiting the upregulation of HDACs in spinal glial cells. *J Neuroinflamm.* 2017;14(1):213. doi:10.1186/s12974-017-0988-1
- Xu F, Li Y, Li S, et al. Complete Freund's adjuvant-induced acute inflammatory pain could be attenuated by triptolide via inhibiting spinal glia activation in rats. *J Surg Res.* 2014;188(1):174–182. doi:10.1016/j.jss.2013.11.1087
- Mei Z, Wu Q, Hu S, Lib X, Yang X. Triptolide loaded solid lipid nanoparticle hydrogel for topical application. *Drug Dev Ind Pharm.* 2005;31(2):161–168. doi:10.1081/DDC-200047791
- Tang X, Wang C, Hsieh Y, et al. Triptolide induces toxicity in inner ear stem cells via promoting DNA damage. *Toxicol in Vitro.* 2019;61:104597. doi:10.1016/j.tiv.2019.104597
- Xi C, Peng S, Wu Z, Zhou Q, Zhou J. Toxicity of triptolide and the molecular mechanisms involved. *Biomed Pharmacother.* 2017;90:531–541. doi:10.1016/j.biopha.2017.04.003
- Li J, Shen F, Guan C, et al. Activation of Nrf2 protects against triptolide-induced hepatotoxicity. *PLoS One.* 2014;9(7):e100685. doi:10.1371/journal.pone.0100685
- Vliegthart ADB, Wei C, Buckley C, et al. Characterization of triptolide-induced hepatotoxicity by imaging and transcriptomics in a novel zebrafish model. *Toxicol Sci.* 2017;159(2):380–391. doi:10.1093/toxsci/kfx144
- Hu DD, Chen XL, Xiao XR, et al. Comparative metabolism of triptolide and triptonide using metabolomics. *Food Chem Toxicol.* 2018;115:98–108. doi:10.1016/j.fct.2018.03.009
- Xiang S, Zhao Z, Zhang T, et al. Triptonide effectively suppresses gastric tumor growth and metastasis through inhibition of the oncogenic Notch1 and NF-kappaB signaling pathways. *Toxicol Appl Pharmacol.* 2019;114870.
- Wang SS, Lv Y, Xu XC, et al. Triptonide inhibits human nasopharyngeal carcinoma cell growth via disrupting Lnc-RNA THOR-IGF2BP1 signaling. *Cancer Lett.* 2019;443:13–24. doi:10.1016/j.canlet.2018.11.028
- Zhang B, Meng M, Xiang S, et al. Selective activation of tumor-suppressive MAPK signaling pathway by triptonide effectively inhibits pancreatic cancer cell tumorigenicity and tumor growth. *Biochem Pharmacol.* 2019;166:70–81. doi:10.1016/j.bcp.2019.05.010
- Zhang M, Tan S, Yu D, et al. Triptonide inhibits lung cancer cell tumorigenicity by selectively attenuating the Shh-Gli1 signaling pathway. *Toxicol Appl Pharmacol.* 2019;365:1–8. doi:10.1016/j.taap.2019.01.002
- Zhao Q, Zhang J-L, Li F. Application of metabolomics in the study of natural products. *Nat Prod Bioprospect.* 2018;8(4):321–334. doi:10.1007/s13659-018-0175-9
- Jiang BC, Liu T, Gao YJ. Chemokines in chronic pain: cellular and molecular mechanisms and therapeutic potential. *Pharmacol Ther.* 2020;212:107581. doi:10.1016/j.pharmthera.2020.107581
- Ji RR, Chamesian A, Zhang YQ. Pain regulation by non-neuronal cells and inflammation. *Science (New York, N Y).* 2016;354(6312):572–577. doi:10.1126/science.aaf8924
- Gao Y, Bai L, Zhou W, et al. PARP-1-regulated TNF- α expression in the dorsal root ganglia and spinal dorsal horn contributes to the pathogenesis of neuropathic pain in rats. *Brain Behav Immun.* 2020;88:482–496. doi:10.1016/j.bbi.2020.04.019
- Nadeau S, Filali M, Zhang J, et al. Functional recovery after peripheral nerve injury is dependent on the pro-inflammatory cytokines IL-1 β and TNF: implications for neuropathic pain. *J Neurosci.* 2011;31(35):12533–12542. doi:10.1523/JNEUROSCI.2840-11.2011
- Fang D, Kong LY, Cai J, et al. Interleukin-6-mediated functional upregulation of TRPV1 receptors in dorsal root ganglion neurons through the activation of JAK/PI3K signaling pathway: roles in the development of bone cancer pain in a rat model. *Pain.* 2015;156(6):1124–1144.
- Liu Y, Tong C, Tang Y, et al. Tanshinone IIA alleviates blast-induced inflammation, oxidative stress and apoptosis in mice partly by inhibiting the PI3K/Akt/FoxO1 signaling pathway. *Free Radic Biol Med.* 2020;152:52–60. doi:10.1016/j.freeradbiomed.2020.02.032
- Han X, Li B, Ye X, et al. Dopamine D2 receptor signalling controls inflammation in acute pancreatitis via a PP2A-dependent Akt/NF-kappaB signalling pathway. *Br J Pharmacol.* 2017;174(24):4751–4770. doi:10.1111/bph.14057
- Jiang S-P, Zhang Z-D, Kang L-M, Wang Q-H, Zhang L, Chen H-P. Celecoxib reverses oxaliplatin-induced neuropathic pain through inhibiting PI3K/Akt2 pathway in the mouse dorsal root ganglion. *Exp Neurol.* 2016;275:11–16. doi:10.1016/j.expneurol.2015.11.001
- Guo J-R, Wang H, Jin X-J, Jia D-L, Zhou X, Tao Q. Effect and mechanism of inhibition of PI3K/Akt/mTOR signal pathway on chronic neuropathic pain and spinal microglia in a rat model of chronic constriction injury. *Oncotarget.* 2017;8(32):52923. doi:10.18632/oncotarget.17629

30. Xu JT, Zhao JY, Zhao XL, et al. Opioid receptor-triggered spinal mTORC1 activation contributes to morphine tolerance and hyperalgesia. *J Clin Invest.* **2014**;124(2):592–603. doi:10.1172/JCI70236
31. Kim MJ, Lee TH, Kim SH, Choi YJ, Heo J, Kim YH. Triptolide inactivates Akt and induces caspase-dependent death in cervical cancer cells via the mitochondrial pathway. *Int J Oncol.* **2010**;37(5):1177–1185. doi:10.3892/ijo.00000769
32. He X-H, Zang Y, Chen X, et al. TNF- α contributes to up-regulation of Nav1.3 and Nav1.8 in DRG neurons following motor fiber injury. *PAIN®.* **2010**;151(2):266–279. doi:10.1016/j.pain.2010.06.005
33. Copray J, Mantingh I, Brouwer N, et al. Expression of interleukin-1 beta in rat dorsal root ganglia. *J Neuroimmunol.* **2001**;118(2):203–211. doi:10.1016/S0165-5728(01)00324-1
34. Wei X-H, Na X-D, Liao G-J, et al. The up-regulation of IL-6 in DRG and spinal dorsal horn contributes to neuropathic pain following L5 ventral root transection. *Exp Neurol.* **2013**;241:159–168. doi:10.1016/j.expneurol.2012.12.007
35. Wang W, Mei XP, Chen L, et al. Triptolide prevents and attenuates neuropathic pain via inhibiting central immune response. *Pain Physician.* **2012**;15(6):E995–1006.
36. Zhuang Z-Y, Xu H, Clapham DE, Ji -R-R. Phosphatidylinositol 3-kinase activates ERK in primary sensory neurons and mediates inflammatory heat hyperalgesia through TRPV1 sensitization. *J Neurosci.* **2004**;24(38):8300–8309. doi:10.1523/JNEUROSCI.2893-04.2004
37. Chen S-P, Zhou Y-Q, Liu D-Q, et al. PI3K/Akt pathway: a potential therapeutic target for chronic pain. *Curr Pharm Des.* **2017**;23(12):1860–1868. doi:10.2174/1381612823666170210150147
38. Li D, Chen H, Luo X-H, Sun Y, Xia W, Xiong Y-C. CX3CR1-mediated Akt1 activation contributes to the paclitaxel-induced painful peripheral neuropathy in rats. *Neurochem Res.* **2016**;41(6):1305–1314. doi:10.1007/s11064-016-1827-y
39. Cook AD, Christensen AD, Tewari D, McMahon SB, Hamilton JA. Immune cytokines and their receptors in inflammatory pain. *Trends Immunol.* **2018**;39(3):240–255. doi:10.1016/j.it.2017.12.003
40. Matsuoka Y, Yamashita A, Matsuda M, Kawai K, Sawa T, Amaya F. The NLRP2 inflammasome in dorsal root ganglion as a novel molecular platform that produces inflammatory pain hypersensitivity. *Pain.* **2019**;160(9):2149–2160. doi:10.1097/j.pain.0000000000001611
41. Wu W, Peden DB, McConnell R, Fruin S, Diaz-Sanchez D. Glutathione-S-transferase M1 regulation of diesel exhaust particle-induced pro-inflammatory mediator expression in normal human bronchial epithelial cells. *Part Fibre Toxicol.* **2012**;9(1):31. doi:10.1186/1743-8977-9-31
42. Hwang S-Y, Kim J-Y, Kim K-W, et al. IL-17 induces production of IL-6 and IL-8 in rheumatoid arthritis synovial fibroblasts via NF- κ B-and PI3-kinase/Akt-dependent pathways. *Arthritis Res Ther.* **2004**;6(2):R120. doi:10.1186/ar1038

Journal of Pain Research

Dovepress

Publish your work in this journal

The Journal of Pain Research is an international, peer reviewed, open access, online journal that welcomes laboratory and clinical findings in the fields of pain research and the prevention and management of pain. Original research, reviews, symposium reports, hypothesis formation and commentaries are all considered for publication. The manuscript

management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-pain-research-journal>