Utilizing a Minimum of Four Antenatal Care Visits and Associated Factors in Debre Berhan Town, North Shewa, Amhara, Ethiopia, 2020

This article was published in the following Dove Press journal:
Risk Management and Healthcare Policy

Michael Amara Tizazu
Eyosiyas Yeshialem Asefa
Mikyas Arega Muluneh
Asalif Beyene Haile

Department of Midwifery, Health Science
College, Debre Berhan University, Debre
Berhan, Ethiopia

Background: Receiving a minimum of four antenatal care, as per the World Health Organization recommendation, enhances the possibility of utilizing effective maternal health care during antenatal follow-up. Hence, this study aimed to identify the level of a minimum of 4 ANC usage and factors associated with Debre Berhan town.

Methods: A community-based cross-sectional study was employed from February 20 to March 10/2020 among women who gave birth in the last 6 months before the data collection period in Debre Berhan town. Data were collected using face-to-face interviews with the pre-tested tool by trained data collectors from randomly selected participants. After entering the data into Epi-data software version 3.1, it was further processed using SPSS version 23 statistical package. Both bivariate and multivariable Logistic regressions were fitted to identify the determinants for the utilization of a minimum of 4 antenatal care. A significance level of 0.05 was used to decide the significance of statistical tests.

Results: A total of 390 women were interviewed giving a response rate of 99.2%. In this study, 78.5% (95% CI: 74.4–82.6) of women received a minimum of four antenatal care visits. Maternal educational status, initiation of antenatal visit at ≤ 16 weeks of gestation, and husband support/partner involvement were identified predictors of utilization of minimum 4 antenatal visits.

Conclusion and Recommendations: Use of at least four antenatal visits in Debre Berhan town was found to be good. Health promotion programs targeting mothers with no education and lower educational level are important to increase their awareness about the importance of antenatal services. Male partner involvement during care related to pregnancy and promoting early initiation of antenatal care is important to overcome challenges towards a minimum of 4 antenatal visits in Debre Berhan town.

Keywords: minimum four antenatal, utilization, Ethiopia

Background

Antenatal care (ANC) is routine care given by a trained health care provider for a woman during pregnancy starting from the time of conception up to the delivery of the baby. ANC from trained health care providers is important to monitor pregnancy and to reduce complications and risks for the mother and child during pregnancy, delivery, and after giving birth.^{1,2}

Although significant progress has been made in the last two decades, about 295 000 women died during and following pregnancy and childbirth in 2017. About 810 Women approximately die every day from preventable causes related to pregnancy

Correspondence: Eyosiyas Yeshialem
Asefa

Department of Midwifery, Health Science
College, Debre Berhan University, Debre
Berhan, Ethiopia
Tel +251 913836869
Email eyosi143@gmail.com

and childbirth and 94% of all maternal deaths occur in low and lower-middle-income countries.³

Pregnancy-related preventable morbidity and mortality remain one of the global most critical challenges irrespective of significant progress and ambitions to end by 2030. The world will face difficulties to achieve this target by more than one million lives with the current pace of progress.^{1,4,5}

The world health organization (WHO) visions a world where each pregnant woman and newborn accessed quality care throughout the whole stage of pregnancy. ANC gives a platform for significant health care functions including, screening and diagnosis, health promotion, and disease prevention. Implementing important evidence-based practices on time, ANC can improve lives and provides the chance to communicate with and support pregnant women, families, and societies at an important time in the course of a woman's life.^{4,6}

Antenatal care can support women making ready for delivery and understand danger signs during pregnancy and childbirth. It can be a source of micronutrient provision, care of hypertension to prevent eclampsia, tetanus immunization, and prevention of mother-to-child transmission of HIV/AIDS. In the malaria-endemic area, the health care provider can also give medication insecticide-treated mosquito nets to a pregnant woman.^{1,6,7}

Globally, about 86% of pregnant women access antenatal care from trained health personnel at least once, but only two in three (65%) receive at least four antenatal visits. In Sub-Saharan Africa and South Asia, about half of them received at least four antenatal visits (52% and 48%) respectively.⁶ The Ethiopian Min Demographic and Health Survey (EMDHS) showed that four in 10 women (43%) had four or more ANC visits for their recent live births. Antenatal care coverage has increased in Ethiopia, from 28% in 2005 to 74% during 2019.^{2,8–10}

Place of residence, region, mothers' education level, household wealth index, desire for pregnancy, frequency of reading newspaper, frequency of listening to the radio, and frequency of watching TV was associated with the utilization of a minimum of four ANC services.^{11,12}

Unlike other parts of the world, there is no available data about the level of at least 4 ANC utilization and related factors in our study area. Therefore; this study was aimed to determine the level and factors associated with the utilization of a minimum of four antenatal care services in Debre Berhan town in 2020.

Methods

Study Area

The study was conducted in Debre Birhan town, located in the North Shewa Zone of Amhara region, 130 kilometers far from the capital city, Addis Ababa. According to the 2019 report of Debre Berhan Town administration's mayor office, the Town consists of nine kebeles (the smallest administrative unit) with an estimated total population of 114,652 (Male=51,843 and Female=62,809). Of the 62,809 female population, 39,066 were within the age group of 15–49 years. The town has four public (one referral hospital and three health centers) and three private health institutions. All the above health institutions had maternal health services. Health extension workers are community health workers responsible for identifying and registering all pregnant women at a time in their respective kebeles additional to the provision of tasks related to maternal health like provision of antenatal care, clean and safe deliveries, postnatal care, family planning, immunization and nutritional advice.¹³ According to the health extension worker's report; a total of 771 women gave birth in the last 6 months in Debre Berhan town.

Study Design and Period

A cross-sectional study at the community level study was conducted from February 20 to March 10/2020 among individuals randomly selected from the registration list of health extension workers.

Source Population

The source population for this study was all women who gave birth in the last 6 months before the data collection period in Debre Berhan town.

Study Population

This study was done on randomly selected women who gave birth in the last 6 months in Debre Berhan town. Women who reside less than six months at the study area during the time of data collection and who were unable to respond during data collection were excluded from the study.

Sample Size Determination

The sample size for this study was calculated using the assumptions for single population proportion with estimated proportion of completion of at least four antenatal care 36.6% from a cross-sectional study which was

analyzed from the 2016 Ethiopian Demographic and Health Survey (EDHS) data,¹¹ confidence level of 95%, 5% degree of precision, and 10% non-response rate, using the following formula.

$$n = \frac{Z_{a/2}^2 p(1-p)}{d^2}$$

Where,

n= sample size,

$Z_{a/2}$ = the value under the normal standard table for the given confidence level,

p=estimated of the population proportion,

d= margin of error, by using the above equation the sample size was calculated as follows:

$$n = \frac{(1.96)^2 * 0.366(1 - 0.366)}{(0.05)^2},$$

n = 356.568~357 by considering the non-response rate the last sample included became 393.

Sampling Technique

We found a total of 771 women who gave birth in the last 6 months from the registry book. Then after excluding those using the exclusion criteria we reached 769 women as a frame. Finally, we selected a total of 393 samples to be included in this study using the lottery method.

Data Collection methods and Procedures

Data was collected using the interviewer guide pre-tested tool using trained 2 BSc midwives data collectors and one MSc midwife as a supervisor. The data collection questionnaire was adapted from the Ethiopian Demographic and Health Survey (EDHS) and other literature. The tool incorporates socio-demographic variables, obstetrics related variables, and maternal health service-related variables. The data collectors and supervisors accessed the house of study subjects with the help of health development army (HDA) leaders.

Variables

Dependent Variable

“Utilization of a minimum of 4 ANC services”

Explanatory Variables

Socio-demographic (age, marital status, educational status, occupation, monthly income), reproductive and obstetrics related (number of pregnancy, planned/unplanned pregnancy, number of children, number of ANC visit, months

of first ANC visit) and maternal health care service-related (counseling on pregnancy complication, birth preparedness, performing urinalysis and blood checkups during ANC, TT vaccination).

Operational Definitions

Antenatal care utilization (at least four visits): utilization of at least 4 antenatal care visits of a woman of reproductive age group with a live birth during her pregnancy time.^{1,6}

Birth preparedness: Well prepared for birth and its complications were considered when women reported that they have implemented five or more components of birth preparedness and complication readiness (BPCR) otherwise considered as “not well prepared”. The components of BPCR considered in this study were identified birthplace, identified sign of labor, identify supplies needed during labor/delivery, saving money for an emergency, identified emergency transportation, people to support during/after birth, and identified potential blood donors as needed.¹⁴

Planned pregnancy: is only if they had met four key criteria. Intending to become pregnant, stopping contraception, partner agreement, and reaching the right time in terms of lifestyle/life stage were also necessary.¹⁵

Partner support: measured if one of the following three criteria were practiced. Reminded his partner's ANC follow-up schedule; covered medical/transport costs of partner's ANC follow-up accompanied partner to ANC clinic at least once.

Data Quality Assurance

A one-day training was given for data collectors and supervisors on the objective of the study, and the content of the tool. Back and forth questionnaire translation was done to check its consistency with the original meaning. Before commencing data collection, a pre-test was conducted among 10% of the study sample and, the required adjustments were made on unclear questions. Additionally, we did a reliability test and we found Cronbach's alpha (reliability coefficient) equals 0.78 showings acceptable level of consistencies. Daily meetings and supervision were performed throughout the data collection period.

Data Processing and Analysis

After Checking the completeness and consistency, the collected data were coded and entered into Epi-data software version 3.1 and then exported to SPSS version 23 statistical package for analysis. Descriptive statistics were done to

quantify the use of at least 4 ANC visits and other characteristics. Variables with a P value of less than 0.25 on bivariate logistic regression were selected as a candidate for multivariate logistic regression. Finally, multivariable Logistic regressions were used to assess the relative impact of explanatory variables on dependent variables and to select important predictors of at least 4 ANC visits. The fitness of the model was measured by Hosmer and Lemeshow goodness of test and multicollinearity between the independent variables was checked by the variance inflation factor ($VIF > 10$). Adjusted Odds Ratio (AOR) and their 95% Confidence Interval (CI) was used to determine the level of association. A significance level of 0.05 was used to declare the significance of statistical tests. Finally, the finding was presented using narration and tables.

Ethical Consideration

A letter of Ethical clearance was secured from the Institutional Review Board (IRB) of health Science College, Debre Berhan University. A formal letter of permission was obtained from the Debre Berhan town health office. Informed written consent was obtained from study subjects. Additionally, after doing the witness attests that the consent information was accurately explained for those who cannot read and write, that the subject apparently understood the information, and informed consent was given freely. Participants had the right to self-determination regarding participation in research, both initially and during the research. For confidentiality purposes, the names of the participant were not included in the questionnaire. The collected data were kept confidential and used only for the study. This study was conducted following the Declaration of Helsinki.

Results

Socio-Demographic Characteristics of the Study Participants

From a total of 393 participants, 390 were interviewed. The majority 269 (69%) of respondents were aged 25–34 years and the mean age was 28.9 ± 4.55 years. Most women 370 (94.9%) were married. While 37 (9.5%) had no education, 167 (42.8%) had college and above education. Almost 97% of study participants were Amhara by ethnicity. Nearly half (48.3%) were housewives. Most women 349 (89.5%) were ever heard about maternal health services. Over half 209 (53.6%) of participant's partners attained tertiary education (college and above). By occupation, 158 (40.5%) of their partner were government employee (Table 1).

Table 1 Socio-Demographic Characteristics of the Study Participants in Debre Berhan Town, Ethiopia 2020 (n=390)

Characteristics		Frequency	Percentage
Age of women in years	19–24	65	16.7
	25–34	269	69.0
	≥35	56	14.4
Religion	Orthodox	342	87.8
	Muslim	18	4.6
	Protestant	26	6.7
	Others	4	1.0
Ethnicity	Amhara	378	96.9
	Oromo	3	0.8
	Tigre	2	0.5
	Guragie	5	1.3
	Others	2	0.5
Marital status	Married	370	94.9
	Divorced/separated	16	4.1
	Widowed	1	0.3
	Single	3	0.8
Educational status	No education	37	9.5
	Primary education ^{1–8}	125	32.1
	Secondary education ^{9–12}	61	15.6
	College and above	167	42.8
Occupation	Farmer	4	1.0
	Housewives	9	2.3
	Private employed	188	48.2
	Gov't employed	46	11.8
	Merchant	107	27.4
	Others	29	7.4
		7	1.8
Family monthly income(ETB)	≤500	7	1.8
	501–1500	23	5.9
	1501–2500	49	12.6
	≥2501	311	79.7
Educational status of partner	No education	37	9.5
	Primary education ^{1–8}	46	11.8
	Secondary education ^{9–12}	98	25.1
	College and above	209	53.6
Occupation of partner	Farmer	20	5.1
	Private employed	100	25.6
	Gov't employed	158	40.5
	Merchant	70	17.9
	Daily laborer	27	6.9
	Others	15	3.8
Ever Heard maternal health services	Yes	349	89.5
	No	41	10.5
Source of maternal health information (n=349)	Mass-media(tv/radio)	122	35.0
	Health care provider	214	61.3
	Others	13	3.7

Reproductive and Maternal Health Services-Related Characteristics of Study Participants

Among study participants, 140 (35.9%) had their first pregnancy, whereas, 41 (10.5%) had four and more pregnancies. Almost 97% of the participant's recent pregnancy was planned. Three hundred thirty-seven (86.4%) of women had initiated their first antenatal visit at or less than 16 weeks of gestation and 306 (78.5%) had four or more antenatal visits throughout their pregnancy. Out of all respondents, 228 (58.5%) of women had received their antenatal care at public health centers. Nearly 90% had their husband's support during maternal health care visit. Regarding the services provided during the pregnancy period; 378 (96.9%) had blood pressure measurement, 370 (94.9%) received nutritional counseling, 260 (66.7%) had taken vaccination for tetanus, 359 (92.1%) informed about danger signs, 352 (90.3%) counseled about birth preparedness and complication readiness plan and 353 (90.5%) had received iron and folate (Table 2).

Factors Associated with the Utilizing a Minimum of 4 Antenatal Visits

This study identified some important factors that are associated with the utilization of a minimum of 4 ANC visits in Debre Berhan town. Maternal educational status, initiation of ANC visit ≤ 16 weeks of gestation, and husband support/partner involvement were statistically significant association with the outcome variable.

The odds of utilization of a minimum of 4 ANC visits were three times higher among women who had college and above education (AOR: 3.09, 95% CI: 1.16–8.27) than those who had no education. The odds of utilization of minimum 4 ANC visits was eight times higher for those who had first ANC visit at ≤ 16 weeks of gestation than their counterparts (AOR: 8.30, 95% CI: 4.01–17.18). Similarly, The odds of utilization of a minimum of 4 ANC visits were two times higher for those who had husband support than their counterparts (AOR: 2.28, 95% CI: 1.03–5.06) (Table 3).

Discussion

The result of the study revealed that the utilization of a minimum of four ANC visits was 78.5% (95% CI: 74.4–82.6) in Debre Berhan town. This value was greater than the finding of previous studies in Ethiopia 33%,¹⁶ 36.2%,¹¹ and 36.78%.¹⁷ This could be because; previous studies were conducted in both urban and rural settings,

Table 2 Reproductive and Maternal Health Services-Related Characteristics of the Study Participants in Debre Berhan Town Ethiopia 2020 (n=390)

Characteristics		Frequency	Percentage
Number of pregnancy	1	140	35.9
	2–3	209	53.6
	≥ 4	41	10.5
Pregnancy planned	Yes	378	96.9
	No	12	3.1
Number of children	1	142	36.4
	2–3	213	54.6
	≥ 4	35	9.0
Number of ANC visit for the last pregnancy	< 4	84	21.5
	≥ 4	306	78.5
Timing of first ANC visit for the last pregnancy	≤ 16 weeks	337	86.4
	> 16 weeks	53	13.6
Place of ANC visit	Gov't hospital	109	27.9
	Gov't health center	228	58.5
	Private hospital	30	7.7
	Private clinic	23	5.9
Partner support	Yes	345	88.5
	No	45	11.5
Informed key pregnancy danger signs	Yes	359	92.1
	No	31	7.9
Blood pressure measured at ANC	Yes	378	96.9
	No	12	3.1
Urine sample taken	Yes	368	94.4
	No	22	5.6
Blood sample taken	Yes	382	97.9
	No	8	2.1
Nutritional counseling	Yes	370	94.9
	No	20	5.1
TT vaccine at ANC	Yes	260	66.7
	No	130	33.3
Number of TT vaccine (n=260)	One	87	33.5
	Two or more	173	66.5
Iron tablet taken	Yes	353	90.5
	No	37	9.5
Deworming during ANC	Yes	140	90.5
	No	250	9.5
Informed about Birth preparedness plan	Yes	352	90.3
	No	38	9.7
Status of Birth preparedness	Well prepared	199	51.0
	Not well prepared	191	49.0

Table 3 Factors Associated with the Utilization of a Minimum 4 Antenatal Visits in Debre Berhan Town Ethiopia 2020 (n=390)

Variables	Utilization of Minimum 4 ANC		COR(95% CI)	AOR(95% CI)
	Yes	No		
Educational status of the mother				
No education	21	16	1	1
Primary education	92	33	2.124(0.991–4.553)	1.89(0.788–4.524)
Secondary education	45	16	2.143(0.902–5.090)	1.88(0.708–4.965)
College and above	148	19	5.935(2.648–13.300)	3.79(1.539–9.351)*
Number of pregnancy				
One	115	25	2.385(1.097–5.188)	1.61(0.567–4.562)
Two-three	164	45	1.890(0.915–3.901)	1.18(0.459–3.011)
Four or more	27	14	1	1
Was the pregnancy planned				
Yes	301	77	5.473(1.691–17.715)	2.08(0.440–9.851)
No	5	7	1	1
Timing of first ANC visit				
≤16 weeks	288	49	11.429(6.001–21.764)	8.87(4.478–17.558)*
>16 weeks	18	35	1	1
Husband support				
Yes	283	62	4.366(2.289–8.329)	2.29(1.076–4.877)*
No	23	22	1	1
Informed key pregnancy danger signs				
Yes	288	71	2.930(1.371–6.259)	1.10(0.388–3.146)
No	18	13	1	1
Nutritional counseling during pregnancy				
Yes	293	77	2.049(0.790–5.311)	1.17(0.318–4.318)
No	13	7	1	1
TT vaccination during pregnancy				
Yes	215	45	2.048(1.250–3.355)	1.73(0.979–3.057)
No	91	39	1	1
Iron and folate taken				
Yes	286	67	3.628(1.803–7.300)	1.18(0.450–3.096)
No	20	17	1	1
Informed about BPCR				
Yes	281	71	2.058(1.003–4.224)	1.29(0.519–3.224)
No	25	13	1	1
Status of BPCR				
Well-prepared	168	31	2.081(1.266–3.422)	1.45(0.840–2.676)
Not well-prepared	138	53	1	1

Note: *p-value < 0.05.

which includes women with lower access to antenatal care visits but the current study was conducted in the urban areas. This explanation is maintained by previous studies conducted in Ethiopia, where urban women have better access to antenatal care than those women living in rural areas.^{7,18} Similarly, utilization of at least 4 ANC visits in

our study area was much higher than a study in Northern Ethiopia,¹⁹ Vietnam,²⁰ Nigeria,²¹ and Timor-Leste.²² This discrepancy might be because of the difference in the study setting like time and study area (rural areas).

On the other hand, the finding was found to be lower than a study done in Ghana where About 95% of women

had a minimum of four antenatal visits.²³ This variation might be from the difference in sociodemographic variables between the study population.

The odds of usage of at least 4 ANC visits were higher among women with higher education compared to uneducated women. This finding was consistent with the findings of a prior study done in Ethiopia,^{11,16,18} and finding from Benin republic,²⁴ Tanzania,²⁵ Vietnam,²⁰ and Timor-Leste²² which showed that being educated or having a higher level of education had a positive impact on increasing the chance of at least 4 ANC visits. The reason might be; education creates a better chance to access through different sources, understand and utilize pieces of information²¹ regarding maternal and child health care as well as a better understanding of the importance of completing ANC visits and the consequence of danger signs on pregnancy. It is also true that education helps the mothers to engage in different activities and programs that scale up their understanding of ANC services.^{7,26} A possible explanation also might be that educated women might have a better chance of having their income sources and autonomy to decide on maternal health-care seeking. This is also plausible with the fact that women's autonomy to decide on health care service utilization increases their ANC visits/coverage.¹⁷

Women who had an early antenatal visit were 9 times more likely to utilize a minimum of 4 ANC visits compared to late started women. This finding was found to be consistent with a study in Tanzania²⁵ showing initiation of ANC after the first 4 weeks of pregnancy increases the chance of utilization of at least 4 ANC visits. Early antenatal care initiation is a key strategy to improve the utilization of maternal health care visits during pregnancy; WHO's recommendations to women attain adequate antenatal care visits.¹ This might be due to early initiation helps the women to get early counseling about ANC visits schedules and establish a client-provider relationship, this, in turn, helps them to complete at least 4 ANC visits.

The chance of completing ≥ 4 ANC visits was higher among pregnant women who had their husbands support/approval for antenatal visits compared to those counterparts. The above result was supported by a prior study done in Yem special wereda/district, Hadiya/South Ethiopia, Northern Ethiopia,^{19,27,28} Jayapura (Indonesia),²⁹ and Vietnam.²⁰ The possible reason might be due to women feels confident and increase their health-seeking behavior, these encourage them to attend more antenatal care visits. Women having a big problem in permission and support (not getting money) to

visit health facility could have a lower number of the visit of ANC during pregnancy,^{22,21}

On contrary; Number of pregnancy, Informed key pregnancy danger signs, TT vaccination during pregnancy, Iron and folate taken, Informed about Birth Preparedness Complication Readiness, and Status of Birth Preparedness Complication Readiness were those variables which were not significantly associated with utilization of at least 4 ANC visits in our study while these were found to have significant association on different studies.^{21,25} This discrepancy might be from the difference in the study setting.

Conclusion and Recommendation

The utilization of 4 ANC visits in Debre Berhan town is better compared to the national data (43%). Additionally; this study revealed that early initiation of ANC visits, male partner support, and higher educational level was significantly associated with the utilization of a minimum of 4 ANC services in Debre Berhan town. Promoting partner involvement during maternal health care services utilization and encouraging early initiation of first ANC visits are very important to scale up their utilization of at least 4 ANC services. In addition to increasing, universal access to education for women in the long run; Health promotion programs focusing on women with no education/lower educational status are important to increase their awareness about the importance of completing full antenatal services. Therefore, a collaboration of different stakeholders and an integrated effort should be applied to sustain the current positive outcome and to scale up the antenatal care service utilization.

Abbreviations

ANC, antenatal care; AOR, Adjusted Odds Ratio; BSc, Bachelor of Science; CI, Confidence Interval; COR, Crude Odds Ratio; AOR, adjusted odds ratio; SD, Standard Deviation; WHO, World Health Organization; SPSS- statistical package for social science; HSTP, Health sector transformation plan; HAD, health development army, RMNCAH, reproductive maternal newborn child and adolescent health; EDHS, Ethiopian demographic health survey; HIV, human immune-deficiency virus.

Data Sharing Statement

The data utilized to support the result of this study will be available from the corresponding author based on reasonable request.

Consent for Publication

Not applicable

Acknowledgment

First of all, we would like to thank God for giving us the courage and wisdom to do our work. Then, we would like to mention and pass our acknowledgment to Debre Berhan University, friends, and colleagues. We finally thank data collectors and supervisors for their commitment during the data collection process.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis, and interpretation, or in all these areas; took part in drafting, revising, or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted, and agree to be accountable for all aspects of the work.

Funding

The whole required (material and humanitarian) cost for this research work was covered by Debre Berhan University.

Disclosure

The authors of this paper declare that we have no conflict of interests.

References

1. WHO. *WHO Recommendations on Antenatal Care for a Positive Pregnancy Experience*. World Health Organization; 2016.
2. Institute EPH, ICF. *Ethiopia Mini Demographic and Health Survey 2019: Key Indicators*. Rockville, Maryland, USA: EPHI and ICF; 2019.
3. WHO. Maternal health global situation 2020 2020. Available from: https://www.who.int/health-topics/maternal-health#tab=tab_1. Accessed November 25, 2020.
4. WHO. *Strategies Towards Ending Preventable Maternal Mortality (EPMM)*. World Health Organization; 2015.
5. WHO. *Trends in Maternal Mortality 2000 to 2017: Estimates by WHO*. UNICEF, UNFPA, World Bank Group and the United Nations Population Division; 2019.
6. UNICEF. *UNICEF Data: Monitoring The Situation Of Children And Women*, in *2e State of the World's Children Report*. New York, NY, USA: UNICEF; 2019.
7. Yeneneh A, Alemu K, Dadi AF, Alamirrew A. Spatial distribution of antenatal care utilization and associated factors in Ethiopia: evidence from Ethiopian demographic health surveys. *BMC Pregnancy Childbirth*. 2018;18(1):1–12. doi:10.1186/s12884-018-1874-2
8. Agency CS, ICF. *Ethiopia Demographic and Health Survey 2016: Key Indicators Report*. Addis Ababa, Ethiopia, and Rockville, Maryland, USA: CSA and ICF; 2016.
9. Central Statistical Agency I. *Ethiopia Demographic and Health Survey 2011: Key Indicators Report*. Addis Ababa, Ethiopia, and Rockville, Maryland, USA: CSA and ICF; 2011.
10. Central Statistical Agency I. *Ethiopia Demographic and Health Survey 2005: Key Indicators Report*. Addis Ababa, Ethiopia, and Rockville, Maryland, USA: CSA and ICF; 2005.
11. Basha GW. Factors affecting the utilization of a minimum of four antenatal care services in Ethiopia. *Obstet Gynecol Int*. 2019;2019.
12. Tsegaye B, Ayalew M. Prevalence and factors associated with antenatal care utilization in Ethiopia: an evidence from demographic health survey 2016. *BMC Pregnancy Childbirth*. 2020;20(1):1–9. doi:10.1186/s12884-020-03236-9
13. Medhanyie A, Spigt M, Kifle Y, et al. The role of health extension workers in improving utilization of maternal health services in rural areas in Ethiopia: a cross sectional study. *BMC Health Serv Res*. 2012;12(1):352. doi:10.1186/1472-6963-12-352
14. Belda SS, Gebremariam MB. Birth preparedness, complication readiness and other determinants of place of delivery among mothers in Goba District, Bale Zone, South East Ethiopia. *BMC Pregnancy Childbirth*. 2016;16(1):73. doi:10.1186/s12884-016-0837-8
15. Barrett G, Wellings K. What is a 'planned' pregnancy? Empirical data from a British study. *Soc Sci Med*. 2002;55(4):545–557. doi:10.1016/S0277-9536(01)00187-3
16. Muchie KF. Quality of antenatal care services and completion of four or more antenatal care visits in Ethiopia: a finding based on a demographic and health survey. *BMC Pregnancy Childbirth*. 2017;17(1):1–7. doi:10.1186/s12884-017-1488-0
17. Tegegne TK, Chojenta C, Getachew T, Smith R, Loxton D. Antenatal care use in Ethiopia: a spatial and multilevel analysis. *BMC Pregnancy Childbirth*. 2019;19(1):399. doi:10.1186/s12884-019-2550-x
18. Ousman SK, Mdala I, Thorsen VC, Sundby J, Magnus JH. Social determinants of antenatal care service use in Ethiopia: changes over a 15-year span. *Front Public Health*. 2019;7:161. doi:10.3389/fpubh.2019.00161
19. Ftwi M, Gebretsadik GG-E, Berhe H, Haftu M, Gebremariam G, Tesfau YB. Coverage of completion of four ANC visits based on recommended time schedule in Northern Ethiopia: a community-based cross-sectional study design. *PLoS One*. 2020;15(8):e0236965. doi:10.1371/journal.pone.0236965
20. Ha BT, Tac PV, Duc DM, Duong DT, Thi LM. Factors associated with four or more antenatal care services among pregnant women: a cross-sectional survey in eight south central coast provinces of Vietnam. *Int J Women's Health*. 2015;7:699.
21. Adewuyi EO, Auta A, Khanal V, et al. Prevalence and factors associated with underutilization of antenatal care services in Nigeria: a comparative study of rural and urban residences based on the 2013 Nigeria demographic and health survey. *PLoS One*. 2018;13(5):e0197324. doi:10.1371/journal.pone.0197324
22. Khanal V, da Cruz JLNB, Mishra SR, Karkee R, Lee AH. Underutilization of antenatal care services in Timor-Leste: results from Demographic and Health Survey 2009–2010. *BMC Pregnancy Childbirth*. 2015;15(1):211. doi:10.1186/s12884-015-0646-5
23. Yeji F, Shibanuma A, Oduro A, et al. Continuum of care in a maternal, newborn and child health program in Ghana: low completion rate and multiple obstacle factors. *PLoS One*. 2015;10(12):e0142849. doi:10.1371/journal.pone.0142849
24. Dansou J, Adekunle AO, Arowojolu AO. Factors associated with antenatal care services utilisation patterns amongst reproductive age women in Benin Republic: an analysis of 2011/2012 Benin Republic's demographic and health survey data. *Nigerian Postgrad Med J*. 2017;24(2):67. doi:10.4103/npmj.npmj_16_17

25. Gupta S, Yamada G, Mpembeni R, et al. Factors associated with four or more antenatal care visits and its decline among pregnant women in Tanzania between 1999 and 2010. *PLoS One*. 2014;9(7):e101893. doi:10.1371/journal.pone.0101893
26. Ayalew TW, Nigatu AM. Focused antenatal care utilization and associated factors in Debre Tabor Town, northwest Ethiopia, 2017. *BMC Res Notes*. 2018;11(1):819. doi:10.1186/s13104-018-3928-y
27. Tewodros B, Dibaba Y. Factors affecting antenatal care utilization in Yem special woreda, southwestern Ethiopia. *Ethiop J Health Sci*. 2009;19:1.
28. Abosse Z, Woldie M, Ololo S. Factors influencing antenatal care service utilization in hadiya zone. *Ethiop J Health Sci*. 2010;20:2.
29. Kamo B, Msen Y, Rantetampang A, Mallongi A. The factors affecting with four visited at public health centre Sub Province Mimika Papuan Province. *Education*. 2018;76(11):6.

Risk Management and Healthcare Policy

Dovepress

Publish your work in this journal

Risk Management and Healthcare Policy is an international, peer-reviewed, open access journal focusing on all aspects of public health, policy, and preventative measures to promote good health and improve morbidity and mortality in the population. The journal welcomes submitted papers covering original research, basic science, clinical & epidemiological studies, reviews and evaluations,

guidelines, expert opinion and commentary, case reports and extended reports. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/risk-management-and-healthcare-policy-journal>