Resveratrol Ameliorates Systemic Sclerosis via Suppression of Fibrosis and Inflammation Through Activation of SIRT1/mTOR Signaling

This article was published in the following Dove Press journal:
Drug Design, Development and Therapy

Qicen Yao^{1,*}
Qingchao Wu^{2,*}
Xiayu Xu¹
Yixi Xing¹
Jin Liang¹
Qianqi Lin¹
Meiqiong Huang¹
Yiling Chen¹
Bo Lin³
Weifei Chen¹

¹Department of Rheumatology and Immunology, The Second Affiliated Hospital of Hainan Medical University, Haikou, People's Republic of China;

²Department of Rheumatology and Immunology, Minda Hospital of Hubei Minzu University, Enshi, Hubei Province, People's Republic of China; ³Department of Pharmacy, The Second Affiliated Hospital of Hainan Medical University, Haikou, Hainan Province, People's Republic of China

*These authors contributed equally to this work

Correspondence: Weifei Chen
Department of Rheumatology and Immunology, The Second Affiliated Hospital of Hainan Medical University, 368 Yehai Boulevard, Haikou City, Hainan Province, People's Republic of China
Email chenwf130@163.com

Bo Lin
Department of Pharmacy, The Second Affiliated Hospital of Hainan Medical University, 368 Yehai Boulevard, Haikou City, Hainan Province, People's Republic of China
Email linbo13518072160@163.com

Purpose: Resveratrol (Res) is a natural polyphenolic compound found in several plants and reported as a promising biological molecule with effective anti-fibrosis and anti-inflammatory activities. However, the underlying mechanism of Res on systemic sclerosis (SSc) remains unclear. In the study, we identified the key cellular signaling pathways involved in the Res regulatory process on SSc.

Methods: Res-targeted genes interaction network was constructed using the STITCH database, and the shared Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways involved in both SSc and Res-targeted genes were then identified. The top five enriched KEGG pathways were visualized by GOplot. KEGG pathways associated with Res-targeted genes were established by Pathway Builder Tool 2.0. Quantitative real-time PCR (qRT-PCR) was used to assess the expression of sirtuin 1 (SIRT1), mammalian targeted of rapamycin (mTOR), and cytokines.

Results: Enrichment analysis of Res-targeted genes showed 79 associated pathways, 27 of which were also involved in SSc. Particularly, SIRT1/mTOR signaling was found as one of the crucial regulatory pathways. In vitro results suggested that SIRT1-mediated mTOR degradation ameliorated bleomycin (BLM)-induced fibrosis and inflammation. Res was capable of elevating the SIRT1 level in fibroblasts and partially reversing mTOR-dependent induction of fibrosis and inflammation.

Conclusion: These results indicated that Res is a feasible and effective choice for SSc and therapeutic target of mTOR could be a potential alternative for treatment of SSc.

Keywords: resveratrol, systemic sclerosis, SIRT1, mTOR, signaling

Introduction

Systemic sclerosis (SSc) is an autoimmune disease characterized by microvascular damage, innate and adaptive immune dysfunction, and multiple organ fibrosis, seriously affecting the quality of a patient's life and placing a heavy burden on society.¹ A previous study reported that the quality of a patient's life was significantly lower than the control group using a SF-36 health questionnaire.² The functional damage of patients is gradually aggravated over the prolongation of time.³ However, the specific pathological mechanism of SSc and its etiological factors remain unclear. Thus, it is an urgent time to investigate the underlying mechanism of SSc and explore a feasible and effective drug to ameliorate the pathological progress of fibrosis and inflammation in SSc patients.

Resveratrol (Res, 3-4'-5-trihydroxystilbene), a non-flavonoid polyphenolic compound, is an antitoxin produced when plants are stimulated. It can be synthesized in

grape leaves and is a bioactive ingredient in grape wine and grape juice.⁴ Previous in vitro and in vivo experiments suggested that Res has antioxidant, anti-inflammatory, anti-cancer and cardiovascular protective effects.⁵ Currently, accumulative attention has been attracted to the therapeutic effect of Res on SSc due to its anti-fibrotic and anti-inflammatory activities.⁶ For example, it was recently reported that Res could alleviate SSc-related pulmonary fibrosis through suppression of pro-inflammatory and pro-fibrotic processes.⁷ Recent research also indicated that Res-induced SIRT1 inhibits transforming growth factor- β (TGF- β) induced cell proliferation and fibroblast differentiation.⁸ mTOR is a serine/threonine protein kinase which has been shown to be involved in the initiation of inflammatory cascades and fibroblast activation.⁹

Microarrays, a lab technique to detect thousands of gene expressions, have been widely applied to the investigation of the pathogenic processes governing the development of a variety of diseases. Integrating microarray and bioinformatic tools makes it an invaluable method for functional genomic studies.¹⁰ In the present study, Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways related to both Res-targeted genes and SSc development were analyzed. The mammalian target of rapamycin (mTOR) signaling was identified as a crucial KEGG pathway related to Res-targeted genes. Therefore, we investigated the effects of Res on SSc while focusing on the mTOR signaling in the present study.

Materials and Methods

Reagent

Res was purchased from Sigma-Aldrich Co. (St Louis, MO, USA). Bleomycin (BLM) was obtained from Enzo Life Sciences, Inc. (Farmingdale, NY, USA). qRT-PCR kit was purchased from Thermo Fisher Scientific Co. (Waltham, MA, USA). Small interfering RNA (siRNA) mTOR and the siRNA control were designed by Applied Biological Materials Inc. (Richmond, BC, Canada). Plasmid SIRT1 was purchased from the Public Protein/Plasmid Library (PPL, #BC012499). High-glucose Dulbecco's modified eagle medium was purchased from Gibco BRL (Thermo Fisher).

Ethical Statement

Protocols of the present study were approved by the Ethics Committee of The Second Affiliated Hospital of Hainan Medical University (Haikou, China; 2019-019-23), and informed written consent was achieved from all participants.

All experiments involved with human samples were conducted in compliance with the Declaration of Helsinki.

Skin Sample Collection

From August 2018 to May 2020, skin biopsy specimens of the affected skin were collected both from SSc patients, and the specimens fulfilled the American College of Rheumatology criteria for SSc. At the same time, skin biopsy specimens were obtained from normal controls who had no history of autoimmune and other dermal diseases.

Culture of Human Skin Fibroblasts (HSFs)

High-glucose Dulbecco's modified eagle medium was used to culture HSFs (FuHeng Biology, Shanghai, China). HSFs were cultured at 37°C with 5% CO₂ and 95% humidity. HSFs treated with BLM (100 mmol/mL) for 24 hours were selected for further research. To knockdown endogenous mTOR, siRNA mTOR transfection was performed by Lipofectamine RNAiMAX (Life Technologies) according to the manufacturer's protocol. qRT-PCR and ELISA analyses were performed 24 hours post-lipofection. The plasmid SIRT1 was purchased from the Public Protein/Plasmid Library (PPL, #BC012499). qRT-PCR and ELISA analyses were performed 24 hours after plasmid transfection.

qRT-PCR Analysis

The total RNA was isolated from cells and skin samples using TRIzol[®] reagent (Thermo Fisher Scientific). The reverse-transcription process was performed using ReverTra Ace[®] qPCR RT Master Mix (Toyobo Life Science, Osaka, Japan) to synthesize cDNA according to the manufacturer's protocol. The relative mRNA levels were normalized to GAPDH levels (the internal control) and were calculated according to the $2^{-\Delta\Delta C_t}$ approach. All experiments were conducted in triplicate, and the primer sequences of mRNAs are shown in Table 1.

Enzyme-Linked Immunosorbent Assay (ELISA)

The concentrations of IL-1 β and IL-6 in cell supernatants were measured with specific ELISA kits (Shanghai Biotechnology, Shanghai, China) in accordance with the manufacturer's instructions.

Res-Targeted Genes Interaction Network Construction

Search Tool for Interacting Chemicals (STITCH) is a database that integrates over 430,000 chemicals into

Table 1 microRNAs and mRNA Primer Sequences

microRNA or Gene Names	Primer Sequence (5' to 3')
hsa-mTOR-Forward	TCCAGGGCTTCTTCCGTT
hsa-mTOR-Reverse	GGGCTGTCGTGGTAGACTTAG
hsa-SIRT1-Forward	CGGAACAATACCTCCACCTGA
hsa-SIRT1-Reverse	GAAGTCTACAGCAAGGCGAGCA
hsa- α -SMA-Forward	TGTATGTGGCTATCCAGGCG
hsa- α -SMA-Reverse	AGAGTCCAGCACGATGCCAG
hsa-Col-1 α 1-Forward	CCTCCAGGGCTCCAACGAG
hsa-Col-1 α 1-Reverse	TCTATCACTGTCTTGCCCCA
hsa-Col-1 α 2-Forward	GATGTTGAACCTGTTGCTGAGC
hsa-Col-1 α 2-Reverse	TCTTTCCCCATTCAATTGTCTT
hsa-IL-1 β -Forward	AGCTACGAATCTCCGACCAC
hsa-IL-1 β -Reverse	CGTTATCCCATGTGTCGAAGAA
hsa-IL-6-Forward	ATGAACCTCTTCTCCACAAGC
hsa-IL-6-Reverse	AGGCGGTGCTTGTTCTCTCA
hsa-GAPDH-Forward	ACCCACTCTCCACCTTTGA
hsa-GAPDH-Reverse	CTGTTGCTGTAGCCAAATTCGT

a single resource. Res-targeted genes were obtained in STITCH based on the following settings: first shell: 0 and up to 10; second shell: 0 and up to 10; third shell: 0 and up to 10. The first shell is directly associated with resveratrol. The second shell is secondarily associated with resveratrol through the first shell. The interaction network of Res and its targeted genes were constructed in STITCH online tool. To obtain the degree, betweenness, closeness of each gene in the network and better display the three layers of genes, the Res-targeted gene symbols were imported into Cytoscape 3.7.2 and calculated. Degree represented the degree of association of one node with all other nodes in the network. Closeness represented the closeness between a node and other nodes in the network. Betweenness was the number of times a node acted as the shortest bridge between two other nodes.

Shared KEGG Pathways Involved Both in SSc and Res-Targeted Genes

miRWalk2.0 is a comprehensive archive, supplying the largest available collection of predicted and experimentally verified genes and related information. miRWalk2.0 was used to retrieve human SSc KEGG pathways. Some functional associations such as gene class, protein class, KEGG, and WIKI pathways can be obtained from the database. The KEGG pathways information was downloaded. Database for Annotation, Visualization, and Integrated Discovery

(DAVID) bioinformatics resources consists of an integrated biological knowledgebase and analytic tools aimed at systematically extracting biological meaning from large gene/protein lists. DAVID was used to generate the Res-targeted genes enriched KEGG pathways. Shared KEGG pathways of SSc and Res-targeted genes were identified using Venn Diagram (<http://www.ehbio.com/ImageGP/index.php/Home/Index/VennDiagram.html>).

Identification of the Hub Genes

The enrichment information of the top KEGG pathways with $p < 0.01$ was presented with GOplot, an R package for visually combining expression data with functional analysis. Genes involved in all the top three shared KEGG pathways were considered as hub genes.

Retrieval of the KEGG Pathways Related to Res-Targeted Genes

The top 3 shared KEGG pathways with smallest p values were selected. The KEGG pathways related to the Res-targeted genes were established with the ScienceSlides. Hub genes and their action mechanisms were shown in the schematic diagrams.

Statistical Analysis

GraphPad Prism 8.0 was used to perform the statistical analyses. Results were shown as the mean $\pm$ SD. Student's t -test was applied to compare two groups, and one-way

Figure 1 The interaction network of the resveratrol-targeted genes. Red is first shell. Yellow is second shell. Green is third shell.

analysis of variance with Tukey's post hoc test was used to compare three or more groups. $P < 0.05$ was considered as statistically significant.

Results

Res-Targeted Genes and the Interaction Network

In total, 30 Res-targeted genes were obtained in STITCH with the limitation of three shells. Then the interaction of Res-targeted genes was reconstructed in Cytoscape (Figure 1). *TP53*, *SIRT1*, *PTGS1*, *SIRT3*, *ESR1*, *PPARG*, *NOS3*, *AKT1*, *SIRT5*, *PTGS2* were involved in the first shell. They were directly associated with resveratrol. *ATM*, *BRCA1*, *FOXO1*, *MTOR*, *EP300*, *RICTOR*, *FOXO3*, *CDKN1A*, *KAT2B*, *MDM2* were involved in the second shell. *HSP90AA1*, *HIPK2*,

NCOA3, *CDKN2A*, *MAPK8*, *SRC*, *USP7*, *RCHY1*, *CREBBP*, *SP1* were involved in the third shell.

Identification of the Shared KEGG Pathways of SSc and Res-Targeted Genes

After the enrichment analysis of the Res-targeted genes in DAVID, 57 KEGG pathways were identified. 79 KEGG pathways involved in human SSc were retrieved using miRWalk2.0. 27 shared KEGG pathways of SSc and Res-targeted genes were identified (Figure 2 and Table S1). The top nine KEGG pathways with $p < 0.01$ were Prostate cancer pathway, Pathway in cancer, Glioma pathway, p53 signaling pathway, Cell cycle pathway, Bladder cancer pathway, Melanoma pathway, Chronic myeloid leukemia pathway, and ErbB signaling pathway. The rich factor, $-\log(p)$, and

Figure 2 Common KEGG pathways of resveratrol-target genes and CCs. There are 57 resveratrol-target genes related KEGG pathways and 79 SSc related KEGG pathways. 27 shared KEGG pathways were obtained.

Figure 3 Enrichment information of KEGG pathways with $p < 0.01$. hsa05220: Chronic myeloid leukemia pathway, hsa05219: Bladder cancer pathway, hsa05218: Melanoma pathway, hsa05215: Prostate cancer pathway, hsa05214: Glioma pathway, hsa05200: Pathways in cancer, hsa04115: p53 signaling pathway, hsa04110: Cell cycle signaling pathway, hsa04012: ErbB signaling pathway.

gene number of the nine pathways are demonstrated in Figure 3. The detailed information of the top 3 KEGG pathways are shown in Table 2.

Identification of the Hub Genes

Among the 30 Res-targeted genes, *AKT1*, *CDKN1A*, *TP53*, *MDM2*, *MTOR*, *HSP90AA1*, *EP300*, *CREBBP*, *FOXO1*, *PTGS2*, *PPARG*, *CDKN2A*, *MAPK8*, *RCHY1*, and *ATM* were involved in the top three shared KEGG pathways. Degree, betweenness, and closeness information of all 30 Res-targeted genes are shown in Figure 4. The enrichment information of the top 9 KEGG pathways is shown in

Figure 5. *AKT1*, *CDKN1A*, *TP53*, *MDM2*, *MTOR* are involved in all of the top three KEGG pathways and were identified as hub genes.

Retrieval of the KEGG Pathways Related to Res-Targeted Genes

The Res-targeted genes related parts of the top three shared KEGG pathways are shown in Figure 6. They are associated with the cell proliferation, survival, tumor growth, impaired G1 and G2 arrest, reduced apoptosis, genomic instability, cell cycle progression, apoptosis inhibition, evading apoptosis, G1/S progression, uncontrolled

Table 2 The Information of Top 3 KEGG Pathways

KEGG Pathway	Count	FDR	Genes
hsa05215: Prostate cancer	9	2.1E-8	<i>CDKN1A</i> , <i>HSP90AA1</i> , <i>CREBBP</i> , <i>MDM2</i> , <i>EP300</i> , <i>AKT1</i> , <i>TP53</i> , <i>FOXO1</i> , <i>MTOR</i>
hsa05200: Pathways in cancer	13	1.4E-7	<i>CDKN1A</i> , <i>CREBBP</i> , <i>HSP90AA1</i> , <i>CDKN2A</i> , <i>PTGS2</i> , <i>FOXO1</i> , <i>MTOR</i> , <i>MAPK8</i> , <i>MDM2</i> , <i>AKT1</i> , <i>EP300</i> , <i>PPARG</i> , <i>TP53</i>
hsa05214: Glioma	6	4.2E-5	<i>CDKN1A</i> , <i>CDKN2A</i> , <i>MDM2</i> , <i>AKT1</i> , <i>TP53</i> , <i>MTOR</i>

Figure 4 Degree, betweenness, closeness of resveratrol-targeted genes.

proliferation. It can be found they may have biological action through PI3K-Akt signaling pathway, p53 signaling pathway, and mTOR signaling pathway.

Res Inhibits Fibrosis and Inflammation Through Reduction of the Expression of mTOR

As seen in Figure 6, mTOR signaling was identified as a critical KEGG pathway related to Res-targeted genes.

Thus, we next examined the association of Res and mTOR and their regulatory roles in fibrosis and inflammation. First, the gene expression of mTOR in skin tissues from both SSc patients and the healthy controls was measured by qRT-PCR. The result showed that mTOR level in the skin sample of SSc patients was significantly higher than the normal controls (Figure 7A). In vitro, BLM was used to treat HSFs to create a cellular model of fibroblast activation. The in vitro results indicated that Res was able to markedly reverse the high level of mTOR induced by BLM. 10 μ M was shown as the

Figure 5 The resveratrol-targeted genes enrichment analysis information.

optimum concentration of Res to exert its effective regulation (Figure 7B). Furthermore, the effect of Res on fibrosis was investigated in vitro and the results suggested that Res could significantly reverse the BLM-induced high expression of

fibrotic genes, including α -SMA, Col 1a1, and Col 1a2 (Figure 7C–E). Similarly, the qRT-PCR results also indicated that Res could significantly reverse the BLM-induced expression of inflammatory cytokines including IL-1 β and

Figure 7 Res inhibits fibrosis and inflammation through reduction of the expression of mTOR. (A) The expression of mTOR in the skin samples from SSc patients and the normal controls was measured by qRT-PCR analysis, $n = 10$, per group; (B) The expression of mTOR in HSFs with different treatments was measured by qRT-PCR analysis; (C–E) The expression of fibrotic genes in the different groups was measured by qRT-PCR analysis; (F, G) The expression of inflammatory cytokines IL-1 β and IL-6 in the different groups was measured by qRT-PCR analysis; (H) The expression of mTOR in HSFs with different treatments was measured by qRT-PCR analysis; (I–K) The expression of fibrotic genes in the different groups was measured by qRT-PCR analysis; (L, M) The expression of inflammatory cytokines IL-1 β and IL-6 in the different groups was measured by qRT-PCR analysis. Data are the means $\pm$ SDs of three independent experiments.

for the generation and secretion of fibrotic and inflammatory cytokines, and play crucial roles in the pathogenesis of SSc.^{19,20} In the present study, we identified *mTOR* signaling as an important KEGG pathway regarding the

Res-targeted genes, and many prior studies have demonstrated that *mTOR* is an important factor in SSc.^{21,22} Previous literature have also shown the *mTOR* signaling which may be regulated by the *SIRT1* expression.²³ For

Figure 8 Res ameliorates SSc via regulation of SIRT1/mTOR signaling. (A) The expression of SIRT1 in HSFs with different treatments was measured by qRT-PCR analysis; (B) The expression of SIRT1 in HSFs after the SIRT1 plasmid inference was measured by qRT-PCR analysis; (C) The expression of mTOR in HSFs after the SIRT1 plasmid inference was measured by qRT-PCR analysis; (D–F) The expression of fibrotic genes in the different groups was measured by qRT-PCR analysis; (G, H) The expression of inflammatory cytokines IL-1 β and IL-6 in the different groups was measured by qRT-PCR analysis. Data are the means $\pm$ SDs of three independent experiments.

example, a recent study pointed out that *SIRT1/mTOR* signaling was the potential mechanism of the lycopene's effect on the improved pain experience in the dorsal horn of burn injury rats.²⁴ Our present study indicated that the Res-dependent overexpression of *SIRT1* could markedly suppress the *mTOR* expression and exert a strong anti-fibrosis and anti-inflammation in vitro. Thus, we speculate that *SIRT1/mTOR* axis be the underlying mechanism of Res on SSc.

In this study, not only *SIRT1* but also *SIRT3* and *SIRT5* were found directly targeted by Res. *SIRT1* is localized to both the nucleus and cytoplasm, deacetylating intracellular signaling proteins, including histones, and regulate gene expression.^{25–27} *SIRT3* and *SIRT5* are exclusively localized to the mitochondria and regulate energy metabolism in response to mitochondrial stress.^{27–30} Whether *SIRT3* and *SIRT5* play a role in

promoting SSc by Res needs to be investigated in future studies.

Conclusion

Our study determined that Res exhibited protective effects on SSc, as evidenced by suppressing the increase in fibrotic and inflammatory cytokines observed in vitro. The mainly involved mechanism was the anti-fibrosis and anti-inflammatory role of Res through *SIRT1/mTOR* axis. This study provided a potential candidate natural compound for the treatment of SSc.

Acknowledgments

This study was supported by the Research and Development fund of the Second Affiliated Hospital of Hainan Medical University (Hospital Fund Number: 2018–19).

Author Contributions

WC and BL designed the study. QY and QW collated the data, analyzed the data and produced the initial draft of the manuscript. QY, QW, XX, YX, JL, QL, MH, YC, BL and WC revised the figures and tables. QY, QW, BL and WC helped to draft the manuscript. All authors contributed to data analysis, drafting or revising the article, have agreed on the journal to which the article will be submitted, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Disclosure

All authors declared no conflicts of interest.

References

- Hoffmann-Vold AM, Molberg Ø. Detection, screening, and classification of interstitial lung disease in patients with systemic sclerosis. *Curr Opin Rheumatol*. 2020;32(6):497–504. doi:10.1097/BOR.0000000000000741
- Lo LMP, Taylor BV, Winzenberg T, et al. Estimating the relative contribution of comorbidities in predicting health-related quality of life of people with multiple sclerosis. *J Neurol*. 2020. doi:10.1007/s00415-020-10195-w
- Geraldes R, Esiri MM, Perera R, et al. Vascular disease and multiple sclerosis: a post-mortem study exploring their relationships. *Brain*. 2020;143(10):2998–3012. doi:10.1093/brain/awaa255
- Rawat D, Chhonker SK, Naik RA, et al. Modulation of antioxidant enzymes, SIRT1 and NF-κB by resveratrol and nicotinamide in alcohol-aflatoxin B1-induced hepatocellular carcinoma. *J Biochem Mol Toxicol*. 2020:e22625. doi:10.1002/jbt.22625
- Fonseca-Santos B, Chorilli M. The uses of resveratrol for neurological diseases treatment and insights for nanotechnology based-drug delivery systems. *Int J Pharm*. 2020;589:119832. doi:10.1016/j.ijpharm.2020.119832
- Ahmad I, Hoda M. Attenuation of diabetic retinopathy and neuropathy by resveratrol: review on its molecular mechanisms of action. *Life Sci*. 2020;245:117350. doi:10.1016/j.lfs.2020.117350
- Chu H, Jiang S, Liu Q, et al. Sirtuin1 protects against systemic sclerosis-related pulmonary fibrosis by decreasing proinflammatory and profibrotic processes. *Cell Mol Biol*. 2018;58:28–39.
- Nowosad A, Besson A. CDKN1B/p27 regulates autophagy via the control of regulator and MTOR activity in amino acid-deprived cells. *Autophagy*. 2020;1–2. doi:10.1080/15548627.2020.1831217
- Sharma D, Malik A, Balakrishnan A, et al. RIPK3 promotes mefv expression and pyrin inflammasome activation via modulation of mtor signaling. *J Immunol*. 2020;ji2000244. doi:10.4049/jimmunol.2000244
- Xiong Y, Cao F, Chen L, et al. Identification of key microRNAs and targeted genes for the diagnosis of bone nonunion. *Mol Med Rep*. 2020;21:1921–1933. doi:10.3892/mmr.2020.10996
- Komorowska J, Wątroba M, Szukiewicz D. Review of beneficial effects of resveratrol in neurodegenerative diseases such as alzheimer's disease. *Adv Med Sci*. 2020;65(2):415–423. doi:10.1016/j.advms.2020.08.002
- Viana-Mattioli S, Cineaglia N, Bertozzi-Matheus M, et al. SIRT1-dependent effects of resveratrol and grape juice in an in vitro model of preeclampsia. *Biomed Pharmacother*. 2020;131:110659. doi:10.1016/j.biopha.2020.110659
- Giancchetti E, Fierabracci A. Insights on the effects of resveratrol and some of its derivatives in cancer and autoimmunity: a molecule with a dual activity. *Antioxidants (Basel)*. 2020;9. doi:10.3390/antiox9020091
- Zerr P, Palumbo-Zerr K, Huang J, et al. Sirt1 regulates canonical TGF-β signaling to control fibroblast activation and tissue fibrosis. *Ann Rheum Dis*. 2016;75(1):226–233. doi:10.1136/annrheumdis-2014-205740
- Jiang Y, Hu F, Li Q, et al. Tanshinone IIA ameliorates the bleomycin-induced endothelial-to-mesenchymal transition via the Akt/mTOR/p70S6K pathway in a murine model of systemic sclerosis. *Int Immunopharmacol*. 2019;77:105968. doi:10.1016/j.intimp.2019.105968
- Chen C, Akiyama K, Wang D, et al. mTOR inhibition rescues osteopenia in mice with systemic sclerosis. *J Exp Med*. 2015;212(1):73–91. doi:10.1084/jem.20140643
- Mutlu E, Gibbs ST, South N, et al. Comparative toxicokinetics of trans-resveratrol and its major metabolites in harlan sprague dawley rats and B6C3F1/N mice following oral and intravenous administration. *Toxicol Appl Pharmacol*. 2020;394:114962. doi:10.1016/j.taap.2020.114962
- Cao X, Tian S, Fu M, et al. Resveratrol protects human bronchial epithelial cells against nickel-induced toxicity via suppressing p38 MAPK, NF-κB signaling, and NLRP3 inflammasome activation. *Environ Toxicol*. 2020;35(5):609–618. doi:10.1002/tox.22896
- Trapiella-Martínez L, Díaz-López JB, Caminal-Montero L, et al. Very early and early systemic sclerosis in the Spanish scleroderma registry (RESCLE) cohort. *Autoimmun Rev*. 2017;16(8):796–802. doi:10.1016/j.autrev.2017.05.013
- Gordon SM, Stitt RS, Nee R, et al. Risk factors for future scleroderma renal crisis at systemic sclerosis diagnosis. *J Rheumatol*. 2019;46(1):85–92. doi:10.3899/jrheum.171186
- Qi Q, Mao Y, Tian Y, et al. Geniposide inhibited endothelial-mesenchymal transition via the mTOR signaling pathway in a bleomycin-induced scleroderma mouse model. *Am J Transl Res*. 2017;9:1025–1036.
- Suto T, Karonitsch T. The immunobiology of mTOR in autoimmunity. *J Autoimmun*. 2020;110:102373. doi:10.1016/j.jaut.2019.102373
- Peng C, Fuchao C, Jiexin L, et al. Activation of the miR-34a-mediated SIRT1/mTOR signaling pathway by urolithin a attenuates d-galactose-induced brain aging in mice. *Neurotherapeutics*. 2019;16(4):1269–1282. doi:10.1007/s13311-019-00753-0
- Qin Y, Jin-Feng W, Xiao-Hua X, et al. Effect of lycopene on pain facilitation and the SIRT1/mTOR pathway in the dorsal horn of burn injury rats. *Eur J Pharmacol*. 2020; 889:173365.
- Chang H-C, Guarente L. SIRT1 and other sirtuins in metabolism. *Trends Endocrinol Metab*. 2014;25(3):138–145. doi:10.1016/j.tem.2013.12.001
- Guarente L. Calorie restriction and sirtuins revisited. *Genes Dev*. 2013;27(19):2072–2085. doi:10.1101/gad.227439.113
- Lee H-C, Chathuranga K, Lee J-S. Mechanisms and disease implications of sirtuin-mediated autophagic regulation. *Exp Mol Med*. 2019;51(12):1–11. doi:10.1038/s12276-019-0299-y
- Choi J-E, Mostoslavsky R. Sirtuins, metabolism, and DNA repair. *Curr Opin Genet Dev*. 2014;26:24–32. doi:10.1016/j.gde.2014.05.005
- Yang W, Nagasawa K, C. M, et al. Mitochondrial sirtuin network reveals dynamic SIRT3-dependent deacetylation in response to membrane depolarization. *Cell*. 2016;167(4):985–1000.e21. doi:10.1016/j.cell.2016.10.016
- van de Ven Robert AH, Santos D, Haigis Marcia C. Mitochondrial sirtuins and molecular mechanisms of aging. *Trends Mol Med*. 2017;23(4):320–331. doi:10.1016/j.molmed.2017.02.005

Drug Design, Development and Therapy**Dovepress****Publish your work in this journal**

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which has also

been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/drug-design-development-and-therapy-journal>