Immunohistochemical Analysis of PGC-1 α and ERR α Expression Reveals Their Clinical Significance in Human Ovarian Cancer

This article was published in the following Dove Press journal:
OncoTargets and Therapy

Xiqi Huang ¹
Guanyu Ruan ^{1,2}
Guifen Liu ¹
Yuqin Gao¹
Pengming Sun ^{1,2}

¹Laboratory of Gynecologic Oncology, Fujian Provincial Maternity and Children's Health Hospital, Affiliated Hospital of Fujian Medical University, Fuzhou, People's Republic of China; ²Key Laboratory of Women and Children's Critical Diseases Research, Fujian Provincial Maternity and Children's Health Hospital, Affiliated Hospital of Fujian Medical University, Fuzhou, People's Republic of China

Purpose: Peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1 α) and estrogen-related receptor alpha (ERR α) play a vital role in various human cancers. The purpose of this study was to investigate whether the PGC-1 α /ERR α axis could serve as an effective prognostic marker in ovarian cancer (OC).

Patients and Methods: We investigated the expression of both PGC-1 α and ERR α in 42 ovarian cancer and 31 noncancerous ovarian samples by immunohistochemistry (IHC). The relationship between the expression of PGC-1 α and ERR α in OC and the clinical characteristics of patients was evaluated. In addition, data from the Human Protein Atlas (HPA) database were collected to validate the prognostic significance of PGC-1 α and ERR α mRNA expression in OC.

Results: PGC-1 α and ERR α showed notably higher expression in OC tissues than in noncancerous tissues ($P=0.0059$, $P=0.002$). Moreover, in patients with OC, high ERR α and PGC-1 α /ERR α expression significantly correlated with tumor differentiation ($P=0.027$; $P=0.04$), lymph node status ($P=0.023$; $P=0.021$), CA125 ($P=0.036$; $P=0.021$), and HE4 ($P=0.021$; $P=0.05$), while high PGC-1 α expression was only significantly associated with tumor differentiation ($P=0.029$). The combined analysis of high PGC-1 α and ERR α expression revealed a tendency towards poor cancer-specific survival ($P=0.1276$).

Conclusion: PGC-1 α and ERR α are overexpressed in OC and might be significant prognostic factors for this cancer.

Keywords: prognostic significance, estrogen-related receptor alpha, immunohistochemistry, ovarian malignance, peroxisome proliferator-activated receptor gamma coactivator 1-alpha

Introduction

Ovarian cancer is one of the most fatal malignancies in women and poses a serious threat to women's health. Despite some progress achieved with extensive clinical and basic research, the etiology and tumorigenesis of OC are not fully understood. Tumor heterogeneity leads to different stages and characteristics of disease development. Different molecular drivers may exist in patients with OC at the same stage, leading to heterogeneous treatment responses and prognosis.¹ It is therefore urgent to investigate the molecular markers underlying ovarian cancer and identify novel therapeutic targets for OC treatment.

Estrogen-related receptor alpha (ESRRA, also known as ERR α), a member of the ligand-independent orphan nuclear receptor superfamily, is expressed primarily in tissues with high metabolic demand and controls the expression of genes involved in

Correspondence: Pengming Sun
Key Laboratory of Women and Children's Critical Diseases Research, Fujian Provincial Maternity and Children's Health Hospital, Affiliated Hospital of Fujian Medical University, No. 18, Daoshan Road, Gulou District, Fuzhou, 350001 Fujian, People's Republic of China
Tel +86 591 8755 8732
Fax +86 591 8755 1247
Email sunfemy@hotmail.com

cellular energy metabolism.^{2–4} Notably, *ERRα* has been reported to be overexpressed in breast, colon, and endometrial cancers, and its overexpression is related to poor prognosis.^{5–8} Our prior studies have demonstrated that the mRNA levels of *ERRα* increase with the clinical stage of ovarian cancer, thus suggesting *ERRα* as a prognostic factor for ovarian cancer.⁷ *ERRα* also has potential diagnostic and therapeutic value in OC. *ERRα* activity highly relies on the presence of coregulatory proteins, most notably that of peroxisome proliferator-activated receptor gamma coactivator 1- α (*PPARGC1A*, also known as *PGC-1α*).^{9,10} The *PGC-1α/ERRα* axis has been implicated in controlling the expression of metabolic gene networks and mitochondrial biogenesis.^{11,12} Accumulating evidence suggests that the *PGC-1α/ERRα* axis plays a vital role in various cancer development and progression, which are accompanied by metabolic dysfunction.^{10,13} Our recent study indicated that *PGC-1α* and *ERRα* positively correlate with more advanced myometrial invasion in endometrial cancer, and were robust predictors for myometrial invasion.¹⁴

Currently, the suitability of *PGC-1α* and *ERRα* as biomarkers in OC tissues has not been thoroughly examined. To investigate whether the *PGC-1α/ERRα* axis acts as an effective prognostic marker for patients with OC, we detected the expression of these proteins in human OC tissues using immunohistochemistry and investigated their clinical significance in OC.

Patients and Methods

The HPA Database

First, we explored mRNA expression of *PGC-1α* and *ERRα* in normal tissues and ovarian cancer tissues from the HPA database (<https://www.proteinatlas.org/>). The HPA database provides abundant transcriptome and proteome data in specific human tissues through RNA-sequencing analysis and immunohistochemistry analysis.^{15,16} In addition, we analyzed the prognostic significance of *PGC-1α* and *ERRα* mRNA in patients with OC.

Patients and Tissue Samples

A total of 42 OC and 31 noncancer ovarian samples with related clinical data were obtained for immunohistochemistry from patients who received surgical therapy between September 2012 and April 2019. None of the patients received preoperative chemotherapy or radiotherapy. This study was approved by the Ethics Committee of Fujian Maternity and Child Health Hospital affiliated with Fujian

Medical University (approval number 2013-004), and performed in compliance with the Declaration of Helsinki. Informed consent was obtained from all patients. The patients were followed up postoperatively by their surgeons at 3-month intervals for 5 years and yearly thereafter. Overall survival (OS) was defined as the interval between surgery and mortality or the last follow-up (censored data for living patients). At the end of the follow-up period, seven patients (16.7%) died of OC.

Immunohistochemistry

To examine the expression of *PGC-1α* and *ERRα* in OC, we performed a tissue microarray constructed by Shanghai Zhuoli Biotechnology Co., Ltd (Zhuoli Biotechnology Co, Shanghai, China). Rabbit polyclonal anti-*ERRα* (ab93173, Abcam) and rabbit polyclonal anti-*PGC-1α* (ab191838, Abcam) antibodies were used. In each case, 1–2 μ m-thick sections from paraffin tissue blocks were cut, dewaxed, pretreated, and transferred to glass slides using an adhesive tape transfer system, in order to conduct ultraviolet cross linkage. All reactions were performed using an automated staining device. Two pathologists independently evaluated the quantitation of immunostaining for *PGC-1α* and *ERRα*, who were blinded to patient details. The expression of *PGC-1α* and *ERRα* in tumor parenchyma was semi-quantified by immunoreactivity score (IR score) based on intensity and heterogeneity. The IR score was determined as the sum of heterogeneity and intensity. The percentage of positive cells was scored as 0 points (0%), 1 point (1–25%), 2 points (26–50%), 3 points (51–75%), and 4 points (76–100%). Positive staining intensity was scored as 0 points (none), 1 point (low), 2 points (medium), and 3 points (high).

Statistical Analysis

Frequency and percentage were calculated for the classified variables. We used the Chi-square test to evaluate the differences in *PGC-1α* and *ERRα* expression between the two groups of clinicopathological features. The comparison between IR score in cancer foci and in noncancer lesions was analyzed by Student's *t*-test. Cancer-specific survival curves were obtained using the Kaplan–Meier method and verified by the log-rank (Mantel–Cox) test. All statistical analyses were performed using SPSS 19.0 software (SPSS Inc., Chicago, IL, USA) and GraphPad Prism Version 8.0 software (GraphPad Software, Inc., La Jolla, CA, USA). All *P*-values in the statistical analysis were two-tailed, and *P* < 0.05 was considered statistically significant.

Result

Expression and Prognostic Significance of PGC-1 α and ERR α mRNA in the HPA Database

The mRNA expression profiles retrieved from HPA (<https://www.proteinatlas.org/ENSG00000109819-PPARGC1A/tissue>, <https://www.proteinatlas.org/ENSG00000173153-ESRRA/tissue>) revealed low mRNA expression of PGC-1 α

and ERR α in normal ovarian tissues (Figure 1A and B). We then explored the association between mRNA expression of PGC-1 α and ERR α and survival outcome in OC from the HPA database (<https://www.proteinatlas.org/ENSG00000109819-PPARGC1A/pathology>, <https://www.proteinatlas.org/ENSG00000173153-ESRRA/pathology>). As shown in Figure 1C, the 5-year OS rates in patients with high and low ERR α expression were 25% and 34%, respectively. Patients with high ERR α expression had

Figure 1 Expression and prognostic significance of PGC-1 α and ERR α mRNA in HPA database. (**A** and **B**) PGC-1 α and ERR α mRNA expression in different normal human tissues. (**C**) Patients with high PGC-1 α mRNA expression tend to have a poorer overall rate compared with those with low PGC-1 α mRNA expression in OC ($P = 0.17$). (**D**) Patients with high ERR α mRNA expression exhibited a poorer overall rate compared with those with low ERR α mRNA expression in OC ($P = 0.035$). All the pictures were downloaded from HPA database.

Abbreviations: ERR α , estrogen-related receptor α ; HPA, Human Protein Atlas; OC, ovarian cancer; PGC-1 α , peroxisome proliferator-activated receptor gamma coactivator 1- α .

significantly lower OS rates than those with low *ERRα* expression ($P=0.035$; [Figure 1D](#)). The 5-year OS rates in patients with high and low *PGC-1α* expression were 27% and 39%, respectively. However, while patients with high *PGC-1α* expression tended to show poor cancer-specific survival rates, this association was not significant ($P = 0.17$; [Figure 1C](#)).

Expression of *PGC-1α* and *ERRα* in OC Tissues

Next, we determined the protein expression of *PGC-1α* and *ERRα* in 42 OC tissues and 31 noncancerous ovarian tissues by IHC. Statistical analysis showed that OC tissues exhibited significantly higher *PGC-1α* and *ERRα* expression than noncancerous ovarian tissues ($P=0.0059$, $P=0.002$; [Figure 2A](#) and [B](#)). As shown in [Figure 2C](#), *ERRα* was primarily detected in the nuclei of tumor cells, while *PGC-1α* was mainly detected in the cytoplasm. IHC also revealed that the expression of *PGC-1α* and *ERRα* was low in noncancerous tissues, but elevated in OC tissues.

Association Between *PGC-1α/ERRα* Expression and Clinical Parameters in OC

To investigate the clinical significance of *PGC-1α* and *ERRα* in ovarian cancer, we analyzed the association between expression of these proteins and clinical characteristics of 42 patients with OC ([Table 1](#)). Almost all tumor foci showed IR scores ≥ 3 and ≥ 4 for *PGC-1α* and *ERRα*, respectively ([Figure 2A](#) and [B](#)). Thus, we defined an IR score of 4 and 7 as the cut-off for high *PGC-1α* and *ERRα*, respectively, in order to identify a potential correlation between *PGC-1α/ERRα* expression and patient clinical characteristics. Significant associations were identified between high *ERRα* and *PGC-1α/ERRα* expression and tumor differentiation ($P=0.027$; $P=0.04$), lymph node status ($P=0.023$; $P=0.021$), CA125 ($P=0.036$; $P=0.021$), and HE4 ($P=0.021$; $P=0.05$). However, high *PGC-1α* expression was only significantly associated with tumor differentiation ($P=0.029$).

Prognostic Significance of *PGC-1α* and *ERRα* in OC

The Kaplan-Meier estimator model was employed to evaluate the prognostic significance of *PGC-1α* and *ERRα* expression ([Figure 3](#)). Seven (16.7%) patients died of ovarian cancer during the follow-up period. No significant

relationship was observed between expression of *PGC-1α* and *ERRα* and cancer-specific survival rate ($P=0.4736$, $P=0.3119$; [Figure 3A](#) and [B](#)). However, combined analysis of high *PGC-1α* and *ERRα* expression revealed a tendency towards poor cancer-specific survival ($P=0.1276$; [Figure 3C](#)).

Discussion

Increasing evidence has suggested that *ERRα* promotes tumor cell proliferation, angiogenesis, metastasis, and drug resistance in various cancers.^{10,17} Willy et al¹⁸ and Chisamore et al¹⁹ revealed that suppression of *ERRα* blocks the cell cycle in the G1/S transition, hindering the growth of breast tumor cell lines and their tumorigenicity in vivo. Epithelial-mesenchymal transition (EMT) plays an essential role in OC cell invasion and metastasis.^{20,21} Wang et al²² reported that suppression of *ERRα* expression inhibits tumor metastasis and migration in OC cell lines by restraining mitochondrial activity and preventing EMT in vitro. Moreover, in an orthotopic model of OC, Lam et al²³ found that inhibition of *ERRα* dramatically reduces tumor burden, ascites formation, and metastatic peritoneal nodules in vivo. Our recent research also demonstrated that inhibition of *ERRα* reduces EMT phenotypes, thereby significantly inhibiting invasion and migration in endometrial cancer cells.¹⁴ *ERRα* expression has also been associated with negative outcome in various human cancers. For example, in breast cancer, the mRNA and protein expression of *ERRα* positively correlates with node status, increased risk of recurrence, and metastatic status.^{24,25} Our previous studies have demonstrated that a high mRNA level of *ERRα* is associated with advanced FIGO stage and histological grade, and is thus associated with poor prognosis and shorter median overall survival time in OC.⁷

However, *ERRα* activity is primarily controlled by its coactivators, especially *PGC-1α*. The *PGC-1α/ERRα* axis also has been implicated in regulating several genes involved in energy metabolism, and increased mRNA and protein levels of *ERRα* in tissues are accompanied by high expression of *PGC-1α*.^{26,27} Expression of *PGC-1α* and *ERRα* is sensitive to physiological and pathological changes, which strengthens their crucial role in energy homeostasis in health and disease.⁹ Compounds affecting either the coactivator or the receptor could regulate signaling activity.^{9,10} Moreover, several studies indicate that *PGC-1α/ERRα* efficiently induce vascular endothelial growth factor (VEGF), and promote angiogenesis, mitochondrial biogenesis, and OXPHOS, leading to tumor angiogenesis, invasion, and metastasis.^{28–30}

Figure 2 The expression of PGC-1 α and ERR α in OC by IHC. (**A** and **B**) Ovarian tissues exhibited significantly higher PGC-1 α and ERR α expression compared with noncancer ovarian tissues ($P=0.0059$, $P=0.002$). (**C**) PGC-1 α and ERR α presented low expression in noncancer tissues while elevated PGC-1 α and ERR α expression was recorded in the OC tissues. Original magnification, x200.

Note: ** $p<0.01$.

Abbreviations: ERR α , estrogen-related receptor α ; IHC, Immunohistochemistry; IR score, immunoreactivity score; OC, ovarian cancer; PGC-1 α , peroxisome proliferator-activated receptor gamma coactivator 1- α .

Numerous studies have focused on the potential mechanism of action of PGC-1 α /ERR α ,^{31–33} and multiple potential novel agonists have been identified, such as genistein, apigenin, resveratrol, daidzein, flavone, and cholesterol.^{18,34–36} Our

latest research detected ERR α levels in uterine tumors by IHC and found that high expression of ERR α is associated with myometrial invasion. In addition, we identified a novel role for PGC-1 α and ERR α as positive regulators of EMT,

Table I Association Between PGC-1 α /ERR α Expression and Clinicopathological Parameters in Patients with Ovarian Cancer

Clinical Features	Total (n)	PGC-1 α Expression		χ^2	P	ERR α Expression		χ^2	P	PGC-1 α /ERR α Expression		χ^2	P
		High	Low			High	Low			High/High	Neither High		
Age, years <55 ≥55	42	17 12	8 5	0.032	0.859	15 11	10 6	0.095	0.758	11 7	14 10	0.033	0.856
Tumor size, cm <10 ≥10	41	21 8	5 7	3.459	0.063	17 9	9 6	0.119	0.73	13 5	13 10	1.073	0.3
Tumor differentiation Well, moderate Poor	37	6 18	8 5	4.786	0.029	6 19	8 4	6.275 ^a	0.027	3 14	11 9	5.451 ^a	0.04
FIGO stage I–II III–IV	41	14 15	5 7	0.149	0.699	10 15	9 7	1.036	0.309	7 11	12 11	0.717	0.397
Lymph node status Absent Present	32	13 9	8 2	1.332 ^a	0.425	10 10	11 1	5.772 ^a	0.023	5 8	16 3	7.161 ^a	0.021
CA125, U/mL <78 ≥78	42	8 21	5 8	0.497	0.481	5 21	8 8	4.388	0.036	2 16	11 13	5.802 ^a	0.021
CA153, U/mL <14.8 ≥14.8	38	5 20	6 7	2.844	0.092	5 19	6 8	2.085	0.149	2 14	9 13	3.635 ^a	0.078
HE4, U/mL <75.5 ≥75.5	34	6 19	3 6	0.296	0.67	2 17	7 8	5.625 ^a	0.025	1 13	8 12	4.568 ^a	0.05

Notes: ^aFisher test was performed. The rest of the scores were from Chi-square test.

Abbreviations: ERR α , estrogen-related receptor α ; PGC-1 α , peroxisome proliferator-activated receptor gamma coactivator 1- α .

and showed that disruption of PGC-1 α /ERR α signaling could reverse EMT and inhibit endometrial cancer invasion and migration.¹⁴

Currently, there are no reports in the literature on the analysis of PGC-1 α and ERR α expression in OC tissues by IHC. To verify the clinical significance of PGC-1 α /ERR α expression in OC, we first performed analyzed the mRNA levels of PGC-1 α and ERR α in OC tissues using data from the public database HPA. We found that patients with high ERR α mRNA expression had significantly lower OS rates compared with patients with low ERR α expression, which supported our prior observations.⁷ Similar trends were observed between PGC-1 α mRNA levels and patients with

OC in HPA. According to the central dogma of molecular biology,³⁷ the sequential information transfer residue-by-residue, thus mRNA expression correlates to protein expression. Consistent with these results, immunohistochemical staining indicated that PGC-1 α and ERR α were remarkably overexpressed in OC tissues rather than in non-cancer ovarian tissues. We further investigated the correlation between PGC-1 α and ERR α expression and the clinical features of patients with OC. High ERR α expression was significantly associated with tumor differentiation, lymph node status, CA125, and HE4, whereas high PGC-1 α expression only significantly correlated with tumor differentiation. Moreover, Kaplan-Meier survival analysis revealed that the

Figure 3 The prognostic significance of PGC-1 α and ERR α in OC. (A and B) No significant relation was observed between PGC-1 α and ERR α expression and the cancer-specific survival rate ($P=0.4736$, $P=0.3119$). (C) Combined analysis of high PGC-1 α and ERR α expression tended to show poor cancer-specific survival ($P=0.1276$).

Abbreviations: ERR α , estrogen-related receptor α ; OC, ovarian cancer; PGC-1 α , peroxisome proliferator-activated receptor gamma coactivator 1-alpha.

prognostic value of PGC-1 α and ERR α in OC was not significant. Interestingly, combined analyses of the expression of the two proteins enhanced their clinical significance in OC as compared with the analysis of their expression alone. Poor differentiation and positive lymph node predict poor outcomes in OC. CA125 and HE4 are crucial biomarkers in the diagnostic and therapeutic monitoring phase.³⁸ These data enhance the role of PGC-1 α and ERR α in the development and progression of ovarian cancer, as with what we observed in the experiments of cell lines and animals.^{22,23}

This study has several limitations. Firstly, this is a retrospective study and might have selection bias. Secondly, our current study was based on small sample size and lack of adequate samples for further validation of protein biomarkers. Thirdly, future studies are necessary to address the mechanism that regulates the development and progression of OC.

Conclusion

In conclusion, the present study showed the expression and clinical significance of PGC-1 α and ERR α in human OC. The combined analysis of PGC-1 α and ERR α expression could be a useful prognostic indicator of OC.

Abbreviations

ERR α , estrogen-related receptor alpha; HPA, Human Protein Atlas; IHC, Immunohistochemistry; OC, ovarian cancer; PGC-1 α , peroxisome proliferator-activated receptor gamma coactivator 1-alpha.

Acknowledgments

This study was supported in part by grant YCXZ18-01 from Fujian Provincial Maternity and Children's Health Hospital, China.

Disclosure

The authors report no conflicts of interest in this work.

References

- Kopper O, de Witte CJ, Löhmußsaar K, et al. An organoid platform for ovarian cancer captures intra- and interpatient heterogeneity. *Nat Med*. 2019;25(5):838–849. doi:10.1038/s41591-019-0422-6
- Crevet L, Vanacker J-M. Regulation of the expression of the estrogen related receptors (ERRs). *Cell Mol Life Sci*. 2020;77:4573–4579. doi:10.1007/s00018-020-03549-0
- Cai Q, Lin T, Kamarajugadda S, Lu J. Regulation of glycolysis and the Warburg effect by estrogen-related receptors. *Oncogene*. 2013;32(16):2079–2086. doi:10.1038/ncr.2012.221
- Tennessen JM, Baker KD, Lam G, Evans J, Thummel CS. The Drosophila estrogen-related receptor directs a metabolic switch that supports developmental growth. *Cell Metab*. 2011;13(2):139–148. doi:10.1016/j.cmet.2011.01.005
- Vernier M, McGuirk S, Dufour CR, et al. Inhibition of DNMT1 and ERR α crosstalk suppresses breast cancer via derepression of IRF4. *Oncogene*. 2020;39(41):6406–6420. doi:10.1038/s41388-020-01438-1
- Zhou S, Xia H, Xu H, et al. ERR α suppression enhances the cytotoxicity of the MEK inhibitor trametinib against colon cancer cells. *J Exp Clin Cancer Res*. 2018;37(1):218. doi:10.1186/s13046-018-0862-8
- Sun P, Sehoul J, Denkert C, et al. Expression of estrogen receptor-related receptors, a subfamily of orphan nuclear receptors, as new tumor biomarkers in ovarian cancer cells. *J Mol Med (Berl)*. 2005;83(6):457–467. doi:10.1007/s00109-005-0639-3
- Gao M, Sun P, Wang J, Zhao D, Wei L. Expression of estrogen receptor-related receptor isoforms and clinical significance in endometrial adenocarcinoma. *Int J Gynecol Cancer*. 2006;16(2):827–833. doi:10.1111/j.1525-1438.2006.00527.x
- Lynch C, Zhao J, Huang R, et al. Identification of estrogen-related receptor α agonists in the Tox21 compound library. *Endocrinology*. 2018;159(2):744–753. doi:10.1210/en.2017-00658
- Deblois G, St-Pierre J, Giguère V. The PGC-1/ERR signaling axis in cancer. *Oncogene*. 2013;32(30):3483–3490. doi:10.1038/ncr.2012.529
- Chang C-Y, McDonnell DP. Molecular pathways: the metabolic regulator estrogen-related receptor α as a therapeutic target in cancer. *Clin Cancer Res*. 2012;18(22):6089–6095. doi:10.1158/1078-0432.CCR-11-3221
- Villena JA, Kralli A. ERR α : a metabolic function for the oldest orphan. *Trends Endocrinol Metab*. 2008;19(8):269–276. doi:10.1016/j.tem.2008.07.005

13. Audet-Walsh É, Papadopoulos DJ, Gravel S-P, et al. The PGC-1 α /ERR α axis represses one-carbon metabolism and promotes sensitivity to anti-folate therapy in breast cancer. *Cell Rep*. 2016;14(4):920–931. doi:10.1016/j.celrep.2015.12.086
14. Chen L, Mao X, Huang M, Lei H, Xue L, Sun P. PGC-1 α and ERR α in patients with endometrial cancer: a translational study for predicting myometrial invasion. *Aging (Albany NY)*. 2020;12(17):16963–16980. doi:10.18632/aging.103611
15. Uhlen M, Fagerberg L, Hallström BM, et al. Proteomics. Tissue-based map of the human proteome. *Science*. 2015;347(6220):1260419. doi:10.1126/science.1260419
16. Pontén F, Jirstrom K, Uhlen M. The human protein atlas—a tool for pathology. *J Pathol*. 2008;216(4):387–393. doi:10.1002/path.2440
17. Liu G, Sun P, Dong B, Sehoul J. Key regulator of cellular metabolism, estrogen-related receptor α , a new therapeutic target in endocrine-related gynecological tumor. *Cancer Manag Res*. 2018;10:6887–6895. doi:10.2147/CMAR.S182466
18. Willy PJ, Murray IR, Qian J, et al. Regulation of PPAR γ coactivator 1 α (PGC-1 α) signaling by an estrogen-related receptor α (ERR α) ligand. *Proc Natl Acad Sci U S A*. 2004;101(24):8912–8917. doi:10.1073/pnas.0401420101
19. Chisamore MJ, Cunningham ME, Flores O, Wilkinson HA, Chen JD, Bauer JA. Characterization of a novel small molecule subtype specific estrogen-related receptor α antagonist in MCF-7 breast cancer cells. *PLoS One*. 2009;4(5):e5624. doi:10.1371/journal.pone.0005624
20. Vergara D, Merlot B, Lucot J-P, et al. Epithelial-mesenchymal transition in ovarian cancer. *Cancer Lett*. 2010;291(1):59–66. doi:10.1016/j.canlet.2009.09.017
21. Loret N, Denys H, Tummers P, Berx G. The role of epithelial-to-mesenchymal plasticity in ovarian cancer progression and therapy resistance. *Cancers (Basel)*. 2019;11(6):838. doi:10.3390/cancers11060838
22. Wang C-W, Hsu W-H, Tai C-J. Antimetastatic effects of cordycepin mediated by the inhibition of mitochondrial activity and estrogen-related receptor α in human ovarian carcinoma cells. *Oncotarget*. 2017;8(2):3049–3058. doi:10.18632/oncotarget.13829
23. Lam SS, Mak AS, Yam JW, Cheung AN, Ngan HY, Wong AS. Targeting estrogen-related receptor α inhibits epithelial-to-mesenchymal transition and stem cell properties of ovarian cancer cells. *Mol Ther*. 2014;22(4):743–751. doi:10.1038/mt.2014.1
24. Zhao Y, Li Y, Lou G, et al. MiR-137 targets estrogen-related receptor α and impairs the proliferative and migratory capacity of breast cancer cells. *PLoS One*. 2012;7(6):e39102. doi:10.1371/journal.pone.0039102
25. Stein RA, Gaillard S, McDonnell DP. Estrogen-related receptor α induces the expression of vascular endothelial growth factor in breast cancer cells. *J Steroid Biochem Mol Biol*. 2009;114(1–2):106–112. doi:10.1016/j.jsbmb.2009.02.010
26. Deblois G, Giguère V. Functional and physiological genomics of estrogen-related receptors (ERRs) in health and disease. *Biochim Biophys Acta*. 2011;1812(8):1032–1040. doi:10.1016/j.bbdis.2010.12.009
27. Eichner LJ, Giguère V. Estrogen related receptors (ERRs): a new dawn in transcriptional control of mitochondrial gene networks. *Mitochondrion*. 2011;11(4):544–552. doi:10.1016/j.mito.2011.03.121
28. LeBleu VS, O'Connell JT, Gonzalez Herrera KN, et al. PGC-1 α mediates mitochondrial biogenesis and oxidative phosphorylation in cancer cells to promote metastasis. *Nat Cell Biol*. 2014;16(10). doi:10.1038/ncb3039
29. Arany Z, Foo S-Y, Ma Y, et al. HIF-independent regulation of VEGF and angiogenesis by the transcriptional coactivator PGC-1 α . *Nature*. 2008;451(7181):1008–1012. doi:10.1038/nature06613
30. Dwyer MA, Joseph JD, Wade HE, et al. WNT11 expression is induced by estrogen-related receptor α and β -catenin and acts in an autocrine manner to increase cancer cell migration. *Cancer Res*. 2010;70(22):9298–9308. doi:10.1158/0008-5472.CAN-10-0226
31. Yan M, Audet-Walsh É, Manteghi S, et al. Chronic AMPK activation via loss of FLCN induces functional beige adipose tissue through PGC-1 α /ERR α . *Genes Dev*. 2016;30(9):1034–1046. doi:10.1101/gad.281410.116
32. Choi YK, Park JH, Baek -Y-Y, et al. Carbon monoxide stimulates astrocytic mitochondrial biogenesis via L-type Ca channel-mediated PGC-1 α /ERR α activation. *Biochem Biophys Res Commun*. 2016;479(2):297–304. doi:10.1016/j.bbrc.2016.09.063
33. Craige SM, Kröller-Schön S, Li C, et al. PGC-1 α dictates endothelial function through regulation of eNOS expression. *Sci Rep*. 2016;6:38210. doi:10.1038/srep38210
34. Teng CT, Hsieh J-H, Zhao J, et al. Development of novel cell lines for high-throughput screening to detect estrogen-related receptor α modulators. *SLAS Discov*. 2017;22(6):720–731. doi:10.1177/2472555216689772
35. Wei W, Schwaid AG, Wang X, et al. Ligand activation of ERR α by cholesterol mediates statin and bisphosphonate effects. *Cell Metab*. 2016;23(3):479–491. doi:10.1016/j.cmet.2015.12.010
36. Teng CT, Beames B, Alex Merrick B, Martin N, Romeo C, Jetten AM. Development of a stable cell line with an intact PGC-1 α /ERR α axis for screening environmental chemicals. *Biochem Biophys Res Commun*. 2014;444(2):177–181. doi:10.1016/j.bbrc.2014.01.033
37. Strasser BJ. A world in one dimension: Linus Pauling, Francis Crick and the central dogma of molecular biology. *Hist Philos Life Sci*. 2006;28(4):491–512.
38. Potenza E, Parpinel G, Laudani ME, Macchi C, Fuso L, Zola P. Prognostic and predictive value of combined HE-4 and CA-125 biomarkers during chemotherapy in patients with epithelial ovarian cancer. *Int J Biol Markers*. 2020;1724600820955195. doi:10.1177/1724600820955195

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic

agents and protocols on patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/oncotargets-and-therapy-journal>

Dovepress