Characteristics, Preferences and Health Care Utilization in Patients Using a Dietary Supplement for Improving Sleeping Disturbances: Results from an Explorative Online Survey

This article was published in the following Dove Press journal:
Patient Preference and Adherence

Stefania Di Gangi

Stefan Markun

Thomas Rosemann

Andreas Plate

Institute of Primary Care, University and
University Hospital of Zürich, Zürich,
Switzerland

Purpose: Sleeping disturbances are highly prevalent in the general population, and pharmacological drug treatment harbours the risk of serious side effects. Many affected persons use dietary supplements for self-treating their symptoms, but little is known about the specific characteristics and preferences of these patients. Even less evidence exists about the consequences of a specific dietary supplement usage on health care utilization. The aim of this study was to explore characteristics, preferences and the impact on health care utilization in patients using a specific over-the-counter dietary supplement, which is promoted for improving sleeping disturbances.

Patients and Methods: We conducted a structured survey and invited a sample of 297 customers of a specific dietary supplement to participate. The survey was open between June and September 2020. Participants were invited by email. All participants accepted an informed consent.

Results: A total of 127 customers participated in the survey (participation rate: 42.8%). Of them, 87.7% were female and the mean age 50.5 years. Participants reported an overall good improvement of symptoms (7.66 on a ten-point Likert-scale) and showed a strong belief in the effectiveness of supplements in general; 67% of participants reported that the intake led to fewer physician encounters, and 48.3% reported that they could stop the intake of other pharmaceutical sleeping drugs.

Conclusion: A significant proportion of participants reported a substantial reduction in pharmaceutical drug use and health care utilization. While these self-reports lack an adequate control, they are still real consumer experiences, and the large beneficial effects – whether placebo or not – explain the popularity of such supplements and their therapeutic potential in sleeping disorders.

Keywords: supplement use, sleeping disorders, usage patterns, drugs, health care utilization

Introduction

The use of dietary supplements is common, and it is estimated that up to three-quarters of adults use dietary supplements.^{1,2} In addition to being used to achieve a healthier lifestyle, dietary supplements are used as remedies for a variety of complaints and unspecific symptoms. Sleeping disorders are highly prevalent, and insufficient sleeping quality/quantity is associated with a wide range of negative health outcomes.³ Self-treatment with herbals, over-the-counter medication or

Correspondence: Stefania Di Gangi
Institute of Primary Care, University of
Zurich, Pestalozzistrasse 24, Zürich 8091,
Switzerland
Tel +41 44 255 75 02
Fax +41 44 255 90 97
Email stefania.digangi@usz.ch

prescribed medications is common.⁴ Classical pharmacological treatments of sleeping disorders (for example benzodiazepines or hypnotics) are associated with serious side effects and the risk for abuse or dependence.⁵ Consequently, a growing demand and offer of dietary supplements tackling sleeping disorders has been observed for many years. There is some evidence¹ that dietary supplement users, aiming for a healthier lifestyle, are more frequently females with higher education and higher incomes compared to non-users. However, little is known about the characteristics, attitudes and beliefs of dietary supplements users who use the dietary supplements for sleeping disorders. In addition, there is no evidence on the impact of self-treatment on patients' health care utilization, as sleeping disorders are a common reason for physician encounters.^{6–8} To gain a more detailed insight into this topic, we conducted an explorative survey among dietary supplement users using a specific over-the-counter dietary supplement, which is promoted for improving sleeping disturbances. The goal of the present explorative study was to determine characteristics, preferences and the impact of use on their health care utilization.

Patients and Methods

Methods

Data for this study were obtained by an anonymous online survey. The survey was conducted in Switzerland, and 297 customers of Sero-Night,⁹ a dietary supplement advertised for improving sleeping quality, were invited by email to participate in the survey. The dietary supplement's main ingredients were L-tryptophan, ashwagandha extract, inositol, L-methionin as well as vitamins B 5/6/12. The survey was open for 11 weeks beginning from mid-June 2020, and customers received up to two reminders by email. We used a commercial survey software (Survey Monkey Inc., San Mateo, California, USA). For subgroup analysis, we used adaptive questioning depending on the previous answers of the customers. The survey had a total length of 13 screens, with a mean of 2.8 questions on each screen. All participants were able to review their answers. Software's function was used to prevent multiple entries from the same device. The survey was fully anonymous, and no questions were mandatory. Questions were closed-ended (ie education, field of employment), open-ended (ie type of medication, further supplements), multiple choice (ie reasons for intake) and Likert-scale (ie overall satisfaction). A checklist for reporting results of internet e-surveys

(CHERRIES)¹⁰ is provided in the supplemental material ([Supplemental Table 1](#)).

Statistics

Results are presented as means (standard deviations [SD]) for continuous variables and as number *N* (%), or only %, for categorical variables. The range (min–max) was reported for age and all variables defined in Likert-scale points. When necessary, we reported the missing values. In case of incomplete questionnaires, we analysed only the available data, without imputation. All statistical analyses were carried out using statistical package R, R Core Team (2016). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

Ethics

Fully anonymous surveys do not fall under the scope of the Swiss national Human Research Act, and thus no ethics approval is necessary to perform health surveys. Participation to the survey was voluntary, and each participant signed an informed consent, which contained all information about the purpose of the study. All participants agreed to a scientific analysis and the publication of the data.

Results

A total of 127 customers participated in the survey, and the participation rate was 42.8% (127/297). Basic demographic and health data of the participants are reported in [Table 1](#): 87.7% of participants were female, and the mean age was 50.5 (SD 10.3) years (range: 26–76 years). Furthermore, 33% of the participants had an education level equal to or higher than a Master degree, and 60% had an education level equal to or higher than a Bachelor degree. The most common fields of employment were economy, management (27%), education, social issues (23.5%) and health (17.4%). Mean self-reported wellbeing, on a ten-point Likert-scale, was 7.15 (SD 1.5). Active smoking was reported by 4.3% of participants, and one-fifth of participants reported no additional comorbidities. The most common reported comorbidities were sleeping disorders (31.5%) and postmenopausal complaints (24.4%). Participants reported a mean of 3.5 (SD 4.8) physician consultations within the previous twelve months.

The specific user preferences of supplement use are reported in [Table 2](#). The overall satisfaction of supplements, on a ten-point Likert-scale, was on average 7.59

Table 1 Basic Demographic and Health Data of 127 Participants

Variables	Results	Answer Available, N (%)
Demographic data		
Age (mean, SD), range	50.47 (10.33), 26–76	N=122 (96.1)
Female sex	107 (87.7)	N=122 (96.1)
Highest educational level		N=127 (100)
≤ 9 years of school education	2 (1.6)	
≤ 13 years of school education	4 (3.1)	
Apprenticeship certificate	35 (27.6)	
Bachelor degree	34 (26.8)	
Master degree	37 (29.1)	
PhD or higher degree	5 (3.9)	
No answer	10 (7.9)	
Field of employment		N=115 (90.5)
Nature	2 (1.7)	
Hospitality industry	3 (2.6)	
Aesthetics, sports	5 (4.3)	
Design, arts	5 (4.3)	
Printing	1 (0.9)	
Construction	1 (0.9)	
Wood, interior construction	1 (0.9)	
Motor vehicles	1 (0.9)	
Electrical engineering	2 (1.7)	
Metal, engineering	1 (0.9)	
Development, construction	1 (0.9)	
Sales	6 (5.2)	
Economy, management	31 (27.0)	
Traffic, logistics	3 (2.6)	
Computer science	2 (1.7)	
Culture	3 (2.6)	
Health	20 (17.4)	
Education, social issues	27 (23.5)	
Health data		
General wellbeing*, mean (SD), range	7.15 (1.5), 3–10	N=123 (96.7)
Smoking	5 (4.3)	N=116 (91.3)
Comorbidities:		N=127 (100)
None	28 (22.0)	
Hypertension	10 (7.9)	
Dyslipidaemia	9 (7.1)	
Diabetes mellitus	1 (0.8)	
Depression	11 (8.7)	
Sleeping disorders	40 (31.5)	
Postmenopausal complaints	31 (24.4)	
Other	42 (33.1)	

(Continued)

Table 1 (Continued).

Variables	Results	Answer Available, N (%)
Medical consultations in the last 12 months		
Any physician ⁺ consultation	114	N=122 (96.1)
Mean number of consultations	3.53 (4.81)	
Any medical consultation	119	N=122 (96.1)
Mean number of consultations	11.76 (14.95)	
Type of consultation, mean number of visits (SD), range		Number of participants
Family physician	2.04 (2.26), 0–12	N=89
Other physician/specialist	2.59 (4.53), 0–45	N=103
Alternative practitioners	2.16 (3.89), 0–20	N=51
Traditional Chinese Medicine (TCM)	2.54 (4.16), 0–16	N=50
Physiotherapy, ergotherapy	7.60 (11.10), 0–40	N=60
Other	6.62 (8.78), 0–35	N=53

Notes: Results are presented as absolute numbers and percentages (in parenthesis) unless else stated. *Likert scale: 0 (worst health status) to 10 (best health status). ⁺Including categories visiting “family physician” and “other physician/specialist”.

Abbreviations: N, number of participants; SD, standard deviation.

(SD 2.9). Participants reported a mean duration of intake of 13.5 (SD 10.3) months with a mean of 5.9 (SD 1.8) intakes per week. Specialists or physicians different from family doctors generally recommended the supplements, in nearly 80% of the cases. More than 80% of the participants used the supplement in order to optimize their sleeping patterns. However, mood disorders (mood swings or depressive mood) were common additional reasons for the supplement use in up to one-third of patients. Regarding the improvement of symptoms, the mean satisfaction score, on a ten-point Likert-scale, was 7.66 (SD 2.1). Just over 70% of the participants who used the supplement to optimize their sleeping patterns reported improvement in staying asleep during the night. Generally, more than 81% of the participants did not experience side effects during the use of supplements. Almost 90% of all participants reported the use of a mean of 3.35 (SD 1.9) further supplements. Participants reported the intake of various

Table 2 Use of Supplements: Specific Characteristics and Preferences

Question	Results	Answer Available (%)
Overall satisfaction, mean (SD), range Very unsatisfied (score 0) Almost unsatisfied (score 1–5) Almost satisfied (score 6–9) Very satisfied (score 10)	7.59 (2.94), 0–10 9 (7.2) 12 (9.6) 60 (48.0) 44 (35.2)	N=125 (98.4)
Time of intake (months), mean (SD), range	13.53 (10.26), 1–48	N=127 (100)
Source of supply Family physician Other physician/specialist Online shop Other	17 (13.4) 56 (44.1) 84 (66.1) 4 (3.1)	N=127 (100)
Number of purchased packages with the last year, mean (SD), range	3.13 (1.97), 0–9	N=125 (98.4)
Initial recommendation by Family physician Other physician/specialist Friends, relatives Own research on the topic Advertising	15 (11.8) 100 (78.7) 10 (7.9) 1 (0.8) 1 (0.8)	N=127 (100)
Weekly intake (in days), mean (SD), range	5.91 (1.79), 0–7	N=125 (98.4)
Reasons for intake Optimizing sleeping patterns (fall asleep, sleeping the whole night) Reduce intake of other sleeping pills Performance increase (due to better sleeping) Reducing postmenopausal complaints Improvement of memory Mood swings Depressive mood Other	105 (82.7) 15 (11.8) 39 (30.7) 28 (22.0) 17 (13.4) 42 (33.1) 29 (22.8) 8 (6.3)	N=127 (100)
Improvement of symptoms (in general), mean (SD), range	7.66 (2.08), 0–10	N=126 (99.2)
Improvement of symptoms (refers only to “Optimizing sleeping patterns”) I can fall asleep faster I can stay asleep during the night I can sleep longer I need fewer sleeping pills Sleeping quality improves I have more power during the day I fell less tired during the day General mood improved Feeling less tight Feeling less anxious Other	42 (40.0) 74 (70.5) 15 (14.3) 14 (13.3) 54 (51.4) 36 (34.3) 27 (25.7) 34 (32.4) 21 (20.0) 6 (5.7) 8 (7.6)	N=105 (100)
Experience of side effects? No Dizziness Nausea, vomiting Diarrhoea	104 (81.9) 4 (3.1) – 2 (1.6)	N=127 (100)

(Continued)

Table 2 (Continued).

Question	Results	Answer Available (%)
Tiredness	9 (7.1)	
Increasing weight	2 (1.6)	
Other	7 (5.5)	
Use of further supplements		
Concomitant use of other supplements	111 (89.5)	N=124 (97.6)
Number of further supplements, mean (SD), range	3.35 (1.86), 1–10	N=111 (100)
Years of intake, mean (SD), range	6.92 (7.55), 0–47	N=111 (100)
Weekly intake (in days), mean (SD), range	6.38 (1.29), 1–7	N=111 (100)
Type of supplements		N=111 (100)
Multivitamins	41 (36.9)	
Vitamin D	77 (69.4)	
Vitamin B12	45 (40.5)	
Other*	80 (72.1)	
Reason for concomitant use		N=111 (100)
Improvement of general health	72 (64.9)	
Strengthening of bones/joints	24 (21.6)	
Compensation for bad eating habits	21 (18.9)	
Improvement of the cardiovascular system	9 (8.1)	
Improvement of immune system	38 (34.2)	
Improvement of hair, nails or skin	21 (18.9)	
Regulation of gastrointestinal complaints	22 (19.8)	
Improvement of mental health	36 (32.4)	
Improvement of concentration	38 (34.2)	
Improvement of tiredness	50 (45.0)	
Other	19 (17.1)	
Did further household members use supplements?	27 (21.4)	N=126 (99.2)
Number of household members using supplements?		N=27 (100)
1	22 (81.5)	
2	5 (18.5)	
Kind of supplements		N=27 (100)
Same dietary supplement	9 (24.3)	
Others	28 (75.7)	

Notes: Results are presented as absolute numbers and percentages (in parenthesis) unless else stated. *This group consists of dietary products with various combinations of vitamins, electrolytes, herbals and amino acids.

Abbreviations: N, number of patients; SD, standard deviation.

combinations of vitamins, electrolytes, herbals and amino acids, with vitamin D the most commonly reported single supplement.

Results about the effects of supplements on co-mediations and health care utilization are reported in [Table 3](#). A total of 87.1% of participants reported a previous health care utilization because of index symptoms in the past, and more than two-third of them (67%)

reported that the use of the supplement led to reduced physician consultations. Over one-half of participants reported the use of other medications (for example pharmaceutical drugs [56.4%] or herbals [49%]), with antidepressants, hypnotics and melatonin the most frequently used drugs and sleeping pills the most frequently used herbal medication ([supplemental Table 2](#)). In the subgroup of participants with other medications, 84.5% reported that

the use of the supplement led to a stop or decrease in the uptake of additional medications (Table 3).

Participants' attitudes and beliefs regarding the effectiveness of dietary supplements for various health conditions are presented in Figure 1. Whereas nearly half of the participants (47.5%) disagreed with the statement that dietary supplements are a possibility to compensate for bad living habits in general (answer one [red] and two [yellow] on a five-point Likert-scale), most of the participants reported that they believed in the effectiveness of dietary supplements for various health complaints. An exception was cardiovascular health, where belief in the effectiveness of dietary supplements was considerably low.

Discussion

In this study, we investigated characteristics, beliefs and preferences of supplement users taking a specific dietary supplement to improve sleeping quality. We found that users were predominantly highly educated females, experienced in supplement use and that the intake of the dietary supplement led to self-perceived reduced health care utilization and reduced consumption of pharmaceutical drugs.

Sleeping disorders are a common complaint in developed countries,³ and inadequate sleeping quality is associated with various adverse outcomes (ie depression,¹¹ risk of falling,¹² diabetes mellitus¹³). Several studies reported

Table 3 Influence of Supplement Use on Co-Medications and Health Care Utilization

Question	N (%)	Answer Available (%)
Previous physician consultation because of symptoms?	108 (87.1)	N=124 (97.6)
Did the intake lead to fewer physician consultations?	61 (67.0)	N=91 (84.3)
Was there an intake of other medication for the symptoms?	58 (47.5)	N=122 (96.1)
Pharmaceutical drugs	31 (56.4)	N=55 (94.8)
Herbals	27 (49.0)	
Amino acids	2 (3.6)	
Electrolytes	3 (5.4)	
Other	4 (7.3)	
Influence of starting SKE Vital on other medications		N=58 (100)
Stopped other medications	28 (48.3)	
Reduced intake of other medications	21 (36.2)	
No change in co medications	8 (13.8)	
Other	1 (1.7)	

the association between sleeping disorders and health care utilization,^{14–16} but the impact of dietary supplements on improving sleeping disorders is unknown. For some

Figure 1 Attitudes and beliefs in effectiveness of dietary supplements. % of participants, for each point Likert-scale are shown. **(A)**: Results from a five point Likert-scale (pink: totally agree to red: totally disagree). Bad habits: Dietary supplements as a compensation for bad life habits. **(B)**: Results from a five point Likert-scale (pink: very effective to red: not effective). General: Improvement of general health; Bones/joints: Strengthening of bones/joints; Bad eating: compensation for bad eating habits; CVS: Improvement of the cardiovascular system; Immune system: improvement of the immune system; Hair/nails: improvement of hair, nails or skin; Gastrointestinal: regulation of gastrointestinal complaints; Mental: improvement of mental health; Concentration: improvement of concentration; Tiredness: improvement of tiredness.

ingredients of the studied dietary supplement, there is little/moderate evidence of a clinical effect in patients with sleeping disorders: the essential amino acid L-tryptophan, which is a precursor of the endogenous melatonin and serotonin synthesis, is often used to treat insomnia and depression,^{17,18} and sparse evidence suggests that ashwagandha extract can improve insomnia and anxiety.^{19,20} However, there are no comparative studies reported in the literature, and general transferability of study results is hardly feasible because ingredients and composition of the studied dietary supplements differ.

Basic Characteristics

Basic characteristics of our participants were similar to the characteristics reported in larger scale population-based surveys on supplement use.^{1,21} Most participants were highly educated females, and the overall wellbeing was high. With the exception of cardiovascular health, participants reported strong beliefs in the effectiveness of dietary supplements on various health conditions. Therefore, the majority of participants reported a long experience in supplement use, taking more than one supplement almost every day. Moreover, the low proportion of smokers in our patient group (4.3% vs 27.1% in the general population²²) could reveal that the use of supplements is perceived as a way to help a healthier lifestyle.¹

Health Care Utilization

Sleeping disorders are associated with health care utilization^{14–16} and are a common reason for physician encounters.^{6–8} Two large population-based surveys from Canada⁴ and Australia²³ reported that between 11% and 13% of the population sought medical care because of sleeping disturbances. In patients diagnosed with an insomnia disorder, the proportion was up to 40%.⁴ The characteristics of our participants resembled the patterns described in the literature. Almost 90% of all participants reported a physician consultation because of symptoms, but the mean consultation numbers were close to the mean consultation numbers in the overall population (3.5 vs 4.3 per year²⁴). However, two-thirds of participants in our study reported that the use of dietary supplement reduced the physician consultations. Available, sparse evidence showed no positive impact of supplement use in general and on health care costs,²⁵ but its association in specific users is still unknown. Though we reported explorative data from a highly selected population of experienced supplement users, given the high prevalence

of sleeping disorders, a relative reduction of consultations in two-third of patients, or in a subgroup of patients, could have an economic impact, in terms of health cost reduction, which should be confirmed in future studies.

Self-Medication of Sleeping Disorders

We found a self-perceived reduction in sleeping disorder-specific drug consumption. Sleeping disorders are a common indication for self-medication.^{26,27} Herbals and pharmaceutical drugs are the most commonly used medications, and their use differs internationally and depends on particular types of sleeping disorder.^{4,5} Accordingly, the participants of our survey reported the use of various herbals and pharmaceutical drugs. Major issues with pharmaceutical drugs are potential serious side effects, abuse or dependence.⁵ In addition, evidence on effectiveness of some commonly used drugs (for example antihistamines or melatonin) is limited.^{28,29} A reduction of these drugs would be beneficial as a kind of harm reduction. On the other hand, the consequences of excessive dietary supplement use is unknown,³⁰ and still dietary supplements harbour a significant risk of serious side effects.³¹ Especially if multiple supplements are used, the recommended daily intake is often surpassed.³² In our patient group, half of the participants reported the use of other medication for improving symptoms before starting the supplement. In the subgroup of pharmaceutical drug users, the majority of participants reported to either have stopped or reduced the drug after starting the dietary supplement. We cannot know if a real effect based on the ingredients or a placebo effect of the supplement use led to a stop or reduction of the pharmaceutical drugs, or whether these patients would have stopped the drugs by themselves. However, taking into account that only 22% of this subgroup started the dietary supplement with the intention to reduce the intake of their other sleeping medications, there might be a beneficial effect, which should be confirmed and evaluated in further trials too.

Strengths and Limitations

Our study has some limitations. First, we reported results from a survey among a specific group of dietary supplement users, and second, our study lacked a control group. Therefore, it must be acknowledged that our results merely represent beliefs of this specific population, and our results must not be misunderstood for specific effects of the supplement. However, our findings highlight the beliefs

of users and their perceptions of the supplement despite the paucity of supporting evidence available. Given the high prevalence of sleeping disorders in the general population there is a potential for relevant supplement use of unknown benefit, costs or even harm that needs to be better understood. In addition, given the fact that millions of people use supplements, it is of medical importance to understand patients' attitudes, beliefs and expectations. Lastly, we are aware about a potential non-response bias: those with positive experiences associated with supplement intake may have been more likely to respond. However, available evidence suggested that the impact of the non-response bias in satisfaction surveys is limited.^{33,34}

Conclusion

A significant proportion of participants reported a substantial reduction in pharmaceutical drug use and health care utilization. While these self-reports lack an adequate control, they are still real consumer experiences and beliefs. The large beneficial effects – whether placebo effect or not – explain the popularity of such supplements and their therapeutic potential in sleeping disorders.

Funding

The study was sponsored by SKE Vital. The sponsor had no influence on conceptualization of the study and analysing and interpreting of the obtained data.

Disclosure

The authors report no conflicts of interest in this work.

References

- Dickinson A, MacKay D. Health habits and other characteristics of dietary supplement users: a review. *Nutr J*. 2014;13(1):14. doi:10.1186/1475-2891-13-14
- Knudsen VK, Rasmussen LB, Haraldsdóttir J, et al. Use of dietary supplements in Denmark is associated with health and former smoking. *Public Health Nutr*. 2002;5(3):463–468. doi:10.1079/PHN2001276
- Chattu VK, Manzar MD, Kumary S, Burman D, Spence DW, Pandi-Perumal SR. The Global Problem of Insufficient Sleep and Its Serious Public Health Implications. *Healthcare*. 2018;7:1.
- Morin CM, LeBlanc M, Daley M, Gregoire JP, Mérette C. Epidemiology of insomnia: prevalence, self-help treatments, consultations, and determinants of help-seeking behaviors. *Sleep Med*. 2006;7(2):123–130. doi:10.1016/j.sleep.2005.08.008
- Lie JD, Tu KN, Shen DD, Wong BM. Pharmacological Treatment of Insomnia. *Peer Rev J Formulary Manage*. 2015;40(11):759–771.
- Fukushi M, Ishibashi Y, Nago N. Final diagnoses and probability of new reason-for-encounter at an urban clinic in Japan: A 4-year observational study. *J Clin Med*. 2017;96(22):e6999.
- Liu Y, Chen C, Jin G, et al. Reasons for encounter and health problems managed by general practitioners in the rural areas of Beijing, China: A cross-sectional study. *PLoS One*. 2017;12(12):e0190036. doi:10.1371/journal.pone.0190036
- Senthilvel E, Auckley D, Dasarathy J. Evaluation of sleep disorders in the primary care setting: history taking compared to questionnaires. *J Clin Sleep Med*. 2011;7(1):41–48.
- Vital SKE. 2020; <https://www.ske-vital.ch/produktseite/sero-night>. Accessed September 2020.
- Eysenbach G. Improving the quality of Web surveys: the Checklist for Reporting Results of Internet E-Surveys (CHERRIES). *J Med Internet Res*. 2004;6(3):e34. doi:10.2196/jmir.6.3.e34
- Morphy H, Dunn KM, Lewis M, Boardman HF, Croft PR. Epidemiology of insomnia: a longitudinal study in a UK population. *Sleep*. 2007;30(3):274–280.
- Stone KL, Ancoli-Israel S, Blackwell T, et al. Actigraphy-measured sleep characteristics and risk of falls in older women. *Arch Intern Med*. 2008;168(16):1768–1775. doi:10.1001/archinte.168.16.1768
- Gottlieb DJ, Punjabi NM, Newman AB, et al. Association of sleep time with diabetes mellitus and impaired glucose tolerance. *Arch Intern Med*. 2005;165(8):863–867. doi:10.1001/archinte.165.8.863
- Kaufmann CN, Canham SL, Mojtabai R, et al. Insomnia and health services utilization in middle-aged and older adults: results from the Health and Retirement Study. *J Gerontol A Biol Sci Med Sci*. 2013;68(12):1512–1517. doi:10.1093/gerona/glt050
- Kapur VK, Redline S, Nieto FJ, Young TB, Newman AB, Henderson JA. The relationship between chronically disrupted sleep and healthcare use. *Sleep*. 2002;25(3):289–296.
- Novak M, Mucsi I, Shapiro CM, Rethelyi J, Kopp MS. Increased utilization of health services by insomniacs—an epidemiological perspective. *J Psychosom Res*. 2004;56(5):527–536. doi:10.1016/j.jpsychores.2004.02.007
- Lieberman HR, Agarwal S, Fulgoni VL. Tryptophan Intake in the US Adult Population Is Not Related to Liver or Kidney Function but Is Associated with Depression and Sleep Outcomes. *J Nutr*. 2016;146(12):2609S–2615S. doi:10.3945/jn.115.226969
- Simper T, Gilmartin M, Allwood D, Taylor L, Chappell A. The effects of a sleep/recovery supplement: 'Night Time Recharge' on sleep parameters in young adults. *Nutr Health*. 2019;25(4):265–274. doi:10.1177/0260106019875911
- Langade D, Kanchi S, Salve J, Debnath K, Efficacy AD. Safety of Ashwagandha (*Withania somnifera*) Root Extract in Insomnia and Anxiety: A Double-blind, Randomized, Placebo-controlled Study. *Cureus*. 2019;11(9):e5797.
- Lopresti AL, Smith SJ, Malvi H, Kodgule R. An investigation into the stress-relieving and pharmacological actions of an ashwagandha (*Withania somnifera*) extract: A randomized, double-blind, placebo-controlled study. *Medicine*. 2019;98(37):e17186. doi:10.1097/MD.00000000000017186
- Satia-Abouta J, Kristal AR, Patterson RE, Littman AJ, Stratton KL, White E. Dietary supplement use and medical conditions: the VITAL study. *Am J Prev Med*. 2003;24(1):43–51. doi:10.1016/S0749-3797(02)00571-8
- Zahlen BAG & Fakten: tabak. 2020; <https://www.bag.admin.ch/bag/de/home/zahlen-und-statistiken/zahlen-fakten-zu-sucht/zahlen-fakten-zu-tabak.html>. Accessed September 2020.
- Bartlett DJ, Marshall NS, Williams A, Grunstein RR. Predictors of primary medical care consultation for sleep disorders. *Sleep Med*. 2008;9(8):857–864. doi:10.1016/j.sleep.2007.09.002
- Schweizerische Gesundheitsbefragung 2017. 2018; <https://www.bfs.admin.ch/bfs/de/home/statistiken/gesundheit/erhebungen/sgb.assetdetail.6426300.html>. Accessed September 2020.
- Kilonzo MM, Vale LD, Cook JA, Milne AC, Stephen AI, Avenell A. A cost-utility analysis of multivitamin and multimineral supplements in men and women aged 65 years and over. *Clin nutrition*. 2007;26(3):364–370. doi:10.1016/j.clnu.2006.11.002

26. Shehnaz SI, Agarwal AK, Khan N. A systematic review of self-medication practices among adolescents. *J Adolescent Health*. 2014;55(4):467–483. doi:10.1016/j.jadohealth.2014.07.001
27. Lund HG, Reider BD, Whiting AB, Prichard JR. Sleep patterns and predictors of disturbed sleep in a large population of college students. *J Adolescent Health*. 2010;46(2):124–132. doi:10.1016/j.jadohealth.2009.06.016
28. Ramar K, Olson EJ. Management of common sleep disorders. *Am Fam Physician*. 2013;88(4):231–238.
29. Brzezinski A, Vangel MG, Wurtman RJ, et al. Effects of exogenous melatonin on sleep: a meta-analysis. *Sleep Med Rev*. 2005;9(1):41–50. doi:10.1016/j.smrv.2004.06.004
30. Chen F, Du M, Blumberg JB, et al. Association among dietary supplement use, nutrient intake, and mortality among u.s. adults: a cohort study. *Ann Intern Med*. 2019;170(9):604–613. doi:10.7326/M18-2478
31. Geller AI, Shehab N, Weidle NJ, et al. Emergency department visits for adverse events related to dietary supplements. *N Engl J Med*. 2015;373(16):1531–1540. doi:10.1056/NEJMsa1504267
32. Panel NS-o-tS. National institutes of health state-of-the-science conference statement: multivitamin/mineral supplements and chronic disease prevention. *Am J Clin Nutr*. 2007;85(1):257S–264S. doi:10.1093/ajcn/85.1.257S
33. Lasek RJ, Barkley W, Harper DL, Rosenthal GE. An evaluation of the impact of nonresponse bias on patient satisfaction surveys. *J Int Med*. 1997;35(6):646–652.
34. Perneger TV, Chamot E, Bovier PA. Nonresponse bias in a survey of patient perceptions of hospital care. *Med Care*. 2005;43(4):374–380. doi:10.1097/01.mlr.0000156856.36901.40

Patient Preference and Adherence

Dovepress

Publish your work in this journal

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focusing on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize clinical outcomes for existing disease

states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/patient-preference-and-adherence-journal>