

IgA and FcαRI: Versatile Players in Homeostasis, Infection, and Autoimmunity

This article was published in the following Dove Press journal:
ImmunoTargets and Therapy

Melissa Maria Johanna
van Gool^{1,2}
Marjolein van Egmond¹⁻³

¹Department of Molecular Cell Biology and Immunology, Amsterdam UMC, Amsterdam, Vrije Universiteit Amsterdam, Amsterdam, Netherlands; ²Amsterdam institute for Infection and Immunity, Amsterdam UMC, Amsterdam, Netherlands; ³Department of Surgery, Amsterdam UMC, Vrije Universiteit Amsterdam, Amsterdam, Netherlands

→ Video abstract



Point your Smartphone at the code above. If you have a QR code reader the video abstract will appear. Or use:
<https://youtu.be/xlijxy5W0xA>

Abstract: Mucosal surfaces constitute the frontiers of the body and are the biggest barriers of our body for the outside world. Immunoglobulin A (IgA) is the most abundant antibody class present at these sites. It passively contributes to mucosal homeostasis via immune exclusion maintaining a tight balance between tolerating commensals and providing protection against pathogens. Once pathogens have succeeded in invading the epithelial barriers, IgA has an active role in host-pathogen defense by activating myeloid cells through diverse receptors, including its Fc receptor, FcαRI (CD89). To evade elimination, several pathogens secrete proteins that interfere with either IgA neutralization or FcαRI-mediated immune responses, emphasizing the importance of IgA-FcαRI interactions in preventing infection. Depending on the IgA form, either anti- or pro-inflammatory responses can be induced. Moreover, the presence of excessive IgA immune complexes can result in continuous FcαRI-mediated activation of myeloid cells, potentially leading to severe tissue damage. On the one hand, enhancing pathogen-specific mucosal and systemic IgA by vaccination may increase protective immunity against infectious diseases. On the other hand, interfering with the IgA-FcαRI axis by monovalent targeting or blocking FcαRI may resolve IgA-induced inflammation and tissue damage. This review describes the multifaceted role of FcαRI as immune regulator between anti- and pro-inflammatory responses of IgA, and addresses potential novel therapeutic strategies that target FcαRI in disease.

Keywords: neutrophil, CD89, mucosa, infection, inflammation, autoimmunity

Introduction

The immune system is a central player in protecting the host against infectious diseases. Synergy between both innate and adaptive immunity is essential to induce effective immune responses against invading microbes. Immunoglobulins are major players of the adaptive immune response, and contribute to both immune defense and maintaining homeostasis.¹ Based on structure and effector functions, five major immunoglobulin isotypes can be distinguished, ie, IgM, IgD, IgG, IgA, and IgE that differ in the Fc tail. Immunoglobulins can mediate neutralization, thereby preventing invasion of pathogens or toxins. Additionally, immunoglobulins constitute a bridge between pathogens and the innate immune system facilitating complement activation as well as inducing effector cell functions by immune cells.^{2,3} Binding of the immunoglobulin Fc tail to their cognate Fc receptor, which can be distinguished in receptors for IgG (FcγRs), IgE (FcεRI), IgA (FcαRI), IgM (FcμR), and IgA/IgM (Fcα/μR), induces cellular activation.⁴

Mucosal surfaces like the respiratory-, urogenital- and gastrointestinal tracts are continuously exposed to environmental factors and therefore considered as the

Correspondence: Marjolein van Egmond
Tel +31204448080
Email m.vanegmond@amsterdamumc.nl

frontiers of the body. Tolerating harmless antigens while providing protection against pathogens is a challenging feature of mucosal immunity. IgA is the predominant immunoglobulin at mucosal surfaces and in external secretions. It contributes to mucosal homeostasis by neutralizing toxins and viruses, preventing colonization and invasion of pathogenic bacteria, clearing unwanted particles, and promoting sampling of luminal antigens.^{5,6} In serum, IgA is the second most abundant antibody after IgG. Nonetheless, the exact functions of serum IgA are relatively unexplored and ill understood.⁷

Binding of IgA to its Fc receptor, Fc α RI, can initiate either pro- or anti-inflammatory responses. It was demonstrated that interaction of monomeric serum IgA with Fc α RI induces inhibitory signals (Figure 1A).⁸ As such, it is suggested that IgA and Fc α RI contribute to homeostatic conditions.⁹ By contrast, IgA immune complexes (eg, IgA-opsonized bacteria) induce pro-inflammatory responses by cross-linking of Fc α RI, which is important in controlling infections (Figure 1B).^{10,11} The presence of excessive IgA immune complexes or IgA-opsonized

bacteria can however lead to uncontrolled and disproportionate Fc α RI-mediated immune cell activation, resulting into severe tissue damage as observed during chronic inflammation and autoimmunity.¹² Increased serum IgA levels or IgA autoantibodies have been reported in multiple diseases including rheumatoid arthritis, IgA nephropathy, IgA vasculitis, dermatitis herpetiformis, celiac disease, inflammatory bowel disease, Sjögren's syndrome, ankylosing spondylitis, alcoholic liver cirrhosis, and acquired immunodeficiency syndrome.^{13–20} The role of Fc α RI-mediated inflammation in pathology is still poorly understood. This review summarizes the different functions of Fc α RI and its ligand IgA during homeostasis, infection, chronic inflammation, or autoimmunity, and addresses the possibilities of targeting Fc α RI for therapeutic strategies.

Fc α RI and IgA: The Basics

Fc α RI Structure and Expression

Human Fc α RI is a member of the Fc receptor immunoglobulin superfamily. Nevertheless, Fc α RI has some distinct features compared to other Fc receptors. The Fc α RI gene




Figure 1 Inhibitory and activating signaling via Fc α RI after ligand binding. **(A)** Monomeric IgA (not complexed to an antigen) does not induce Fc α RI cross-linking resulting in partial phosphorylation of immunoreceptor tyrosine-based activation motifs (ITAMs) and recruitment of Src homology region 2 domain-containing phosphatase-1 (SHP-1). This results in inhibition of ITAM signaling, and impairs phosphorylation of spleen tyrosine kinase (Syk), LAT and ERK, which is initiated through signaling via other activating Fc receptors (like IgG-mediated Fc γ receptor activation). The exact binding of free dIgA to Fc α RI, and concomitant signaling, has not yet been resolved. **(B)** IgA immune complexes (eg IgA-coated *Escherichia coli*) induce cross-linking of Fc α RI, resulting in ITAM phosphorylation of the associated Fc γ -chain. Phosphorylated ITAMs subsequently function as a docking site for signaling molecules such as Syk. Syk plays an essential role in initiating signaling pathways, including the Ras/Raf/MEK/MAPK pathway. Activation of signaling pathways results in pro-inflammatory cellular functions such as phagocytosis, antibody-dependent cellular cytotoxicity, respiratory burst, degranulation, antigen presentation, and release of NETs, cytokines and inflammatory mediators. Created with BioRender.com.

(FCAR) is located on chromosome 19 (19q13.4) within the leukocyte receptor cluster (LRC), whereas other human FcR genes are located on chromosome 1.^{21,22} LRC also encodes leukocyte Ig-like receptors and natural killer cell immunoglobulin-like receptors. The amino acid (aa) sequence of FcαRI resembles LRC encoded receptors more closely than other Fc receptors.²³ FCAR consists of five exons that encode for the leader peptide (S1; 34 base pairs (bp), and S2; 36 bp), two extracellular Ig-like domains (EC1 and EC2; 291 bp and 288 bp) and a combined transmembrane and cytoplasmic region (TM/C; 215 bp).²⁴ FCAR encodes a transmembrane receptor, which consists of two extracellular domains (EC1 and EC2; each 206 aa) that are folded with an angle of approximately 90° to each other.²⁵ The transmembrane region (19 aa) is crucial for association with FcR γ-chain,²⁶ and FcαRI has a short cytoplasmic tail (41 aa). FcαRI is expressed on the surface of myeloid cells, including neutrophils, eosinophils, monocytes, macrophages, Kupffer cells, and human platelets.^{12,27,28} Additionally, low FcαRI levels were observed on in vitro cultured immature monocyte-derived dendritic cells (DCs) as well as on monocyte-derived CD103⁺ DCs, which resemble human epithelial interstitial-type DCs.^{29,30} The molecular weight of FcαRI ranges between 50 and 75 kilodalton (kDa) due to differences in N glycosylation, with the exception of FcαRI on eosinophils, which is between 70 and 100 kDa.³¹ Orthologues of human FcαRI have been identified in several monkey species, horses, cattle, hamsters, gerbils, and rats, but not in mice due to a gene translocation.^{32,33}

FcαRI is constitutively expressed and independent of its ligand. Expression can be modulated by several mediators, such as lipopolysaccharide (LPS), chemoattractants, cytokines, or adapter protein binding to the intracellular domain of FcαRI.^{33,34} Upregulation of FcαRI on neutrophils occurs rapidly by either transport from an intracellular pool to the cell surface or via de novo synthesis and is induced by N-formylmethionyl-leucyl-phenylalanine (fMLP), interleukin (IL)-8, tumor necrosis factor-α (TNF-α), LPS, and granulocyte-macrophage colony-stimulating factor (GM-CSF).^{35–38} On monocytes and monocyte-like cell lines FcαRI expression was enhanced by calcitriol, LPS, TNF-α, GM-CSF, and IL-1β.^{39,40} Downregulation occurs in the presence of transforming growth factor-β (TGF-β), interferon-γ (IFN-γ), or by ligand binding due to FcαRI aggregation and internalization.^{41,42}

IgA Binding to FcαRI

Humans express two closely related IgA subclasses, ie, IgA1 and IgA2, whereas most mammals, except for rabbits and certain primates, express only a single IgA subclass that resembles IgA2.^{43–45} In serum, IgA1 is the predominant subclass with an IgA1:IgA2 ratio of 9:1, while IgA2 is mainly found in the colon. In other mucosal tissues, IgA1 and IgA2 are more evenly distributed. IgA1 and IgA2 differ in their hinge region and number of glycosylation sites.⁴⁶ IgA1 contains a hinge region that is 13 amino acids longer compared to IgA2, which results in enhanced antigen recognition capacity but also in increased susceptibility for proteolytic cleavage by bacterial proteases.⁴⁷ Furthermore, IgA1 contains three to six O-linked glycans in the hinge region, while IgA2 is devoid of O-linked glycosylation.⁴⁸ O-linked glycans of IgA1 in external secretions can interact with bacterial adhesion molecules, contributing to mucosal homeostasis.⁴⁹ Altered O-linked glycosylation can cause IgA1 conformational changes resulting in increased immune complex formation, which is a key pathogenic factor in diseases like IgA nephropathy.⁵⁰

In humans, IgA is expressed in three different forms: ie, monomeric IgA, dimeric IgA (dIgA), and secretory IgA (SIgA). Serum IgA is mostly monomeric and produced by plasma cells in the bone marrow, spleen, and lymph nodes. By contrast, IgA at mucosal sites is predominantly dimeric and produced by local plasma cells in the lamina propria.^{51,52} Dimeric IgA (dIgA) is composed of two monomers that are linked tail-to-tail with a joining (J-) chain via Cys471-mediated disulfide bonds forming a boomerang-like structure.⁵³ It is transported across the epithelium by binding to the polymeric Ig receptor (pIgR), which is expressed on the basolateral membrane of epithelial cells.^{53,54} At the luminal side, pIgR is cleaved and a part of this receptor, referred to as secretory component (SC), remains attached to dIgA by binding both Fc-tails and J-chain across the ~50° gap between the two monomers, thereby forming SIgA (Figure 2A).^{55,56}

All (iso)forms of IgA are ligands for FcαRI, although binding of SIgA to FcαRI is (partially) hampered due to steric hindrance of SC.⁵⁷ Furthermore, binding of SIgA requires the presence of macrophage-1 antigen (Mac-1, CD11b/CD18).⁵⁸ Monomeric IgA and dIgA bind with moderate affinity ($K_a = 10^6 \text{ M}^{-1}$), whereas IgA immune complexes bind with higher avidity and induce crosslinking of FcαRI.³³ Optimal binding of IgA immune complexes occurs with five to six molecules of IgA per complex.⁵⁹ In particular residues Pro440-Phe443 and




Figure 2 Roles of IgA and FcαRI in homeostasis and infection at mucosal sites. **(A)** Local plasma cells in the lamina propria produce dimeric IgA (dIgA), which is transported across the epithelium into luminal secretions by binding to the polymeric Ig receptor (pIgR). At the luminal side it is released as secretory IgA (SIgA) where it can neutralize pathogens and toxins. **(B)** On route of being secreted, dIgA can intercept viruses, which have infected epithelial cells and redirect them into the lumen. **(C)** Invading pathogens and antigens in the lamina propria are opsonized by dIgA and transported back into the lumen. **(D)** Microbes that are opsonized with SIgA are shuttled via microfold (M) cells to dendritic cells (DCs) in Peyer's patches for sampling. Additionally, DCs can extend dendrites through the epithelial layer for sampling of the luminal content. **(E)** During infection, dIgA-opsonized pathogens are taken up by FcαRI-expressing DCs, and presented to T cells. Additionally, phagocytosis of dIgA-opsonized pathogens by neutrophils results in the release of leukotriene B4 (LTB4), which mediates the chemotaxis of more neutrophils to the site of infection, thereby functioning as a self-contained positive feedback loop of immune cell recruitment to clear invading pathogens. Created with BioRender.com.

Leu257-Leu258 in the FcαRI EC1 are essential for IgA binding.⁶⁰ Monomeric IgA binds to FcαRI via its Ca2 and Ca3 domains in a 2:1 stoichiometry (one IgA molecule binds simultaneously two FcαRI molecules).^{25,61} This is in contrast with other Fc receptors since FcεRI and FcγRIII bind their ligands in EC2 in a 1:1 stoichiometry.^{62–65} In theory, dIgA can bind four FcαRI due to its multiplied binding sites, although this may not be possible due to steric hindrance. As such, it still

remains unclear how dIgA exactly binds to FcαRI.⁶⁶ It has been described that pentraxins, including acute phase C reactive protein (CRP) and serum amyloid P (SAP), compete for FcαRI binding since these proteins recognize a similar binding site on FcαRI as IgA.⁶⁷ Mutations in FcαRI outside the IgA-binding site enhanced pentraxin binding to FcαRI with 2-fold whereas IgA binding was unaffected, suggesting additional binding sites for pentraxins.⁶⁷

Fc α RI harbors six N-glycosylation sites which affect the binding affinity of IgA to Fc α RI. Altered glycosylation of Fc α RI due to a specific mutation (Asn58 to Glu58) resulted in a nearly 2-fold increased binding of IgA. Furthermore, removal of sialic acids increased IgA binding with nearly 4-fold.⁶⁸ IgA binding to Fc α RI can also be regulated by intracellular signals that are modulated by cytokine stimulation of cells (also referred to as inside-out signaling), independently of Fc α RI expression levels.⁶⁹ Increased binding of IgA to Fc α RI in transfected cells, eosinophils, and monocytes was reported in the presence of cytokines like GM-CSF, IL-4, and IL-5 without increasing Fc α RI expression levels on the cell surface.^{70,71}

Fc α RI Signaling and Cellular Activation

Fc α RI does not contain any known signaling motifs. In order to initiate effector functions, Fc α RI associates with the FcR γ -chain subunit, which contains an immunoreceptor tyrosine-based activation motif (ITAM) in its intracellular domain.²⁶ In the transmembrane regions, the positively charged arginine on position 209 (R209) in Fc α RI associates with the negatively charged aspartic acid 11 (D11) in FcR γ -chain.⁷² The FcR γ -chain ITAMs contain conserved paired tyrosines and leucines in a consensus sequence (YxxL-x₇₋₁₂-YxxL). Binding of monomeric serum IgA (not complexed with an antigen) leads to partial phosphorylation of FcR γ -chain and involves extracellular signal-related kinases (ERK)-dependent recruitment of tyrosine phosphatase Src homology region 2 domain-containing phosphatase-1 (SHP-1) to sphingolipid-cholesterol-rich membrane domains.⁸ Cytoplasmic clusters referred to as inhibisomes hamper spleen tyrosine kinase (Syk), linker for activation of T cells (LAT), and ERK phosphorylation, thereby inhibiting pro-inflammatory responses that are induced by other activating Fc receptors.⁷³ This process is referred to as inhibitory ITAM (ITAMi) signaling (Figure 1A), which may represent an anti-inflammatory mechanism to prevent uncontrolled release of inflammatory responses (see also below).

By contrast, binding of IgA immune complexes to Fc α RI induces cross-linking of Fc α RI, resulting in pro-inflammatory responses (Figure 1B). After crosslinking, tyrosines in ITAMs are phosphorylated by the Src kinase Fyn, forming docking sites for other tyrosine kinases, including Syk.⁷⁴⁻⁷⁶ Syk plays an essential role in the activation of several proteins, including phosphoinositide 3-kinase (PI3-K), phospholipase C γ (PLC γ), and Src homology and collagen adaptor protein (Shc). These

proteins initiate multiple signaling pathways, including the Ras/Raf/MEK/MAPK pathway and the Rho family GTPase pathway. These pathways are also interconnected and can therefore modulate each other.⁷⁴⁻⁷⁶ Syk additionally induces the release of second messengers such as calcium and diacylglycerol.⁷⁷ Activation of signaling pathways results in remodeling of the actin cytoskeleton and activation of transcription factors such as nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ b), ultimately leading to cellular activation (for detailed description of Fc α RI signaling, see Aleyd et al).¹⁰

Depending on the cell type, Fc α RI activation can result in phagocytosis, degranulation, superoxide generation, release of neutrophil extracellular traps (NETs), antibody-dependent cellular cytotoxicity (ADCC), release of cytokines and chemokines, or antigen presentation.¹² It has been shown that phagocytosis of IgA-coated particles by neutrophils induces increased reactive oxygen species (ROS) production and NET release, which can contribute to pathogen elimination.⁷⁸ Additionally, cross-linking of Fc α RI by serum or dIgA initiates the release of the chemoattractant leukotriene B₄ (LTB₄) with concomitant neutrophil recruitment.⁷⁹ SIgA was able to induce respiratory burst in neutrophils, although less efficiently compared to serum IgA. It did not induce efficient uptake of pathogens by either neutrophils or Kupffer cells.^{27,80} Cross-linking of Fc α RI by IgA immune complexes on immature DCs resulted in antigen presentation through the major histocompatibility complex class II pathway, DC maturation, and production of IL-10.⁸¹⁻⁸³ Since IL-10, together with TGF- β , mediates IgA isotype switching in B cells, Fc α RI-positive DCs are described to be important for the induction of IgA by B cells in secondary lymphoid organs.⁸⁴ Additionally, cross-linking of Fc α RI on in vitro generated human CD103⁺ DCs (resembling human epithelial interstitial-type DCs) resulted in the release of pro-inflammatory cytokines like TNF- α , IL-1 β , IL-6, and IL-23.²⁹ SIgA was internalized by DCs as well, albeit through carbohydrate-recognizing receptors instead of Fc α RI, which did not result in DC maturation.⁸⁵

Fc α RI and IgA: Homeostasis Immune Exclusion and Immune Regulation

In mucosal areas like the respiratory-, urogenital-, and gastrointestinal tracts, SIgA plays a key role in keeping a tight balance between tolerating commensals and harmless antigens while providing protection against harmful pathogens.⁸⁶

Due to its multiple binding sites, SIgA opsonizes bacteria with high avidity thereby interfering with bacterial motility and inducing bacterial agglutination. This can block the entrance of bacteria into the mucosal epithelium.⁸⁷ Additionally, bacterial products like enzymes and toxins are also neutralized by SIgA.⁸⁸ This process, referred to as immune exclusion, is the main described function of SIgA and prevents local and systemic infection (Figure 2A).⁸⁶ Not only bacteria but also viruses can be neutralized by mucosal IgA. On route to be secreted, dIgA has the ability to intercept and disarm viruses, which have infected epithelial cells and redirect them back into the lumen (Figure 2B).¹² Similarly, pathogens that have invaded the epithelial barrier into the lamina propria are likely opsonized by dIgA and transported back into the lumen (Figure 2C). It has been shown that IgA neutralized several viruses, including sendai, influenza, rota, measles, and human immunodeficiency virus 1 (HIV-1).^{89–94} Additionally, SIgA plays an important role in shaping and diversifying the gut microbiome.⁹⁵ Mice lacking IgA showed reduced overall microbial diversity resulting in altered bacterial composition, increased bacterial translocation, and ultimately intestinal inflammation.^{96–98} Similarly, humans with defective mucosal IgA responses, such as Selective IgA-deficiency (SIgAd), showed modest but significant changes in the gut microbiota composition.⁹⁹ High-microbiota binding as well as cross-species reactivity of IgA was shown to promote host-microbiota symbiosis, hereby maintaining intestinal mucosal integrity and homeostasis.^{100–102} Yet, the functional effects of IgA binding to microbiota fitness or physiology remain unclear. IgA production is also microbiota-dependent, as germ-free showed diminished IgA titers compared to specific-pathogen-free mice.^{103,104} IgA-producing plasma cells were absent in most tissues of germ-free mice and significantly reduced in the small intestine, reflecting the importance of microbiota in initiating humoral responses.¹⁰¹

SIgA immune complexes (eg, IgA-opsonized bacteria) can be transported from the lumen into the lamina propria via the transferrin receptor 1 (TfR1 or CD71) on epithelial cells or via Dectin-1 on microfold cells (M cells) (Figure 2D).^{105,106} Reverse transcytosis of SIgA is important for the uptake and delivery of antigens from the intestinal lumen to gut-associated lymphoid tissues (GALTs) influencing immune responses.^{106,107} SIgA immune complexes are taken up by DCs through interaction with Dendritic Cell-Specific Intercellular adhesion molecule-3-Grabbing Non-integrin (DC-SIGN).⁸⁵

In the circulation, serum IgA has likely an immunomodulatory role through inhibitory signals via ITAMi signaling (Figure 1A). It was shown that monovalent targeting of Fc α RI resulted in the inhibition of oxidative burst activity, chemotaxis and IgG-mediated phagocytosis and cytokine production.^{108–111} Furthermore, in transgenic mice that express human Fc α RI on myeloid cells, IgE-mediated asthma was prevented by the binding of soluble IgA to Fc α RI, which inhibited Fc ϵ RI-induced degranulation of mast cells.⁸ ITAMi signaling by Fc α RI is therefore suggested to play a role in maintaining homeostasis and protection against enhanced Fc γ receptor- or Fc ϵ RI-mediated activation during inflammatory diseases and allergies.⁹

Early-Life Immunity

SIgA is the predominant antibody in human colostrum and described to protect offspring from infection when the neonatal immune system is still immature.^{112,113} IgA and IgM are not able to cross the placenta through the neonatal Fc receptor (FcRn) and can therefore only be provided through maternal milk.¹¹⁴ Breastfeeding is important for the development of the neonatal intestinal microbiota and can protect infants from infectious diseases.^{114–116} Studies in mice have shown that colostrum is the only source of SIgA in the first weeks of life since it takes up to approximately 4 weeks for the neonatal intestine to be populated by IgA-secreting B cells.^{116,117} In humans, mucosal IgA in the fetal intestine is absent or rarely present until 10 days after birth.¹¹⁸ IgA-producing B cells become pre-dominant 1–2 months after birth and increase in number until 6 to 11 months of age.¹¹⁹ The highest concentration of SIgA is found in colostrum; however, prolonged breastfeeding resulted in increased IgA in maternal milk as well.^{120,121} Although the opsonic activity of SIgA is poor, early studies demonstrated that SIgA initiated macrophage phagocytosis and neutrophil respiratory burst.¹²² After antigen interaction, SIgA undergoes conformational changes, which enhanced the binding of SIgA immune complexes to Fc α RI.¹²³ In addition to immunoglobulins, maternal milk also facilitates the transfer of leukocytes. IgA-induced cellular effector functions may therefore contribute to early immunity against pathogens in newborns. It has been demonstrated that the majority of leukocytes present in maternal milk have a similar phenotype to blood cells, although leukocyte subsets are present in different frequencies. Myeloid precursors, neutrophils, and immature granulocytes are the main identified cells

present in colostrum.¹²⁴ Colostral neutrophils were able to phagocytose IgA-opsonized bacteria although this did not result in significant bacterial-killing and release of superoxide anion. This is likely due to FcαRI expression without γ subunit association.¹²⁵ The exact contribution of FcαRI-mediated cellular responses initiated by SIgA or leukocytes in maternal milk remains incompletely understood.

FcαRI and IgA: Infection

The production of IgA, with a synthesis rate of 66mg/kg, exceeds that of all other antibodies combined supporting its importance in host-pathogen defense.¹² Pathogen-specific IgA is found at mucosal surfaces and in circulation during several infectious diseases, while the exact function of IgA remains unclear.^{6,11} Passive immunity (via neutralization) as well as immune activating properties of IgA have been described for infections in the respiratory- and reproductive tracts such as *Mycobacterium tuberculosis* (*Mtb*), HIV-1, and more recently severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infections.^{126–128}

Passive Immunity

Respiratory infections often occur via inhalation of air-borne droplets containing pathogens or via direct contact with respiratory secretions.^{129,130} The nasopharyngeal mucosa is furthermore a natural reservoir for several pathogens like *Streptococcus pneumoniae* (*S. pneumoniae*) and *Neisseria meningitidis* (*N. meningitidis*).¹³¹ In individuals who lack protective antibody titers, infections can occur causing various pathologies like pneumonia, meningitis, bacteremia, and sepsis.^{132,133} In mice, SIgA against *Mtb* or *S. pneumoniae* has been associated with protection against tuberculosis (TB) or pneumococcal disease respectively.^{134,135} The neutralizing capacities of IgA are not only described for bacteria since IgA against HIV-1 or rotavirus was able to neutralize either virus. Monomeric IgA1 or dIgA2 against envelope proteins (Env) of HIV-1 showed neutralizing activity in vitro.¹³⁶ Moreover, mucosally applied Env-specific dIgA1 and dIgA2 protected rhesus macaques against an intrarectal challenge with simian-human immunodeficiency virus (SHIV).^{137,138} IgA against rotavirus (RV-IgA) neutralized both virus in solution and virus that had bound to epithelial cells in vitro.¹³⁹ Furthermore, it was shown that RV-IgA was produced in all rotavirus animal models (horse, cow, sheep, gnotobiotic piglet, rat, rabbit, and mouse) which correlated with clearance of infection and protective immunity.^{140–142}

Generation of specific IgA in coronavirus disease 2019 (COVID-19) was described as well.^{143–146} It was suggested that SIgA antibodies against SARS-CoV-2 were able to neutralize the spike protein or nucleocapsid protein thereby providing protective mucosal immunity.¹⁴⁷ It is unknown whether serum IgA anti-SARS-CoV-2 antibodies contribute to protection against COVID-19. On the one hand, it was demonstrated that severe COVID-19 illness was significantly associated with increased total serum IgA but not with total serum IgG levels.¹⁴⁸ Furthermore, IgA levels correlated with disease score in critically ill patients.¹⁴⁹ On the other hand, the presence of IgA and IgG antibodies against SARS-CoV-2 spike protein subunit 1 (S1) showed an inverse correlation with viral load. Additionally, enhanced titers of specific anti-S1 IgA in serum correlated with significant increased survival in COVID-19 patients 28 days post intensive care unit admission.¹⁵⁰ Passive antibody therapy using convalescent plasma from recovered COVID-19 patients resulted in a reduction in viral load and better disease outcomes, most notably in less severely ill COVID-19 patients.^{151–153} Nonetheless, since IgA-specific SARS-CoV-2 neutralizing antibodies were not determined in convalescent plasma that was transferred to COVID-19 patients, the exact contribution of IgA neutralizing antibodies in COVID-19 remains unknown.^{154–156} Longitudinal investigation of potentially protective functions of mucosal and systemic IgA is needed to determine their role in COVID-19.

Active Immunity

Although few FcαRI-positive cells are observed in mucosal areas in homeostatic conditions, FcαRI-positive neutrophils are the first cells that are recruited after infection.^{79,157} Once pathogens have successfully invaded the epithelial barrier, they can become opsonized by dIgA in the lamina propria. It was shown that dIgA induced efficient phagocytosis by neutrophils.⁸⁰ As such, a pro-inflammatory role for dIgA in eliminating infiltrating pathogens was proposed.³³ Enhanced phagocytosis of *Escherichia coli*, *Streptococcus pneumoniae*, *Staphylococcus aureus*, *Porphyromonas gingivalis*, *Candida albicans*, *Bordetella pertussis*, and *Neisseria meningitidis* by neutrophils after targeting FcαRI has been demonstrated in vitro.^{27,79,80,158–161} Specific IgA antibodies against meningococcal capsular polysaccharides were shown to induce moderate phagocytosis by neutrophils but initiated respiratory burst more potently than IgG.⁸⁰ Cross-linking of FcαRI on neutrophils furthermore results in the release of LTB₄, which is a potent neutrophil

chemoattractant, and additionally mediates the migration of monocytes and monocyte-derived DCs.^{79,162,163} LTB₄ release after neutrophil activation through FcαRI has therefore been proposed to function as a self-contained positive feedback loop of immune cell recruitment to clear invading pathogens and maintain homeostasis (Figure 2E).³³ Alveolar macrophages, which express an alternatively spliced variant of FcαRI, also contribute to host-pathogen defense by clearance of inhaled antigens that were opsonized by IgA.¹⁶⁴ A recent study showed that antibodies from patients with latent TB induced enhanced macrophage killing of intracellular *Mtb*.¹⁶⁵ Since IgA levels against TB antigens distinguished patients with pulmonary TB, patients with latent TB, and non-infected individuals, a role for FcαRI-mediated effector functions in *Mtb* control is predicted, although this is formally not yet established.¹⁶⁶ Interestingly, a recent study showed that FcαRI also acts as an innate receptor, by binding bacteria directly independent of its ligands IgA or CRP. Binding of bacteria to FcαRI induced phagocytosis by CD11c⁺ DCs and monocytes/macrophages. Moreover, FcαRI transgenic mice were protected against two different sepsis-induced models, identifying FcαRI as a first-line innate receptor for bacterial clearance.¹⁶⁷

The contribution of antibody-dependent cellular phagocytosis (ADCP) in protective immune responses against viruses remains debatable. Nonetheless, there is substantial evidence that supports the involvement of ADCP in protection against several types of viruses and reduction of disease.¹⁶⁸ It has been described that human NK cells and neutrophils perform ADCC and ADCP of HIV-1 gp120-pulsed target CEM-NKr cells or gp120-coated beads, respectively, in the presence of polyclonal antibodies from different HIV-positive subject groups.¹⁶⁹ Moreover, HIV-1 gp41 envelope-specific IgA induced FcαRI-mediated ADCC of HIV-1 Clade A- and B-infected target cells by monocytes.¹⁷⁰ HIV-1 gp41 envelope-specific IgA additionally triggered ADCP of HIV-1 infected CD4⁺ T cells by monocytes and neutrophils more efficiently than anti-gp41 IgGs.¹⁷¹ FcαRI-mediated neutrophil phagocytosis initiated by vaccine-induced IgA was associated with reduced risk of infection against simian immunodeficiency virus (SIV) after immunization via the nasal, but not the intramuscular route in non-human primates.¹⁷² These studies may provide new insights into FcαRI-mediated immune responses against HIV-1 infected cells and address the potential functions of IgA during viral infections. By contrast, the RV144 HIV-1 vaccine trial

showed that IgA interfered with IgG-mediated ADCC by NK cells of HIV-1 infected cells.¹⁷³ It was proposed that Env-specific IgA competed with IgG for NK cell-mediated ADCC of HIV-1 infected cells, while enhancing ADCP by FcαRI-expressing cells like neutrophils, which represent the dominant phagocyte population in tissues from the lower female reproductive tract.

The severity of infections increases when invading pathogens from the respiratory- or gastrointestinal tract enter the bloodstream. In mice it was shown that serum IgA directed against commensal bacteria protected mice against lethal sepsis when the intestinal barrier was damaged, suggesting that serum IgA provided protection against systemic infection.¹⁰³ Mice lacking pIgR and SIgA have epithelial barrier disruption, enhanced numbers of IgA-secreting plasma cells, and increased levels of serum IgG and IgA.¹⁷⁴ It was furthermore demonstrated that in human FcαRI-transgenic mice IgA-opsonized bacteria in the circulation were phagocytosed by Kupffer cells, which express FcαRI.²⁷ FcαRI cross-linking by serum IgA immune complexes and cross-talk with pathogen recognition receptors (PRRs) on Kupffer cells initiated the release of pro-inflammatory cytokines such as TNF-α, IL-1β, and IL-6.^{175,176} As such, clearance of serum IgA-coated bacteria by Kupffer cells may act as a systemic line of defense, through elimination of invasive bacteria that have escaped mucosal immune responses (Figure 3).

Bacterial Evasion Mechanisms

Pathogens have evolved and developed strategies to either disarm IgA or evade FcαRI-mediated activation of immune cells, emphasizing the importance of IgA-FcαRI interactions in the elimination of pathogens and prevention of infection.¹⁷⁷ Pathogens that colonize the oral and upper respiratory mucosa like *S. pneumoniae* and *N. meningitidis* produce specific proteases cleaving both mucosal and serum IgA1 at the hinge region, which abrogates its protective effects and interferes with host antibacterial immunity.^{178,179} Binding of pneumococcal surface protein (SpsA) to SIgA is suggested to recruit SIgA to the bacterial surface to promote its degradation by IgA proteases, hinder SIgA from clearing bacteria, or block the interaction between SIgA and FcαRI thereby preventing immune responses.⁵³ Additionally, IgA1 proteases were shown to degrade lysosomal-associated membrane protein 1 (LAMP-1) promoting intracellular bacterial survival in epithelial cells in vitro.¹⁸⁰ Furthermore, it was described that serine proteases do not only cleave IgA1, which prevents FcαRI-mediated immune responses but also reduce the binding




Figure 3 Systemic protection by serum IgA and Fc α RI-expressing Kupffer cells. Serum IgA-opsonized bacteria in the circulation are transported to the liver through the portal vein and phagocytosed by Kupffer cells (KC), which express Fc α RI, providing protection against systemic infection. Cross-talk between Fc α RI and pathogen recognition receptors (PRRs) may break the tolerance of Kupffer cells to bacterial structures. Created with BioRender.com.

avidity of IgA1 thereby reducing the neutralizing ability of IgA1.¹⁸¹ Closely related strains of *N. meningitides*, as well as *S. pneumoniae* that lack IgA1 proteases are considered nonvirulent.¹⁸² Furthermore, *Staphylococcus aureus* (*S. aureus*) and group A and B streptococci developed evasion mechanisms to circumvent Fc α RI-mediated elimination by secreting decoy proteins that inhibit binding of IgA to Fc α RI. Decoy proteins Sir22, Arp4, and an unrelated β protein from group B streptococci, as well as staphylococcal superantigen-like protein seven that is produced by *S. aureus*, bind to specific Fc residues in the Ca2 and Ca3 domains of IgA.^{183,184} The ability of bacteria to escape Fc α RI-mediated immunity enhances survival in mucosal environments and may contribute to systemic infections.

Fc α RI and IgA: Chronic Inflammation and Autoimmunity

Because Fc α RI potently activates immune cells, aberrant IgA responses may contribute to pathogenesis in inflammatory and/or autoimmune diseases. The presence of IgA autoantibodies, aberrant IgA glycosylation, or excessive IgA immune complexes can contribute to chronic inflammation and tissue damage. The detrimental role of IgA and Fc α RI-mediated immune responses has been proposed in

several inflammatory diseases like IgA nephropathy, rheumatoid arthritis, IgA vasculitis, dermatitis herpetiformis, linear IgA bullous disease, and inflammatory bowel disease.

IgA Nephropathy

IgA nephropathy (IgAN) is the most common IgA-mediated autoimmune disease. It is characterized by primary glomerulonephritis, resulting in chronic renal failure in 30% of the patients.^{185,186} Infections or microbiome dysbiosis prior to IgAN have been suggested to initiate abnormal IgA1 glycosylation as glycosyltransferases are regulated by bacterial products.¹⁸⁷ Elevated synthesis of galactose-deficient IgA1 (gd-IgA1) triggers the production of anti-glycan antibodies. IgA immune complexes can deposit in the glomeruli inducing mesangial proliferation and matrix expansion, which eventually initiates renal injury.¹⁸⁸ The release of pro-inflammatory cytokines such as TNF- α , IL-6, and TGF- β by mesangial cells after IgA immune complex binding was suggested to induce inflammation and glomerulosclerosis.^{187,189} In mice, gd-IgA1 immune complexes were not cleared from the circulation and deposited in the kidney. Shedding of Fc α RI in transgenic mice however resulted in the deposition of soluble

FcαRI-IgA immune complexes in the mesangium, which induced glomerular and interstitial macrophage infiltration, hematuria, mesangial matrix expansion, and mild proteinuria.^{190–192} Soluble FcαRI-IgA complexes additionally mediated kidney inflammation by interacting with Tfr1 on mesangial cells, resulting in the release of pro-inflammatory mediators.¹⁹⁰ Human FcαRI transgenic mice developed IgAN after injection of serum IgA of IgAN patients. Furthermore, human IgA1 knock-in mice that had been crossed with FcαRI transgenic mice already developed IgAN in 6 weeks.¹⁹³ These data clearly support the contribution of IgA and FcαRI-mediated inflammation in IgAN.

Rheumatoid Arthritis

Rheumatoid arthritis (RA) is a systemic and chronic autoimmune disorder characterized by inflammation in the joints, which frequently leads to joint destruction and disability.¹⁹⁴ Although the etiology of RA is unknown, the presence of autoantibodies is a distinctive feature of RA. In particular, IgM rheumatoid factor (RF) and IgG anti-citrullinated protein antibodies (ACPA) are commonly used for diagnosing and classifying RA.¹⁹⁵ Interestingly, elevated levels of IgA RF, but not of IgM- or IgG RF, correlated with worse disease prognosis and extra-articular manifestations in RA.¹⁹⁶ Additionally, high levels of IgA RF were associated with a poor response rate to TNF-α blockers as therapeutic agents in advanced RA.¹⁹⁷ In IgM RF negative patients it was shown that IgA ACPA was associated with disease severity based on disease activity score.¹⁵ RF or ACPA can be elevated in the serum years prior to the onset of clinical disease, referred to as pre-clinical RA.^{198,199} An increased number of IgA⁺ plasma-blasts was identified in preclinical RA without a history of or current inflammatory arthritis.²⁰⁰ In both clinical and preclinical RA, IgA ACPA was highly specific for RA.^{201,202} IgA ACPA and RF immune complexes furthermore increased the release of pro-inflammatory cytokines such as TNF-α, IL-1β, and IL-6 by myeloid cells via cross-linking of FcαRI, whereas blocking of FcαRI on macrophages resulted in reduced levels of TNF-α.²⁰³ Additionally, increased NET release by neutrophils in the presence of plasma of RA patients containing IgA RF was demonstrated, which was dependent on FcαRI.⁷⁸ Altogether, this supports that IgA contributes to disease, as activation of FcαRI-expressing neutrophils and macrophages in the joint via IgA auto-immune complexes likely enhances inflammation and pathology of RA.

IgA Vasculitis

IgA vasculitis, formerly known as Henoch-Schönlein purpura, is the most common form of vasculitis involving the small vessels of the joints, kidneys, gastrointestinal tract, and the skin.²⁰⁴ Although the etiology of IgA vasculitis remains unknown, the disease is characterized by IgA1 immune deposits and neutrophil infiltrates damaging the small vessels. Abnormal glycosylation of the hinge region of IgA1 is suggested to cause aggregation into macromolecular immune complexes.²⁰⁴ FcαRI-mediated cross-linking of neutrophils induces inflammatory processes like ROS production, NET formation, and LTB4 release. As such, it is hypothesized that FcαRI-mediated activation of neutrophils results in vessel damage and leakage of red blood cells into the skin causing typical cutaneous hemorrhages.¹⁴ Antigen recognition sites of accumulated IgA1 in IgA vasculitis have not yet been identified. However, it was proposed that these antibodies can recognize epitopes on endothelial cells as serum IgA from IgA vasculitis patients was shown to bind to human but not bovine glomerular endothelial cells in vitro.²⁰⁵ Notably, sera of IgA vasculitis patients had increased levels of soluble FcαRI-IgA complexes, which was associated with decreased FcαRI expression on monocytes.²⁰⁶ A positive feedback loop of inflammation in IgA vasculitis was proposed, since binding of IgA1 antibodies to endothelial cells induced the release of IL-8, thereby attracting more neutrophils that can be activated through FcαRI.¹⁶

Skin Blistering Diseases

Dermatitis herpetiformis (DH) is an autoimmune disease that often occurs in combination with gluten-sensitive enteropathy (celiac disease). Anti-transglutaminase IgA antibodies (mainly anti-epidermal transglutaminase 3; TGase3) play a key role in disease pathogenesis of DH as it was shown that serum anti-epidermal transglutaminase IgA positively correlated with disease activity.²⁰⁷ IgA anti-tissue transglutaminase antibodies form immune complexes and deposit in the dermis of patients with DH.¹² In the skin, transglutaminase/IgA immune complexes induce fibrinolysis, which directly contributes to blister formation in DH, and results in chemotaxis of neutrophils and monocytes.^{208,209} Accumulating FcαRI-positive neutrophils in the skin of DH had increased ability to bind IgA. As such, they presumably bind transglutaminase/IgA immune complexes and induce cellular activation contributing to tissue damage.^{210,211} Released proteolytic enzymes by activated neutrophils can induce sub-epidermal split by cleaving adhesion molecules

such as collagen VII.²¹² Circulating epidermal transglutaminase/IgA immune complexes have been detected in patients with DH which, together with disease manifestations, decrease over time after following a gluten-free diet.²¹³ Some DH patients also showed IgA immune complex deposition in the kidney, which ultimately can lead to IgA nephropathy (IgAN).²¹³

Similar to DH, linear IgA bullous disease (LABD) is a skin blistering disease characterized by IgA autoantibodies against collagen XVII and Fc α RI-mediated neutrophil activation.²¹⁴ It was shown in an ex vivo skin model that activated neutrophils in the presence of serum of LABD patients (containing anti-collagen XVII IgA) caused the separation of dermis from epidermis (Figure 4). This separation, reflecting blister formation in LABD patients, was prevented by blocking IgA-Fc α RI interaction on neutrophils.²¹⁵ Additionally, eosinophil influx has also been observed in the skin of LABD, which may contribute to LABD pathology through Fc α RI-mediated respiratory burst activity.^{214,215}

Inflammatory Bowel Disease

Inflammatory bowel disease (IBD) is characterized by chronic inflammation of the gastrointestinal tract and disruption of the epithelium. IBD can be subdivided into two major forms, ulcerative colitis (UC) and Crohn's disease (CD).²¹⁶ The etiology of IBD is ill-understood, but likely

involves derailed immune responses against commensal bacteria. Fecal bacteria of patients with IBD, that were highly opsonized with IgA, induced colitis in germ-free mice.²¹⁷ Furthermore, fecal bacteria in IBD patients showed increased opsonization with IgA compared to healthy individuals, suggesting that leakage of serum IgA or dIgA through the disrupted epithelial layer contributes to enhanced opsonization.^{217,218} In line with this, it was shown that IBD patients have increased levels of specific IgA against microbiota in their serum.²¹⁹ The percentage of IgA-opsonized bacteria in CD strongly correlated with multiple clinical indexes of disease activity and may therefore be useful for monitoring CD.²²⁰ Due to the disruption of the intestinal epithelial barrier in these patients, bacteria are able to invade the sub-epithelial lamina propria and are likely opsonized with dIgA. This may subsequently induce neutrophil activation via Fc α RI cross-linking, since uptake of IgA complexes by neutrophils was observed in the mucosa of patients with UC.⁷⁹ Fc α RI-mediated neutrophil activation induces migration through the release of LTB₄. It was therefore hypothesized that in UC patients a continuous neutrophil recruitment loop is initiated. Newly recruited neutrophils will be activated by IgA-opsonized bacteria and initiate a feedback loop of neutrophil chemotaxis, which eventually results in tissue damage.⁷⁹ Moreover, cross-talk of Fc α RI and toll-like receptor 4 (TLR4) on neutrophils resulted in enhanced release of pro-inflammatory TNF- α , which plays a central role in IBD pathogenesis.^{221,222} Excessive influx of neutrophils causing major tissue damage was demonstrated in UC patients.²²² As such, a role for IgA and Fc α RI is proposed in especially UC, but their exact roles still need to be established.




Figure 4 IgA-Fc α RI induced pathology in skin blistering diseases. Cross-linking of Fc α RI on neutrophils with IgA immune complexes results in the release of leukotriene B₄ (LTB₄) inducing enhanced neutrophil influx, causing tissue damage in a variety of inflammatory- and autoimmune diseases. An example is the blister formation in linear IgA bullous disease (LABD). Created with BioRender.com.

Fc α RI and IgA: Therapeutic Opportunities

IgA Vaccination Strategies

Inducing pathogen-specific mucosal and systemic IgA through passive or active immunization may be an attractive strategy to combat bacterial and viral infections.²²³ Licensed oral and nasal vaccines have demonstrated to induce mucosal SIgA responses as well as system responses contributing to protection.^{223,224} Furthermore, mucosal application of the TB protein subunit vaccine H56/CAF01 followed by intramuscular priming induced high levels of antigen-specific lung mucosal and systemic IgA.²²⁵ *Mtb*-specific IgA may contribute to prevention of TB, since

mucosally administered human IgA antibodies against *Mtb* in mice provided passive protection.²²⁶ Additionally, vaccine-induced pulmonary SIgA in mice was associated with passive protection against TB.^{134,227} Recombinant monoclonal antibodies (mAb), generated from single isolated B cells of untreated adult patients with acute pulmonary TB and from MTB-exposed healthcare workers, revealed that IgA against *Mtb* inhibited mycobacterial infection of human epithelial-like and macrophage-like cells, whereas IgG promoted infection.¹²⁶ Thus, it is suggested that IgA may be particularly important for protection against TB infection.²²⁸ This is supported by experiments with human FcαRI transgenic mice, which had a lower *Mtb* infection rate compared to control WT mice after inoculation of human IgA mAbs, implicating an important role for FcαRI in clearing *Mtb* infection.²²⁶ Nasal immunization with pneumococcal surface protein A (PspA) and cholera toxin as adjuvant elicited PspA-specific mucosal IgA and systemic IgG antibody responses, which provided protection against colonization of *S. pneumoniae* and pneumococcal infection in mice.²²⁹ PspA-specific SIgA effectively neutralized colonized *S. pneumoniae*, diminished nasal carriage and prevented subsequent infection with *S. pneumoniae*. IgA-deficient mice failed to prevent nasal colonization by *S. pneumoniae*, in spite of having functional anti-PspA IgG antibodies in the nasal cavity, stating the necessity of specific IgA to prevent infection.²³⁰

Vaccine-induced IgA also provides effective protection against viral infections. Both the inactivated and the live-attenuated oral poliovirus vaccines, which played a tremendous role in the elimination of the poliovirus, induced potent-specific IgA responses.²³¹ Oral vaccines consisting of live attenuated rotavirus (RV) mediate significantly reduced disease. Serum rotavirus-specific IgA (RV-IgA) correlated with vaccine efficacy and protective immunity.²³² IgA knockout, as well as J-chain knockout mice, showed delayed viral clearance and absence of protective immunity, supporting the critical role of IgA in RV immunity.^{233,234} Interestingly, in both humans and animals, RV-IgA can persist for a long time, suggesting the presence of long-lived IgA⁺ plasma cells.²³⁵ Similar to RV, it has been shown that vaccination with live attenuated poliovirus induced long-lived IgA memory immune responses in elderly.²³⁶ Intravenous administration of polymeric IgA against influenza showed protection against infection in mice. Inhibition of influenza virus shedding by IgA was 10 times more effective than IgG.²³⁷ Additionally, IgA antibodies against influenza A virus hemagglutinin,

purified from respiratory tract washings of hemagglutinin immunized mice, were able to protect naïve mice from influenza infection.²³⁸ Similar protective effects of IgA were shown in mice that received passive oral administration of IgA against reovirus-specific adhesion molecule sigma1, which prevented entry of reovirus into the Peyer's patches. Since IgG mAbs against sigma1 did not prevent Peyer's patch infection by reovirus, this effect was IgA specific.²³⁹ Thus, inducing systemic and mucosal IgA responses through vaccination may play an important role in either passive or active protection against pathogens (Figure 5). Notably, the adjuvant alum, which is currently used in many injectable vaccines does not effectively induce IgA class-switching.²⁴⁰ Newly developed mucosal adjuvants like TLR agonists and toxin derivatives (ADP-ribosyltransferase enterotoxins and adenylate cyclase toxins) are suggested to improve IgA responses in vaccination strategies.²⁴¹ Thus, inducing enhanced IgA responses through vaccination represents a promising therapeutic strategy for future vaccine development.




Figure 5 IgA and FcαRI as therapeutic targets. Enhanced IgA complexes or auto-antibodies result in excessive activation of immune cells contributing to chronic inflammation and tissue damage in autoimmune diseases. Blocking IgA-FcαRI interactions by either monoclonal antibodies or peptides may reduce inflammation and tissue damage in these diseases. Treatment with monomeric IgA or anti-FcαRI Fabs may dampen immune responses by inducing inhibitory ITAM signaling and inhibiting IgG-induced phagocytosis and IgE-mediated allergic diseases. To combat bacterial and viral infections, inducing pathogen-specific IgA via passive or active vaccination may result in enhanced protective immunity. Enhancing pro-inflammatory responses by FcαRI-expressing immune cells via IgA monoclonal antibody therapy, bi-specific antibodies, or cross-isotype hybrid antibodies may result in efficient killing of tumor cells and represent a promising therapeutic opportunity for cancer patients. Created with BioRender.com.

Promoting Fc α RI-Mediated Anti-Tumor Immunity

IgA or Fc α RI bispecific antibodies have been proposed as novel drugs to treat cancer by enhancing activation of Fc α RI-expressing immune cells.^{242–245} Tumor cell killing by neutrophils was superior after targeting Fc α RI, compared with targeting to IgG Fc receptors, which was demonstrated for several tumor antigens such as EGFR, HER2, EpCAM, HLA-II, CD20, CD30, and carcinoembryonic antigen.^{242,245,246} Treatment of Fc α RI transgenic mice with anti-tumor IgA mAbs against CD20, HER2, or EGFR resulted in enhanced migration of Fc α RI-expressing immune cells and significantly increased anti-tumor cytotoxicity, which was mostly mediated by Fc α RI-expressing macrophages.^{243,247,248} However, in contrast to IgG, IgA mAbs cannot activate natural killer cells, which do not express Fc α RI. Moreover, IgA is a poor complement activator and has a shorter half-life compared to IgG. IgA therapeutic antibodies that are produced in non-human systems contain different glycosylation profiles, which may also enhance immunogenicity and are likely to be cleared from the human circulation.¹² Alternatively, the efficacy of Fc α RI bispecific antibodies targeting both tumor antigens and Fc α RI has been investigated. Fc α RI bispecific antibodies efficiently induced neutrophil migration *in vitro*, which was not observed after targeting Fc γ receptors.²⁴⁹ Additionally, tumor cell killing was more effective.^{250,251} Engineering of a cross-isotype antibody that contains merged Fc domain residues from IgG1 and IgA, combining the effector functions of both isotypes, mediated increased complement-dependent cytotoxicity and Her2⁺ tumor cell killing by both neutrophils and macrophages.²⁵² Furthermore, it was shown that anti-epidermal growth factor receptor-2 IgG1/IgA2 cross-isotype antibodies induced increased recruitment of neutrophils, resulting in enhanced ADCC of human breast cancer cells by neutrophils.²⁵³ Pharmacokinetics of IgG and IgA cross-isotype antibodies were similar to the parental IgG antibodies and may therefore augment IgG-based antibody therapies.²⁵³ Thus, antibody-based strategies that target Fc α RI represent a promising therapeutic opportunity for cancer patients (Figure 5).

Inhibiting Fc α RI-Mediated Immunity

Naturally occurring serum IgA was shown to dampen immune responses by inducing ITAMi signaling via Fc α RI (Figure 1A).⁸ Enhancing ITAMi signaling by monovalent targeting of Fc α RI has been proposed as a promising strategy

to inhibit IgG-induced phagocytosis and IgE-mediated allergic diseases (Figure 5).²⁵⁴ Monovalent targeting of Fc α RI with the anti-Fc α RI mAb A77 inhibited degranulation of RBL-2H3 transfected cells.⁸ In *in vivo* models for allergic asthma and arthritis, Fc α RI transgenic mice had reduced airway inflammation or resolution of arthritis after monovalent targeting of Fc α RI.^{254,255} Additionally, targeting of Fc α RI by mAb A77 suppressed inflammation in transgenic mice with IgG immune complex-induced glomerulonephritis and obstructive nephropathy.²⁵⁶ Renal inflammation induced by CpG dinucleotides (TLR9 agonist) in Fc α RI transgenic mice was downregulated by monomeric targeting of Fc α RI as well.²⁵⁷ Thus, monomeric targeting of Fc α RI was shown to induce anti-inflammatory properties, which could be used as treatment of inflammatory diseases with involvement of myeloid cells (Figure 5). Alternatively, blocking Fc α RI with mAb MIP8a inhibited cytokine production, leukocyte recruitment, and inflammation in a lupus nephritis model.²⁵⁸ Fc α RI blocking also reduced NET release by neutrophils that had been stimulated with IgA immune complexes obtained from serum and synovial fluid of RA patients.²⁵⁹ Similarly, in an *ex vivo* LABD skin model, Fc α RI blocking with MIP8a prevented IgA-induced neutrophil-mediated blister formation.²¹⁵ In a recent study, peptides targeting the interaction sites of IgA and Fc α RI were suggested as novel therapeutic strategy for IgA-mediated skin autoimmune diseases, as these peptides were able to penetrate into human skin *ex vivo* and reduced IgA-induced neutrophil migration.²⁶⁰ As such, blocking Fc α RI-IgA interactions represents a promising therapeutic strategy for IgA-associated inflammatory diseases and autoimmunity (Figure 5).

Concluding Remarks and Future Perspectives

IgA and Fc α RI-mediated cell activation is important for maintaining homeostasis and preventing infections. SIgA contributes to passive immunity at mucosal surfaces by neutralizing pathogens, whereas active immunity is provided by dIgA and serum IgA through induction of Fc α RI-mediated activation of myeloid cells. Future research regarding vaccination and infectious diseases should therefore also include serological and functional studies of IgA and Fc α RI-mediated immune responses. It is becoming clear that the presence of overabundant IgA complexes or autoantibodies can result in excessive activation and recruitment of immune cells contributing to tissue damage and the development of

chronic inflammation in multiple diseases. A better understanding of the IgA-FcαRI axis in health and disease can provide multiple options for therapeutic interventions. On the one hand, immune responses against pathogens or cancer cells may be enhanced by vaccination or passive transfer of therapeutic IgA. On the other hand, initiating inhibitory ITAM signaling by monomeric targeting of FcαRI or blocking IgA-FcαRI interactions with mAbs or peptides can dampen inflammation and disease.

Highlights

- SIgA contributes to mucosal homeostasis and passive immunity through immune exclusion and by diversifying the gut microbiome
- Monomeric IgA induces inhibitory signaling, whereas active immunity is provided by complexed dIgA and serum IgA through cross-linking of FcαRI on myeloid cells
- Pathogen-specific IgA is associated with protection against infections
- Altered IgA glycosylation or excessive IgA immune complexes contribute to chronic inflammation and autoimmunity through FcαRI-mediated immune activation
- Understanding the IgA-FcαRI axis during infection and autoimmunity will provide new therapeutic opportunities.

Disclosure

The authors report no conflicts of interest for this work.

References

1. Belkaid Y, Harrison OJ. Homeostatic immunity and the microbiota. *Immunity*. 2017;46(4):562–576. doi:10.1016/j.immuni.2017.04.008
2. Lu LL, Suscovich TJ, Fortune SM, Alter G. Beyond binding: antibody effector functions in infectious diseases. *Nat Rev Immunol*. 2018;18(1):46–61. doi:10.1038/nri.2017.106
3. Dimitrov JD, Lacroix-Desmazes S. Noncanonical functions of antibodies. *Trends Immunol*. 2020;41(5):379–393. doi:10.1016/j.it.2020.03.006
4. Bruhns P, Jonsson F. Mouse and human FcR effector functions. *Immunol Rev*. 2015;268(1):25–51. doi:10.1111/imr.12350
5. Macpherson AJ, Yilmaz B, Limenitakis JP, Ganai-Vonarburg SC. IgA function in relation to the intestinal microbiota. *Annu Rev Immunol*. 2018;36(1):359–381. doi:10.1146/annurev-immunol-042617-053238
6. Pabst O, Slack E. IgA and the intestinal microbiota: the importance of being specific. *Mucosal Immunol*. 2020;13(1):12–21. doi:10.1038/s41385-019-0227-4
7. Leong KW, Ding JL. The unexplored roles of human serum IgA. *DNA Cell Biol*. 2014;33(12):823–829. doi:10.1089/dna.2014.2639
8. Pasquier B, Launay P, Kanamaru Y, et al. Identification of FcαRI as an inhibitory receptor that controls inflammation: dual role of FcRgamma ITAM. *Immunity*. 2005;22(1):31–42. doi:10.1016/j.immuni.2004.11.017
9. Mkaddem SB, Christou I, Rossato E, Berthelot L, Lehuen A, Monteiro RC. IgA, IgA receptors, and their anti-inflammatory properties. *Curr Top Microbiol Immunol*. 2014;382:221–235. doi:10.1007/978-3-319-07911-0_10
10. Alelyd E, Heineke MH, van Egmond M. The era of the immunoglobulin A Fc receptor FcαRI; its function and potential as target in disease. *Immunol Rev*. 2015;268(1):123–138. doi:10.1111/imr.12337
11. Davis SK, Selva KJ, Kent SJ, Chung AW. Serum IgA Fc effector functions in infectious disease and cancer. *Immunol Cell Biol*. 2020;98(4):276–286. doi:10.1111/imcb.12306
12. Breedveld A, van Egmond M. IgA and FcαRI: pathological roles and therapeutic opportunities. *Front Immunol*. 2019;10:553. doi:10.3389/fimmu.2019.00553
13. Sallustio F, Curci C, Di Leo V, Gallone A, Pesce F, Gesualdo L. A new vision of IgA nephropathy: the missing link. *Int J Mol Sci*. 2019;21(1):189. doi:10.3390/ijms21010189
14. Heineke MH, Ballering AV, Jamin A, Ben Mkaddem S, Monteiro RC, Van Egmond M. New insights in the pathogenesis of immunoglobulin A vasculitis (Henoch-Schönlein purpura). *Autoimmun Rev*. 2017;16(12):1246–1253. doi:10.1016/j.autrev.2017.10.009
15. Karimifar M, Moussavi H, Babaei M, Akbari M. The association of immunoglobulin A, immunoglobulin G and anti-cyclic citrullinated peptide antibodies with disease activity in seronegative rheumatoid arthritis patients. *J Res Med Sci*. 2014;19(9):823–826.
16. Yang YH, Huang YH, Lin YL, et al. Circulating IgA from acute stage of childhood Henoch-Schönlein purpura can enhance endothelial interleukin (IL)-8 production through MEK/ERK signalling pathway. *Clin Exp Immunol*. 2006;144(2):247–253. doi:10.1111/j.1365-2249.2006.03076.x
17. Di Niro R, Mesin L, Zheng NY, et al. High abundance of plasma cells secreting transglutaminase 2-specific IgA autoantibodies with limited somatic hypermutation in celiac disease intestinal lesions. *Nat Med*. 2012;18(3):441–445. doi:10.1038/nm.2656
18. Mercieca C, van der Horst-bruinsma IE, Borg AA. Pulmonary, renal and neurological comorbidities in patients with ankylosing spondylitis; implications for clinical practice. *Curr Rheumatol Rep*. 2014;16(8):434. doi:10.1007/s11926-014-0434-7
19. Torruellas C, French SW, Medici V. Diagnosis of alcoholic liver disease. *World J Gastroenterol*. 2014;20(33):11684–11699. doi:10.3748/wjg.v20.i33.11684
20. Lopez E, Shattock RJ, Kent SJ, Chung AW. The multifaceted nature of immunoglobulin A and its complex role in HIV. *AIDS Res Hum Retroviruses*. 2018;34(9):727–738. doi:10.1089/aid.2018.0099
21. Kremer EJ, Kalatzis V, Baker E, Callen DF, Sutherland GR, Maliszewski CR. The gene for the human IgA Fc receptor maps to 19q13.4. *Hum Genet*. 1992;89(1):107–108.
22. Nimmerjahn F, Ravetch JV. Fcγ receptors: old friends and new family members. *Immunity*. 2006;24(1):19–28. doi:10.1016/j.immuni.2005.11.010
23. Monteiro RC, Van De Winkel JG. IGAFC receptors. *Annu Rev Immunol*. 2003;21(1):177–204. doi:10.1146/annurev.immunol.21.120601.141011
24. de Wit TP, Morton HC, Capel PJ, van de Winkel JG. Structure of the gene for the human myeloid IgA Fc receptor (CD89). *J Immunol*. 1995;155(3):1203–1209.
25. Herr AB, Ballister ER, Bjorkman PJ. Insights into IgA-mediated immune responses from the crystal structures of human FcαRI and its complex with IgA1-Fc. *Nature*. 2003;423(6940):614–620. doi:10.1038/nature01685
26. Pfefferkorn LC, Yeaman GR. Association of IgA-Fc receptors (FcαRI) with FcεRI γ subunits in U937 cells. Aggregation induces the tyrosine phosphorylation of γ2. *J Immunol*. 1994;153(7):3228–3236.

27. van Egmond M, van Garderen E, van Spriel AB, et al. Fc α R1-positive liver kupffer cells: reappraisal of the function of immunoglobulin A in immunity. *Nat Med*. 2000;6(6):680–685. doi:10.1038/76261
28. Qian K, Xie F, Gibson AW, Edberg JC, Kimberly RP, Wu J. Functional expression of IgA receptor Fc α R1 on human platelets. *J Leukoc Biol*. 2008;84(6):1492–1500. doi:10.1189/jlb.0508327
29. Hansen IS, Krabbendam L, Bernink JH, et al. Fc α R1 co-stimulation converts human intestinal CD103(+) dendritic cells into pro-inflammatory cells through glycolytic reprogramming. *Nat Commun*. 2018;9(1):863. doi:10.1038/s41467-018-03318-5
30. Heystek HC, Moulon C, Woltman AM, Garonne P, van Kooten C. Human immature dendritic cells efficiently bind and take up secretory IgA without the induction of maturation. *J Immunol*. 2002;168(1):102–107.
31. Monteiro RC, Cooper MD, Kubagawa H. Molecular heterogeneity of Fc alpha receptors detected by receptor-specific monoclonal antibodies. *J Immunol*. 1992;148(6):1764–1770.
32. Reljic R. In search of the elusive mouse macrophage Fc-alpha receptor. *Immunol Lett*. 2006;107(1):80–81. doi:10.1016/j.imlet.2006.04.014
33. Bakema JE, van Egmond M. The human immunoglobulin A Fc receptor Fc α R1: a multifaceted regulator of mucosal immunity. *Mucosal Immunol*. 2011;4(6):612–624. doi:10.1038/mi.2011.36
34. Bakema JE, Hiemstra IH, Bakker J, et al. c-Jun activating binding protein 1 binds to the IgA receptor and modulates protein levels of Fc α R1 and Fc γ 3-chain. *Eur J Immunol*. 2010;40(7):2035–2040. doi:10.1002/eji.200939985
35. Hostoffer RW, Krukavets I, Berger M. Increased Fc alpha R expression and IgA-mediated function on neutrophils induced by chemoattractants. *J Immunol*. 1993;150(10):4532–4540.
36. Nikolova EB, Russell MW. Dual function of human IgA antibodies: inhibition of phagocytosis in circulating neutrophils and enhancement of responses in IL-8-stimulated cells. *J Leukoc Biol*. 1995;57(6):875–882. doi:10.1002/jlb.57.6.875
37. Yin N, Peng M, Xing Y, Zhang W. Intracellular pools of Fc α R (CD89) in human neutrophils are localized in tertiary granules and secretory vesicles, and two Fc α R isoforms are found in tertiary granules. *J Leukoc Biol*. 2007;82(3):551–558. doi:10.1189/jlb.0207112
38. Wehrli M, Schneider C, Cortinas-Elizondo F, et al. IgA triggers cell death of neutrophils when primed by inflammatory mediators. *J Immunol*. 2020.
39. Maliszewski CR, Shen L, Fanger MW. The expression of receptors for IgA on human monocytes and calcitriol-treated HL-60 cells. *J Immunol*. 1985;135(6):3878–3881.
40. Shen L, Collins JE, Schoenborn MA, Maliszewski CR. Lipopolysaccharide and cytokine augmentation of human monocyte IgA receptor expression and function. *J Immunol*. 1994;152(8):4080–4086.
41. Reterink TJ, Levarht EW, Klar-Mohamad N, Van Es LA, Daha MR. Transforming growth factor-beta 1 (TGF-beta 1) down-regulates IgA Fc-receptor (CD89) expression on human monocytes. *Clin Exp Immunol*. 1996;103(1):161–166.
42. Grossetete B, Launay P, Lehuen A, Jungers P, Bach JF, Monteiro RC. Down-regulation of Fc alpha receptors on blood cells of IgA nephropathy patients: evidence for a negative regulatory role of serum IgA. *Kidney Int*. 1998;53(5):1321–1335. doi:10.1046/j.1523-1755.1998.00885.x
43. Kawamura S, Saitou N, Ueda S. Concerted evolution of the primate immunoglobulin alpha-gene through gene conversion. *J Biol Chem*. 1992;267(11):7359–7367.
44. Pinheiro A, de Sousa-pereira P, Strive T, et al. Identification of a new European rabbit IgA with a serine-rich hinge region. *PLoS One*. 2018;13(8):e0201567.
45. de Sousa-pereira P, Woof JM. IgA: structure, function, and developability. *Antibodies (Basel)*. 2019;8(4). doi:10.3390/antib8040057
46. Steffen U, Koeleman CA, Sokolova MV, et al. IgA subclasses have different effector functions associated with distinct glycosylation profiles. *Nat Commun*. 2020;11(1):120. doi:10.1038/s41467-019-13992-8
47. Kilian M, Reinholdt J, Lomholt H, Poulsen K, Frandsen EV. Biological significance of IgA1 proteases in bacterial colonization and pathogenesis: critical evaluation of experimental evidence. *APMIS*. 1996;104(5):321–338. doi:10.1111/j.1699-0463.1996.tb00724.x
48. Tarelli E, Smith AC, Hendry BM, Challacombe SJ, Pouria S. Human serum IgA1 is substituted with up to six O-glycans as shown by matrix assisted laser desorption ionisation time-of-flight mass spectrometry. *Carbohydr Res*. 2004;339(13):2329–2335. doi:10.1016/j.carres.2004.07.011
49. Royle L, Roos A, Harvey DJ, Wormald MR, van Gijlswijk-janssen D. Secretory IgA N- and O-glycans provide a link between the innate and adaptive immune systems. *J Biol Chem*. 2003;278(22):20140–20153. doi:10.1074/jbc.M301436200
50. Novak J, Barratt J, Julian BA, Renfrow MB. Aberrant glycosylation of the IgA1 molecule in IgA nephropathy. *Semin Nephrol*. 2018;38(5):461–476. doi:10.1016/j.semnephrol.2018.05.016
51. Kerr MA. The structure and function of human IgA. *Biochem J*. 1990;271(2):285–296. doi:10.1042/bj2710285
52. Bunker JJ, Bendelac A. IgA responses to microbiota. *Immunity*. 2018;49(2):211–224. doi:10.1016/j.immuni.2018.08.011
53. Wang Y, Wang G, Li Y, et al. Structural insights into secretory immunoglobulin A and its interaction with a pneumococcal adhesin. *Cell Res*. 2020;30(7):602–609. doi:10.1038/s41422-020-0336-3
54. Johansen FE, Kaetzel CS. Regulation of the polymeric immunoglobulin receptor and IgA transport: new advances in environmental factors that stimulate pIgR expression and its role in mucosal immunity. *Mucosal Immunol*. 2011;4(6):598–602. doi:10.1038/mi.2011.37
55. Brandtzaeg P, Prydz H. Direct evidence for an integrated function of J chain and secretory component in epithelial transport of immunoglobulins. *Nature*. 1984;311(5981):71–73. doi:10.1038/311071a0
56. Kumar N, Arthur CP, Ciferri C, Matsumoto ML. Structure of the secretory immunoglobulin A core. *Science*. 2020;367(6481):1008–1014. doi:10.1126/science.aaz5807
57. Herr AB. Secret(ory) revealed: the long-awaited structures of secretory IgA. *Cell Res*. 2020;30(7):558–559. doi:10.1038/s41422-020-0351-4
58. Van Spriel AB, Leusen JH, Vile H. Mac-1 (CD11b/CD18) as accessory molecule for Fc alpha R (CD89) binding of IgA. *J Immunol*. 2002;169(7):3831–3836. doi:10.4049/jimmunol.169.7.3831
59. Reterink TJ, van Zandbergen G, van Egmond M, et al. Size-dependent effect of IgA on the IgA Fc receptor (CD89). *Eur J Immunol*. 1997;27(9):2219–2224. doi:10.1002/eji.1830270915
60. Pleass RJ, Dehal PK, Lewis MJ, Woof JM. Limited role of charge matching in the interaction of human immunoglobulin A with the immunoglobulin A Fc receptor (Fc α R1) CD89. *Immunology*. 2003;109(3):331–335. doi:10.1046/j.1365-2567.2003.01677.x
61. Posgai MT, Tondast-Navaei S, Jayasinghe M, Ibrahim GM, Stan G, Herr AB. Fc α R1 binding at the IgA1 CH2-CH3 interface induces long-range conformational changes that are transmitted to the hinge region. *Proc Natl Acad Sci U S A*. 2018;115(38):E8882–E91. doi:10.1073/pnas.1807478115
62. Zhang Y, Boesen CC, Radaev S, et al. Crystal structure of the extracellular domain of a human Fc gamma RIII. *Immunity*. 2000;13(3):387–395. doi:10.1016/S1074-7613(00)00038-8

63. Keown MB, Henry AJ, Ghirlando R, Sutton BJ, Gould HJ. Thermodynamics of the interaction of human immunoglobulin E with its high-affinity receptor Fc epsilon RI. *Biochemistry*. 1998;37(25):8863–8869. doi:10.1021/bi972354h
64. Keown MB, Ghirlando R, Mackay GA, Sutton BJ, Gould HJ. Basis of the 1:1 stoichiometry of the high affinity receptor Fc epsilon RI-IgE complex. *Eur Biophys J*. 1997;25(5–6):471–476. doi:10.1007/s002490050062
65. Sondermann P, Huber R, Oosthuizen V, Jacob U. The 3.2-A crystal structure of the human IgG1 Fc fragment-Fc gammaRIII complex. *Nature*. 2000;406(6793):267–273. doi:10.1038/35018508
66. Bonner A, Furtado PB, Almogren A, Kerr MA, Perkins SJ. Implications of the near-planar solution structure of human myeloma dimeric IgA1 for mucosal immunity and IgA nephropathy. *J Immunol*. 2008;180(2):1008–1018. doi:10.4049/jimmunol.180.2.1008
67. Lu J, Marjon KD, Mold C, Marnell L, Du Clos TW, Sun P. Pentraxins and IgA share a binding hot-spot on Fc alphaRI. *Protein Sci*. 2014;23(4):378–386. doi:10.1002/pro.2419
68. Xue J, Zhao Q, Zhu L, Zhang W. Deglycosylation of Fc alphaR at N58 increases its binding to IgA. *Glycobiology*. 2010;20(7):905–915. doi:10.1093/glycob/cwq048
69. Bakema JE, Bakker A, de Haij S, Honing H, Bracke M, Koenderman L. Inside-out regulation of Fc alpha RI (CD89) depends on PP2A. *J Immunol*. 2008;181(6):4080–4088. doi:10.1038/s41467-017-00294-0
70. Weisbart RH, Kacena A, Schuh A, Golde DW. GM-CSF induces human neutrophil IgA-mediated phagocytosis by an IgA Fc receptor activation mechanism. *Nature*. 1988;332(6165):647–648. doi:10.1038/332647a0
71. Bracke M, Dubois GR, Bolt K, et al. Differential effects of the T helper cell type 2-derived cytokines IL-4 and IL-5 on ligand binding to IgG and IgA receptors expressed by human eosinophils. *J Immunol*. 1997;159(3):1459–1465.
72. Morton HC, van den Herik-oudijk IE, Vossebeeld P, et al. Functional association between the human myeloid immunoglobulin A Fc receptor (CD89) and FcR gamma chain. Molecular basis for CD89/FcR gamma chain association. *J Biol Chem*. 1995;270(50):29781–29787. doi:10.1074/jbc.270.50.29781
73. Pfirsch-Maisonnas S, Aloulou M, Xu T, et al. Inhibitory ITAM signaling traps activating receptors with the phosphatase SHP-1 to form polarized “inhibisome” clusters. *Sci Signal*. 2011;4(169):ra24. doi:10.1126/scisignal.2001309
74. Lang ML, Chen YW, Shen L, et al. IgA Fc receptor (FcalphaR) cross-linking recruits tyrosine kinases, phosphoinositide kinases and serine/threonine kinases to glycolipid rafts. *Biochem J*. 2002;364(Pt 2):517–525. doi:10.1042/bj20011696
75. Gulle H, Samstag A, Eibl MM, Wolf HM. Physical and functional association of Fc alpha R with protein tyrosine kinase Lyn. *Blood*. 1998;91(2):383–391. doi:10.1182/blood.V91.2.383
76. Mkaddem SB, Murua A, Flament H, et al. Lyn and Fyn function as molecular switches that control immunoreceptors to direct homeostasis or inflammation. *Nat Commun*. 2017;8(1):246.
77. Getahun A, Cambier JC. Of ITIMs, ITAMs, and ITAMis: revisiting immunoglobulin Fc receptor signaling. *Immunol Rev*. 2015;268(1):66–73. doi:10.1111/imr.12336
78. Aleyd E, van Hout MW, Ganzevles SH, et al. IgA enhances NETosis and release of neutrophil extracellular traps by polymorphonuclear cells via Fc alpha receptor I. *J Immunol*. 2014;192(5):2374–2383. doi:10.4049/jimmunol.1300261
79. van der Steen L, Tuk CW, Bakema JE, et al. Immunoglobulin A: Fc(alpha)RI interactions induce neutrophil migration through release of leukotriene B4. *Gastroenterology*. 2009;137(6):2018–29e1–3. doi:10.1053/j.gastro.2009.06.047
80. Vidarsson G, van Der Pol WL, van Den Elsen JM, et al. Activity of human IgG and IgA subclasses in immune defense against neisseria meningitidis serogroup B. *J Immunol*. 2001;166(10):6250–6256. doi:10.4049/jimmunol.166.10.6250
81. Geissmann F, Launay P, Pasquier B, et al. A subset of human dendritic cells expresses IgA Fc receptor (CD89), which mediates internalization and activation upon cross-linking by IgA complexes. *J Immunol*. 2001;166(1):346–352. doi:10.4049/jimmunol.166.1.346
82. Chen YW, Lang ML, Wade WF. Protein kinase C-alpha and -delta are required for Fc alphaR (CD89) trafficking to MHC class II compartments and Fc alphaR-mediated antigen presentation. *Traffic*. 2004;5(8):577–594. doi:10.1111/j.1600-0854.2004.00202.x
83. Pasquier B, Lepelletier Y, Baude C, Hermine O, Monteiro RC. Differential expression and function of IgA receptors (CD89 and CD71) during maturation of dendritic cells. *J Leukoc Biol*. 2004;76(6):1134–1141. doi:10.1189/jlb.0204101
84. Fayette J, Dubois B, Vandenabeele S, et al. Human dendritic cells skew isotype switching of CD40-activated naive B cells towards IgA1 and IgA2. *J Exp Med*. 1997;185(11):1909–1918. doi:10.1084/jem.185.11.1909
85. Baumann J, Park CG, Mantis NJ. Recognition of secretory IgA by DC-SIGN: implications for immune surveillance in the intestine. *Immunol Lett*. 2010;131(1):59–66. doi:10.1016/j.imlet.2010.03.005
86. Cortes B. Multi-faceted functions of secretory IgA at mucosal surfaces. *Front Immunol*. 2013;4:185. doi:10.3389/fimmu.2013.00185
87. Rogier EW, Frantz AL, Bruno ME, Kaetzel CS. Secretory IgA is concentrated in the outer layer of colonic mucus along with gut bacteria. *Pathogens*. 2014;3(2):390–403. doi:10.3390/pathogens3020390
88. Mantis NJ, Rol N, Cortes B. Secretory IgA's complex roles in immunity and mucosal homeostasis in the gut. *Mucosal Immunol*. 2011;4(6):603–611. doi:10.1038/mi.2011.41
89. Yan H, Lamm ME, Bjorling E, Huang YT. Multiple functions of immunoglobulin A in mucosal defense against viruses: an in vitro measles virus model. *J Virol*. 2002;76(21):10972–10979. doi:10.1128/JVI.76.21.10972-10979.2002
90. Mazanec MB, Kaetzel CS, Lamm ME, Fletcher D, Nedrud JG. Intracellular neutralization of virus by immunoglobulin A antibodies. *Proc Natl Acad Sci U S A*. 1992;89(15):6901–6905. doi:10.1073/pnas.89.15.6901
91. Mantis NJ, Palaia J, Hessel AJ, et al. Inhibition of HIV-1 infectivity and epithelial cell transfer by human monoclonal IgG and IgA antibodies carrying the b12 V region. *J Immunol*. 2007;179(5):3144–3152. doi:10.4049/jimmunol.179.5.3144
92. Wright A, Lamm ME, Huang YT. Excretion of human immunodeficiency virus type 1 through polarized epithelium by immunoglobulin A. *J Virol*. 2008;82(23):11526–11535. doi:10.1128/JVI.01111-08
93. Kulkarni V, Ruprecht RM. Mucosal IgA responses: damaged in established HIV infection-yet, effective weapon against HIV transmission. *Front Immunol*. 2017;8:1581. doi:10.3389/fimmu.2017.01581
94. Blutt SE, Conner ME. The gastrointestinal frontier: IgA and viruses. *Front Immunol*. 2013;4:402. doi:10.3389/fimmu.2013.00402
95. Sutherland DB, Suzuki K, Fagarasan S. Fostering of advanced mutualism with gut microbiota by immunoglobulin A. *Immunol Rev*. 2016;270(1):20–31. doi:10.1111/imr.12384
96. Kawamoto S, Maruya M, Kato LM, et al. Foxp3(+) T cells regulate immunoglobulin A selection and facilitate diversification of bacterial species responsible for immune homeostasis. *Immunity*. 2014;41(1):152–165. doi:10.1016/j.immuni.2014.05.016

97. Wijburg OL, Uren TK, Simpfendorfer K, Johansen FE, Brandtzaeg P, Strugnell RA. Innate secretory antibodies protect against natural salmonella typhimurium infection. *J Exp Med*. 2006;203(1):21–26. doi:10.1084/jem.20052093
98. Reikvam DH, Derrien M, Islam R, et al. Epithelial-microbial crosstalk in polymeric Ig receptor deficient mice. *Eur J Immunol*. 2012;42(11):2959–2970. doi:10.1002/eji.201242543
99. Catanzaro JR, Strauss JD, Bielecka A, et al. IgA-deficient humans exhibit gut microbiota dysbiosis despite secretion of compensatory IgM. *Sci Rep*. 2019;9(1):13574. doi:10.1038/s41598-019-49923-2
100. Kabbert J, Benckert J, Rollenske T, et al. High microbiota reactivity of adult human intestinal IgA requires somatic mutations. *J Exp Med*. 2020;217(11):11. doi:10.1084/jem.20200275
101. Bunker JJ, Erickson SA, Flynn TM, et al. Natural polyreactive IgA antibodies coat the intestinal microbiota. *Science*. 2017;358(6361):6361. doi:10.1126/science.aan6619
102. Nakajima A, Vogelzang A, Maruya M, et al. IgA regulates the composition and metabolic function of gut microbiota by promoting symbiosis between bacteria. *J Exp Med*. 2018;215(8):2019–2034. doi:10.1084/jem.20180427
103. Wilmore JR, Gaudette BT, Gomez Atria D, et al. Commensal microbes induce serum IgA responses that protect against polymicrobial sepsis. *Cell Host Microbe*. 2018;23(3):302–11 e3. doi:10.1016/j.chom.2018.01.005
104. Kozakova H, Schwarzer M, Tuckova L, et al. Colonization of germ-free mice with a mixture of three lactobacillus strains enhances the integrity of gut mucosa and ameliorates allergic sensitization. *Cell Mol Immunol*. 2016;13(2):251–262. doi:10.1038/cmi.2015.09
105. Matysiak-Budnik T, Moura IC, Arcos-Fajardo M, et al. Secretory IgA mediates retrotranscytosis of intact gliadin peptides via the transferrin receptor in celiac disease. *J Exp Med*. 2008;205(1):143–154. doi:10.1084/jem.20071204
106. Rochereau N, Drocourt D, Perouzel E, et al. Dectin-1 is essential for reverse transcytosis of glycosylated SIgA-antigen complexes by intestinal M cells. *PLoS Biol*. 2013;11(9):e1001658. doi:10.1371/journal.pbio.1001658
107. Alpan O, Rudomen G, Matzinger P. The role of dendritic cells, B cells, and M cells in gut-oriented immune responses. *J Immunol*. 2001;166(8):4843–4852. doi:10.4049/jimmunol.166.8.4843
108. Van Epps DE, Williams RC Jr. Suppression of leukocyte chemotaxis by human IgA myeloma components. *J Exp Med*. 1976;144(5):1227–1242. doi:10.1084/jem.144.5.1227
109. Wilton JM. Suppression by IgA of IgG-mediated phagocytosis by human polymorphonuclear leucocytes. *Clin Exp Immunol*. 1978;34(3):423–428.
110. Wolf HM, Fischer MB, Pühringer H, Samstag A, Vogel E, Eibl MM. Human serum IgA downregulates the release of inflammatory cytokines (tumor necrosis factor- α , interleukin-6) in human monocytes. *Blood*. 1994;83(5):1278–1288. doi:10.1182/blood.V83.5.1278.1278
111. Olas K, Butterweck H, Teschner W, Schwarz HP, Reipert B. Immunomodulatory properties of human serum immunoglobulin A: anti-inflammatory and pro-inflammatory activities in human monocytes and peripheral blood mononuclear cells. *Clin Exp Immunol*. 2005;140(3):478–490. doi:10.1111/j.1365-2249.2005.02779.x
112. Rogier EW, Frantz AL, Bruno ME, et al. Secretory antibodies in breast milk promote long-term intestinal homeostasis by regulating the gut microbiota and host gene expression. *Proc Natl Acad Sci U S A*. 2014;111(8):3074–3079. doi:10.1073/pnas.1315792111
113. Hurley WL, Theil PK. Perspectives on immunoglobulins in colostrum and milk. *Nutrients*. 2011;3(4):442–474. doi:10.3390/nu3040442
114. Gopalakrishna KP, Hand TW. Influence of maternal milk on the neonatal intestinal microbiome. *Nutrients*. 2020;12(3):823. doi:10.3390/nu12030823
115. Jones G, Steketee RW, Black RE, Bhutta ZA, Morris SS. Bellagio child survival study G. How many child deaths can we prevent this year? *Lancet*. 2003;362(9377):65–71. doi:10.1016/S0140-6736(03)13811-1
116. Gopalakrishna KP, Macadangdang BR, Rogers MB, et al. Maternal IgA protects against the development of necrotizing enterocolitis in preterm infants. *Nat Med*. 2019;25(7):1110–1115. doi:10.1038/s41591-019-0480-9
117. Rognum TO, Thrane S, Stoltenberg L, Vege A, Brandtzaeg P. Development of intestinal mucosal immunity in fetal life and the first postnatal months. *Pediatr Res*. 1992;32(2):145–149. doi:10.1203/00006450-199208000-00003
118. Brandtzaeg P, Nilssen DE, Rognum TO, Thrane PS. Ontogeny of the mucosal immune system and IgA deficiency. *Gastroenterol Clin North Am*. 1991;20(3):397–439.
119. Blanco E, Perez-Andres M, Arriba-Mendez S, et al. Age-associated distribution of normal B-cell and plasma cell subsets in peripheral blood. *J Allergy Clin Immunol*. 2018;141(6):2208–19 e16. doi:10.1016/j.jaci.2018.02.017
120. Miller EM, McConnell DS. Milk immunity and reproductive status among Ariaal women of northern Kenya. *Ann Hum Biol*. 2015;42(1):76–83. doi:10.3109/03014460.2014.941398
121. Fujita M, Wander K, Paredes Ruvalcaba N, Brindle E. Human milk sIgA antibody in relation to maternal nutrition and infant vulnerability in northern Kenya. *Evol Med Public Health*. 2019;2019(1):201–211. doi:10.1093/emph/eoz030
122. Gorter A, Hiemstra PS, Leijh PC, et al. IgA- and secretory IgA-opsonized *S. aureus* induce a respiratory burst and phagocytosis by polymorphonuclear leucocytes. *Immunology*. 1987;61(3):303–309.
123. Duc M, Johansen F-E, Cortes B. Antigen binding to secretory immunoglobulin A results in decreased sensitivity to intestinal proteases and increased binding to cellular Fc receptors. *J Biol Chem*. 2010;285(2):953–960. doi:10.1074/jbc.M109.059220
124. Trend S, de Jong E, Lloyd ML, et al. Leukocyte populations in human preterm and term breast milk identified by multicolour flow cytometry. *PLoS One*. 2015;10(8):e0135580. doi:10.1371/journal.pone.0135580
125. Honorio-Franca AC, Launay P, Carneiro-Sampaio MM, Monteiro RC. Colostral neutrophils express Fc alpha receptors (CD89) lacking gamma chain association and mediate noninflammatory properties of secretory IgA. *J Leukoc Biol*. 2001;69(2):289–296.
126. Zimmermann N, Thormann V, Hu B, et al. Human isotype-dependent inhibitory antibody responses against *Mycobacterium tuberculosis*. *EMBO Mol Med*. 2016;8(11):1325–1339. doi:10.15252/emmm.201606330
127. Magri G, Cerutti A. IgA summons IgG to take a hit at HIV-1. *Cell Host Microbe*. 2020;27(6):854–856. doi:10.1016/j.chom.2020.05.017
128. Chao YX, Rotzschke O, Tan EK. The role of IgA in COVID-19. *Brain Behav Immun*. 2020;87:182–183. doi:10.1016/j.bbi.2020.05.057
129. Riley RL, Mills CC, Nyka W, et al. Aerial dissemination of pulmonary tuberculosis. A two-year study of contagion in a tuberculosis ward. 1959. *Am J Epidemiol*. 1995;142(1):3–14. doi:10.1093/oxfordjournals.aje.a117542
130. Bijlsma MW, Bekker V, Brouwer MC, Spanjaard L, van de Beek D, van der Ende A. Epidemiology of invasive meningococcal disease in the Netherlands, 1960–2012: an analysis of national surveillance data. *Lancet Infect Dis*. 2014;14(9):805–812. doi:10.1016/S1473-3099(14)70806-0

131. Zhang Q, Finn A. Mucosal immunology of vaccines against pathogenic nasopharyngeal bacteria. *J Clin Pathol.* 2004;57(10):1015–1021. doi:10.1136/jcp.2004.016253
132. Tuomanen EI, Austrian R, Masure HR. Pathogenesis of pneumococcal infection. *N Engl J Med.* 1995;332(19):1280–1284. doi:10.1056/NEJM199505113321907
133. Musher DM, Groover JE, Reichler MR, et al. Emergence of antibody to capsular polysaccharides of *Streptococcus pneumoniae* during outbreaks of pneumonia: association with nasopharyngeal colonization. *Clin Infect Dis.* 1997;24(3):441–446. doi:10.1093/clinids/24.3.441
134. Wu M, Zhao H, Li M, Yue Y, Xiong S, Xu W. Intranasal vaccination with mannoseylated chitosan formulated DNA vaccine enables robust IgA and cellular response induction in the lungs of mice and improves protection against pulmonary mycobacterial challenge. *Front Cell Infect Microbiol.* 2017;7:445. doi:10.3389/fcimb.2017.00445
135. Ferreira DM, Darrieux M, Silva DA, et al. Characterization of protective mucosal and systemic immune responses elicited by pneumococcal surface protein PspA and PspC nasal vaccines against a respiratory pneumococcal challenge in mice. *Clin Vaccine Immunol.* 2009;16(5):636–645. doi:10.1128/CVI.00395-08
136. Jia M, Liberatore RA, Guo Y, et al. VSV-displayed HIV-1 envelope identifies broadly neutralizing antibodies class-switched to IgG and IgA. *Cell Host Microbe.* 2020;27(6):963–75 e5. doi:10.1016/j.chom.2020.03.024
137. Watkins JD, Sholukh AM, Mukhtar MM, et al. Anti-HIV IgA isotypes: differential virion capture and inhibition of transcytosis are linked to prevention of mucosal R5 SHIV transmission. *AIDS.* 2013;27(9):F13–20. doi:10.1097/QAD.0b013e328360eac6
138. Sholukh AM, Watkins JD, Vyas HK, et al. Defense-in-depth by mucosally administered anti-HIV dimeric IgA2 and systemic IgG1 mAbs: complete protection of rhesus monkeys from mucosal SHIV challenge. *Vaccine.* 2015;33(17):2086–2095. doi:10.1016/j.vaccine.2015.02.020
139. Johansen K, Svensson L. Neutralization of rotavirus and recognition of immunologically important epitopes on VP4 and VP7 by human IgA. *Arch Virol.* 1997;142(7):1491–1498. doi:10.1007/s007050050175
140. To TL, Ward LA, Yuan L, Saif LJ. Serum and intestinal isotype antibody responses and correlates of protective immunity to human rotavirus in a gnotobiotic pig model of disease. *J Gen Virol.* 1998;79(Pt 11):2661–2672.
141. Crawford SE, Patel DG, Cheng E, et al. Rotavirus viremia and extraintestinal viral infection in the neonatal rat model. *J Virol.* 2006;80(10):4820–4832. doi:10.1128/JVI.80.10.4820-4832.2006
142. Feng N, Burns JW, Bracy L, Greenberg HB. Comparison of mucosal and systemic humoral immune responses and subsequent protection in mice orally inoculated with a homologous or a heterologous rotavirus. *J Virol.* 1994;68(12):7766–7773. doi:10.1128/JVI.68.12.7766-7773.1994
143. Iyer AS, Jones FK, Nodoushani A, et al. Persistence and decay of human antibody responses to the receptor binding domain of SARS-CoV-2 spike protein in COVID-19 patients. *Sci Immunol.* 2020;5(52):52. doi:10.1126/sciimmunol.abe0367
144. Kowitdamrong E, Puthanakit T, Jantarabenjakul W, et al. Antibody responses to SARS-CoV-2 in patients with differing severities of coronavirus disease 2019. *PLoS One.* 2020;15(10):e0240502. doi:10.1371/journal.pone.0240502
145. Sterlin D, Mathian A, Miyara M, et al. IgA dominates the early neutralizing antibody response to SARS-CoV-2. *medRxiv.* 2020;2020.06.10.20126532.
146. Xue M, Zhang T, Hu H, et al. Predictive effects of IgA and IgG combination to assess pulmonary exudation progression in COVID-19 patients. *J Med Virol.* 2020. doi:10.1002/jmv.26437
147. Ejemel M, Li Q, Hou S, et al. A cross-reactive human IgA monoclonal antibody blocks SARS-CoV-2 spike-ACE2 interaction. *Nat Commun.* 2020;11(1):4198. doi:10.1038/s41467-020-18058-8
148. Hasan Ali O, Bomze D, Risch L, et al. Severe COVID-19 is associated with elevated serum IgA and antiphospholipid IgA-antibodies. *Clin Infect Dis.* 2020. doi:10.1093/cid/ciaa1496
149. Yu HQ, Sun BQ, Fang ZF, et al. Distinct features of SARS-CoV-2-specific IgA response in COVID-19 patients. *Eur Respir J.* 2020;56(2):2001526. doi:10.1183/13993003.01526-2020
150. Fourati S, Hue S, Pawlotsky JM, Mekontso-Dessap A, de Prost N. SARS-CoV-2 viral loads and serum IgA/IgG immune responses in critically ill COVID-19 patients. *Intensive Care Med.* 2020;46(9):1781–1783. doi:10.1007/s00134-020-06157-5
151. Casadevall A, Pirofski LA. The convalescent sera option for containing COVID-19. *J Clin Invest.* 2020;130(4):1545–1548. doi:10.1172/JCI138003
152. Li L, Zhang W, Hu Y, et al. Effect of convalescent plasma therapy on time to clinical improvement in patients with severe and life-threatening COVID-19: a randomized clinical trial. *JAMA.* 2020;324(5):460–470. doi:10.1001/jama.2020.10044
153. Liu STH, Lin HM, Baine I, et al. Convalescent plasma treatment of severe COVID-19: a propensity score-matched control study. *Nat Med.* 2020;26(11):1708–1713. doi:10.1038/s41591-020-1088-9
154. Ye M, Fu D, Ren Y, et al. Treatment with convalescent plasma for COVID-19 patients in Wuhan, China. *J Med Virol.* 2020;92(10):1890–1901. doi:10.1002/jmv.25882
155. Salazar E, Perez KK, Ashraf M, et al. Treatment of coronavirus disease 2019 (COVID-19) patients with convalescent plasma. *Am J Pathol.* 2020;190(8):1680–1690. doi:10.1016/j.ajpath.2020.05.014
156. Perotti C, Baldanti F, Bruno R, et al. Mortality reduction in 46 severe Covid-19 patients treated with hyperimmune plasma. A proof of concept single arm multicenter trial. *Haematologica.* 2020. doi:10.3324/haematol.2020.261784
157. Nauseef WM, Borregaard N. Neutrophils at work. *Nat Immunol.* 2014;15(7):602–611. doi:10.1038/ni.2921
158. Hellwig SM, van Sriel AB, Schellekens JF, Mooi FR, van de Winkel JG. Immunoglobulin A-mediated protection against bordetella pertussis infection. *Infect Immun.* 2001;69(8):4846–4850. doi:10.1128/IAI.69.8.4846-4850.2001
159. van Sriel AB, van den Herik-oudijk IE, van Sorge NM, Vile HA, van Strijp JA, van de Winkel J. Effective phagocytosis and killing of *Candida albicans* via targeting FcγRI (CD64) or FcαRI (CD89) on neutrophils. *J Infect Dis.* 1999;179(3):661–669. doi:10.1086/314643
160. van der Pol W, Vidarsson G, Vile HA, van de Winkel JG, Rodriguez ME. Pneumococcal capsular polysaccharide-specific IgA triggers efficient neutrophil effector functions via FcαRI (CD89). *J Infect Dis.* 2000;182(4):1139–1145. doi:10.1086/315825
161. Kobayashi T, Yamamoto K, Sugita N, et al. Effective in vitro clearance of *Porphyromonas gingivalis* by Fcα receptor I (CD89) on gingival crevicular neutrophils. *Infect Immun.* 2001;69(5):2935–2942. doi:10.1128/IAI.69.5.2935-2942.2001
162. Shin EH, Lee HY, Bae YS. Leukotriene B4 stimulates human monocyte-derived dendritic cell chemotaxis. *Biochem Biophys Res Commun.* 2006;348(2):606–611. doi:10.1016/j.bbrc.2006.07.084
163. Subramanian BC, Majumdar R, Parent CA. The role of the LTB4-BLT1 axis in chemotactic gradient sensing and directed leukocyte migration. *Semin Immunol.* 2017;33:16–29. doi:10.1016/j.smim.2017.07.002
164. Patry C, Sibille Y, Lehuen A, Monteiro RC. Identification of Fcα receptor (CD89) isoforms generated by alternative splicing that are differentially expressed between blood monocytes and alveolar macrophages. *J Immunol.* 1996;156(11):4442–4448.

165. Lu LL, Chung AW, Rosebrock TR, et al. A functional role for antibodies in tuberculosis. *Cell*. 2016;167(2):433–43 e14. doi:10.1016/j.cell.2016.08.072
166. Legesse M, Ameni G, Medhin G, et al. IgA response to ESAT-6/CFP-10 and Rv2031 antigens varies in patients with culture-confirmed pulmonary tuberculosis, healthy Mycobacterium tuberculosis-infected and non-infected individuals in a tuberculosis endemic setting, Ethiopia. *Scand J Immunol*. 2013;78(3):266–274. doi:10.1111/sji.12080
167. de Tymowski C, Heming N, Correia MDT, et al. CD89 is a potent innate receptor for bacteria and mediates host protection from sepsis. *Cell Rep*. 2019;27(3):762–75 e5. doi:10.1016/j.celrep.2019.03.062
168. Tay MZ, Wiehe K, Pollara J. Antibody-dependent cellular phagocytosis in antiviral immune responses. *Front Immunol*. 2019;10:332. doi:10.3389/fimmu.2019.00332
169. Ackerman ME, Mikhailova A, Brown EP, et al. Polyfunctional HIV-specific antibody responses are associated with spontaneous HIV control. *PLoS Pathog*. 2016;12(1):e1005315. doi:10.1371/journal.ppat.1005315
170. Duchemin M, Khamassi M, Xu L, Tudor D, Bomsel M. IgA targeting human immunodeficiency virus-1 envelope gp41 triggers antibody-dependent cellular cytotoxicity cross-clade and cooperates with gp41-specific IgG to increase cell lysis. *Front Immunol*. 2018;9:244. doi:10.3389/fimmu.2018.00244
171. Duchemin M, Tudor D, Cottignies-Calamarte A, Bomsel M. Antibody-dependent cellular phagocytosis of HIV-1-infected cells is efficiently triggered by IgA targeting HIV-1 envelope subunit gp41. *Front Immunol*. 2020;11:1141. doi:10.3389/fimmu.2020.01141
172. Ackerman ME, Das J, Pittala S, et al. Route of immunization defines multiple mechanisms of vaccine-mediated protection against SIV. *Nat Med*. 2018;24(10):1590–1598. doi:10.1038/s41591-018-0161-0
173. Rerks-Ngarm S, Pitisuttithum P, Nitayaphan S, et al. Vaccination with ALVAC and AIDSVAX to prevent HIV-1 infection in Thailand. *N Engl J Med*. 2009;361(23):2209–2220. doi:10.1056/NEJMoa0908492
174. Uren TK, Johansen FE, Wijburg OL, Koentgen F, Brandtzaeg P, Strugnell RA. Role of the polymeric Ig receptor in mucosal B cell homeostasis. *J Immunol*. 2003;170(5):2531–2539. doi:10.4049/jimmunol.170.5.2531
175. Hansen IS, Hoepel W, Zaat SAJ, Baeten DLP, den Dunnen J. Serum IgA immune complexes promote proinflammatory cytokine production by human macrophages, monocytes, and kupffer cells through FcαRI-TLR cross-talk. *J Immunol*. 2017;199(12):4124–4131. doi:10.4049/jimmunol.1700883
176. Hansen IS, Baeten DLP, den Dunnen J. The inflammatory function of human IgA. *Cell Mol Life Sci*. 2019;76(6):1041–1055. doi:10.1007/s00018-018-2976-8
177. Woof JM. Immunoglobulins and their receptors, and subversion of their protective roles by bacterial pathogens. *Biochem Soc Trans*. 2016;44(6):1651–1658. doi:10.1042/BST20160246
178. Fasching CE, Grossman T, Corthesy B, Plaut AG, Weiser JN, Janoff EN. Impact of the molecular form of immunoglobulin A on functional activity in defense against Streptococcus pneumoniae. *Infect Immun*. 2007;75(4):1801–1810. doi:10.1128/IAI.01758-06
179. Janoff EN, Rubins JB, Fasching C, et al. Pneumococcal IgA1 protease subverts specific protection by human IgA1. *Mucosal Immunol*. 2014;7(2):249–256. doi:10.1038/mi.2013.41
180. Lin L, Ayala P, Larson J, et al. The Neisseria type 2 IgA1 protease cleaves LAMP1 and promotes survival of bacteria within epithelial cells. *Mol Microbiol*. 1997;24(5):1083–1094. doi:10.1046/j.1365-2958.1997.4191776.x
181. Vidarsson G, Overbeeke N, Stemerding AM, et al. Working mechanism of immunoglobulin A1 (IgA1) protease: cleavage of IgA1 antibody to neisseria meningitidis PorA requires de novo synthesis of IgA1 protease. *Infect Immun*. 2005;73(10):6721–6726. doi:10.1128/IAI.73.10.6721-6726.2005
182. Polissi A, Pontiggia A, Feger G, et al. Large-scale identification of virulence genes from Streptococcus pneumoniae. *Infect Immun*. 1998;66(12):5620–5629. doi:10.1128/IAI.66.12.5620-5629.1998
183. Pleass RJ, Areschoug T, Lindahl G, Woof JM. Streptococcal IgA-binding proteins bind in the Calpha 2-Calpha 3 interdomain region and inhibit binding of IgA to human CD89. *J Biol Chem*. 2001;276(11):8197–8204. doi:10.1074/jbc.M009396200
184. Ramsland PA, Willoughby N, Trist HM, et al. Structural basis for evasion of IgA immunity by staphylococcus aureus revealed in the complex of SSL7 with Fc of human IgA1. *Proc Natl Acad Sci U S A*. 2007;104(38):15051–15056. doi:10.1073/pnas.0706028104
185. D'Amico G. Natural history of idiopathic IgA nephropathy and factors predictive of disease outcome. *Semin Nephrol*. 2020;12(3):823–896. doi:10.1016/j.semnephrol.2004.01.001
186. McGrogan A, Franssen CF, de Vries CS. The incidence of primary glomerulonephritis worldwide: a systematic review of the literature. *Nephrol Dial Transplant*. 2011;26(2):414–430. doi:10.1093/ndt/gfq665
187. Novak J, Julian BA, Tomana M, Mestecky J. IgA glycosylation and IgA immune complexes in the pathogenesis of IgA nephropathy. *Semin Nephrol*. 2019;10(1):78–87. doi:10.1016/j.semnephrol.2007.10.009
188. Chen A, Yang SS, Lin TJ, Ka SM. IgA nephropathy: clearance kinetics of IgA-containing immune complexes. *Semin Immunopathol*. 2018;40(6):539–543. doi:10.1007/s00281-018-0708-7
189. Gutierrez E, Carvaca-Fontan F, Luzardo L, Morales E, Alonso M, Praga M. A personalized update on IgA nephropathy: a new vision and new future challenges. *Nephron*. 2020;144(11):555–571. doi:10.1159/000509997
190. Berthelot L, Papista C, Maciel TT, et al. Transglutaminase is essential for IgA nephropathy development acting through IgA receptors. *J Exp Med*. 2012;209(4):793–806. doi:10.1084/jem.20112005
191. Launay P, Grossetete B, Arcos-Fajardo M, et al. FcαRI (CD89) mediates the development of immunoglobulin A (IgA) nephropathy (Berger's disease). Evidence for pathogenic soluble receptor-IgA complexes in patients and CD89 transgenic mice. *J Exp Med*. 2000;191(11):1999–2009. doi:10.1084/jem.191.11.1999
192. Rifai A, Fadden K, Morrison SL, Chintalacharuvu KR. The N-glycans determine the differential blood clearance and hepatic uptake of human immunoglobulin (Ig)A1 and IgA2 isotypes. *J Exp Med*. 2020;27(6):963–975.e5. doi:10.1084/jem.191.12.2171
193. Xu L, Li B, Huang M, et al. Critical role of kupffer cell CD89 expression in experimental IgA nephropathy. *PLoS One*. 2016;11(7):e0159426.
194. Aletaha D, Neogi T, Silman AJ, et al. Rheumatoid arthritis classification criteria: an American college of rheumatology/European league against rheumatism collaborative initiative. *Arthritis Rheum*. 2010;62(9):2569–2581. doi:10.1002/art.27584
195. Moez S, John P, Bhatti A. Anti-citrullinated protein antibodies: role in pathogenesis of RA and potential as a diagnostic tool. *Rheumatol Int*. 2013;33(7):1669–1673. doi:10.1007/s00296-012-2635-6
196. Jonsson T, Arinbjarnarson S, Thorsteinsson J, et al. Raised IgA rheumatoid factor (RF) but not IgM RF or IgG RF is associated with extra-articular manifestations in rheumatoid arthritis. *Scand J Rheumatol*. 1995;24(6):372–375. doi:10.3109/03009749509095183

197. Bobbio-Pallavicini F, Caporali R, Alpini C, et al. High IgA rheumatoid factor levels are associated with poor clinical response to tumour necrosis factor alpha inhibitors in rheumatoid arthritis. *Ann Rheum Dis*. 2007;66(3):302–307. doi:10.1136/ard.2006.060608
198. Nielen MM, van Schaardenburg D, Reesink HW, et al. Specific autoantibodies precede the symptoms of rheumatoid arthritis: a study of serial measurements in blood donors. *Arthritis Rheum*. 2004;50(2):380–386. doi:10.1002/art.20018
199. Rantapaa-Dahlqvist S, de Jong BA, Berglin E, et al. Antibodies against cyclic citrullinated peptide and IgA rheumatoid factor predict the development of rheumatoid arthritis. *Arthritis Rheum*. 2003;48(10):2741–2749. doi:10.1002/art.11223
200. Kinslow JD, Blum LK, Deane KD, et al. Elevated IgA plasma-blast levels in subjects at risk of developing rheumatoid arthritis. *Arthritis Rheumatol*. 2016;68(10):2372–2383. doi:10.1002/art.39771
201. Svard A, Kastbom A, Soderlin MK, Reckner-Olsson A, Skogh T. A comparison between IgG- and IgA-class antibodies to cyclic citrullinated peptides and to modified citrullinated vimentin in early rheumatoid arthritis and very early arthritis. *J Rheumatol*. 2011;38(7):1265–1272. doi:10.3899/jrheum.101086
202. Kokkonen H, Mullazehi M, Berglin E, et al. Antibodies of IgG, IgA and IgM isotypes against cyclic citrullinated peptide precede the development of rheumatoid arthritis. *Arthritis Res Ther*. 2011;13(1):R13. doi:10.1186/ar3237
203. Anquetil F, Clavel C, Offer G, Serre G, Sebbag M. IgM and IgA rheumatoid factors purified from rheumatoid arthritis sera boost the Fc receptor- and complement-dependent effector functions of the disease-specific anti-citrullinated protein autoantibodies. *J Immunol*. 2015;194(8):3664–3674. doi:10.4049/jimmunol.1402334
204. Du L, Wang P, Liu C, Li S, Yue S, Yang Y. Multisystemic manifestations of IgA vasculitis. *Clin Rheumatol*. 2020. doi:10.1007/s10067-020-05166-5
205. Yang YH, Wang SJ, Chuang YH, Lin YT, Chiang BL. The level of IgA antibodies to human umbilical vein endothelial cells can be enhanced by TNF-alpha treatment in children with Henoch-Schonlein purpura. *Clin Exp Immunol*. 2002;130(2):352–357. doi:10.1046/j.1365-2249.2002.01964.x
206. Berthelot L, Jamin A, Viglietti D, et al. Value of biomarkers for predicting immunoglobulin A vasculitis nephritis outcome in an adult prospective cohort. *Nephrol Dial Transplant*. 2018;33(9):1579–1590. doi:10.1093/ndt/gfx300
207. Sardy M, Karpati S, Merkl B, Paulsson M, Smyth N. Epidermal transglutaminase (TGase 3) is the autoantigen of dermatitis herpetiformis. *J Exp Med*. 2002;195(6):747–757. doi:10.1084/jem.20011299
208. Taylor TB, Schmidt LA, Meyer LJ, Zane JJ. Transglutaminase 3 present in the IgA aggregates in dermatitis herpetiformis skin is enzymatically active and binds soluble fibrinogen. *J Invest Dermatol*. 2015;135(2):623–625. doi:10.1038/jid.2014.368
209. Davalos D, Akassoglou K. Fibrinogen as a key regulator of inflammation in disease. *Semin Immunopathol*. 2012;34(1):43–62.
210. Smith AD, Streilein RD, Hall RP 3rd. Neutrophil CD11b, L-selectin and Fc IgA receptors in patients with dermatitis herpetiformis. *Br J Dermatol*. 2002;147(6):1109–1117. doi:10.1046/j.1365-2133.2002.05004.x
211. Gornowicz-Porowska K, Seraszek-Jaros A, Bowszyc-Dmochowska M, Kaczmarek E, Dmochowski M. Immunorexpression of IgA receptors (CD89, CD71) in dermatitis herpetiformis. *Folia Histochem Cytobiol*. 2017;55(4):212–220. doi:10.5603/FHC.a2017.0024
212. Russo V, Klein T, Lim DJ, et al. Granzyme B is elevated in autoimmune blistering diseases and cleaves key anchoring proteins of the dermal-epidermal junction. *Sci Rep*. 2018;8(1):9690. doi:10.1038/s41598-018-28070-0
213. Coppo R, Amore A, Roccatello D. Dietary antigens and primary immunoglobulin A nephropathy. *J Am Soc Nephrol*. 1992;2(10 Suppl):S173–80.
214. Sitaru C, Zillikens D. Mechanisms of blister induction by autoantibodies. *Exp Dermatol*. 2005;14(12):861–875. doi:10.1111/j.1600-0625.2005.00367.x
215. van der Steen LP, Bakema JE, Sesarman A, et al. Blocking Fcalpha receptor I on granulocytes prevents tissue damage induced by IgA autoantibodies. *J Immunol*. 2012;189(4):1594–1601. doi:10.4049/jimmunol.1101763
216. Park JH, Peyrin-Biroulet L, Eisenhut M, Shin JI. IBD immunopathogenesis: a comprehensive review of inflammatory molecules. *Autoimmun Rev*. 2017;16(4):416–426. doi:10.1016/j.autrev.2017.02.013
217. Palm NW, de Zoete MR, Cullen TW, et al. Immunoglobulin A coating identifies colitogenic bacteria in inflammatory bowel disease. *Cell*. 2014;158(5):1000–1010. doi:10.1016/j.cell.2014.08.006
218. van der Waaij LA, Kroese FG, Visser A, et al. Immunoglobulin coating of faecal bacteria in inflammatory bowel disease. *Eur J Gastroenterol Hepatol*. 2004;16(7):669–674. doi:10.1097/01.meg.0000108346.41221.19
219. Mitsuyama K, Niwa M, Takedatsu H, et al. Antibody markers in the diagnosis of inflammatory bowel disease. *World J Gastroenterol*. 2016;22(3):1304–1310. doi:10.3748/wjg.v22.i3.1304
220. Rengarajan S, Vivio EE, Parkes M, et al. Dynamic immunoglobulin responses to gut bacteria during inflammatory bowel disease. *Gut Microbes*. 2020;11(3):405–420. doi:10.1080/19490976.2019.1626683
221. Bakema JE, Tuk CW, van Vliet SJ, et al. Antibody-opsonized bacteria evoke an inflammatory dendritic cell phenotype and polyfunctional Th cells by cross-talk between TLRs and FcRs. *J Immunol*. 2015;194(4):1856–1866. doi:10.4049/jimmunol.1303126
222. Maloy KJ, Powrie F. Intestinal homeostasis and its breakdown in inflammatory bowel disease. *Nature*. 2011;474(7351):298–306. doi:10.1038/nature10208
223. Miquel-Clopes A, Bentley EG, Stewart JP, Carding SR. Mucosal vaccines and technology. *Clin Exp Immunol*. 2019;196(2):205–214. doi:10.1111/cei.13285
224. Vela Ramirez JE, Sharpe LA, Peppas NA. Current state and challenges in developing oral vaccines. *Adv Drug Deliv Rev*. 2017;114:116–131. doi:10.1016/j.addr.2017.04.008
225. Thakur A, Rodriguez-Rodriguez C, Saatchi K, et al. Dual-isotope SPECT/CT imaging of the tuberculosis subunit vaccine H56/CAF01: induction of strong systemic and mucosal IgA and T-cell responses in mice upon subcutaneous prime and intrapulmonary boost immunization. *Front Immunol*. 2018;9:2825. doi:10.3389/fimmu.2018.02825
226. Balu S, Reljic R, Lewis MJ, et al. A novel human IgA monoclonal antibody protects against tuberculosis. *J Immunol*. 2011;186(5):3113–3119. doi:10.4049/jimmunol.1003189
227. Williams A, Reljic R, Naylor I, et al. Passive protection with immunoglobulin A antibodies against tuberculous early infection of the lungs. *Immunology*. 2004;111(3):328–333. doi:10.1111/j.1365-2567.2004.01809.x
228. Casadevall A. Antibodies to mycobacterium tuberculosis. *N Engl J Med*. 2017;376(3):283–285. doi:10.1056/NEJMcibr1613268
229. Briles DE, Ades E, Paton JC, et al. Intranasal immunization of mice with a mixture of the pneumococcal proteins PsaA and PspA is highly protective against nasopharyngeal carriage of *Streptococcus pneumoniae*. *Infect Immun*. 2000;68(2):796–800. doi:10.1128/IAI.68.2.796-800.2000
230. Fukuyama Y, King JD, Kataoka K, Kobayashi R, Gilbert RS, Oishi K. Secretory-IgA antibodies play an important role in the immunity to *Streptococcus pneumoniae*. *J Immunol*. 2010;185(3):1755–1762. doi:10.4049/jimmunol.1000831

231. Parker EP, Molodecky NA, Pons-Salort M, O'Reilly KM, Grassly NC. Impact of inactivated poliovirus vaccine on mucosal immunity: implications for the polio eradication endgame. *Expert Rev Vaccines*. 2015;14(8):1113–1123. doi:10.1586/14760584.2015.1052800
232. Patel M, Glass RI, Jiang B, Santosham M, Lopman B, Parashar U. A systematic review of anti-rotavirus serum IgA antibody titer as a potential correlate of rotavirus vaccine efficacy. *J Infect Dis*. 2013;208(2):284–294. doi:10.1093/infdis/jit166
233. Blutt SE, Miller AD, Salmon SL, Metzger DW, Conner ME. IgA is important for clearance and critical for protection from rotavirus infection. *Mucosal Immunol*. 2012;5(6):712–719. doi:10.1038/mi.2012.51
234. McNeal MM, Stone SC, Basu M, et al. Protection against rotavirus shedding after intranasal immunization of mice with a chimeric VP6 protein does not require intestinal IgA. *Virology*. 2006;346(2):338–347. doi:10.1016/j.virol.2005.11.016
235. McNeal MM, Ward RL. Long-term production of rotavirus antibody and protection against reinfection following a single infection of neonatal mice with murine rotavirus. *Virology*. 1995;211(2):474–480. doi:10.1006/viro.1995.1429
236. Buisman AM, Abbink F, Schepp RM, Sonsma JA, Herremans T, Kimman TG. Preexisting poliovirus-specific IgA in the circulation correlates with protection against virus excretion in the elderly. *J Infect Dis*. 2008;197(5):698–706. doi:10.1086/527487
237. Renegar KB, Small PA Jr, Boykins LG, Wright PF. Role of IgA versus IgG in the control of influenza viral infection in the murine respiratory tract. *J Immunol*. 2004;173(3):1978–1986. doi:10.4049/jimmunol.173.3.1978
238. Tamura S, Funato H, Hirabayashi Y, et al. Cross-protection against influenza A virus infection by passively transferred respiratory tract IgA antibodies to different hemagglutinin molecules. *Eur J Immunol*. 1991;21(6):1337–1344. doi:10.1002/eji.1830210602
239. Hutchings AB, Helander A, Silvey KJ, et al. Secretory immunoglobulin A antibodies against the sigma1 outer capsid protein of reovirus type 1 Lang prevent infection of mouse Peyer's patches. *J Virol*. 2004;78(2):947–957. doi:10.1128/JVI.78.2.947-957.2004
240. Boyaka PN. Inducing mucosal IgA: a challenge for vaccine adjuvants and delivery systems. *J Immunol*. 2017;199(1):9–16. doi:10.4049/jimmunol.1601775
241. Sterlin D, Gorochoy G. When therapeutic IgA antibodies might come of age. *Pharmacology*. 2020;1–11. doi:10.1159/000510251
242. Heemskerk N, van Egmond M. Monoclonal antibody-mediated killing of tumour cells by neutrophils. *Eur J Clin Invest*. 2018;48(Suppl 2):e12962. doi:10.1111/eci.12962
243. Lohse S, Loew S, Kretschmer A, Jansen JHM, Meyer S, Ten Broeke T. Effector mechanisms of IgA antibodies against CD20 include recruitment of myeloid cells for antibody-dependent cell-mediated cytotoxicity and complement-dependent cytotoxicity. *Br J Haematol*. 2018;181(3):413–417. doi:10.1111/bjh.14624
244. Dechant M, Valerius T. IgA antibodies for cancer therapy. *Crit Rev Oncol Hematol*. 2001;39(1–2):69–77. doi:10.1016/S1040-8428(01)00105-6
245. Valerius T, Stockmeyer B, van Spriell AB, et al. FcalphaRI (CD89) as a novel trigger molecule for bispecific antibody therapy. *Blood*. 1997;90(11):4485–4492. doi:10.1182/blood.V90.11.4485
246. Bakema JE, van Egmond M. Immunoglobulin A: a next generation of therapeutic antibodies? *MAbs*. 2011;3(4):352–361. doi:10.4161/mabs.3.4.16092
247. Boross P, Lohse S, Nederend M, et al. IgA EGFR antibodies mediate tumour killing in vivo. *EMBO Mol Med*. 2013;5(8):1213–1226. doi:10.1002/emmm.201201929
248. Meyer S, Nederend M, Jansen JH, et al. Improved in vivo anti-tumor effects of IgA-Her2 antibodies through half-life extension and serum exposure enhancement by FcRn targeting. *MAbs*. 2016;8(1):87–98.
249. Otten MA, Bakema JE, Tuk CW, et al. Enhanced FcalphaRI-mediated neutrophil migration towards tumour colonies in the presence of endothelial cells. *Eur J Immunol*. 2012;42(7):1815–1821.
250. Otten MA, Rudolph E, Dechant M, et al. Immature neutrophils mediate tumor cell killing via IgA but not IgG Fc receptors. *J Immunol*. 2005;174(9):5472–5480. doi:10.4049/jimmunol.174.9.5472
251. Bakema JE, Ganzevles SH, Fluitsma DM, et al. Targeting FcalphaRI on polymorphonuclear cells induces tumor cell killing through autophagy. *J Immunol*. 2011;187(2):726–732. doi:10.4049/jimmunol.1002581
252. Kelton W, Mehta N, Charab W, et al. IgGA: a “cross-isotype” engineered human Fc antibody domain that displays both IgG-like and IgA-like effector functions. *Chem Biol*. 2014;21(12):1603–1609. doi:10.1016/j.chembiol.2014.10.017
253. Borrok MJ, Luheshi NM, Beyaz N, et al. Enhancement of antibody-dependent cell-mediated cytotoxicity by endowing IgG with FcalphaRI (CD89) binding. *MAbs*. 2015;7(4):743–751. doi:10.1080/19420862.2015.1047570
254. Ben Mkaddem S, Rossato E, Heming N, Monteiro RC. Anti-inflammatory role of the IgA Fc receptor (CD89): from autoimmunity to therapeutic perspectives. *Autoimmun Rev*. 2013;12(6):666–669. doi:10.1016/j.autrev.2012.10.011
255. Rossato E, Ben Mkaddem S, Kanamaru Y, et al. Reversal of arthritis by human monomeric IgA through the receptor-mediated SH2 domain-containing phosphatase 1 inhibitory pathway. *Arthritis Rheumatol*. 2015;67(7):1766–1777. doi:10.1002/art.39142
256. Kanamaru Y, Pfirsch S, Aloulou M, et al. Inhibitory ITAM signaling by Fc alpha RI-FcR gamma chain controls multiple activating responses and prevents renal inflammation. *J Immunol*. 2008;180(4):2669–2678. doi:10.4049/jimmunol.180.4.2669
257. Watanabe T, Kanamaru Y, Liu C, et al. Negative regulation of inflammatory responses by immunoglobulin A receptor (FcalphaRI) inhibits the development of toll-like receptor-9 signalling-accelerated glomerulonephritis. *Clin Exp Immunol*. 2011;166(2):235–250. doi:10.1111/j.1365-2249.2011.04452.x
258. Liu C, Kanamaru Y, Watanabe T, et al. Targeted IgA Fc receptor I (FcalphaRI) therapy in the early intervention and treatment of pristane-induced lupus nephritis in mice. *Clin Exp Immunol*. 2015;181(3):407–416. doi:10.1111/cei.12647
259. Aleyd E, Al M, Tuk CW, van der Laken CJ, van Egmond M. IgA complexes in plasma and synovial fluid of patients with rheumatoid arthritis induce neutrophil extracellular traps via FcalphaRI. *J Immunol*. 2016;197(12):4552–4559. doi:10.4049/jimmunol.1502353
260. Heineke MH, van der Steen LPE, Korthouwer RM, et al. Peptide mimetics of immunoglobulin A (IgA) and FcalphaRI block IgA-induced human neutrophil activation and migration. *Eur J Immunol*. 2017;47(10):1835–1845. doi:10.1002/eji.201646782

ImmunoTargets and Therapy

Dovepress

Publish your work in this journal

ImmunoTargets and Therapy is an international, peer-reviewed open access journal focusing on the immunological basis of diseases, potential targets for immune based therapy and treatment protocols employed to improve patient management. Basic immunology and physiology of the immune system in health, and disease will be also covered. In addition, the journal will focus on the impact of management

programs and new therapeutic agents and protocols on patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/immunotargets-and-therapy-journal>