


Protection of the Eye from Ultraviolet Radiation Damage Among Adults in Addis Zemen Town, Northwest Ethiopia

This article was published in the following Dove Press journal:
Clinical Optometry

Gizachew Tilahun Belete 
Kalkidan Getahun Tolessa
Mohammed Seid Hussien

University of Gondar, College of
Medicine and Health Science,
Department of Optometry, Gondar,
Ethiopia

Background: Ultraviolet radiation is electromagnetic radiation or light having a wavelength of less than 400 nm but greater than 100 nm. Ultraviolet radiation, majorly from sunlight, can potentially damage any organ that is exposed to any part of its spectrum. Aside from the skin, the organ most susceptible to sunlight-induced damage is the eye. Ultraviolet radiation is reported to be the cause of multiple ocular problems ranging from benign conditions like pterygium and pinguecula to ocular malignancies such as basal cell carcinoma which finally leads to visual impairment and blindness. Protection practice is mandatory to avoid the burden of diseases caused by ultraviolet exposure and maximizing the protection measures implementation is important. To maximize this, it is essential to know the current practice and the associated factors affecting the usage of protective devices in the study area.

Methods: Community-based cross-sectional study was conducted from April to May 2019 on 453 study participants. The study participants were selected through a systematic random sampling method. Data were collected using a pretested structured questionnaire. The analyzed result was summarized and presented using texts, tables and charts. A Chi-square test was applied to assess the significant association.

Results: A total of 430 study subjects were participated and completed the questionnaire with a response rate of 94.92%. The mean age of the study participants was 35.3 (SD±6.68). Two hundred forty-three (56.5%) study participants were males. About 228 (53%) of participants were married, and 356 (82.8%) were Christian in religion. From the total study participants, 173 (40.23%) had good practice in protecting the eye from ultraviolet radiation damages while the remaining 59.77% had poor practice. A significant association was found between sex and protection practice of the eye from ultraviolet radiation damages.

Conclusion: The majority of the study participants had poor protection practice of the eye from ultraviolet radiation damages. Improving awareness and protection practice are vital to reduce the burden of ocular abnormalities due to excessive exposure to ultraviolet radiation.

Keywords: ultraviolet radiation, protection practice, Addis Zemen Town, Ethiopia

Introduction

Ultraviolet radiation is an electromagnetic spectrum with a short wavelength between 100 and 400nm. Prolonged exposure to the sun increases exposure to this radiation which can cause chronic skin,¹ ocular and immune-related health problems.^{2,3}

The sensitivity of the eye to ultraviolet radiation (UVR) has been documented in both animal and human studies.⁴ Ultraviolet radiation is reported to be the cause of multiple ocular problems ranging from benign conditions like pterygium and pinguecula to ocular malignancies such as basal cell carcinoma.⁴⁻¹²

Correspondence: Gizachew Tilahun Belete
University of Gondar, College of Medicine
and Health Science, Department of
Optometry, Gondar, Ethiopia
Tel +251 (0)912684323
Email gizachewtilahun1@gmail.com

Even though the contribution of UV radiation to the global burden of disease has not yet been quantified, 25–33% of the global disease burden can be attributed to environmental risk factors. According to estimates by the World Health Organization, Worldwide, 12 to 15 million people become blind from cataracts annually, of which up to 20% may be caused or enhanced by sun exposure.³ To avoid the consequences of ultraviolet radiation on the eye there are several protective measures to be taken. Wearing wide-brimmed hat,¹³ sunglasses,^{13–15} wearing glasses with anti-reflective coatings,¹⁶ avoiding mid-day sun and provision of shades can reduce the ocular dose of ultraviolet radiation.¹³

Worldwide, some studies suggest that there has been an increase in the awareness of adverse effects of ultraviolet radiation exposure on the skin^{1,5,13,17–19} and even lesser studies on the eyes, however having the awareness and knowledge does not guarantee the use of the appropriate protective measures.^{5,6,12,18,20}

The few studies conducted show that there is poor practice of implementing the protective measures against ultraviolet exposure of the eyes,^{5,6,21,22} and since the usage of protection practice is mandatory to avoid the burden of diseases caused by the ultraviolet exposure, maximizing the protection measures implementation is important. In order to maximize this, it is essential to know the current practice and the associated factors affecting the usage of protective devices. In the study area, there is no evidence suggesting the current practice creating a knowledge gap and affecting which measures to take for the future in reducing the burden of ultraviolet exposure-related ocular morbidities. This study aims to fill the gap and provide baseline data on the current protection practice and associated factors of ultraviolet radiation hazards to the eye.

Methods and Materials

Study Design, Study Area and Period

A community-based cross-sectional study was conducted in Addis Zemen town, Northwest Ethiopia, from April to May 2019. Data obtained from Addis Zemen town administration's statistical office indicated that Addis Zemen city is located ~650 kilometers from the capital city of Ethiopia, Addis Ababa. It is a town in northwest Ethiopia with a latitude and longitude of 12°07'N 37°47'E and an elevation of about 1975 meters above sea level. The place is classified as humid subtropical bordering with subtropical high lands where ultraviolet radiation from sunlight could have a hazardous effect. According to

the 2007 Ethiopian national census, the city has a total population of 37,347 with four Kebeles (the smallest administrative unit) which hosts approximately 14,232 households. There are one governmental hospital, one health center and few private clinics.

Study Population, Sample Size and Sampling Procedure

The study population was all adults living in Addis Zemen town. The sample size was determined by using the single population proportion formula with the following assumptions. Level of significance (α) = 5% (with a confidence level of 95%), marginal error (w) = 4%, $P=22\%$ (0.22) (taken from similar study done in Australia by considering regular and frequent use of sunglass as a good practice in protecting ultraviolet radiation),⁵ Z-value of 1.96 at 95% for a confidence level. (n = sample size, P = proportion, w = marginal error).

$$n = \frac{Z^2 \alpha / 2p(1-p)}{w^2}$$

$$= \frac{(1.96)^2 (0.22)(0.78)}{(0.04)^2}$$

=412

Considering 10% for non-respondents, the final sample size determined was 453.

A simple random sampling method was used to select the first household and then systematic random sampling with an interval of 31 ($k=31$) was applied to select the remaining households. A simple random method (lottery) has been employed in a household when there was more than one participant in the selected household. Participants with a serious illness that makes them unable to communicate directly or through the translator to the interviewer were excluded. Individuals who stay indoors or who were not exposed to any other sources of ultraviolet radiation for the last one year were not also included in the study.

Data Collection Tool and Procedure

The data were collected using a structured and standardized questionnaire. The questionnaire comprised 8 socio-demographic questions, 10 major and 4 elaborative questions regarding the protection practice of the eyes from ultraviolet radiation damages. The questionnaire was first prepared in the English language, then translated to Amharic (local language), and later back-translated to the English language to maintain its consistency. The data were collected by trained optometrists through face to face interviews.

Protection practice was also assessed by using 14 questions. Each correct response was scored as 1 and each wrong response had a score of 0. The sum of scores varied from 0 to 14 points. Finally, the overall protection practice of the eyes from ultraviolet radiation damage was categorized as good or poor using the mean score as a cutoff point. Participants who scored the mean and above were considered as doing good practice while any score below the mean was grouped under poor practice.

Data Analysis

After the data were checked for completeness and cleared, it was coded and entered into Epi-data version 3.1 and analyzed by SPSS version 20. The descriptive statistics parameters were summarized using measures of central tendency and dispersion. A Chi-square test was applied to identify possible associations of factors for the protection practice of the eye from ultraviolet radiation damages. The analyzed data were organized and presented in a tabular form.

Results

Socio-Demographic Characteristics of Study Participants

In this study, a total of 430 study subjects were participated and completed the questionnaire with a response rate of 94.92%. The mean age of the study participants was 35.3 (SD±6.68). Two hundred forty-three (56.5%) study participants were males while 228 (53%) of participants were married and, 356 (82.8%) were Christian in religion. Almost half (48.6%) of the study participants had completed either primary or secondary school. Regarding the study participants' occupation, 123 (28.60%) was employed followed by merchants, 116 (27%) (See Table 1).

Protection Practice of the Eye from Ultraviolet Radiation Damage

From the total study participants, 173 (40.23%) had good practice in protecting the eye from ultraviolet radiation damages while the remaining 59.77% had poor practice (See Figure 1).

Wearing Time and Mechanism of Protection of the Eye from Ultraviolet Radiation

Among ultraviolet protection users, 112 (26.04%) used either sunglasses or photochromic lenses followed by

Table 1 Socio-Demographic Characteristics of Study Participants in Addis Zemen Town, Northwest Ethiopia, 2019

Variables	Frequency (n=430)	Percent (%)
Sex		
Male	243	56.51
Female	187	43.49
Age (Years)		
< 25	126	29.30
26–30	98	22.79
31–43	101	23.49
>43	105	24.42
Religion		
Christian	356	82.79
Muslim	74	17.21
Marital status		
Single	161	37.44
Married	228	53.03
Divorced	27	6.28
Widowed	14	3.25
Educational status		
No education	91	21.16
Non-formal education	63	14.65
Primary school	107	24.88
Secondary school	102	23.72
College and above	67	15.58
Occupation		
Employed	123	28.60
Merchant	116	26.98
Student	103	23.95
Housewife	64	14.88
Other	24	5.58
Household income		
<2500 ETB	183	42.56
≥2500 ETB	247	57.44

antireflection coated glasses. A significant number of participants, 123 (28.6%) did not use any type of protection.

More than 70% of protection users wore their protection device between 10 AM and 2 PM when ultraviolet radiation reaches its peak. The majority of study participants used protective device to protect ultraviolet radiation from sunlight (see Table 2).

Association Between Protection Practice and Socio-Demographic Characteristics

Among participants who had poor protective practice, males account for 35%. Fifty percent of participants




Figure 1 Protection practice of the eye from ultraviolet radiation among adults in Addis Zemen town, Northwest Ethiopia, 2019.

with poor protective practice are Christians in their religion.

By applying the chi-square test, a significant association has been found between sex and protection practice of the eye from ultraviolet radiation damages ($P=0.03$) but no other significant association was obtained between socio-demographic variables and

Table 2 Wearing Time, Protective Device and Sources of Ultraviolet Radiation Among Adults in Addis Zemen Town, Northwest Ethiopia, 2019

Variables	Frequency	Percent (%)
Protection materials/means (n=430)		
Sunglass/photochromic lens	112	26.04
Combined ARC and tint glasses	57	13.25
Occupational safety glass	38	8.83
Brimmed hat	35	8.13
Clothes	21	4.88
Umbrella	29	6.74
No protection	123	28.60
Others	15	3.49
Protection wearing time (n=307)		
12 AM to 10 AM	35	11.40
10 AM to 2 PM	221	71.97
2 PM to 12 AM	51	16.61
Protection of eye from (UVR sources) (n=307)		
Sunlight	170	45.95
Welding arc	38	10.27
Visual display units emissions	52	14.05
Reflection from snow/ocean/sea	18	4.86
Reflection from buildings/roads	21	5.68
Others	7	1.89

protection practice of the eye from UVR damages (See Table 3).

Discussion

In this study, the proportion of good practice to protect the eye from ultraviolet radiation damage was 40.23% (95% CI: 36.62–48.91%) which was in line with studies done in Canada, Brazil and Malta.^{23–26} This level of protection practice is categorized as low which needs prevention and protection campaigns to reduce the damages of ultraviolet radiation.

The proportion of protective practice in this study is lower than studies done in different states of America (69% and 80%) and Australia (71% and 61%).^{6,27–29}

Table 3 Distribution of Protection Practice Among Socio-Demographic Characteristics in Addis Zemen, Northwest Ethiopia, 2019

Variables	Protection Practice		
	Good	Poor	p-value
Sex			
Male	94	149	0.03
Female	79	108	
Age			
<25	50	76	0.52
26–30	39	59	
31–43	41	60	
>43	43	62	
Religion			
Christian	143	213	0.82
Muslim	30	44	
Educational status			
No education	37	54	2.75
Non formal education	25	38	
Primary school	43	64	
Secondary school	41	61	
College and above	27	40	
Occupation			
Employed	49	74	0.16
Merchant	47	69	
Student	41	62	
Housewife	26	38	
Others	10	14	
Household income (ETB)			
<2500	74	109	0.21
≥2500	99	148	

Abbreviations: ETB, Ethiopian Birr; household income, average monthly household or family income.

This difference may be due to variations in the nature of the profession, time spent recreation, geographical location and level of knowledge. Study subjects (over 80%) in the USA who use ultraviolet protective devices were lifeguards in their profession. Because of the nature of the profession, these subjects are forced to use the protective device but none of the participants in this study were lifeguards. The other study in Galveston, USA (69% use protective device) includes participants who spent the majority of their at the beach area. The study subjects in western Sydney (Australia-61% use protective devices) were outdoor workers. Both in the American and Australian studies, the proportion of awareness and knowledge about the protection of UVR is high which leads to a better use of the protective device to avoid its damage. In the study area, there are only a few awareness campaigns and low eye care service utilization. Additionally, the study settings both in Australia and America are at higher risk of UVR exposure which enforces individuals to use the protective device.

However, the protective practice of the eye from ultraviolet radiation damages found in this study was higher than the studies done in South Australia, Brazil, South Africa and California (America).^{5,30-32} The difference is likely due to the differences in the study population characteristics. The study subjects in South Australia (1-18 years) and Brazil (university students with a mean age of 22 years) were younger than this study (mean age 35.3 with a range of 18-85 years). There were also differences in sampling technique and sample size (large sample size in the case of South African study). Another possible explanation for this difference could be the criteria or base of classification as some of the studies used median and other methods to classify protective practice than the mean value. Likewise, the possible source of differences includes; study design and settings, socio-economic variations, level of awareness and health care seeking and utilization.

Sex was found to have a significant association with the protective practice of the eye from ultraviolet radiation damages ($P=0.03$). This finding was evidenced and supported by other studies.^{27,31} This could be explained in either way as most males are outdoor workers where they are at a state of exposure to ultraviolet radiation. This frequent exposure may enforce subjects to use some type of protective device to prevent the damaging effects of ultraviolet radiation on the eyes. On the other side, females are more conscious of their beauty and usually protecting

their bodies from sunlight exposure by sunscreens and umbrellas.

As a limitation of the study, there were limited works of literature done in the area which makes the comparison a bit difficult. Most of the published researches reviewed in this study were more on skin protection. Additionally, works of literature did not label the habit of protection practice but focused on the type of protection methods and devices which make direct comparison and explanation were not adequate.

Conclusion

The majority of the study participants had poor protection practice of the eye from ultraviolet radiation damages. A significant association was found between sex and protection practice of the eye from ultraviolet radiation damages. Improving awareness and protection practice are vital to reduce the burden of ocular abnormalities due to excessive exposure to ultraviolet radiation.

Data Sharing Statement

The data sets generated and analyzed for the current study are available from the corresponding author and accessed upon reasonable request.

Ethical Approval

Ethical clearance and approval were obtained from the Ethical Review Committee, University of Gondar. Both the study and verbal informed consent were also approved by the Ethical Review Committee, University of Gondar. The study was conducted in accordance with the Declaration of Helsinki. The objective of the study was clearly described for each participant. Then, oral informed consent was received from each study participant. The survey participants were granted the full right to cease or refuse to take part. Confidentiality was assured through keeping records and omitting any participants' identifiers.

Author Contributions

All authors contributed to data analysis, drafting or revising the article, have agreed on the journal to which the article will be submitted, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Disclosure

The authors declare that we have no conflicts of interest regarding this work or the publication of this article.

References

- Grandahl K, Ibler KS, Laier GH, Mortensen OS. Skin cancer risk perception and sun protection behavior at work, at leisure, and on sun holidays: a survey for Danish outdoor and indoor workers. *Environ Health Prev Med*. 2018;23(1):47. doi:10.1186/s12199-018-0736-x
- Robyn L, Tony M, Wayne S, Bruce A. *Solar Ultraviolet Radiation: Global Burden of Disease from Solar Ultraviolet Radiation*. Geneva: World Health Organization; 2006.
- WHO. *Intersun - The Global UV Project: A Guide and Compendium*. Geneva: World Health Organization; 2003.
- Rosenthal FS, Safran M, Taylor H. The ocular dose of ultra violet radiation from sunlight exposure. *Photochem Photobiol*. 1985;42(2):163–171. doi:10.1111/j.1751-1097.1985.tb01555.x
- Pakroui N, Casson R, Fung S, Ferdowsi N, Lee G, Selva D. South Australian adolescent ophthalmic sun protective behaviours. *Eye*. 2006;22:808–814. doi:10.1038/sj.eye.6702619
- Lee GA, Hirst LW, Sheehan M. Knowledge of sunlight effects on the eyes and protective behaviors in adolescents. *Ophthalmic Epidemiol*. 1999;6(3):171–180. doi:10.1076/oep.6.3.171.1501
- Rosenthal Aeb FS, Taylor H. The effect of prescription eyewear on ocular exposure to ultraviolet radiation. *AJPH*. 1986;76:10.
- Informal Consultation on the Effects of Solar UVRotE, Blindness WHOPftPo, World Health Organization. Office of G, Integrated Environmental H. *The Effects of Solar UV Radiation on the Eye: Report of an Informal Consultation, Geneva, 30 August - 3 September 1993*. Geneva: World Health Organization; 1994.
- Lindgren G, Diffey BL, Larko O. Basal cell carcinoma of the eyelids and solar ultraviolet radiation exposure. *BRJ Ophthalmol*. 1998;82(12):1412–1415. doi:10.1136/bjo.82.12.1412
- Sliney DH. Ocular exposure to environmental light and ultraviolet - the impact of lid opening and sky conditions. *Dev Ophthalmol*. 1997;27:63–75.
- Young RW. The family of sunlight related eye disease. *Optom Vis Sci*. 1994;71(2):125–144. doi:10.1097/00006324-199402000-00013
- Lee Graham A, Hirst Lawrence W, Sheehan M. Knowledge of sunlight effects on the eyes and protective behaviors in the general community. *Ophthalmic Epidemiol*. 1994;1(2):67–84. doi:10.3109/09286589409052363
- Peter H. Protection against solar ultraviolet radiation. *Mutat Res*. 1998;15–22.
- Frank S, Bakalian AE, Lou CQ, Taylor HR. The effect of sunglasses on ocular exposure to ultraviolet radiation. *Am J Public Health*. 1988;78(1):72–74. doi:10.2105/AJPH.78.1.72
- Sliney DH. Photoprotection of the eye – UV radiation and sunglasses. *J Photochem Photobiol B*. 2001;64(2–3):166–175. doi:10.1016/S1011-1344(01)00229-9
- Citek K. Anti-reflective coatings reflect ultraviolet radiation. *Optometry*. 2008;79(3):143–148. doi:10.1016/j.optm.2007.08.019
- Diffey BL. Sunscreens as a preventative measure in melanoma: an evidence-based approach or the precautionary principle? *Br J Dermatol*. 2009;161:25–27. doi:10.1111/j.1365-2133.2009.09445.x
- Parkin DM, Mesher D, Sasieni P. Cancers attributable to solar (ultraviolet) radiation exposure in the UK in 2010. *Br J Cancer*. 2011;105(S2):S66–S9. doi:10.1038/bjc.2011.486
- Cook BE, Bartley GB. Treatment options and future prospects for the management of eyelid malignancies. *Ophthalmology*. 2001;108(11):11. doi:10.1016/S0161-6420(01)00796-5
- Thomas-Gavelan E, Sáenz-Anduaga E, Ramos W, Sánchez-Saldaña L, Sialer M. Knowledge, attitudes and practices about sun exposure and photoprotection in outpatients attending dermatology clinics at four hospitals in Lima, Peru. *An Bras Dermatol*. 2011;86(6):1122–1128. doi:10.1590/S0365-05962011000600009
- Wright CY, Reeder AI, Albers PN. Knowledge and practice of sun protection in schools in South Africa where no national sun protection programme exists. *Health Educ Res*. 2016;31(2):247–259. doi:10.1093/her/cyw005
- Janjani H, Nedjat S. Sun exposure and health safety practices of high school students in an urban population of Iran. *BMC Public Health*. 2019;19(1):19. doi:10.1186/s12889-019-8100-7
- Campbell HS, Birdsall JM. Knowledge, beliefs, and sun protection behaviors of Alberta adults. *Prev Med*. 1994;23(2):160–166. doi:10.1006/pmed.1994.1022
- Dallazem LND, Benvegnú AM, Stramari JM, Beber AAC, Chemello RML, Beck M. Knowledge and habits of sun exposure in university students: a cross-sectional study in Southern Brazil. *An Bras Dermatol*. 2019;94(2):172–181.
- Benvenuto-Andrade C, Zen B, Fonseca G, De Villa D, Cestari T. Sun exposure and sun protection habits among high-school adolescents in Porto Alegre, Brazil. *Photochem Photobiol*. 2005;81(3):630. doi:10.1562/2005-01-25-RA-428.1
- Scerri L, Aquilina S, Gauci AA, Dalmás M. Sun awareness and sun protection practices in Malta. *J Eur Acad Dermatol Venereol*. 2002;16(1):47–52. doi:10.1046/j.1468-3083.2002.00376.x
- Houghtaling PM, Foster T, Heiner DJ, Uchida T, Wagner JR. Eyelid protection from ultraviolet radiation injury at the beach. *HPC Suppl Focus Sun Care*. 2010;3:12.
- Cioffi J, Wilkes L, O'Brien J. Outdoor workers and sun protection: knowledge and behaviour. *Constr Econ Build*. 2002;2(2):10–14. doi:10.5130/AJCEB.v2i2.2896
- Hall DM, McCarty F, Elliott T, Glanz K. Lifeguards' sun protection habits and sunburns: association with sun-safe environments and skin cancer prevention program participation. *Arch Dermatol*. 2009;145(2):139–144. doi:10.1001/archdermatol.2008.553
- Castilho IG, Sousa M, Leite R. Photoexposure and risk factors for skin cancer: an evaluation of behaviors and knowledge among university students. *An Bras Dermatol*. 2010;85(2):173–178. doi:10.1590/S0365-05962010000200007
- Oduntan O, Carelson A, Clarke-Farr P, Hansraj R. South African university student knowledge of eye protection against sunlight. *African Vis Eye Health*. 2009;68(1):25–31. doi:10.4102/aveh.v68i1.149
- Salas R, Mayer JA, Hoerster KD. Sun-protective behaviors of California farmworkers. *J Occup Environ Med*. 2005;47(12):1244–1249. doi:10.1097/01.jom.0000177080.58808.3b

Clinical Optometry

Publish your work in this journal

Clinical Optometry is an international, peer-reviewed, open access journal publishing original research, basic science, clinical and epidemiological studies, reviews and evaluations on clinical optometry. All aspects of patient care are addressed within the journal as well as the practice of optometry including economic and business analyses. Basic and clinical research papers are published that cover

all aspects of optics, refraction and its application to the theory and practice of optometry. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-optometry-journal>

Dovepress