Burnout and Associated Factors Among Medical Students in a Public University in Uganda: A Cross-Sectional Study

This article was published in the following Dove Press journal:
Advances in Medical Education and Practice

Jonathan Kajjimu ¹
Mark Mohan Kaggwa ²
Felix Bongomin ³

¹Faculty of Medicine, Mbarara University of Science and Technology, Mbarara, Uganda; ²Department of Psychiatry, Faculty of Medicine, Mbarara University of Science and Technology, Mbarara, Uganda; ³Department of Medical Microbiology and Immunology, Faculty of Medicine, Gulu University, Gulu, Uganda

Background: Burnout is a well-recognized phenomenon that may manifest with feelings of stress, fatigue, or exhaustion. It is a common and emerging problem among healthcare workers. Medical students may be at increased risk of burnout given the rigorous nature of their training. However, there is a paucity of data on the burden of burnout among medical students in Africa.

Aim: This study aimed to determine the prevalence of burnout, as assessed using the Maslach Burnout Inventory – Student Survey (MBI-SS) as well as factors associated with the development of burnout among students pursuing Bachelor of Medicine and Bachelor of Surgery (MBChB) degrees at Mbarara University of Science and Technology (MUST), Uganda.

Methods: A single-centre, cross-sectional, online survey was conducted among MBChB students of MUST. Burnout was assessed using the Maslach Burnout Inventory – Student Survey (MBI-SS) tool. Bivariate analysis and backward stepwise logistic regression analysis were performed to assess possible associations between variables related to participants' demography, socioeconomic, personal, learning environment, outside school environment aspects and burnout prevalence scores.

Results: A total of 145 medical students, 102 (70.3%) male, with a median (range) age of 23 (18–40) years were studied. A total of 135 students (93.1%) presented with high levels of emotional exhaustion, 90 (62.1%) students had low levels of professional efficacy scores and 141 (97.2%) of the medical students had high levels of cynicism. Overall, 79 (54.5%) students had burnout, as defined by the MBI-SS tool. Choosing MBChB willingly appears to be an independent predictor of burnout (Adjusted odds ratio: 7.2; 95% CI: 1.4–36.9; $p=0.018$).

Conclusion: More than one-half of medical students questioned at MUST do experience a degree of burnout. Preventative and interventional measures should be considered in the development of the medical curriculum.

Keywords: medical students, burnout, Maslach Burnout Inventory, mental health, well-being; Uganda

Introduction

Burnout is, according to the 11 revision of the International Classification of Diseases (ICD-11),

“A syndrome conceptualized as resulting from chronic workplace stress that has not been successfully managed. It's characterised by three dimensions: feelings of energy

Correspondence: Jonathan Kajjimu
Faculty of Medicine, Mbarara University of Science and Technology, Mbarara, Uganda
Tel +256759719384
Email jonathkebenz37@gmail.com

depletion or exhaustion; increased mental distance from one's job or feelings of negativism or cynicism related to one's job; and reduced personal efficacy."¹

Burnout is an increasingly recognised problem among medical professionals.² Previous studies in medical students found 14.8–79.9% of them experience burnout during their medical training.^{3–7} Its prevalence has been found to increase with medical school academic progression^{8,9} but studies have suggested that by equipping medical students with evidence-based stress management approaches to their education, the incidence of burnout is reduced irrespective of their academic progression.³²

In working-class healthcare professions, burnout has been linked majorly to job- or work-related chronic stress. In medical students, it is most likely related to personal factors such as mental health disorder, physical activity, personal financial satisfaction, high parental expectations, alcohol consumption, and smoking status or individual learning environment.^{1,10} A 2017 systematic review suggested that, on average, one medical student out of two suffered from burnout as defined by the ICD-11. Medical students from the Middle East and Oceania countries were found to have a higher prevalence of burnout than those from countries in other continents, with no African data available at the time of the review.¹¹ Higher levels of stress coupled with poor coping strategies may be the major factors responsible for the development of burnout among medical students,³ due to the cyclic relation between stress and emotional exhaustion.¹² Medical students, when compared to age-matched members of the general public, have been shown to have a poorer psychological quality of life.¹³ A 2015 systematic review showed that medical students displayed a variety of coping mechanisms for burnout such as personal engagement, extracurricular activities including mainly musical and physical activities, student autonomy, mentorship programs, evaluation systems, career counseling and life coaching.³

The prevalence of burnout among medical students at Mbarara University of Science and Technology (MUST) was unknown. We sought to identify the degree of burnout and identify any causal factors such as learning environment, personal life factors and outside school life factors. This cross-sectional study aimed at investigating the prevalence and factors associated with burnout among medical students in their pre-clinical and clinical years of study at MUST.

Materials and Methods

Study Design

An online descriptive cross-sectional study was conducted between 6th September 2020 until 20th September 2020, using a Google form.

Study Setting

The study was carried out at Mbarara University of Science and Technology, commonly called Mbarara University. It is a public university accredited by the National Council for Higher Education, offering both undergraduate and postgraduate programs. MUST was established in 1989 and has two campuses – Kihumuro campus and Mbarara campus. The Faculty of Medicine is located at the Mbarara campus approximately 266 kilometers southwest of Kampala. The MBChB (Bachelor of Medicine and Bachelor of Surgery) program at Mbarara University takes 5 years. There were approximately 423 medical students enrolled for the 2019/2020 academic year in the Faculty of Medicine at MUST.

Study Sample

Undergraduate students from first to final years at MUST, pursuing a medical program of Bachelor of Medicine and Bachelor of Surgery, were surveyed using an enrollment criteria including students being in the official WhatsApp groups of the respective class years and voluntarily consenting to take part in the study (see Figure 1). Representatives of each class were contacted by the principal investigator and informed about the study procedures, ethical issues and data collection. The participants were enrolled using a consecutive sampling technique. Students were recruited into the study through class representatives. The survey tool was then disseminated through class official WhatsApp groups. The inclusion criteria were medical students at MUST enrolled for the academic year 2019/2020. Other health profession students were not eligible for enrollment in this study.

The required sample size was calculated using the Kish–Leslie formula.

$$N = \frac{Z^2 p(1-p)}{d^2}$$

where:

N = Sample size

Z = Standard deviation at 95% Confidence Interval (1.960)

Figure 1 Flow chart for enrollment of participants.

p = Expected prevalence of depression among medical students

d = Precision of the estimate (5%)

The prevalence of burnout was taken to be an average of 0.5 since there had not been any previous study carried out at MUST. This point prevalence, of 50% was also found in a recent study assessing burnout among medical students in Ethiopia.¹⁶

We used the modified Kish–Leslie equation³³ to calculate the available sample size; where K was the estimated overall population of the study population.

$$n = \frac{N}{1 + \frac{(N-1)}{K}}$$

To cater for non-correspondents assuming a response rate of 90%, a total of 10% of the sample size was added. This led to the required sample size of 229 students.

Measurements

We developed a questionnaire that would collect participants' characteristics (demography and socioeconomic factors) and their MBI-SS scores. We also asked about factors that might be associated with burnout among medical students. Participants' characteristics included sex,

age, year of study, religious/spiritual affiliation, source of tuition funding, marital status, family income, agreement with the effectiveness of wellness initiatives at their institution, and family residence.

Burnout was assessed using the Maslach Burnout Inventory – Student Survey tool which was adapted from^{17,18} and had been validated in Brazil.¹⁹ The MBS-SS is a self-administered tool consisting of 15 questions. It is subdivided into three subscales: emotional exhaustion (five items), cynicism (four items), and professional efficacy (six items). All items are assessed by frequency using a Likert scale ranging from never (0) to every day (6).

The tool has an internal consistency of each latent factor greater than 0.7 with values of 0.77, 0.72, and 0.81 for emotional exhaustion, professional efficacy, and cynicism, respectively.²⁰ A test–retest reliability determined at about 4 weeks later, a yielded reliability coefficients of 89, 84 and 67 respectively for exhaustion, cynicism and academic efficacy.²¹ The sensitivity, specificity, positive and negative predictive values of the of the 15-item MBI-SS were 91.9% (95% CI = 82.5–96.5%), 93.2% (95% CI = 87.5–96.4%), 86.4% (95% CI = 76.1–92.7%) and 96.1% (95% CI = 91.2–98.3%) respectively. The positive and negative likelihood ratios were 13.48 (95% CI = 7.15–25.44) and 0.09 (95% CI = 0.04–0.20) respectively.²⁰

There are two validated calculation methods used to compare burnout scores among participants ie, the summation (SUM) and average (AVE) method.¹⁷ Burnout is determined by the total score of the MBI-SS based on the participant's responses by using both the tridimensional and bidimensional diagnostic criteria for burnout syndrome as listed in the ICD-11. On a two-dimensional model, it is identified by high scores on both emotional exhaustion and cynicism subscales. In the three-dimensional model, low-level burnout was determined based on the following scores: emotional exhaustion (0 - 9), cynicism (0 - 1), and professional efficacy <27. Moderate level burnout was determined based on the following scores: emotional exhaustion (10 - 14), cynicism (2 - 6), and professional efficacy (23 - 27). At a high level, burnout was identified based on the following scores: emotional exhaustion >14, cynicism >6, and professional efficacy <23.¹⁸ High scores on emotional exhaustion and cynicism with low scores on professional efficacy are indicative of burnout.²⁰

Factors associated with burnout among medical students were adapted from Haile et al,^{16,18,22} and modified

to suit the institution's low-income setting. Most responses were dichotomised.

The questionnaire was piloted by 23 (10% of the sample size) Gulu University medical students. A Cronbach alpha of 0.8099 was obtained, demonstrating the high reliability of the tool in assessing the study outcomes.

Data Analysis

Data analyses were performed using StataCorp. 2019. Stata: Release 16. Statistical Software. College Station, TX: StataCorp LLC. Medical students' characteristics were summarized with proportions. The prevalence of burnout was determined from the total score of the MBI-SS tool using the tridimensional criteria based on the participants' responses to Likert scale questions related to emotional exhaustion, professional efficacy, and cynicism. Proportions were used to estimate the prevalence of burnout in the student sample. To quantify possible associations between the independent variable and dependent variables, we calculated crude odds ratios. Backward regression analysis was used to assess the predictive relationship between the associated factors obtained from the medical students and their respective burnout prevalence scores.

Results

Participant Characteristics

One hundred and fifty (35.5%) responses were received. We removed one incomplete response and four duplicate responses, leaving behind 145 responses that were analysed (34.3% response rate). The participants included 102 (70.3%) male and 43 (29.7%), female medical students. The majority (49%, $n=71$) of the students were under government sponsorship and over 90% ($n=134$) had a religious affiliation. The majority of the students were aged less than 24 years, and the rest of the sociodemographic characteristics were comparable across the 5 years of studies as shown in Table 1.

Prevalence of Burnout Among Medical Students

Table 2 presents the prevalence of burnout among medical students. More than half (54.5%, $n=79$) of the medical students fitted the definition of burnout using the tridimensional criteria. Of the 145 students, 93.1% ($n=135$) had high emotional exhaustion

scores, 62.1% ($n=90$) had low professional efficacy scores and 97.2% ($n=141$) had high cynicism scores.

Factors Associated with Burnout Among Medical Students

In the unadjusted logistic regression analysis, several factors were found to be independently associated with burnout. These including factors intrinsic to the student such as being optimistic about their future career (OR=2.70, 95% CI 1.11 to 6.55; $p=0.028$), feeling fulfilled and happy at work (OR=2.77, 95% CI 1.31 to 5.86; $p=0.008$), willingly choosing to pursue MBChB (OR=9.44, 95% CI 2.04 to 43.58; $p=0.004$), feeling satisfied with medicine as their career choice (OR=5.07, 95% CI 2.09 to 12.31; $p=0.000$), feeling motivated with their studies (OR=2.43, 95% CI 1.08 to 5.45; $p=0.031$), and having ever developed any thoughts of dropping out of school (OR=0.17, 95% CI 0.06 to 0.48; $p=0.001$). Key factors, outside of the students locus included an exhaustive day-to-day routine of studies (OR=0.43, 95% CI 0.19 to 0.96; $p=0.040$), the challenge of commuting long distances (OR=0.30, 95% CI 0.13 to 0.68; $p=0.004$), insufficient leisure time (OR=2.2, 95% CI 1.12 to 4.30; $p=0.021$), and considering one's self important to their family member (OR=6.08, 95% CI 1.27 to 29.22; $p=0.024$). More details in Table 3.

In multiple logistic backward regression only one factor, choosing MBChB willingly (OR =7.20, 95% CI 1.40 to 36.93; $p=0.018$) was found to be independently associated with burnout. Table 4 contains more details.

Discussion

Our cross-sectional study looked at the prevalence of burnout among medical students at MUST, as well as possible associations. More than half of the students described burnout. Willingly choosing a career in medicine was the only identified associated factor. The high prevalence of burnout in our study is consistent with a similar study in Egypt which reported a burnout prevalence of 79.9%.⁷ One possible reason could be that medical students were unaware of the issue of burnout. A 2010 systematic review, suggested that there existed an association between a lower level of tolerance for uncertainty or ambiguity and self-reported burnout among medical students.²³ Studies conducted among medical students in Ethiopia and Spain found low levels of burnout, 34% and 14.8%, respectively.^{6,16} Such discrepancies in the burnout prevalence in various studies may be attributed to the

Table I Participants' Characteristics

Characteristics	Year 1 (n = 22)	Year 2 (n= 28)	Year 3 (n= 38)	Year 4 (n= 26)	Year 5 (n= 31)	Total N = 145	P-value
Sex, n (%)							0.414
Male	10 (23.3)	8 (18.6)	29 (28.4)	20 (19.6)	21 (20.6)	102 (70.3)	
Female	10 (22.2)	8 (17.7)	9 (20.9)	6 (14.0)	10 (23.3)	43 (29.7)	
Age (years)							0.000*
18–23	19 (20.9)	25 (27.5)	32 (35.2)	8 (8.8)	7 (7.7)	91 (62.8)	
24–40	3 (5.6)	3 (5.6)	6 (11.1)	18 (33.3)	24 (44.4)	54 (37.2)	
Religious/spiritual affiliation n (%)							0.255
Yes	22 (16.4)	26 (19.4)	36 (26.9)	24 (17.9)	26 (19.4)	134 (92.4)	
No	0	2 (18.1)	2 (18.1)	2 (18.1)	5 (45.5)	11 (7.6)	
How is your tuition funded?, n (%)							0.066
Government funded	11 (15.5)	12 (16.9)	21 (29.6)	15 (21.1)	12 (16.9)	71 (49.0)	
NGO funded	0	2 (25.0)	1 (12.5)	3 (37.5)	2 (25.0)	8 (5.5)	
Private funded	10 (18.2)	8 (14.6)	14 (25.5)	6 (10.9)	17 (30.9)	55 (37.9)	
Self-funded	1 (9.1)	6 (54.6)	2 (18.2)	2 (18.2)	0	11 (7.6)	
Relationship status, n (%)							0.790
Married	1 (20.0)	2 (40.0)	1 (20.0)	1 (20.0)	0	5 (3.5)	
In a relationship with a steady partner	3 (11.1)	2 (7.4)	9 (33.3)	4 (14.8)	9 (33.3)	27 (18.6)	
In a relationship without a steady partner	1 (16.7)	1 (16.7)	2 (33.3)	1 (16.7)	1 (16.7)	6 (4.1)	
Single	17 (15.9)	23 (21.5)	26 (24.3)	20 (18.7)	21 (19.6)	107 (73.8)	
How is your family income wage (minimum wage)?, n (%)							0.113
Unsatisfactory	15 (19.7)	18 (23.7)	15 (19.7)	11 (14.5)	17 (22.4)	76 (52.4)	
Satisfactory	7 (10.1)	10 (14.5)	23 (33.3)	15 (21.7)	14 (20.3)	69 (47.6)	
How would you gauge wellness initiatives at your institution?, n (%)							0.657
Not effective	13 (13.4)	19 (19.6)	24 (24.7)	17 (17.5)	24 (24.7)	97 (66.9)	
Effective	9 (18.8)	9 (18.8)	14 (29.7)	9 (18.8)	7 (14.6)	48 (33.1)	
What's your family residence?, n(%)							0.773
Urban	14 (15.1)	17 (18.3)	23 (24.7)	16 (17.2)	23 (24.7)	93 (64.1)	
Rural	8 (15.4)	11 (21.2)	15 (28.9)	10 (19.2)	8 (15.4)	52 (35.9)	

Note: *Bolded to represent a significant p-value.

different burnout measurement tools used. Additionally, students in different countries are exposed to different socio-economic, political, health, and conflict-related stress factors.³

Burnout was higher among males (70.9%) compared to females (29.1%), a consistent finding with a systematic review that found male medical students to have a higher burnout prevalence.¹⁵ This could be due to the many male medical student participants as indicated in most studies. However, whether these many male survey responders are a true representative of their various institutions' male medical student population,

remains a question to answer to come up with a conclusive finding. It has been proposed, by some, that female medical students are at a high risk of burnout than their male counterparts.¹⁴ Female students are more likely to suffer from stress factors than male medical students,⁴ with more demands placed on them outside of the university.⁸

Third and fourth-year medical students had the highest level of burnout in this study. This correlates with prior studies that showed a higher degree of burnout in senior medical students.^{16,17} The transition from preclinical years to clinical years of training appears to be a risk factor for

Table 2 Prevalence of Burnout Among Medical Students

	n = 145	%
Burnout prevalence, subscales and levels with scores*	79	54.5
Emotional Exhaustion		
Low (0–9)	0	0
Moderate (10–14)	10	6.9
High >14	135	93.1
Professional Efficacy		
Low >27	42	29.0
Moderate (23–27)	13	9.0
High <23	90	62.1
Cynicism		
Low (0–1)	0	0
Moderate (2–6)	4	2.8
High >6	141	97.2

Note: *Scores based on Ef et al.¹⁸

academic stress in medical school.^{5,24} The low burnout prevalence of fifth years may be due to them having learned some coping strategies such as life coaching, reflective practice, self-care, musical and physical activities.

One interesting finding in our study was that willingly choosing to undertake MBChB was found to be significantly associated with an increased risk of burnout. This is contrary to a Cameroonian study that reported that students who regretted choosing their medical studies had higher burnout scores than their peers.²⁵ Medical schools in countries like America have established wellness programs that could be adapted to address the “bio-psycho-social-spiritual burnout” epidemic among medical students.^{23,26} In Australasia, several wellness initiatives have been trialed. These include the integration of a well-being curriculum alongside the traditional medical curriculum such as the SAFE-DRS program at the University of Auckland³⁰ and the Health Enhancement Program at Monash University Medical School.³¹ A U.S based study evaluating the success of a student-led program teaching stress management and self-care skills to first-year medical students, found it to be helpful (based on longevity, student participation, and student feedback).²⁷ Additionally, the GMC – UK has provided guidance to medical schools on the creation of peer support networks, group sessions on stress management, and mentoring schemes.²⁹

Medical schools promote competition rather than cooperation. Training-related stress factors including

annual examinations, an inability to cope with academic demands, and pressure with an obligation to succeed all compound the problem. Additionally, a sense of helplessness coupled with sleep deprivation, financial challenges, and the emotional burden of witnessed suffering stress medical students.^{5,24} Other studies have found associations between burnout in medical students with not seeing coursework as a source of pleasure, feeling uncomfortable with course activities, not acquiring skills necessary to become a doctor, poor social support, being less than satisfied with the education system, self-perception of health, optimism, and motivation in studies.^{16,18,22} Our study did not find any such associations.

Burnout is not only harmful to the medical students, their relatives, medical schools, and peers but to also the patients they care for. There is thus a heightened need to address burnout among medical students to ensure that delivery of compassionate patient care is kept paramount.^{4,5,28}

Limitations

Our study had several limitations. First, this was a cross-sectional study. It could not allow us to study the variation of burnout among medical students throughout their entire 5-year study period in medical school at MUST. Second, the small response rate might have led to a type 2 error. Census involving these students would reveal more reliable findings. Recall bias may also have occurred amongst students since, by the time the study was conducted, students were off-campus due to the cessation of educational activities due to the COVID-19 pandemic. This, in itself, may have also contributed.

Implications

Our study findings should be generalizable to many other low- and middle-income countries. We used a standardised burnout measurement tool to increase the reliability and validity of our study findings. This has been the first study to be conducted among medical students in Uganda and it has revealed high levels of emotional exhaustion, cynicism, and low levels of professional efficacy. There needs to be an investment in the provision of mental health support to medical students through interventions to raise awareness, understanding, and prevention of the development of psychological distress, burnout, and mental health disorders like depression and anxiety among medical students. The efficacy

Table 3 Distribution of MUST Medical Students According to Variables Related to Participant Characteristics, Personal Aspects, Learning Environment, Outside School Environment and Prevalence of Burnout

Variables	n (%)	Burnout (%)	Crude OR (CI 95%)	P-value
All responses (n= 145)	79 (100.0)	54.5		
Sex				
Male (n= 102)	56 (54.9)	70.9	1.06 (0.52–2.16)	0.876
Female (n= 43)	23 (53.5)	29.1	1	
Age (years)				
18–23 (n = 91)	51 (56.0)	64.6	1	
23–40 (n = 54)	28 (51.9)	35.4	0.84 (0.43–1.66)	0.624
Year of study				
Year 1 (n= 22)	14 (63.6)	17.7	1	
Year 2 (n= 28)	14 (50.0)	17.7	0.57 (0.18–1.79)	0.337
Year 3 (n= 38)	24 (63.2)	30.4	0.98 (0.33–2.91)	0.970
Year 4 (n= 26)	15 (57.7)	19.0	0.78 (0.24–2.50)	0.675
Year 5 (n= 31)	12 (38.7)	15.2	0.36 (0.12–1.12)	0.077
Religion/spiritual affiliation				
Yes (n= 134)	73 (54.5)	92.4	1 (0.29–3.43)	0.997
No (n= 11)	6 (54.6)	7.6	1	
Tuition funding				
Government funded (n= 71)	38 (53.5)	48.1	1	
NGO funded (n= 8)	3 (37.5)	3.8	0.52 (0.12–2.35)	0.396
Private funded (n= 54)	30 (55.6)	38.0	1.09 (0.53–2.21)	0.821
Self-funded (n= 12)	8 (66.7)	10.1	1.74 (0.48–6.29)	0.401
Marital status				
Married (n= 5)	4 (80.0)	5.1	5 (0.49–50.83)	0.174
In a relationship with a steady partner (n= 27)	12 (44.4)	15.2	1	
In a relationship without a steady partner (n= 6)	3 (50.0)	3.8	1.25 (0.21–7.35)	0.805
Single (n= 107)	60 (56.1)	76.0	1.60 (0.68–3.73)	0.281
Family Income				
Unsatisfactory (n= 76)	36 (47.4)	45.6	1	
Satisfactory (n= 69)	43 (62.3)	54.4	1.84 (0.95–3.57)	0.072
Effectiveness wellness initiatives				
Not effective (n= 97)	55 (56.7)	69.6	1	
Effective (n= 48)	24 (50.0)	30.4	0.76 (0.38–1.53)	0.446
Family residence				
Rural (n= 52)	29 (55.8)	36.7	1.08 (0.54–2.15)	0.816
Urban (n= 93)	50 (53.8)	63.3	1	
What's your health self-perception?				
Bad (n=14)	8 (57.1)	10.1	1	
Good (n= 131)	71 (54.2)	89.9	0.89 (0.29–2.70)	0.834
How do you consider yourself emotionally?				
Calm (n= 119)	68 (57.1)	86.1	1.82 (0.77–4.29)	0.172
Tense (n= 26)	11 (42.3)	13.9	1	
Have you had any mental disorder with prior diagnosis by a psychiatrist?				
Yes (n= 3)	2 (66.7)	2.5	1.69 (0.15–19.04)	0.672
No (n= 142)	77 (54.2)	97.5	1	

(Continued)

Table 3 (Continued).

Variables	n (%)	Burnout (%)	Crude OR (CI 95%)	P-value
Do you have regular WHO recommended physical activity? Yes (n= 54) No (n= 91)	33 (61.1) 46 (50.6)	41.8 58.2	1.54 (0.78–3.05) 1	0.218
Do you have optimism in the future of medicine as your career? Not optimistic (n= 26) Optimistic (n= 119)	9 (34.6) 70 (58.8)	11.4 88.6	1 2.70 (1.11–6.55)	0.028
What's your personal financial satisfaction? Dissatisfied (n=114) Satisfied (n=31)	58 (50.9) 21 (67.7)	73.4 26.6	1 2.07 (0.88–4.69)	0.098
Do you have fulfillment and happiness as a medical student at your institution? No (n= 41) Yes (n= 104)	15 (36.7) 64 (61.5)	19.0 81.0	1 2.77 (1.31–5.86)	0.008
Do you have parents who are doctors? No (n= 139) Yes (n= 6)	76 (54.7) 3 (50.0)	96.2 3.8	1 0.83 (0.16–4.25)	0.822
Do you have high parent expectations? Yes (n= 119) No (n= 26)	65 (54.6) 14 (53.9)	82.3 17.7	1.03 (0.44–2.42) 1	0.943
Do you use any substances like marijuana, shisha, cocaine etc? Yes (n= 8) No (n= 137)	4 (50.0) 75 (54.7)	5.1 94.9	0.83 (0.20–3.44) 1	0.794
How is your alcohol consumption? Low (n= 43) Moderate (n= 15) High (n= 3) I do not consume alcohol and I have never thought of considering it before (n= 69) I do not consume alcohol but I have ever thought of considering it before due to stress. (n=15)	26 (60.5) 10 (66.7) 2 (66.7) 33 (47.8) 8 (53.3)	32.9 12.7 2.53 41.8 10.1	0.76 (0.06–9.10) 1 (0.07–13.87) 1 0.46 (0.04–5.29) 0.57 (0.42–7.74)	0.832 1 0.532 0.674
What's your smoking status? Never smoked and never thought of trying it out (n= 128) Current smoker (n= 5) Former smoker (n= 3) Never smoked but I have thought of trying it out due to stress (n= 9)	68 (53.1) 3(60.0) 3(100.0) 5 (55.6)	86.1 3.8 3.8 6.3	0.76 (0.12–4.68) 1 1 0.83 (0.09–7.68)	0.763 Omitted* 0.872
How are your school activities? Studies and extracurricular (n= 110) Studies alone (n= 35)	64 (58.2) 15 (42.9)	81 19	1.86 (0.86–4.00) 1	0.115
Did you choose MBChB willingly? No (n= 15) Yes (n= 130)	2 (13.3) 77 (59.2)	2.5 97.5	1 9.44 (2.04–43.58)	0.004
What are your average hours at school per day? Up to 12 hours (n= 84) More than 12 hours (n= 61)	49 (58.3) 30 (49.2)	62 38	1 0.69 (0.35–1.34)	0.275

(Continued)

Table 3 (Continued).

Variables	n (%)	Burnout (%)	Crude OR (CI 95%)	P-value
Do you feel you have study overload and academic pressure? Yes (n= 122) No (n= 23)	63 (51.6) 16 (69.6)	79.75 20.25	0.47 (0.18–1.21) I	0.119
Are you satisfied with your medicine career choice? Yes (n= 113) No (n= 32)	71 (62.8) 8 (25.0)	89.9 10.1	5.07 (2.09–12.31) I	0.000
Do you feel motivated for your studies? No (n= 32) Yes (n= 113)	12 (37.5) 67 (59.5)	15.2 84.8	I 2.43 (1.08–5.45)	0.031
Do you consider yourself important to your colleagues? No (n=18) Yes (n= 127)	7 (38.9) 72 (56.7)	8.9 91.1	I 2.06 (0.75–5.65)	0.162
How is the routine of your studies? Exhaustive (n= 109) Not exhaustive (n= 36)	54 (49.5) 25 (69.4)	68.4 31.7	0.43 (0.19–0.96) I	0.040
Do you feel satisfied as a student with your institution's teaching strategies? No (n= 53) Yes (n= 92)	31 (58.5) 48 (52.2)	39.2 60.8	I 0.77 (0.39–1.53)	0.462
Have you experienced any humiliation in your studies before? Yes (n= 85) No (n= 60)	43 (50.6) 36 (60.0)	54.4 45.6	0.68 (0.35–1.33) I	0.263
Have you ever developed any thoughts of dropping out of school? Yes (n= 24) No (n= 121)	5 (20.8) 74 (61.2)	6.3 93.7	0.17 (0.06–0.48) I	0.001
Do you feel you are acquiring skills necessary to become a doctor? No (n= 9) Yes (n= 136)	2 (22.2) 77 (56.6)	2.5 97.5	I 4.57 (0.92–22.8)	0.064
What are your feelings about your course activities? Comfortable (n= 114) Uncomfortable (n= 31)	65 (57) 14 (45.2)	82.3 17.7	1.61 (0.73–3.58) I	0.242
Do you see coursework as a source of pleasure? No (n= 101) Yes (n= 44)	53 (52.5) 26 (59.1)	67.1 32.9	I 1.31 (0.64–2.68)	0.463
What's your experience of searching for mental health support before at your institution? Good and satisfied (n= 10) Unsatisfied (n= 5) Never searched for mental health support in spite of having mental health challenges (n= 40) Never searched for mental health support and never had mental health challenges (n= 90)	6 (60.0) 1 (20.0) 20 (50.0) 52 (57.8)	7.6 1.3 25.3 65.8	I 0.17 (0.01 –2.09) 0.67 (0.16–2.73) 0.91 (0.24–3.46)	0.165 0.573 0.893
Do you have any fears of hurting or harming patients? No (n= 65) Yes (n= 80)	35 (53.9) 44 (55.0)	44.3 55.7	I 1.05 (0.54–2.02)	0.890

(Continued)

Table 3 (Continued).

Variables	n (%)	Burnout (%)	Crude OR (CI 95%)	P-value
Are you in academic competition with your colleagues? Yes (n= 30) No (n= 115)	17 (56.7) 62 (53.9)	21.5 78.5	1.12 (0.50–2.51) I	0.787
Do you have fear of being infected in hospital? Yes (n= 108) No (n= 37)	55 (50.9) 24 (64.9)	69.6 30.4	0.56 (0.26–1.22) I	0.144
Do you feel you have limited medical training time? Yes (n= 71) No (n= 74)	40 (56.3) 39 (52.7)	50.6 49.4	1.16 (0.60–2.23) I	0.660
Do you have inability of answering academic patient questions? Yes (n= 49) No (n= 96)	25 (51.0) 54 (56.3)	31.65 68.35	0.81 (0.41–1.62) I	0.550
Do you have a positive influence of a religious/spiritual life in your studies? No (n= 31) Yes (n= 114)	15 (48.4) 64 (56.1)	19 81	I 1.37 (0.62–3.03)	0.443
Do you have any financial challenges at school which bother you in your studies? Yes (n= 79) No (n= 66)	39 (49.4) 40 (60.6)	49.4 50.6	0.63 (0.33–1.23) I	0.177
Does the distance between campus or hospital and your place of residence/ hostel inconvenience you at all? Yes (n= 34) No (n= 111)	11 (32.4) 68 (61.3)	13.9 86.1	0.30 (0.13–0.68) I	0.004
Do you have frequent family meetings that you attend? No (n= 130) Yes (n= 15)	71 (54.6) 8 (53.3)	89.9 10.1	I 0.95 (0.33–2.77)	0.925
Do you get sufficient leisure time? No (n= 77) Yes (n= 68)	35 (45.5) 44 (64.7)	44.3 55.7	2.2 (1.12–4.30) I	0.021
Do you consider yourself important to your family members? No (n = 11) Yes (n= 134)	2 (18.2) 77 (57.5)	2.5 97.5	I 6.08 (1.27–29.22)	0.024
Have you lost any close family member(s) during your time in medical school whose death affected you negatively in your studies? Yes (n= 28) No (n= 117)	17 (60.7) 62 (53.0)	21.5 78.5	1.37(0.59–3.18) I	0.462
Do you have any close family member(s) battling with a chronic disease currently, whose condition you think may have negatively affected or is negatively affecting your studies? Yes (n= 33) No (n= 112)	13 (39.4) 66 (58.9)	16.5 83.5	0.45 (0.21–1.00) I	0.050
How often do you seek for social support when encountered with emotional challenges? Not often (n= 113) Often (n= 32)	57 (50.4) 22 (68.8)	72.15 27.85	I 2.16 (0.94–4.97)	0.070

(Continued)

Table 3 (Continued).

Variables	n (%)	Burnout (%)	Crude OR (CI 95%)	P-value
Do you have any personal relationship challenges?				
Yes (n= 35)	18 (51.4)	22.8	0.99 (0.42–2.30)	0.974
No (n= 56)	29 (51.8)	36.7	I	
Not in a relationship (n= 54)	32 (59.3)	40.5	1.35 (0.64–2.88)	0.43I
Do you have additional family responsibilities?				
Yes (n= 54)	28 (51.9)	35.4	0.85 (0.43–1.66)	0.624
No (n= 91)	51 (56.0)	64.6	I	

Notes: Bolded to represent a significant p-value. *Omitted represents a p-value that could not be calculated due to having less than 5 observations.

Table 4 Outcome of Logistic Regression Analysis for Related Variables and Burnout in Medical Students

Variables	Crude OR (95% CI)	Adjusted OR (95%)	p-value
Do you have optimism in the future of medicine as your career?			
Not optimism (n= 26)	I	I	
Optimistic (n= 119)	2.70 (1.11–6.55)	1.26 (0.39–4.07)	0.70I
Do you have fulfillment and happiness as a medical student at your institution?			
No (n= 41)	I	I	
Yes (n= 104)	2.77 (1.31–5.86)	1.10 (0.38–3.19)	0.855
Did you choose MBChB willingly?			
No (n= 15)	I	I	
Yes (n= 130)	9.44 (2.04–43.58)	7.20 (1.40–36.93)	0.018*
Are you satisfied with your medicine career choice?			
Yes (n= 113)	5.07 (2.09–12.31)	3.17 (0.93–10.72)	0.064
No (n= 32)	I	I	
Do you feel motivated for your studies?			
No (n= 32)	I	I	
Yes (n= 113)	2.43 (1.08–5.45)	0.63 (0.19–2.09)	0.452
How is the routine of your studies?			
Exhaustive (n= 109)	0.43 (0.19–0.96)	0.54 (0.21–1.38)	0.198
Not exhaustive (n= 36)	I	I	
Have you ever developed any thoughts of dropping out of school?			
Yes (n= 24)	0.17 (0.06–0.48)	0.36 (0.10–1.25)	0.106
No (n= 121)	I	I	
Does the distance between campus or hospital and your place of residence/hostel inconvenience you at all?			
Yes (n= 34)	0.30 (0.13–0.68)	0.44 (0.16–1.16)	0.097
No (n= 111)	I	I	
Do you get sufficient leisure time?			
No (n= 77)	2.2 (1.12–4.30)	1.31 (0.57–2.99)	0.525
Yes (n= 68)	I	I	
Do you consider yourself important to your family members?			
No (n = 11)	I	I	
Yes (n= 134)	6.08 (1.27–29.2)	1.82 (0.29–11.88)	0.530

Note: *Bolded to represent a significant p-value.

of this study could be measured by conducting a multi-centre, prospective longitudinal study.

Some student life activities may be present at MUST but these activities are not united under a comprehensive wellness objective. We hope that this study will help drive the establishment of wellbeing interventions at MUST.

Data Sharing Statement

Data are available upon reasonable request from the first author.

Ethics Approval

The study was performed in accordance with the international ethical standards of the Declaration of Helsinki. This study was approved by the Mbarara University of Science and Technology Research Ethics Committee (Approval number: 19/07–20). We also sought administrative approval from the dean of the Faculty of Medicine at Mbarara University of Science and Technology.

The participant's email and name were requested optionally for entering the participants into a raffle for participatory gift certificates as an incentive for completing the survey. Each study participant who opted to submit their details for enrollment into the raffle was assigned a random number, after which we randomly chose five study participants from the enrolled participants based on those random numbers.

Acknowledgments

We thank all participants for their willingness and the time they took to participate in the study. We thank the following students for serving as class representatives who reached out to their classmates with study link messages during the data collection stage including Viola Nansimbe, Innocent Arinaitwe, Ruth Nakawungu, Janat Nakachwa, Jeremiah Habizana, Taremwa Norman, and Ssewanyana Ernest. We also thank Ronald Olum for his support during data analysis. Specially, we thank Dr. Andrew Tagg (Emergency Physician, Western Health, Melbourne, Australia) for his contributions including copy editing of the study manuscript and provision of financial support for purchasing a license for using the Maslach Burnout Inventory instrument.

Author Contributions

All authors made a significant contribution to the work reported, be it in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all

these areas; took part in drafting, revising, or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

This study was not funded.

Disclosure

The authors declare that they have no conflicts of interest for this work.

References

1. WHO. Burn-out an “occupational phenomenon”: international classification of diseases [Internet]. WHO, World Health Organization; 2019 [cited September 19, 2020]. Available from: http://www.who.int/mental_health/evidence/burn-out/en/. Accessed December 30, 2020
2. Understanding the burnout experience: recent research and its implications for psychiatry - Maslach - 2016 - world psychiatry. Wiley Online Library; 2016 [cited October 7, 2020]. Available from: <https://onlinelibrary.wiley.com/doi/full/10.1002/wps.20311>. Accessed December 30, 2020
3. Fares J, Al Tabosh H, Saadeddin Z, El Mouhayyar C, Aridi H. Stress, burnout and coping strategies in preclinical medical students. *N Am J Med Sci*. 2016;8(2):75–81. doi:10.4103/1947-2714.177299
4. Muzafar Y, Khan HH, Ashraf H, et al. Burnout and its associated factors in medical students of lahore, pakistan. Cureus. [cited September 19, 2020]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4689594/>. Accessed December 30, 2020.
5. Dyrbye LN, Thomas MR, Huntington JL, et al. Personal life events and medical student burnout: a multicenter study. *Acad Med*. 2006;81(4):374–384. doi:10.1097/00001888-200604000-00010
6. Galán F, Sanmartín A, Polo J, Giner L. Burnout risk in medical students in Spain using the Maslach Burnout Inventory-Student Survey. *Int Arch Occup Environ Health*. 2011;84(4):453–459. doi:10.1007/s00420-011-0623-x
7. Atlam SA. Burnout syndrome: determinants and predictors among medical students of Tanta University, Egypt. *Egypt J Community Med*. 2018;36(01):61–73. doi:10.21608/ejcm.2018.6870
8. ElKholy MM, El-Sayed ET, Sedrak AS, Raouf NA. Prevalence and predictors of burnout syndrome among medical students of Cairo University. *Egypt J Community Med*. 2019;37(3):83–92. doi:10.21608/ejcm.2019.43375
9. Melaku L, Mossie A, Negash A. Stress among medical students and its association with substance use and academic performance. *J Biomed Educ*. 2015;2015:1–9. doi:10.1155/2015/149509
10. Dyrbye L, Shanafelt T. A narrative review on burnout experienced by medical students and residents. *Med Educ*. 2016;50(1):132–149. doi:10.1111/medu.12927
11. Frajerman A, Morvan Y, Krebs M-O, Gorwood P, Chaumette B. Burnout in medical students before residency: a systematic review and meta-analysis. *Eur Psychiatry*. 2019;55:36–42. doi:10.1016/j.eurpsy.2018.08.006
12. McManus IC, Keeling A, Paice E. Stress, burnout and doctors' attitudes to work are determined by personality and learning style: a twelve year longitudinal study of UK medical graduates. *BMC Med*. 2004;18(2):29. doi:10.1186/1741-7015-2-29

13. Chang E, Eddins-Folensbee F, Coverdale J. Survey of the prevalence of burnout, stress, depression, and the use of supports by medical students at one school. *Acad Psychiatry*. 2012;36(3):177–182.
14. Vidhukumar K, Hamza M. Prevalence and correlates of burnout among undergraduate medical students - a cross-sectional survey. *Indian J Psychol Med*. 2020;42(2):122–127. doi:10.4103/IJPSYM.IJPSYM_192_19
15. Chunming WM, Harrison R, MacIntyre R, Travaglia J, Balasooriya C. Burnout in medical students: a systematic review of experiences in Chinese medical schools. *BMC Med Educ*. 2017;17(1):217. doi:10.1186/s12909-017-1064-3
16. Haile YG, Senkute AL, Alemu BT, Bedane DM, Kebede KB. Prevalence and associated factors of burnout among Debre Berhan University medical students: a cross-sectional study. *BMC Med Educ*. 2019;19(1):413. doi:10.1186/s12909-019-1864-8
17. Obregon M, Luo J, Shelton J, Blevins T, MacDowell M. Assessment of burnout in medical students using the Maslach Burnout Inventory-Student Survey: a cross-sectional data analysis [Internet]. In review; 2020 [cited Sep 17, 2020]. Available from: <https://www.researchsquare.com/article/rs-30445/v2>. Accessed December 30, 2020.
18. Ef C, Sa S, At S, Ev M, Tm A. Burnout syndrome and associated factors among medical students: a cross-sectional study [Internet]. vol. 67, clinics (Sao Paulo, Brazil). clinics (Sao Paulo); 2012 [cited Sep 16, 2020]. Available from: <https://pubmed.ncbi.nlm.nih.gov/22760894/>.
19. Características psicométricas do Maslach Burnout Inventory. Student Survey (MBI-SS) em estudantes universitários brasileiros. *Psico-USF*. 2017;22(3):569. doi:10.1590/1413-82712017220315
20. Wickramasinghe ND, Dissanayake DS, Abeywardena GS. Clinical validity and diagnostic accuracy of the Maslach burnout inventory-student survey in Sri Lanka. *Health Qual Life Outcomes*. 2018;16(1):220. doi:10.1186/s12955-018-1048-y
21. Zeinab R, Reza AM, Ahmad B. The psychometric characteristics of Maslach Burnout Inventory Student Survey: a study students of Isfahan University. *J Res Med Sci*. 2014;16(9):55–58.
22. Boni RADS, Paiva CE, de Oliveira MA. Burnout among medical students during the first years of undergraduate school: prevalence and associated factors. *PLoS One*. 2018;13(3):e0191746. doi:10.1371/journal.pone.0191746
23. Saadat H, Lin SL, Kain ZN. The role of “wellness” in medical education. *Int Anesthesiol Clin*. 2010;48(3):131–138. doi:10.1097/AIA.0b013e3181e60213
24. Solanky P, Desai B, Kavishwar A, Kantharia S. Study of psychological stress among undergraduate medical students of government medical college, Surat. *Int J Med Sci Public Health*. 2012;1(2):38. doi:10.5455/ijmsph.2012.1.38-42
25. Njim T, Makebe H, Toukam L, et al. Burnout Syndrome amongst medical students in cameroon: a cross-sectional analysis of the determinants in preclinical and clinical students. *J Psychiatry*. 2019;2019:e4157574. doi:10.1155/2019/4157574
26. Drolet BC, Rodgers S. A comprehensive medical student wellness program—design and implementation at Vanderbilt School of Medicine. *Acad Med*. 2010;85(1):103–110. doi:10.1097/ACM.0b013e3181c46963
27. Redwood SK, Pollak MH. Student-led stress management program for first-year medical students. *Teach Learn Med*. 2007;19(1):42–46. doi:10.1080/10401330709336622
28. Wachholtz A, Rogoff M. The relationship between spirituality and burnout among medical students. *J Contemp Med Educ*. 2013;1(2):83–91. doi:10.5455/jcme.20130104060612
29. Supporting medical students with mental health conditions [Internet]. [cited Oct 14, 2020]. Available from: <https://www.gmc-uk.org/education/standards-guidance-and-curricula/guidance/supporting-medical-students-with-mental-health-conditions>.
30. Moir F, Yelder J, Dixon H. SAFE-DRS: 5 years of a health and well-being curriculum in the medical programme at the University of Auckland; 2018 [cited Nov 29, 2020]. Available from: <https://researchspace.auckland.ac.nz/handle/2292/43095>. Accessed December 30, 2020.
31. Hassed C, Sierpina VS, Kreitzer MJ. The health enhancement program at Monash University medical school. *Explore*. 2008;4(6):394–397. doi:10.1016/j.explore.2008.09.008
32. Wild K, Scholz M, Ropohl A, Bräuer L, Paulsen F, Burger PHM. Strategies against burnout and anxiety in medical education – implementation and evaluation of a new course on relaxation techniques (Relacs) for medical students. *PLoS One*. 2014;9(12):e114967. doi:10.1371/journal.pone.0114967
33. Glenn D. Israel determining sample size [Internet]. University of Florida. Available from: <https://www.tarleton.edu/academicassessment/documents/samplesize.pdf>. Accessed December 30, 2020.

Advances in Medical Education and Practice

Publish your work in this journal

Advances in Medical Education and Practice is an international, peer-reviewed, open access journal that aims to present and publish research on Medical Education covering medical, dental, nursing and allied health care professional education. The journal covers undergraduate education, postgraduate training and continuing medical education

Submit your manuscript here: <http://www.dovepress.com/advances-in-medical-education-and-practice-journal>

Dovepress

including emerging trends and innovative models linking education, research, and health care services. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.