Phenotyping Study of Cyclophosphamide 4-Hydroxylation in Malay Cancer Patients

This article was published in the following Dove Press journal:
Drug Design, Development and Therapy

Yesi Ihdina Fityatal Hasanah¹
Yahdiana Harahap^{1,2}
Denni Joko Purwanto³

¹Bioavailability and Bioequivalence Laboratory, Faculty of Pharmacy, Universitas Indonesia, Depok, Indonesia; ²Indonesia Defense University, Bogor 16810, West Java, Indonesia; ³Functional Medical Staff of Surgical Oncology, Dharmas Cancer Hospital, Jakarta, Indonesia

Background: Cyclophosphamide (CP) is an anticancer alkylating group (nitrogen mustard) and a prodrug that will be metabolized to form its active metabolite, 4-hydroxycyclophosphamide (4-OHCP). The various enzymes involved in its bioactivation can cause a wide range of CP expression and activity among patients and ultimately affect the metabolism, efficacy and toxicity of this drug. The effectiveness of CP therapy can be determined by 4-OHCP level in dried blood spot (DBS).

Aim: The purpose of this study was to conduct the phenotyping of CP 4-hydroxylation rate in Malay cancer patients.

Methodology: Phenotyping study of CP 4-hydroxylation rate to 40 subjects of Malay cancer patients was done based on the value of its bioactivity ratio (4-OHCP to CP levels).

Results: The result shown the cyclophosphamide 4-hydroxylation rate of 80% (n=32) subjects as ultrarapid metabolizer (UM) and 20% (n=8) as poor metabolizer (PM).

Conclusion: Phenotyping study of CP 4-hydroxylation in Malay cancer patients can be conducted by quantifying CP bioactivity ratio (4-OHCP to CP level) in dried blood spot. In majority of Malay cancer patients, cyclophosphamide would be bioactivated through 4-hydroxylation in hepar rapidly as indicated by the high value of the bioactivity ratio or the increased CP clearance and 4-OHCP level.

Keywords: 4-hydroxycyclophosphamide, 4-OHCP, 4-hydroxylation rate, cyclophosphamide, CP, dried blood spots, DBS, Malay ethnicity

Introduction

Cyclophosphamide (CP) is a nitrogen mustard, alkylating cytotoxic drug with a broad clinical spectrum for treatment of autoimmune disorders and various types of cancers like chronic lymphocytic leukemia, lymphomas, soft tissue and osteogenic sarcoma, and solid tumours (lung, breast, and ovary).¹ CP is given singly or in combination with other agents, orally or by intravenous injection or infusion. Up to 70–80% of CP is metabolized by the liver microsomal enzymes and yields its active metabolite, 4-hydroxycyclophosphamide (4-OHCP). The drug-metabolizing enzymes (DME) involved in the bioactivation are CYP2A6, CYP2B6, CYP3A4, CYP3A5, CYP2C9, CYP2C18, and CYP2C19, with CYP2B6 providing the greatest activity. In the systemic circulation, 4-OHCP is in equilibrium with its tautomeric, aldophosphamide, which will be hydrolyzed spontaneously by nonenzymatic β -elimination to phosphoramidate mustard and acrolein. Phosphoramidate mustard is cross-linked (covalent) with DNA intra- and interstrand, thus inhibiting DNA replication and causing cell apoptosis.^{2–5}

Several CP detoxification pathways compete with phosphoramidate mustard formation and thus affect phosphoramidate mustard levels. CP can undergo N-dechloroethylation

Correspondence: Yahdiana Harahap
Griya Depok Asri Blok G3 No. 1, Depok
City, West Java 16411, Indonesia
Email yahdiana@farmasi.ui.ac.id

which is catalyzed mainly by CYP3A4 to form 2-dechloroethylcyclophosphamide metabolite which is equimolar with chloroacetaldehyde by-products.³ This minor pathway accounts for approximately 19% of the total CP clearance. Besides, 4-OHCP undergoes secondary metabolism by CYP3A4 to form inactive 4-ketocyclophosphamide. The intermediate product (imino-cyclophosphamide) in this reaction can also undergo glutathione conjugation. Alcohol dehydrogenase and aldo-keto reductase also play a minor role in the elimination of aldophosphamide by reducing them to alcophosphamide. In contrast, the oxidation of aldophosphamide to carboxyphosphamide, catalyzed by *ALDH1A1* and *ALDH3A1*, is an important inactivation pathway. Less than 20% of the CP dose is eliminated in the urine in its entirety. Between 30 and 60% of total CP is eliminated in the kidney in the form of major metabolite and the remains eliminated through feces.^{2,3,6}

The CP chemotherapy effectiveness is determined by the active metabolite, 4-OHCP, level in the blood, which is formed through 4-hydroxylation. The various enzymes involved in CP metabolism can cause a wide range of CP expression and activity among individuals, and ultimately affect the pharmacokinetic, efficacy, and toxicity of this drug. A wide variability of the CP 4-hydroxylation rate between individuals (100 folds) can affect the drug pharmacokinetics (elimination rate and drug level) so that it gives different responses (efficacy and toxicity) among patients. Genetic factors (polymorphism) and environment factors such as diet, smoking, alcohol, drugs, disease, age, and sex can affect the manifestation of CP 4-hydroxylation.⁷

Given the complex scheme of CP metabolism, it is clear that many pathways can affect the interindividual variability of CP response. However, most pharmacogenomic studies have focused on assessing the genetic variant of CYP2B6, an enzyme that catalyzes the initial bioactivation step. The *CYP2B6* gene is located on chromosome 19q13.2, ranges in ~26 kb region, consists of nine exons, and encodes for 491 amino acids.⁸ *CYP2B6* SNP coding region changes the function and expression (activity) of the enzyme produced, which can vary depending on the substrate.^{9,10} *CYP2B6* is highly induced and can be auto-induced by CP within 24 h after continuous infusion or after repeated administration for several days.¹¹ Increased CP activation will increase the 4-OHCP levels and CP clearance.¹² The pharmacogenetics of *CYP2B6* is very complex because SNPs encoding common proteins can exist in various haplotype combinations. This causes

the distribution of allele variant frequencies in each population to be very diverse.

The CYP2B6*6 allele variant with the greatest prevalence was found in the Caucasian population (26%),¹³ Japan (16%),¹⁴ West Africa (42%), and Papua New Guinea (62%).⁸ The CYP2B6 *8 allele was found at low frequency (~1%) in the Caucasian population and was absent in African American, Hispanic, and Asian subjects.¹⁵ In the Timor population (East Nusa Tenggara, Indonesia), *CYP2B6* *9, *4, and *6 were found to be respectively; 40.6; 29.2; and 22.9%.¹⁶ The CP 4-hydroxylation rate by *CYP2B6* was categorized into poor metabolizer (PM) (*6 and/or *7), extensive metabolizer (EM) (*5 and wild type), and ultra rapid metabolizer (UM) (*4).¹⁷ Based on other literature, *CYP2B6* *6/*6 and *6/*7 were classified as PM, *CYP2B6* *1/*1 (wild type), *1/*5, *1/*6, *1/*7, and *5/*5 was classified as EM, and *CYP2B6* *1/*4, *4/*5, and *4/*6 were classified as UM.^{13,18,19} Classification of 4-hydroxylation rate type of *CYP2B6* alleles is still inconsistent with several studies. The Xie et al study found that the *CYP2B6* G516T allele variant (n=8) had an average bioactivity ratio and CP clearance two times greater than the wild type (n=21) with values (0.027±0.0020) and (0.017±0.0014), and (8.13±4.23 L/h) and (5.13±1.67 L/h).²⁰ Allele variants with the G516T mutation are known to be present in *CYP2B6* *6, *7, and *9. From several studies, *CYP2B6* *6 allele variant carrier had greater 4-hydroxylation activity than noncarriers.^{11,20–22} However, in a study on Japanese patients, correlation between the CP 4-hydroxylation rate with *CYP2B6* *6 and *9 alleles was not found.⁶ Differences between study results might be due to several factors including ethnicity, number of subjects, CP dose, or co-medication.

The interindividual variability of CP metabolism influenced the differences on CP toxicity and efficacy profile among patients. The CP metabolism is a manifestation of genetic expression (DME genetic polymorphism factor) combined with the environment factors (diet, lifestyle, co-medication, comorbidity, age, sex, pollutants, etc) so that phenotypes are not always correlated according to the genotype.^{7,11} However, some researchers still consider genetic factors to play a relatively large role in influencing the metabolic phenotype compared to other factors. The genotype is a detailed gene structure in an individual whereas phenotype is more commonly measured using drugs as a marker of a particular DME and measuring the ratio of metabolite levels to their parent drug.²³ Polymorphisms that occur in individuals will continue to be passed down to the next generation so that the

frequency of polymorphisms in each ethnicity or population varies. Therefore, there is a need for personalized medicine and the provision of safe, effective, and rational therapeutic doses in a population. Research on the metabolic activity of CP has been widely carried out but is limited to populations in developed countries such as countries in Europe, USA, Japan, and China.^{13,14,24} However, there have been no studies related to the activity (rate) of CP 4-hydroxylation metabolism in the population in Indonesia until now. This underlies the need for a phenotypic study of the CP 4-hydroxylation rate in cancer patients, especially those of the Malay ethnicity, which is the largest ethnic group in Indonesia. Malay, also known as the Mongoloid ethnicity, is considered to be the ancestors of the Indonesian nation and differentiated into Proto-Malay and Deutro-Malay.^{25,26} In this phenotyping study, the CP 4-hydroxylation will be determined by calculating the CP bioactivity ratio (4-OHCP to CP level) in dried blood spot (DBS) of Malay cancer patients which analyzed using UPLC-MS/MS with internal standard SIL 4-OHCP-d4.

Materials and Methods

Chemicals

Cyclophosphamide (CP) and semicarbazide hydrochloride as derivatization agent were purchased from Sigma-Aldrich (Singapore). 4-hydroxycyclophosphamide (4-OHCP) kit and internal standard (IS) 4-hydroxycyclophosphamide-d4 kit (4-OHCP-d4) were purchased from Toronto Research Chemicals (USA). Methanol (HPLC grade), acetonitrile (HPLC Grade), and formic acid was purchased from Merck (Darmstadt, Germany). Ultrapure water was prepared by using a Millipore Direct-Q™ 5 water system (Millipore, Watford, Hertfordshire, UK). Perkin Elmer 226 DBS card was obtained from PerkinElmer (USA).

Subjects

Dried blood spots (DBS) samples were acquired with written informed consent from cancer patients from “Dharmais” Cancer Hospital, Jakarta, Indonesia. Subjects were interviewed for ethnicity background for three generations background. This study was reviewed and approved by the Research Ethics Committees of “Dharmais” Cancer Hospital, Jakarta 11,420, Indonesia (No. 150/KEPK/VII/2019). Subjects were eligible for this study if they were 18–65 years old cancer patients of Malay ethnicity who received intravenous infusion of CP

as their chemotherapy regimen. Subjects who were contraindicated with CP and had liver and kidney disorders were excluded from the study. DBS samples were collected after complete the CP infuse on by finger-prick through the fingertips using a lancet, collected in a tube, and spotted immediately on DBS card which contain dried semicarbazide derivative agent.

Method Validation

Full validation of CP and 4-OHCP analysis method in DBS was conducted based on the Food and Drug Administration (2018) and European Medicines Agency (2011) guidelines for validation of bioanalysis, by validating LLOQ, linearity, accuracy and precision, selectivity, carryover, recovery, dilution integrity, matrix effect, and stability.^{27,28}

Cyclophosphamide 4-Hydroxylation Assay

Cyclophosphamide 4-hydroxylation was measured by quantifying CP and 4-OHCP in DBS samples with internal standard SIL 4-OHCP-d4 by UPLC-MS/MS. Due to 4-OHCP instability, it carried out on-card DBS derivatization. Initially, the DBS card was spotted by 50 µL semicarbazide 2 M and air-dried for two hours. Blood samples were spotted 30 µL on the DBS card and air-dried for three hours. Blood spots were cut off completely and added by 20 µL internal standard 4-OHCP-d4 1 mg/mL and 1000 µL methanol. The mixture was vortexed for two minutes, sonicated for 10 min, and centrifuged for five minutes at 10,000 rpm. Aliquots of 850 µL of supernatant was evaporated at 60°C for 15 min under N₂ gas flow. The residue was reconstituted by 100 µL of the mobile phase, sonicated for one minute, vortexed for 30 seconds, and centrifuged for five minutes at 10,000 rpm. Afterward, 10 µL of aliquot was injected into the ultra-high-pressure liquid chromatography–tandem mass spectrometry (UPLC-MS/MS) system.

CP and 4-OHCP in DBS were quantified using ESI+ ionization method (Waters Xevo TQD Triple Quadrupole). Chromatographic separation was performed using a UPLC H-Class Acquity column BEH C₁₈ (100×2.1 mm, 1.7 µm) maintained at 50°C. The flow rate was 0.20 mL/min, and the mobile phase was 0.01% formic acid as eluent A and acetonitrile as eluent B. The gradient program was as follows: initial elution with 75% A, followed by a linear gradient to 10% A from 0 to 1.0 min and maintained for 1.0 min, to 75% A from 2.0 to 2.5 min and maintained for 2.5 min until the end of the run. The quantitation was conducted using multiple reaction monitoring (MRM) and the quantitation traces were 261.03 >140.16 (cone

voltage, 34 V; collision voltage, 22 V) for CP, 334.10 >221.04 (cone voltage, 16 V; collision voltage, 14 V) for 4-OHCP-semicarbazone and 338.10 >225.06 (cone voltage, 10 V; collision voltage, 10 V) for IS 4-OHCP-d4-semicarbazone. The optimized parameters for MS were as follows: capillary voltage, 3.5 kV; gas desolvation temperature and flow rate, 350°C; 500 L/h, respectively. Calibration range was 10–40,000 ng/mL for CP and 5–4000 ng/mL for 4-OHCP, at seven levels of concentration for each. Cyclophosphamide 4-hydroxylation rate was determined by bioactivity ratio of CP which can be calculated by dividing 4-OHCP level to CP level ([4-OHCP]:[CP]).

Results and Discussion

Method Validation

Linearity and LLOQ

Calibration curves obtained were linear in the concentration range within 10–40,000 ng/mL for CP and 5–4000 ng/mL for 4-OHCP, with correlation coefficient ($r \geq 0.9980$) and LLOQ 10 and 5 ng/mL, respectively.

Selectivity

Retention times of CP, 4-OHCP, and IS were 3.13; 1.94; and 1.91 min, respectively. There was no significant interfering peak observed around the CP, 4-OHCP, and IS, due to endogenous components or reagents.

Accuracy and Precision

Quality control samples were run a five replicates at four concentration levels for each analyte, which were: 10 ng/mL (LLOQ), 30 ng/mL (QCL), 20,000 ng/mL (QCM), 30,000 ng/mL (QCH) for CP and 5 ng/mL (LLOQ), 15 ng/mL (QCL), 2000 ng/mL (QCM), 3000 ng/mL (QCH) for 4-OHCP by within-run and between-run. They fulfill the requirement if %diff and %CV obtained within 15%, except for LLOQ 20%. The intra-batch accuracy and precision performed on LLOQ, QCL, QCM, and QCH shown in Table 1 were within the acceptable criteria based on FDA 2018 and EMA 2011 guidelines.^{27,28}

Carryover

There was no carryover found in the blank sample after the injection of the highest concentration (ULOQ) sample. The carryover still met the requirements for analytes <20% and <5% for IS, which CP, 4-OHCP, and IS were 17.53, 1.48, and 0.01, respectively.

Table 1 Accuracy and Precision of CP and 4-OHCP

Analyte	QC	Conc. (ng/mL)	Accuracy (%diff)	Precision (%CV)
CP	LLOQ	10	2.57	6.25
	QCL	30	1.72	5.04
	QCM	20,000	8.83	3.32
	QCH	30,000	13.77	4.26
4-OHCP	LLOQ	5	5.19	5.55
	QCL	15	1.22	9.55
	QCM	2000	7.12	3.84
	QCH	4000	14.86	4.22

Recovery and Matrix Effect

The extraction recoveries of CP at three levels of concentration QCL, QCM, and QCH were 85.02–95.69% and 4-OHCP were 75.18–94.55%. The matrix effect of CP, 4-OHCP, and IS obtained were fulfilled the %CV criteria for both matrix factor and internal standard normalized matrix factor, not exceeding $\pm 15\%$. There was no significant matrix effect observed.

Dilution Integrity

Dilution process was conducted in human blank whole blood until QCH and half QCH concentration. Dilution integrity obtained was acceptable and still fulfils accuracy and precision requirements with %diff and %CV not more than 15%.

Stability

Stock solutions storage of CP, 4-OHCP, and IS 4-OHCP-d4 in water at room temperature for 24 h and in the freezer (-80°C) for 30 days (long-term stability) did not alter the analyte of CP, 4-OHCP, and IS 4-OHCP-d4. The stability test results of CP and 4-OHCP in DBS were stable enough during sample preparation, storage conditions and autosampler.

Cyclophosphamide 4-Hydroxylation Assay

All subjects who met the inclusion and exclusion criteria were 40 Malay women. Subject interviews were conducted after signing the informed consent to obtain subject information related to ethnic and tribe backgrounds, lifestyle, and physiological condition. All subjects were patients receiving two-hour prolonged CP infusions. About 200 μL blood sample was taken by finger-prick, after two hours CP administration. The blood was immediately spotted onto the DBS card containing dried semicarbazide

and air-dried for three hours. The blood sampling time at two hours was based on the $t_{\max}$ of active metabolites 4-OHCP (2.1 ± 2.5 h) and CP (1.7 ± 1.3 h) so expected that 4-OHCP has been optimally formed and CP can still be detected in large quantities.²⁹ Blood sampling at the elimination time (four hours) after CP administration was not possible due to patient unwillingness and over hospital operating hours. Blood spotting (30 μ L) on DBS card must be done immediately because 4-OHCP $t_{1/2}$ in blood at 37°C is about four minutes.³⁰

The characteristic data of the subjects is shown in Table 2. A total of 38 subjects (95%) were breast cancer patients while two subjects (5%) were non-Hodgkin's lymphoma cancer patients. Various combination of chemotherapy regimen were given to patients: 27 subjects (67.5%) were given the FAC chemotherapy regimen (fluorouracil+adriamycin+cyclophosphamide), six subjects (15%) AC (adriamycin+cyclophosphamide), four subjects (10%) TAC (docetaxel+adriamycin+cyclophosphamide), two patients (5%) RCHOP (rituximab+cyclophosphamide+adriamycin+vincristine+prednisone), and one subject (2.5%) TC (docetaxel+cyclophosphamide). Subjects

Table 2 Characteristic of Subjects

Characteristic	n=40
Age (years)	
Mean $\pm$ SD	43.6 $\pm$ 9.54
Median (min-max)	42 (23–63)
Weight (kg)	
Mean $\pm$ SD; (min-max)	59.3 $\pm$ 9.81; (38–85)
Cancer Type (n)	
Breast Cancer	38 (95.0%)
LNH	2 (5.0%)
Chemotherapy Regimen (n)	
FAC	27 (67.5%)
AC	6 (15.0%)
TAC	4 (10.0%)
RCHOP	2 (5.0%)
TC	1 (2.5%)
CP Dose (mg)	
Mean $\pm$ SD; (min-max)	804.8 $\pm$ 119.86; (510–1,100)
CP Dose (mg/kg)	
Mean $\pm$ SD; (min-max)	13.72 $\pm$ 1.88; (10–20)
CP Dose (mg/m²)	
Mean $\pm$ SD; (min-max)	508.0 $\pm$ 56.4; (401–691)

Abbreviations: A/H, doxorubicin; C, cyclophosphamide; F, fluorouracil; O, vincristine; P, prednisone; R, rituximab; T, docetaxel.

received an intravenous infusion of CP chemotherapy with doses varying between 510 and 1100 mg; 10–20 mg/kg body weight; or 401–691 mg/m² body area. The average given CP dose was 804.8 $\pm$ 119.86 mg; 13.7 $\pm$ 1.88 mg/kg; or 508.0 $\pm$ 56.40 mg/m². The given CP dose is in accordance with standard intravenous CP chemotherapy protocol, which is between 500 and 1000 mg/m² and is usually done without considering other variables such as age, genetic factors, and comorbid conditions.³¹

The results of the analysis of CP and 4-OHCP levels in DBS can be seen in Table 3. CP and the active metabolite 4-OHCP were found in all DBS samples. CP was found with the lowest levels in SN 01 at 4699 ng/mL, while the highest levels were found in SN 26 at 39,859 ng/mL. The mean CP level was found at 23,790 $\pm$ 11,472.52 ng/mL with 48.22% CV value. The active metabolite 4-OHCP was found with the lowest levels in SN 01 at 281 ng/mL and the highest levels were found in SN 22 at 946 ng/mL. The mean 4-OHCP metabolite level was found at 599 $\pm$ 158.95 ng/mL with 26.54% CV value. The bioactivity ratio is calculated from dividing the 4-OHCP metabolites levels against the CP prodrug levels in DBS and can be seen in Figure 1 and Table 3. The lowest ratio bioactivity ([4-OCP]:[CP]) was found in SN 16 at 0.0112, while the highest ratio was found in SN 09 at 0.0975, and the mean ratio was found at 0.0348 $\pm$ 0.02378 with 68.34% CV value.

Determination of the type of drug metabolism rate can be seen phenotypically based on the bioactivity ratio value,¹¹ but no literature has published for the classification of the three types of CP 4-hydroxylation rate (PM, EM, and UM) with the limiting value of the bioactivity

Table 3 Phenotype Analysis of Subjects

Characteristic	n=40
CP levels (ng/mL)	
Mean $\pm$ SD; (min-max)	23,790 $\pm$ 11,472.5; (4699–39,859)
4-OHCP levels (ng/mL)	
Mean $\pm$ SD; (min-max)	599 $\pm$ 159.0; (281–946)
Phenotype (bioactivity ratio)	
PM (<0,0170; n=8; 20%)	
Mean $\pm$ SD	0.0139 $\pm$ 0.0019
Median (min-max)	0.0141 (0.0112–0.0164)
UM ($\geq$0,0170; n=32; 80%)	
Mean $\pm$ SD	0.0400 $\pm$ 0.0239
Median (min-max)	0.0309 (0.0174–0.0975)

Abbreviations: PM, poor metabolizer; UM, ultrarapid metabolizer.

Figure 1 CP bioactivity ratio [4-OHCP]:[CP] on subjects.

Note: Black dashed-line shows a limit bioactivity ratio of 0.0170.

ratio. Previously it has been agreed in some literature that *CYP2B6* wild type is categorized as extensive metabolizer (EM) type.^{11,13,17–19} Therefore, the (mean) wild type bioactivity ratio of 0.0170,²⁰ can be used as the standard limit for the EM value, so that the CP 4-hydroxylation type is classified into two types, namely PM if the bioactivity ratio value is <0.0170 and the UM for the bioactivity ratio value is ≥ 0.0170 . In this study, the results showed that from all 40 subjects of Malay cancer patients, 80% of subjects ($n=32$) was classified as UM and 20% of subjects ($n=8$) was PM (Table 3). The profile of the distribution of the bioactivity ratio values in the two CP 4-hydroxylation groups, namely poor metabolizers and ultrarapid metabolizers, is shown in Figure 2 and it can be seen that although the data distribution is not symmetrical, extreme data or outliers are not found.

The UM phenotype indicates that CP will be bioactivated through 4-hydroxylation in hepatic rapidly as indicated by the high value of the bioactivity ratio or the increased CP clearance and 4-OHCP level. The prevalence result of CP metabolism phenotypes in this study corresponded with several previous studies in other ethnicities.^{8,11,13,14,16,20–22,24} Based on these other studies, it can be estimated that in Malay ethnicity the majority of the CP 4-hydroxylation phenotype type is obtained as a UM which might be genetically derived from the *CYP2B6* *6 allele variant.

High 4-OHCP level on ultrarapid metabolizer can improve cancer therapy response better but alternatively, it also increases toxicity.¹² Unlike other side effects (myelosuppression, dysuria, nausea, vomiting, alopecia, cardiotoxicity, hypersensitivity, immunosuppression) that can

Figure 2 Box plot of CP bioactivity ratio [4-OHCP]:[CP] in 4-hydroxylation group of poor metabolizer and ultrarapid metabolizer.

also be caused by other anticancer substances, hemorrhagic cystitis is a severe side effect caused especially by the oxazaphosphoric drug class, namely CP and ifosfamide, to be precise by the urotoxic metabolite acrolein. Acrolein is also a spontaneous (nonenzymatic) β -elimination product beside 4-OHCP so that the 4-OHCP level also correlates with the acrolein level. It can be assumed that the CP 4-hydroxylation rate phenotype affects acrolein metabolite levels and the potential for urological side effects, hemorrhagic cystitis. In many cases, side effects of hemorrhagic cystitis generally occur in 10–40% of patients receiving high or long-term CP chemotherapy. Hemorrhagic cystitis cases were also found based on indicators of hematuria

formation with a prevalence of 17.5% (n=40) at “Dharmais” Cancer Hospital.³² It was found that a high number of cases of hemorrhagic cystitis in patients at “Dharmais” Cancer Hospital required several efforts to minimize these side effects. Generally, prevention procedure that can minimize the hemorrhagic cystitis are reducing CP dose, hyperhydration, and administration of a uroprotective drug, mesna, a thiol (sulfhydryl) compound which can neutralize acrolein into a nontoxic thioester compound. Mesna should be used only for chemotherapy with ifosfamide high doses or CP greater than 1000 mg/m².³³ Reducing the CP dose for patients at Dharmais Cancer Hospital is not suggested because the dose has been relatively low (± 500 mg/m²) so it is more advisable to give mesna accompanied by hyperhydration on patients while undergoing chemotherapy.

The clear correlation study of the phenotypes on DME metabolism can facilitate personalized medicine. If the DME metabolism shows a strong phenotype-genotype correlation, then only by analyzing the patient’s DME genetic polymorphisms, the physicians can determine the dose for personalized medicine. Clinically, phenotypic metabolism studies have advantages over genotypes because they can directly and significantly describe DME capacity/activity through blood drug levels which correlate directly with the effectiveness and toxicity of therapy in patients. This is due to genetic factors are only one of several confounding factors that influence the metabolic pharmacokinetics of drugs.^{7,23} The DME metabolic phenotype study itself has limitations because biomatrix sampling can only be carried out within a certain time, namely when $t_{\max}$ of analyte is achieved after drug administration, bioanalysis costs are relatively expensive, and the need for special handling (sampling, transportation, and storage) with the right techniques and conditions, and analysis as soon as possible to maintain the stability of the analyte in the biomatrix. In contrast, practically speaking, the genotyping study of DME metabolism has the advantage because the analysis is faster, easier, simpler, and relatively low-cost than phenotyping study. Genotype studies are also practical because sampling and analysis can be carried out at any time without the need for prior drug administration to the patient. We suggested that this phenotyping study of CP 4-hydroxylation should be continued by doing genotyping study (*CYP2B6* gene polymorphism analysis) and with a larger quantity of subjects to strengthen the data analysis and

obtain a correlation between the phenotype of drug metabolic capacity (activity) and the genetic polymorphism of DME.

Conclusion

The results showed the cyclophosphamide 4-hydroxylation rate of 80% (n=32) subjects as UM and 20% (n=8) as PM. It can be concluded that in Malay cancer patients, cyclophosphamide would be bioactivated through 4-hydroxylation in hepar rapidly as indicated by the high value of the bioactivity ratio or the increased CP clearance and 4-OHCP level.

Abbreviations

4-OHCP, 4-hydroxycyclophosphamide; AC, adriamycin +cyclophosphamide; CP, cyclophosphamide; CV, coefficient of variance; DBS, dried blood spot; diff, difference/bias; EM, extensive metabolizer; EMEA, European Medicines Agency; ESI, electrospray ionization; FAC, fluorouracil +adriamycin+cyclophosphamide; LLOQ, lower limit of quantification; m/z, mass per charge ratio; MRM, multiple reaction monitoring; MS/MS, tandem mass spectrometry; PM, poor metabolizer; QC, quality control; QCH, quality control high; QCL, quality control low; QCM, quality control medium; RCHOP, rituximab+cyclophosphamide+adriamycin+vincristine+prednisone; SN, subject number; TAC, docetaxel+adriamycin+cyclophosphamide; TC, docetaxel +cyclophosphamide; ULOQ, upper limit of quantification; UM, ultrarapid metabolizer; UPLC, ultra high performance liquid chromatography.

Ethics

This study was reviewed and approved by the Research Ethics Committees of “Dharmais” Cancer Hospital, Jakarta 11,420, Indonesia (No. 150/KEPK/VII/2019). This study was conducted in accordance with the Declaration of Helsinki.

Acknowledgments

This study was supported by Directorate of Research and Community Services (DRPM) Universitas Indonesia with grant number: NKB-2095/UN2.RST/HKP.05.00/2020 and Dharmais Cancer Hospital, Jakarta, Indonesia.

Disclosure

The authors report no conflicts of interest in this work.

References

- International Agency for Research on Cancer (IARC). *IARC Monographs - 100A: Cyclophosphamide*. Paris: IARC-WHO; 2012.
- de Jonge ME, van Dam SM, Hillebrand MJX, et al. Simultaneous quantification of cyclophosphamide, 4-hydroxycyclophosphamide, N, N', N''-triethylenethio-phosphoramidate (thiotepa) and N, N', N''-triethylenephosphoramidate (tepa) in human plasma by high-performance liquid chromatography coupled with electrospray ionization tandem mass spectrometry. *J Mass Spectrom*. 2004;39:262–271. doi:10.1002/jms.570
- Ekhart C, Doodeman VD, Rodenhuis S, Smits PH, Beijnen JH, Huitema AD. Influence of polymorphisms of drug metabolizing enzymes (CYP2B6, CYP2C9, CYP2C19, CYP3A4, CYP3A5, GSTA1, GSTP1, ALDH1A1 and ALDH3A1) on the pharmacokinetics of cyclophosphamide and 4-hydroxycyclophosphamide. *Pharmacogenet Genomics*. 2008;18(6):515–523. doi:10.1097/FPC.0b013e3282fe9766
- Emadi A, Jones RJ, Brodsky RA. Cyclophosphamide and cancer: golden anniversary. *Nat Rev Clin Oncol*. 2009;6:638–647.
- Harahap Y, Samuel C, Andalusia R, Syafhan NF. Analysis of 4-hydroxycyclophosphamide in cancer patients plasma for therapeutic drug monitoring of cyclophosphamide. *Int J Pharm Pharm Sci*. 2016;8(9):194–200. doi:10.22159/ijpps.2016v8i9.12918
- Nakajima M, Komagata S, Fujiki Y, et al. Genetic polymorphisms of CYP2B6 affect the pharmacokinetics/pharmacodynamics of cyclophosphamide in Japanese cancer patients. *Pharmacogenet Genomics*. 2007;17(6):431–445. doi:10.1097/FPC.0b013e328045c4fb
- Coleman MD. *Human Drug Metabolism*. 2nd ed. USA: Wiley; 2010.
- Mehlota RK, Ziats MN, Bockarie MJ, Zimmerman PA. Prevalence of CYP2B6 alleles in malaria-endemic populations of West Africa and Papua New Guinea. *Eur J Clin Pharmacol*. 2006;62:267–275.
- Zanger UM, Klein K. Pharmacogenetics of cytochrome P450 2B6 (CYP2B6): advances on polymorphisms, mechanisms, and clinical relevance. *Front Genet*. 2013;4:24. doi:10.3389/fgene.2013.00024
- de Jonge ME, Huitema ADR, Rodenhuis S, Beijnen JH. Clinical pharmacokinetics of cyclophosphamide. *Clin Pharmacokinet*. 2005;44(11):1135–1164. doi:10.2165/00003088-200544110-00003
- Helsby NA, Yong M, van Kan M, de Zoysa JR, Burns KE. The importance of both CYP2C19 and CYP2B6 germline variations in cyclophosphamide pharmacokinetics and clinical outcomes. *Br J Clin Pharmacol*. 2019;1–10.
- Haroun F, Al-Shaar L, Habib RH, et al. Effects of CYP2B6 genetic polymorphisms in patients receiving cyclophosphamide combination chemotherapy for breast cancer. *Cancer Chemother Pharmacol*. 2015;75:207–214. doi:10.1007/s00280-014-2632-4
- Lang T, Klein K, Fischer J, et al. Extensive genetic polymorphism in the human CYP2B6 gene with impact on expression and function in human liver. *Pharmacogenetics*. 2001;11:399–415. doi:10.1097/00008571-200107000-00004
- Hiratsuka M, Takekuma Y, Endo N, et al. Allele and genotype frequencies of CYP2B6 and CYP3A5 in the Japanese population. *Eur J Clin Pharmacol*. 2002;58:417–421. doi:10.1007/s00228-002-0499-5
- Lamba V, Lamba J, Yasuda K, et al. Hepatic CYP2B6 expression: gender and ethnic differences and relationship to CYP2B6 genotype and CAR (constitutive androstane receptor) expression. *J Pharmacol Exp Ther*. 2003;307:906–922. doi:10.1124/jpet.103.054866
- Hananta L, Astuti I, Sadewa AH, et al. The Prevalence of CYP2B6 Gene Polymorphisms in Malaria-endemic Population of Timor in East Nusa Tenggara Indonesia. *Osong Public Health Res Perspect*. 2018;9(4):192–196. doi:10.24171/j.phrp.2018.9.4.08
- Melanson SEF, Stevenson K, Kim H, et al. Allelic variations in CYP2B6 and CYP2C19 and survival of patients receiving cyclophosphamide prior to myeloablative hematopoietic stem cell transplantation. *Am J Hematol*. 2010;85(12):967–971. doi:10.1002/ajh.21889
- Elmaagacli AH, Koldehoff M, Steckel NK, Trensche R, Ottinger H, Beelen DW. Cytochrome P450 2C19 loss of function polymorphism is associated with an increased treatment-related mortality in patients undergoing allogeneic transplantation. *Bone Marrow Transplant*. 2007;40:659–664. doi:10.1038/sj.bmt.1705786
- Zanger UM, Klein K, Saussele T, Blievernicht J, Hofmann M, Schwab M. Polymorphic CYP2B6: molecular mechanisms and emerging clinical significance. *Pharmacogenomics*. 2007;8:743–759. doi:10.2217/14622416.8.7.743
- Xie H, Griskevicius L, Stahle L, et al. Pharmacogenetics of cyclophosphamide in patients with hematological malignancies. *Eur J Pharm Sci*. 2006;27:54–61. doi:10.1016/j.ejps.2005.08.008
- Ariyoshi N, Miyazaki M, Toide K, Sawamura Y-I, Kamataki T. A single nucleotide polymorphism of CYP2B6 found in Japanese enhances catalytic activity by autoactivation. *Biochem Biophys Res Commun*. 2001;281:1256–1260. doi:10.1006/bbrc.2001.4524
- Xie HJ, Yasar U, Lundgren S, et al. Role of polymorphic human CYP2B6 in cyclophosphamide bioactivation. *Pharmacogenomics J*. 2003;3:53–61. doi:10.1038/sj.tpj.6500157
- de Andres F, Teran S, Hernandez F, Teran E, Llerena A. To genotype or phenotype for personalized medicine? CYP450 drug metabolizing enzyme genotype-phenotype concordance and discordance in the ecuadorian population. *OMICS*. 2016;20:699–710. doi:10.1089/omi.2016.0148
- Su G, Min H, Xin L, Xiao C, Eli C, Shu-Feng Z. Intra- and inter-ethnic differences in the allele frequencies of cytochrome P450 2B6 gene in Chinese. *Pharm Res*. 2006;23(9):1983–1990. doi:10.1007/s11095-006-9083-5
- Sofro ASM. *Population Genetic Studies in Indonesia [Dissertation]*. Canberra: Australian National University; 1982.
- Koentjaraningrat. *Metode Penelitian Masyarakat*. Jakarta: Gramedia Pustaka Utama; 1997.
- EMA. *Guideline on Bioanalytical Method Validation*. EMA; 2011.
- FDA. *Bioanalytical Method Validation Guidance*. FDA; 2018.
- Joy M, La M, Wang J, Bridges A, Hu Y, Hogan S. Cyclophosphamide and 4-hydroxycyclophosphamide pharmacokinetics in patients with glomerulonephritis secondary to lupus and small vessel vasculitis. *Br J Clin Pharmacol*. 2012;74(3):445–455. doi:10.1111/j.1365-2125.2012.04223.x
- Hall OM, Peer CJ, Fitzhugh CD, Figg WD. A sensitive and rapid ultrahigh - performance liquid chromatography with tandem mass spectrometric assay for the simultaneous quantitation of cyclophosphamide and the 4-hydroxy-cyclophosphamide metabolite in human plasma. *J Chromatogr B*. 2018;1086:56–62. doi:10.1016/j.jchromb.2018.04.016
- BC Cancer Agency. *Cancer Drug Manual: Cyclophosphamide*. Canada: BC Cancer Agency; 2013.
- Harahap Y, Yanuar A, Cyril M, Melhan M, Purwanto DJ. Quantification of 3-hydroxypropyl mercapturic acid in the urine of patients with breast cancer to monitor cyclophosphamide toxicity. *Ther Drug Monit*. 2020;42(4):548–553. doi:10.1097/FTD.0000000000000737
- London Cancer. *Guidelines for the Administration of Mesna with Ifosfamide and Cyclophosphamide*. London: London Cancer; 2014.

Drug Design, Development and Therapy

Dovepress

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which has also

been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/drug-design-development-and-therapy-journal>