Successful Sequential Treatment for Severe Asthma Coexisting COVID-19 via Budesonide/Glycopyrrolate/Formoterol Fumarate

This article was published in the following Dove Press journal:
International Journal of General Medicine

Yingjian Liang*
Meizhu Chen
Cuiyan Tan
Changli Tu
Xiaobin Zheng
Jing Liu

Department of Pulmonary and Critical Care Medicine (PCCM), The Fifth Affiliated Hospital of Sun Yat-Sen University, Zhuhai, People's Republic of China

*These authors contributed equally to this work

Abstract: Awareness of the management of coronavirus disease 2019 (COVID-19) and airway diseases can effectively help clinical physician during the coronavirus pandemic. Herein, we elucidated a COVID-19 case coexisting with severe asthma. Budesonide/glycopyrrolate/formoterol fumarate (BGF) was used as sequential medicine to systemic glucocorticoids for his persisted symptoms related to bronchospasms. Our case suggests patients with long-term airway diseases like asthma probably attribute their symptoms to COVID-19 instead of primary diseases, which make it more difficult in the symptom control. BGF is able to be an effective and convenient choice as sequential medicine to systemic glucocorticoids in some refractory asthmatic patients complicated with COVID-19.

Keywords: coronavirus disease 2019, asthma, budesonide/glycopyrrolate/formoterol fumarate, sequential treatment

Background

Coronavirus disease 2019 (COVID-19) is caused by a newly notorious coronavirus (severe acute respiratory syndrome coronavirus-2, SARS-CoV-2) and is of great global public health concern as it spiraled into a pandemic.¹ Recently, there has been increasing interest in recognizing the association between COVID-19 and airway diseases including asthma, especially about de-escalation therapy.^{2,3} However, there is no consensus regarding the sequential treatment of COVID-19 coexisting asthma. Herein, we elucidated our successful experience of using budesonide/glycopyrrolate/formoterol fumarate (BGF) in the sequential treatment of asthma coexisted with COVID-19.

Case Presentation

A 54-year-old Guinean man with 25-years smoking history was admitted to our hospital on 17th March 2020 for 3 weeks' non-production cough and sneeze as well as 2 weeks of fever after his travel from COVID-19-ridden part of England and was quarantined as a COVID-19 suspected infection patient. Peripheral ground glass opacity involved in bilateral multi-lobes was found in chest computed tomography scan (Figure 1) and he was confirmed as COVID-19 by real time polymerase chain reaction according to Chinese Center for Disease Control recommended methods. Oxygen support was given after the COVID-19 diagnosis. However, his oxygenation index descends and PaCO₂ ascends (17th March,

Correspondence: Jing Liu; Xiaobin Zheng
The Fifth Affiliated Hospital of Sun Yat-Sen University, 52 East Meihua Road, Zhuhai City, 519000, People's Republic of China
Tel/Fax +86(0)7562528733
Email liujing25@mail.sysu.edu.cn; zhxbn@mail.sysu.edu.cn

Figure 1 Evolution of CT scans. (A–C) represents lesions at admission. (D–F) demonstrates the recovery after out-of-bed activities.

PaCO₂ 30.6mmHg, pH 7.446, BE –3mmol/l; 23th March, PaCO₂ 102mmHg, pH 7.22, BE –5.7mmol/l) persisted even after high-flow nasal cannula (40L/min, 95%) and low-dose methylprednisolone therapy. He was transferred to ICU and received urgent intubation (23th March) for invasive ventilation (A/C, pressure control, Pi 17cmH₂O, PEEP 9cmH₂O, f 26/min, FiO₂ 100%) as abrupt worsening in PaO₂. However, deteriorated progression of his respiratory condition persisted. Due to the supplementary of long-term poor-controlled asthma as well as allergic rhinitis management history was provided from his family and obvious wheezing sound was caught, with a high level of total immunoglobulin E (301IU/mL) in his blood, clear diagnosis of acute asthma attack had been identified. In consideration of it, a large-dose systemic glucocorticoids therapy (intravenous methylprednisolone 4mg/kg/d) was administrated immediately. Respiratory condition improved gradually and tracheal intubation was removed. However, symptoms including chest tightness, dyspnea as well as wheezes still persisted in the course of systemic glucocorticoids converting to inhaled corticosteroid (ICS)/long-acting β 2-agonist (LABA) combination with dry powder inhaler because of his limited cooperation. Triple therapy with BGF with a novel pressure metered dose inhaler was used sequentially to prevents the adverse effect of long-term systematic glucocorticoids therapy. As a dramatic and rapid response to BGF, our patient successfully started out-of-bed activities, with recovery lung image (Figure 1).

Antibiotic was used in the whole course. (17th to 20th March, 5th to 7th, April, moxifloxacin; 20th to 23rd March, ceftriaxone; 23rd March to 5th April, meropenem, 20th to 5th April, linezolid)

Discussion and Conclusion

To our knowledge, it is the first report about the triple therapy with BGF, a combination of ICS, long-acting muscarinic antagonists (LAMA) and LABA with co-suspension delivery technology, used in asthma complicated with COVID-19. Although the occurrence ratio of COVID-19 in asthma were low, several reports have indicated that SARS-CoV-2 commonly trigger the acute attack of asthma.^{2,3} Our case is able to be a potential reminder that asthma can probably increase the diagnosis and treatment difficulty of COVID-19. Two reasons maybe contribute to the diagnosis difficulty according to our clinical experience. Firstly, it is not easy to distinguish COVID-19 with airway diseases because of the similar clinical features like cough, chest tightness and dyspnea.^{4,5} Another is the long-standing asthma patients commonly have poor perception of own symptoms, which make it hard to describe the exact degree of their obstruction in airway and probably attribute their symptoms to COVID-19.

Triple therapy with ICS, LAMA and LABA has become increasingly popular in clinical practice as an effective and convenient treatment. Recently, triple therapy with BGF has been identified effective and convenient among airway diseases as it can result in an improvement

in lung function through distinct and complementary mechanisms.⁶ BGF might be a good choice as sequential medicine to systemic glucocorticoids in some refractory asthma patients. In present case, BGF help our patient relieve the burden of symptoms as well as daily activities, which greatly enhance the doctors' confidence of ICS as well as bronchodilators used in COVID-19 complicated with airway disease. We believe that it may be informative for clinicians who may encounter this type of case in their daily practice and help physicians develop a strategy for optimizing their approach to the concomitant of asthma and COVID-19 management.

Limitation

Our patient cannot provide his pre-exacerbation pulmonary function test (PFT) result as well as the past treatment. In the computed tomography scan, emphysema can be seen in the right lung, which supports the diagnosis of chronic obstructive pulmonary disease. However, our patient did not perform PFT as the precaution of COVID-19 transmission. Thus, we cannot diagnose chronic obstructive pulmonary disease without bronchodilation test result.

Abbreviations

BGF, budesonide/glycopyrrolate/formoterol fumarate; COVID-19, coronavirus disease 2019; SARS-CoV-2; ICS, severe acute respiratory syndrome coronavirus-2; LABA, long-acting β 2-agonist; LAMA, long acting muscarinic antagonists.

Ethics Approval and Consent to Participate

Institutional approval is not applicable or required for publication of this manuscript.

Consent for Publication

Written and informed consent for publication was obtained from the patient. The patient was informed that de-identified data would be used in the scientific research and publications.

Funding

This work was supported by Zhuhai science and technology innovation Bureau (grant number ZH220363022000 21PWC).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Wang C, Horby PW, Hayden FG, Gao GF. A novel coronavirus outbreak of global health concern. *Lancet*. 2020;395(10223):470–473. doi:10.1016/S0140-6736(20)30185-9
2. Li YK, Peng S, Li LQ, et al. Clinical and transmission characteristics of covid-19 - a retrospective study of 25 cases from a single thoracic surgery department. *Curr Med Sci*. 2020;40(2):295–300. doi:10.1007/s11596-020-2176-2
3. Abrams EM, Geert WJ, Yang CL. Asthma and COVID-19. *CMAJ*. 2020.
4. Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA*. 2020;323(11):1061–1069. doi:10.1001/jama.2020.1585
5. Bateman ED, Hurd SS, Barnes PJ, et al. Global strategy for asthma management and prevention: GINA executive summary. *Eur Respir J*. 2008;31(1):143–178. doi:10.1183/09031936.00138707
6. Ishiura Y, Fujimura M, Ohkura N, et al. Triple therapy with budesonide/glycopyrrolate/formoterol fumarate improves inspiratory capacity in patients with asthma-chronic obstructive pulmonary disease overlap. *Int J Chron Obstruct Pulmon Dis*. 2020;15:269–277. doi:10.2147/COPD.S231004

International Journal of General Medicine

Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-general-medicine-journal>

Dovepress

across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.