Prevalence of Early Removal of Long-Acting Contraceptive Methods and Its Associated Factors in Sidama Regional State, Ethiopia

This article was published in the following Dove Press journal:
Open Access Journal of Contraception

Ephrem Geja¹
Fanuel Belayneh²
Deresse Legesse²
Desalegn Tsegaw²
Teshome Abuka²
Addis Gebremariam³
Henok Gebreyohannes⁴
Dawit Jember²
Zewdie Oltaye¹
Alealign Tadele⁵

¹School of Nursing, College of Medicine and Health Sciences, Hawassa University, Hawassa, Sidama Regional State, Ethiopia;

²School of Public Health, College of Medicine and Health Sciences, Hawassa University, Hawassa, Sidama Regional State, Ethiopia; ³Hawassa University Comprehensive Specialized Hospital, Hawassa, Sidama Regional State, Ethiopia;

⁴Yale Global Health Leadership Institute, Yale School of Public Health, Gondar, Ethiopia; ⁵Department of Medical Laboratory Science, Hawassa Health Sciences College, Hawassa, Sidama Regional State, Ethiopia

Background: Long-acting reversible contraceptive methods, IUD and sub-dermal implant, offer women the most effective method to control fertility. Yet, reports on high early removal rates were emerged, prompting concern among service providers and highlighting the need to review removal rates and its reasons. Therefore, this study was conducted to assess the prevalence of early removal rates of LARCs and its associated factors in Sidama Regional State, Southern Ethiopia.

Methods: Community-based cross-sectional study was conducted in Sidama Regional State, Ethiopia from June 1 to June 30, 2019. A multistage sampling technique was used to select 21 administrative units in the first stage. Then, systematic sampling was used to select 475 women who have ever used implants or IUD 3 years preceding the data collection period. Data were entered into Epi Info version 3.4.3 and exported to SPSS version 20 for analysis. Descriptive statistics, bivariate, and multivariate logistic regression were computed. P-value <0.05 was used to declare a significant association.

Results: The mean ($\pm$ SD) of the participant's age was 29.81 ($\pm$ 5.69) years. The prevalence of early removal rate of LARCs was 10.3%, ie, 43 (10.8%) among Implanon/jadelle users and 6 (7.8%) among IUD users. Eleven (22.4%) discontinued within the first six months and 38 (77.6%) utilized for more than six months and discontinued before the 12th month. Women who were not advised about advantage [OR= 2.81 (95% CI: 1.23–6.40)] and effectiveness of contraceptive [OR= 2.70 (95% CI: 1.30–5.60)] and those who were satisfied with the family planning service [OR= 2.24 (95% CI: 1.10–4.57)] were identified as factors.

Conclusions: The prevalence of early removal rate was considerably high among the study subjects. Providing appropriate counseling about the advantage and effectiveness of family planning and improving client satisfaction need to be considered to reduce the early removal rate.

Keywords: IUD, Implanon, jadelle

Introduction

Long-Acting Reversible Contraceptive (LARC) devices which include the Intra-Uterine copper Contraceptive (IUD) and the etonogestrel contraceptive implant are among currently available long-acting contraceptive options in Ethiopia. These methods are classified as long-acting methods due to long-term contraceptive efficacy.¹ Intra-Uterine Devices (IUDs) are T-shaped devices that must be inserted into a woman's uterus by a trained health care provider. Sub-dermal implants are progestin containing rods that are inserted under the skin and can prevent pregnancy for up to five years. Among all reversible modern contraceptives, IUDs are the most

Correspondence: Ephrem Geja
School of Nursing, College of Medicine and Health Sciences Hawassa University,
PO Box 05, Hawassa, Sidama Regional State, Ethiopia
Tel +251-911-066-463
Email ephi.geja@gmail.com

cost-effective and efficient methods in preventing pregnancy for up to 10 years.^{2,3} Implants are also long-acting and extremely effective at preventing pregnancy, with less than one percent of clinical failure rate.¹

Of course, no single method is appropriate for all women. However, despite there being a wide range of contraceptive methods, which are freely available, there are still large numbers of unintended pregnancies ending with abortion every year. For several reasons, many women find it difficult to use contraception consistently and correctly.⁴

LARC methods are commonly discontinued when there is pregnancy while using when a woman has a desire to become pregnant, infrequent sex, husband live away, marital dissolution, or menopause or when a woman is not satisfied with the method or cannot access or afford it.⁵ In different countries, a significant number of women become exposed to the risk of conception after discontinuation. Consequently, the majority of unintended pregnancies ended with abortion or miscarriage.⁶

In Ethiopia, among all women using implants; 2% will discontinue before 6 months and about 11% before 1 year after insertion. Among the IUD users, 27% and 34% will remove the device within 1 and 3 years of insertion.⁷ The study done in three regional states of Ethiopia shows, 15.1% of the respondents had at least one abortion episode.⁸

The controversy between high discontinuation rate and high demand suggests unmet need due to method dissatisfaction. This might increase women's vulnerability to unintended pregnancies-if they do not start another method of contraception. There are inadequate and inconsistent reports indicating high numbers of implant and IUD early removals. This will promote concern among local service providers and highlight the need to review the removal rates and reasons for removal.⁷ Besides, LARC methods are known for their high effectiveness and comprise an increasing share of contraceptive use. Therefore, it is imperative to examine how often and why women discontinue these methods, particularly in low-income countries.⁵ This study was conducted to determine implant and IUD early removal rates and its determinants in Sidama Regional State, Ethiopia.

Methods and Materials

Study Design and Setting

A community-based cross-sectional study was conducted in Sidama Regional State, Southern Ethiopia

from June 1 to 30, 2019. Sidama Regional State is found in the Southern part of Ethiopia. The Region is divided into 36 districts and on average each district has a population of 100,000. According to the Central Statistical Agency (CSA) population projection for 2016, the zone has a total population of 3,590,471 of whom 1,810,134 (50.48%) are men and 1,780,337 (49.52%) women. According to the 2016 Southern Nations Nationalities and Peoples Regional State/SNNPRS health bureau annual report, a total of 601,553 reproductive-age women were used modern family planning in Sidama Regional State. Of whom 92,519 (15.38%) women used LARCs. This research aims to assess the prevalence of early removal rate of long-acting and reversible contraceptives and its associated factors in Sidama Regional State, Southern Ethiopia, from June 1 to 30, 2019.

Population and Sampling

All reproductive age group women who ever used implants or IUDs from the selected 21 kebeles (the smallest administrative unit in Ethiopia) of Sidama Regional State (period of June 1, 2014, up to May 30, 2017) were recruited as the source population of this study. The study population was selected from those women who used either of the above LARCs during the last three years preceding the data collection period. Women who started using LARCs within less than twelve months period preceding the study were excluded.

The required sample size was estimated assuming; 35.0% of women who removed LARCs methods early after insertion (9), 0.05 level of significance (α), and 5% limits of error. Also, a design effect of 1.5 was used to minimize sampling error. Thus, the estimated sample size with the addition of a 10% non-response rate was 525.

A multi-stage sampling technique was implemented. In the first stage, six districts and one city administration were selected by a simple random sampling technique using the lottery method. Then, 21 kebeles were selected similarly. Women who ever used LARCs in the selected kebeles were identified from family planning registration books of health facilities and family planning utilization documents at health posts. Finally, a systematic sampling technique was employed to select study participants from the identified list of women who ever used these methods.

Data Collection Procedure

A structured and pre-tested interview-based questionnaire with both open and closed-ended questions was used to collect data. The questionnaire was developed after reviewing different literature and scientific facts. First, it was prepared in English and translated into Amharic. The Amharic version was retranslated back to English by another person to check whether the translation was consistent with the English version. The data collection tool contains questions about socioeconomic and demographic conditions, past contraception history and Knowledge on LARCs, partner involvement, counseling status, future intention, and information specifically about the use of LARCs. Seven data collectors and two supervisors have collected the data through the home to home visit. Local guides were used to identify the house of the selected women.

Measurement

Early removal of long-acting and reversible contraceptive: To be consistent with previously published data, early discontinuation was defined as the removal of the Implants or IUD within one year of its placement.¹⁰

Data Processing and Analysis

Data were entered into Epi Info version 3.4.3 and then exported to SPSS version 20 for analysis. Summary statistics were computed to determine frequencies and percentages. Bivariate analysis was conducted primarily to check the association of each independent variable with the dependent variable-early removal of LARCs. To control for the possible effect of confounding, variables found to have an association with the dependent variables at a P-value of 0.2 on the bivariate logistic regression analyses were entered into multivariate logistic regression analysis. Those variables which had significant association at p-value less than 0.05 in the multivariate logistic regression were considered to be independent predictors of early removal of LARCs. The final model was diagnosed by Hosmer and Lemeshow goodness of fit tests. Results were presented using tables, charts, and graphs.

Data Quality Control

To ensure the quality of data, the questionnaire was pre-tested on 5% of the total sample size out of the selected kebeles. Possible restructuring and adjustment of the questions were made after the pretest. Orientation was given for data

collectors and supervisors. The data collection process was conducted with close supervision of the assigned supervisors, while completeness and accuracy of collected data were again reviewed and checked by the principal investigators.

Ethical Consideration

Ethical clearance was obtained from the Institution Review Board of Hawassa University, College of Medicine and Health Sciences. Permission and support letter was obtained from the SNNPRS Health Bureau and the Sidama Regional State Health Department. Then an official letter was written to each service delivery point. An information sheet that contains about the benefit and risk of participating in the respondents in this study with verbal informed consent was attached to each questionnaire to brief and obtained the consent of each study participant. Verbal informed consent obtained from the participants was approved by the Institution Review Board of Hawassa University. This study was also conducted in accordance with the Declaration of Helsinki.

Result

Socio-Demographic Characteristics of Study Participants

A total of 475 reproductive age group women were included in the study but 429 participants were participated and make the response rate of 90.4%. The mean ($\pm$ Standard deviation (SD)) age of the respondents was 29.81 ($\pm$ 5.69) years Urban residents account for 408 (85.9%) of women. Nearly half 273 (49.9%) were Protestant by religion followed by orthodox, 203 (43.7%). Four hundred forty-three (93.3%) of the LARCs users were married and 22 (4.6%) were singles. Only 9 (1.95%) of the respondents had no formal education while 443 (93.3%) had formal education. Housewives account for 286 (60.2%), followed by 124 (26.2%) government employees. Thirty-seven (7.8%) of the women were not able to estimate their monthly income and 105 (22.1%) of them reported an estimated monthly income between 2000 and 3000 Ethiopian birrs (Table 1).

Maternal and Reproductive Characteristics

One fourth (25.3%) had two children before using LARC methods. Currently, one among five 91 (19.2%) women has more than four children. One hundred and twenty-seven women (26.7%) had no intention to have children

Table 1 Socio-Demographic Characteristics of Women Who Received LRCMs in Sidama Regional State, Southern Ethiopia, 2019

Characteristics		Frequency	Percent
Age group	≤24	55	11.6
	25–29	203	42.7
	30–34	112	23.6
	35–39	73	15.4
	≥40	32	6.7
Residence	Urban	408	85.9
	Rural	67	14.6
Religion	Protestant	237	49.9
	Orthodox	203	42.7
	Other	35	7.3
Marital status	Single	22	4.6
	Married	443	93.3
	Others	10	2.2
Education	No formal education	32	6.7
	<8 years	149	31.4
	9–12 years	199	41.9
	>12 years	95	20
Occupation	Student	11	2.3
	Merchant	41	8.6
	Government worker	124	26.1
	House wife	286	60.2
	Others	13	2.7
Monthly income	Do not know	37	7.8
	≤1000 birr	32	6.7
	1001–2000 birr	89	18.7
	2001–3000 birr	105	22.1
	3001–4000 birr	66	13.9
	4001–5000 birr	65	13.7
	>5000 birr	81	17.7

in the future, whereas, 52 (10.9%), 110 (23.2%), and 100 (21.1%) had an intention for one, two, and three children in the future, respectively. Women who had a history of at least one abortion were 79 (16.6%) (Table 2).

Family Planning Knowledge and Service

Nearly all, 467 (98.3%) of women have information about injectable contraceptives. Three hundred eighty-seven (81.5%) of them have also responded that they know about oral contraceptive pills. Those participants who reported as having information about Implanon, jadelle, and IUD were 416 (68.8%), 383 (80.6%), and 306 (64.4%)

Table 2 Maternal and Reproductive Characteristics of Women Who Received LRCMs in Sidama Regional State, Southern Ethiopia, 2019

Characteristics	Frequency	Percent
Children before utilization		
No children	31	6.5
1 child	65	13.7
2 children	120	25.3
3 children	97	20.4
4 children	71	14.9
≥ 5 children	91	19.2
Current children		
No children	26	5.5
1 child	52	10.9
2 children	110	23.2
3 children	100	21.1
≥ 4 children	187	39.4
Intention to have more children		
No intention	127	26.7
1 child	61	12.8
2 children	211	44.1
≥ 3 children	76	16
Abortion history		
Yes	79	16.6
No	396	83.4

respectively. Health professionals were the source of information for 412 (86.7%) of women in this study. Duration of pregnancy protection was the most reported known type of information on both methods; intradermal (86.7%) and IUD (82.5%) (Table 3).

Of the total women in this study, 354 (74.5%) received a family planning service for LARC from health centers. Four hundred twenty-three of them reach the health facilities within less than 30 minutes of walking. Almost all (99%) women received counseling before the insertion of the method. The counseling duration was less than 15 minutes for 281 (59.2%) of the clients. The family planning service was satisfying for 349 (73.5%) of the users (Table 4).

Information about the advantage and duration of action of LARC methods were the most frequently addressed topics during counseling. Four hundred (84.2%) of

Table 3 Information on FPM Among Women Who Used LRCMs in Sidama Regional State, 2019

Characteristics	Frequency	Percent
Has information about		
Pills	387	81.5
Injectables	467	98.3
Lactation amenorrhea	61	12.8
Female condom	40	8.4
Male condom	327	68.8
Implanon	416	87.6
Jadelle	383	80.6
IUD	306	64.4
Female Sterilization	49	10.3
Male Sterilization	29	6.1
Information about intradermal methods		
Effectiveness	375	78.9
Side effects	63	13.3
Duration of pregnancy prevention	412	86.7
Benefit	263	55.4
Information about IUD		
Effectiveness	332	69.9
Side effects	262	55.2
Duration of pregnancy prevention	392	82.5
Benefit	280	58.9
Source of information		
Health professional	412	86.7
HEW	193	40.6
Friends or families	108	22.7
Radio/ television	197	41.5
News paper/ magazine	36	7.6
Other printed materials	2	0.4

women were advised about the advantage of these methods and 373 (78.5%) about its duration of action (Figure 1).

Family Planning Methods Utilization and Early Removal

There were 110 (23.2%) women who ever used oral contraceptives and 374 (78.7%) injectable methods. Implanon, Jadelle and IUD ever users were 207 (43.6), 191 (40.2%), and 77 (16.2%) respectively. From the total 398 intradermal (Implanon/Jadelle) contraceptive users, 320 (80.4%) are still using the methods currently. Out of 77 IUD ever users in this study, 55 (71.4%) are using currently (Figure 2).

Table 4 Family Planning Services Provided for LRCMs Users in Sidam Regional State, 2019

Characteristics	Frequency	Percent
Place of insertion		
Health center	354	74.5
Health post	8	1.7
Hospital	75	15.8
NGO health facility	29	6.1
Private health facility	9	1.9
Time to reach the source		
≤30 minutes	423	89.1
>30 minutes	52	10.9
Get counseling before insertion		
Yes	471	99.2
No	4	0.8
Type of counseling		
Individual counseling	364	76.6
Mass counseling	81	17.1
With husband counseling	30	6.3
Duration of counseling		
≤15 minutes	281	59.2
15–30 minutes	189	39.8
>30 minutes	5	1.1
Duration before insertion		
<30 minutes	343	72.2
30 minutes- 1 day	44	9.3
>1 day	88	18.5
Service satisfaction		
Satisfied	349	73.5
Unsatisfied	126	26.5

Of the total participants in this study, 49 discontinued LARC methods within 12 months of use. Based on this, the overall early removal rate of LARCs was 10.3% (95% confidence interval: 7.57–13.06). The proportion of early removal was 10.8% among implant users and 7.8% among IUD users. Eleven (22.4%) of the total early discontinuers used these methods for less than 6 months and the remaining 38 (77.6%) used 6 to 12 months (Figure 3).

The three most common reasons for discontinuation of LARCs were; plan to conceive soon (28.8%), fear of side effect (27.4%), and health concerns together with bodyweight loss (16.4%). Weight gain with health concerns was mentioned by 4.1% of women as a reason for the removal (Figure 4).

Figure 1 Distribution of counseling information given for LRCMs utilizers in Sidama Regional State, Ethiopia, 2019.

Figure 2 Distribution of ever used family planning methods and current LRCMs use among women who received LRCMs in Sidama Regional State, Southern Ethiopia, 2019.

Factors Associated with Early Removal of LRCMs

On the bivariate analysis, age, educational status, intention to have more children, service satisfaction, counseling about advantage, side effect, and effectiveness were selected as a candidate variable for multivariate analysis. On the final model; women's age, educational status, service satisfaction, counseling about advantage, and effectiveness were identified as independent predictors of LARC's early removal.

Women whose age is between 25 and 29 years [OR= 9.86 (95% CI: 2.22–43.74)] and those between 30 and 34 years [OR= 11.5 (95% CI: 2.67–49.56)] were about 10 and

11 times more likely to remove LARCs within one year period than women whose age is above 35 years. When compared to women with no formal education, those who spent ≤ 8 years [OR= 0.32 (95% CI: 0.11–0.93)], 9–12 years [OR= 0.19 (95% CI: 0.06–0.58)] and >12 years [OR= 0.26 (95% CI: 0.26–0.98)] in school are 68%, 81% and 74% less chance of removing LRCMs before one year of use, respectively.

Women who are not advised about the advantage [OR= 2.81 (95% CI: 1.23–6.40)] and effectiveness [OR= 2.70 (95% CI: 1.30–5.60)] of LARCs had three times higher probability of discontinuing the method within 12 months of insertion than those who received pieces of advice on

Figure 3 Distribution of early LRCMs removal and Duration for removal among women who received LRCMs in Sidama Regional State, Southern Ethiopia, 2019.

Figure 4 Reasons for removal of LRCMs mentioned by women who removed the methods in Sidama Regional State, Southern Ethiopia, 2019.

advantage and effectiveness. Unsatisfied women with the family planning service had about two times greater risk of early LARCs removal than those who are satisfied [OR= 2.24 (95% CI: 1.10–4.57)] (Table 5).

Discussion

In this study, early removal of LARC methods was defined as discontinuation of the methods within 12 months of use.

Based on this, the prevalence of LARCs early removal rate was 10.3%. Out of these, 22.4% discontinued within 6 months of use and the remaining 77.6% discontinued within 6 to 12 months of use.

The prevalence of LARC methods early removal rate in this study was higher than the results of the study done in Agarfa District, Bale Zone, South East Ethiopia that reported implants had a discontinuation rate of just 4.0%.¹¹

Table 5 Factors Associated with Early Removal of LRCMs Among Women Who Used These Methods in Sidam Regional State, 2019

Variables (n=475)	LRCM Early Removal		COR (95% CI)	AOR (95% CI)
	Yes	No		
	No(%)	No(%)		
Age group				
≤24	3 (5.5)	52 (94.5)	1.96 (0.38–10.05)	2.37 (0.37–14.93)
25–29	26 (12.8)	177 (87.2)	4.99 (1.47–16.91)*	9.86 (2.22–43.74)**
30–34	17 (15.2)	95 (84.8)	6.08 (1.72–21.42)*	11.5 (2.67–49.56)**
≥35	3 (2.9)	102 (97.1)	1	1
Educational status				
No formal education	9 (28.1)	23 (71.9)	1	1
≤8 years	18 (12.1)	131 (87.9)	0.35 (0.14–0.87)*	0.32 (0.11–0.93)**
9–12 years	16 (8.0)	183 (92.0)	0.22 (0.08–0.56)*	0.19 (0.06–0.58)**
>12 years	6 (6.3)	89 (93.7)	0.17 (0.05–0.533)*	0.26 (0.06–0.98)**
Intention for more children				
No intention	9 (7.1)	118 (92.9)	0.40 (0.16–1.07)	0.51 (0.15–1.70)
1–2 children	28 (10.3)	244 (89.7)	0.61 (0.29–1.27)	0.52 (0.21–1.27)
≥ 3 children	12 (15.8)	64 (84.2)	1	1
Counseled about advantage				
Yes	31 (7.8)	369 (92.2)	1	1
No	18 (24.0)	57 (76.0)	3.75 (1.97–7.16)*	2.81 (1.23–6.40)**
Counseled about side effects				
Yes	6 (19.4)	25 (80.6)	2.23 (0.87–5.75)	2.08 (0.57–7.58)
No	43 (9.7)	401 (90.3)	1	1
Counseled about effectiveness				
Yes	14 (5.8)	226 (94.2)	1	1
No	35 (14.9)	200 (85.1)	2.82 (1.47–5.40)*	2.70 (1.30–5.60)**
Service satisfaction				
Satisfied	26 (7.4)	323 (92.3)	1	1
Unsatisfied	23 (18.3)	103 (82.4)	2.77 (1.51–5.07)*	2.24 (1.10–4.57)**

Note: *P-value<0.05 on COR, **P-value<0.05 on AOR.

This difference might be due to the difference in the age of participants, science, the majority of women in the current study were young, and being young can be associated with a high desire of having more children which in turn leads to a high discontinuation rate. The other possible reason might be inadequate pre-insertion counseling particularly about the advantage and effectiveness of the method.

In contrast, the early LARC removal rate was lower than the contraceptive discontinuation in developing countries in the first year of use; which is 18–63%.¹² It was also lower than the results of studies conducted in Nigeria,

Egypt, Malaysia, and a multi-centered study in other nine countries, which indicated the overall Implanon discontinuation rates of 26.1%, 28.0%, 22.86%, and 20.0%, respectively.^{13–16} The rate was also lower than a 35% national early discontinuation in Ethiopia.⁹ Besides, in the study conducted in Debre Markos, Ethiopia, the rates of discontinuation at 6 months, 12 months, and 24 months were 10.5%, 23.9%, and 38.2%, respectively.¹⁷ These discrepancies could be due to the difference in the composition of study participants as most of these studies are conducted by focusing on the rural population.

Also, the expansion of the health extension program in our study area and the country from time to time might be the reason for the observed difference.

The satisfaction of women with family planning service and improved counseling has a positive impact on the early removal of LARC in our study. This finding was in line with results from the study at the Medical University of South Carolina.¹⁸

In our study, age group, educational status, service satisfaction, counseling about advantage, and effectiveness were identified as independent predictors of early removal of LRCMs. Similarly, the study conducted in Debre Markos Ethiopia showed that attending college and above level education, absence of pre-insertion counseling service, and perceived level of unsatisfying family planning service were significantly associated with Implanon discontinuation.¹⁷

The most common reasons mentioned for early removal by the current study participants were the desire for pregnancy and fear of side effects. This was in line with previous studies, which showed a plan to conceive soon, and the view of side effects was the most frequently given reason.^{19–21}

Despite its strengths, the study has some limitations. Qualitative methods were not included in the study which would support the quantitative results. Besides, some of the women who ever used LARCs outside of the catchment area and whose information about their utilization is inaccessible from health extension workers were not included.

Conclusions and Recommendations

The early removal rate of LARC is in considerable magnitude among women who ever used these methods in Sidama Regional State. The main predictors for early discontinuation of these methods were family planning service-related factors. These include; perceived satisfaction of clients by the family planning service, provision of proper counseling about the advantage, and effectiveness of the methods. Also, socio-demographic factors like; age less than 35 years and educational status are among the significant predictors for early discontinuation of these methods.

Therefore, measures targeted to improve the satisfaction of clients with family planning services and proper counseling on the advantages and effectiveness of these methods should be considered to get LARCs early removal rate lower.

Abbreviations

CSA, Central Statics Agency; FGA, Family Guidance Association; FP, family planning; HEW, health extension

worker; IUDs, intrauterine devices; LARC, long-acting reversible contraceptive; MCH, maternal and child health; REC, Research and Ethical Committee; SRS, Sidama Regional State; SPSS, Statistical Package for Social Science.

Data Sharing Statement

Data supporting the findings of this study are presented in the manuscript.

Ethics Approval and Consent to Participate

Ethical clearance was obtained from the Ethical review committee of the Hawassa University College of Medicine and Health Science & permission was obtained from each participant.

Acknowledgments

We would like to express our gratitude to the study participants for their willingness in providing the necessary information. We would like to acknowledge health Extension workers of the selected kebeles for easing the data collection process. Our gratitude also goes to EzedineMolla, Seife Kibru, AmezenechAyele, Mohammed Ayalew, Fikru Tadesse, BediluDeribe, Sihne Tadesse, Kurabachew Mengistu, Tadesse Yohannes, Sileshi Demelash&Abraham Abate for their unreserved support though out the paper preparation.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis, and interpretation, or in all these areas; took part in drafting, revising, or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

There is no funding to report.

Disclosure

All authors declare that they have no conflicts of interest for this work.

References

1. Ramchandran D, Upadhyay UD. *Implants: The Next Generation*. Population Reports. Baltimore, INFO Project, Johns Hopkins Bloomberg School of Public Health. October 2007. Available from: <http://www.populationreports.org/k7/>. Accessed January 28, 2021.
2. Ali M, Sadler R, Cleland J, Ngo T, Shah I. *Long-Term Contraceptive Protection Discontinuation and Switching Behaviour: Intrauterine Device (IUD) Use Dynamics in 14 Developing Countries*. London: World Health Organization and Marie Stopes International; 2011.
3. Hatcher R, Trussell J, Nelson A. *Contraceptive Technology*. New York, NY: Ardent Media; 2007.
4. Astbury Ward E. Provision of contraception and its influence on abortion. *Nurs Pract*. 2009;2009:47.
5. Staveteig S, Mallick L, Winter R. *Uptake and Discontinuation of Long-Acting Reversible Contraceptives (Larcs) in Low-Income Countries*. DHS Analytical Studies No. 54. Rockville, Maryland, USA. 2015. Available from: <https://www.dhsprogram.com/publications/publication-as54-analytical-studies.cfm>. Accessed January 28, 2021.
6. Frost JJ, Singh S, Finer LB. U.S. women's one-year contraceptive use patterns, 2004. *Perspect Sex Reprod Health*. 2007;38–55.
7. Kidane A, Asefa T, Asres T, et al. *Assessment of Implant and IUCD Removal in Ethiopia International Conference on Family Planning Research and Best Practices Kampala, Uganda November 2009*. Family Health International. 2009.
8. Melkamu Y, Betre M, Tesfaye S. Utilization of post-abortion care services in three regional states of Ethiopia. *Ethiop J Health Dev*. 2010;2010(24 Special Issues 1):123–129.
9. Central Statistical Agency (CSA) [Ethiopia] and ICF. *Ethiopia Demographic and Health Survey 2016*. 2016.
10. Robinson, Register, Ebner, and Orr, Reducing Early Discontinuation Rates of Subdermal Contraception in Your Clinical Practice. In: *Osteopathic Family Physician*. Vol. 1. Addis Ababa, Ethiopia, and Rockville, Maryland, USA: CSA and ICF; 2015:24–28.
11. Bekele T, Gebremariam A, Tura P. Factors associated with contraceptive discontinuation in Agarfa District, Bale Zone, Southeast Ethiopia. *Epidemiology (Sunnyvale)*. 2015;5(2):179. doi:10.4172/2161-1165.1000189
12. Bradley S, Schwandt HM, Khan S. *Levels Trends and Reasons for Contraceptive Discontinuation*. Calverton, Maryland, USA: ICF Macro. 2008.
13. Balogun O, Olaomo N, Adeniran A, Fawole A. Implanon sub-dermal implant: an emerging method of contraception in Ilorin, Nigeria. *J Med Biomed Sci*. 2014;3(1):1–5. doi:10.4314/jmbs.v3i1.1
14. Mastor A, Khaing SL, Omar SZ. Users' perspectives on Implanon in Malaysia, a multicultural Asian country. *Open Access J Contracept*. 2011;Volume 2:79–84. doi:10.2147/OAJC.S9674
15. Said MA-RM, Cairo U. *Implanon Use Pattern Among Ministry of Health and Population Clients 2008–2012*. Cairo: Faculty of Medicine; 2012.
16. Peipert Jeffrey F, Zhao Q, Allsworth JE, et al. Continuation and satisfaction of reversible contraception. *Obstet Gynecol*. 2011;117(5):1105–1113. doi:10.1097/AOG.0b013e31821188ad
17. Siyoum M, Mulaw Z, Abuhay M, Kebebe H. Implanon discontinuation rate and associated factors among women who ever used implanon in the last three years in Debre Markos Town, Northwest Ethiopia, 2016, cross sectional study. *ARC J Public Health Community Med*. 2017;2(1):8–16.
18. Lori M, Vanessa A, Jessica J, et al. Satisfaction, early removal, and side effects associated with long-acting reversible contraception. *Fam Med*. 2013;45(10):701–707.
19. Obijuru L, Bumpus S, Auinger P, et al. Etonogestrel implants in adolescents: experience, satisfaction, and continuation. *J Adolesc Health*. 2016;58(3):284. doi:10.1016/j.jadohealth.2015.10.254
20. Harel Z, Frank M, Kollar L, et al. Adolescents' reasons for and experience after discontinuation of the long-acting contraceptives Depo-Provera and Norplant. *J Adolesc Health*. 1996;19(2):118. doi:10.1016/1054-139X(95)00322-J
21. Lara-Torre E, Spotswood L, Correia N, et al. Intrauterine contraception in adolescents and young women: a descriptive study of use, side effects, and compliance. *J Pediatr Adolesc Gynecol*. 2011;24:39.

Open Access Journal of Contraception

Dovepress

Publish your work in this journal

Open Access Journal of Contraception is an international, peer-reviewed, open access, online journal, publishing original research, reports, reviews and commentaries on all areas of contraception. In addition to clinical research, demographics and health-related aspects, the journal welcomes new findings in animal and preclinical

studies relating to understanding the biological mechanisms and practical development of new contraceptive agents. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/open-access-journal-of-contraception-journal>