

# Evaluation of the Efficacy of Alcohol-Based Hand Sanitizers Sold in Southwest Ethiopia

This article was published in the following Dove Press journal:  
Infection and Drug Resistance

Gizachew Manaye <sup>1</sup>  
Desalegn Muleta<sup>1</sup>  
Andualem Henok<sup>2</sup>  
Abyot Asres <sup>2</sup>  
Yitagesu Mamo<sup>3</sup>  
Desalegn Feyissa <sup>3</sup>  
Fikadu Ejeta<sup>3</sup>  
Wondwossen Niguse <sup>4</sup>

<sup>1</sup>Department of Medical Laboratory Science, College of Medicine and Health Sciences, Mizan-Tepi University, Mizan Teferi, Ethiopia; <sup>2</sup>Department of Public Health, College of Medicine and Health Sciences, Mizan-Tepi University, Mizan Teferi, Ethiopia; <sup>3</sup>Department of Pharmacy, College of Medicine and Health Sciences, Mizan-Tepi University, Mizan Teferi, Ethiopia; <sup>4</sup>Department of Nursing, College of Medicine and Health Sciences, Mizan-Tepi University, Mizan Teferi, Ethiopia

**Background:** Regarding the protection of community and health professionals suffering from a COVID-19 outbreak, currently different alcohol-based hand sanitizers have been distributed. Even though for effective protection effective alcohol-based hand sanitizers are mandatory. Their efficacy was not evaluated. This is the reason why this research was designed to assess the antimicrobial efficacy of hand sanitizers that have been sold in southern parts of Ethiopia.

**Methods:** Six test organisms (*Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Salmonella typhimurium* and *Shigella boydii*) were selected from different clinical specimens. Then seven locally made products of alcohol-based hand sanitizers (MTU, Folium Fine, Epharm, Harego, Taflen and Sheba) were purchased and the disc diffusion, minimum inhibition concentration and minimum bactericidal concentration test were done against selected test organisms. Quality control measures throughout the whole process of the laboratory work were implemented and descriptive parameters were analyzed.

**Results:** The zones of inhibition of the hand sanitizers at their claim concentration were ranged from noninhibition zone (Folium) to 27mm (Sheba). The minimum inhibitory concentration against all selected test organisms was observed at 45%, 55%, 65% and Taflen on undiluted form. The growth of test organisms was decreased across increasing the concentration gradient of different hand sanitizers. MTU except against *E. coli*, Fine, Hargo, Ephra and Sheba hand sanitizers were showed growth below 60% concentrations for all test microorganisms. But a Folium product against all selected test microorganisms and MTU product against *E. coli* were not bactericidal.

**Conclusion:** Fine, Hargo, Ephra, Sheba and Taflen sanitizers were the products that were effective in inhibiting the growth of all the selected test organisms, and they were having effective bactericidal activity in vitro at their claim concentration. But MTU product against *E. coli* and Folium product against all test microorganisms were not having an effective bactericidal activity. So, regulatory authorities and manufacturers should implement strict quality control measures and regular observations throughout the production to ensure the efficacy of hand sanitizers.

**Keywords:** alcohol-based hand sanitizers, hand sanitizers, sanitizers, COVID-19

## Background

Hospital-associated and community-acquired infections being a serious public health problem all over the world and have become a major concern.<sup>1</sup> Hospital-acquired infections are infections developing in hospitalized patients,<sup>2</sup> whereas community-acquired infection on the other hand is an infection acquired anywhere other than a healthcare facility.<sup>1</sup> These infections have considerable impacts on

Correspondence: Gizachew Manaye  
Department of Medical Laboratory Science, College of Medicine and Health Sciences, Mizan-Tepi University, P.O. Box: 206, MizanTefri, Ethiopia  
Tel +251928444333  
Email ayele.gizachew@yahoo.com

individuals such as prolonged hospitalization, disability, increased risk of antimicrobial resistance, huge financial burden and deaths.<sup>3</sup>

The Center for Disease Control and Prevention (CDC) showed that about 2 million people acquire hospital-associated infections every year and about 90,000 of these patients die as a result of their infections.<sup>4</sup> Currently, severe acute respiratory syndrome coronavirus 2 (*SARS-CoV-2*) is a viral infectious agent of both Hospital-associated and community-acquired infections, and it becomes a serious issue in the world. Greater than 2 million people were infected and 139,378 peoples died with it exclusively until this proposal was developed.<sup>5</sup>

Different organizations such as WHO, CDC, and other professional document that practicing hand hygiene is the simplest and preferable technique in the prevention of the spread of the current serious issue of SARS-CoV-2 and other community-acquired and hospital-associated infections.<sup>3,6</sup> Hand hygiene can be done by several methods; the most important method is frequent hand washing with soap and plain water or sanitizing hands with alcohol-based hand sanitizer (ABHS) in the absence of water and soap.<sup>7-9</sup>

The use of ABHS has been reported as the common recommended means of hand hygiene.<sup>10</sup> The most regularly utilized ABHS should have an alcohol concentration of 60 to 85%.<sup>11</sup> The effectiveness of these ABHS depends on the concentration of alcohol and the time of rubbing. Rubbing hands with ABHS (60 to 85% concentrations) for 25–30 seconds was reported to kill 99.99% of microorganisms on hand. For instance, rubbing sanitizers with 70% alcohol is suggested to kill 99.9% of the microorganisms on hands.<sup>12</sup>

Currently, in Ethiopia, different ABHS have been distributed throughout the community and the market, regarding the protection of community and health professionals suffering from a coronavirus disease 2019 (COVID-2019) outbreak. But still now, the efficacy of them was not evaluated even though, for effective protection, effective ABHS is mandatory. This is the reason why this research was designed to evaluate the antimicrobial efficacy of alcohol-based hand sanitizer that has been sold in southern parts of Ethiopia.

The finding of this study will be used for better understanding of status of ABHS that have been sold and distributed in the community. The result will be helpful for ministry of health, policy-makers, and health professionals, other stakeholders and community partners

working on COVID-19 outbreak prevention strategy. It may also help as a baseline data for further related studies and early evaluating of alcohol-based hand sanitizer.

## Methods and Materials

### Test Organisms

Sanitizers containing an alcohol concentration between 60–85% are suggested to kill 99.99% of microorganisms on hands. The sanitizers which are effective on bacteria are effective in virus and vice versa. Therefore, we were forced to assess the efficacy of alcohol-based hand sanitizers only on selected bacteria, but not on virus (*SARS-CoV-2*) due to our laboratory set up constraint for viral isolation and growth. Following this assumption and reviewing related literature, *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Salmonella typhimurium* and *Shigella boydii* were selected at the department of microbiology laboratory at Mizan Tepi University from different left over clinical specimen (Urine, stool and wound) which were collected with laboratory technologists for laboratory test at Mizan Tepi teaching hospital. The microorganisms were isolated from clinical specimens with culture, gram stain, and biochemical test. During the study, clinical specimens were inoculated on chocolate agar, blood agar, MacConkey agar, and Mannitol salt agar, and incubated overnight (35°C - 37°C) for colony isolation. Then on the second day gram stain test was examined to differentiate gram positive from negative bacteria for all colonies. At the 3rd day test organisms were isolated with biochemical tests using their biochemical characteristics after overnight incubation (35°C - 37°C). The isolated test organisms were stored on storage media and kept at 2–8° C. It were refreshed on nutrient agar and used when needed.

### Hand Sanitizers

Seven locally made products of alcohol-based hand sanitizers were purchased from local vendors in Southwest, Ethiopia (Table). Sanitizers that have been sold on markets southern parts of Ethiopia were included and tear or not sealed expired and unlabeled (unknown manufacturer) hand sanitizers were excluded from the study (Table 1).

### McFarland (0.5 Turbidity Standard) Preparation

A 0.5 McFarland turbidity standard was prepared for the standardization of selected test organisms that are needed for the determination of the minimum inhibitory and

**Table I** Hand Sanitizers Used in the Study and Their Ingredients

Hand Sanitizers	Concentration	Product Ingredients
MTU	80% Ethanol alcohol	96%=Ethanol alcohol, 0.125=Glycerin, 1.45=Hydrogen peroxide
Folium	70% Ethanol alcohol	Ethanol alcohol, the other additives unknown
Fine	80% Ethanol alcohol	Ethyl alcohol 96%, Glycerin 98%, Hydrogen peroxide 3%
Epharm	80% Ethanol alcohol	80%=Ethanol alcohol, 0.125=Glycerin, 1.45=Hydrogen peroxide
Harego	80% Ethanol alcohol	80%=Ethanol alcohol, 0.125=Glycerin, 1.45=Hydrogen peroxide
Taflen	Unknown (concentration was unlabeled)	Unknown alcohol 95%, Glycerin 98%, Hydrogen peroxide 3%
Sheba	80% Ethanol alcohol	80%=Ethanol alcohol

bactericidal concentration, for performing the agar diffusion test, and other tests necessary for the study such as biochemical tests. This 0.5 McFarland turbidity standard was prepared into a test tube from the mixture of barium chloride dehydrate ( $\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$ ) solution and sulfuric acid ( $\text{H}_2\text{SO}_4$ ) with verification of the mixture absorbance (0.08–0.10) density precision via a spectrophotometer at a wavelength of 625nm. The prepared tube was stored in a sealed container at room temperature in a dark place.<sup>13</sup>

### Test Organism Standardization

Each selected test organism was standardized before the research was performed by taking a loopful of inoculum using a sterile loop from a pure culture, transferring and suspending it into tubes of sterile normal saline. Then, the suspension density was adjusting by comparing it with the 0.5 McFarland turbidity standards by either adding additional bacteria or sterile saline.<sup>14</sup>

### Agar Diffusion Test

To determine the susceptibility test of selected test organisms for each hand sanitizer well variant agar diffusion method was used.<sup>15</sup> This agar diffusion method was done in duplicates for each hand sanitizer by inoculated sterile

Mueller Hinton agar plates using sterile cotton swab which was immersed into a tube contain standardized test organisms. After Mueller Hinton agar was inoculated, it was allowed to remain at room temperature to dry for only some minutes and 8 equally spaced holes were bored in the agar plate with the 8th hole at the center of the plate with the aid of a sterile 6mm cork borer. The 7 holes were filled with 50µL of the hand sanitizer at the same time as the central hole (8th) was filled with an equal volume of sterile water for control purposes. The Mueller Hinton agar was incubated for 24 hours at 37° C. The zones of inhibition (susceptibility or resistance) of the hand sanitizers to each test organism were examined with the help of a ruler in millimeter by evaluating the average of 2 readings that were found from duplicates of agar diffusion test for each hand sanitizers.

### Minimum Inhibitory Concentration (MIC) Determination

In this study MIC (the lowest concentration of a definite antimicrobial or hand sanitizers required to inhibit the growth of a known test organism *in vitro*<sup>16</sup>) was done on nutrient broth for each hand sanitizer against the selected test organisms. It was done by preparing various concentrations of each hand sanitizers (10%, 20%, 30%, 40%, 50%, 60%, 70%, and 80%). Then, one milliliter from the different concentrations (10%, 20%, 30%, 40%, 50%, 60%, 70%, and 80%) of each sanitizer was introduced into the tube containing equal volumes (1 mL) of nutrient broth inoculated with standardized test organism. A tube, containing only nutrient broth and bacteria without sanitizer and a tube containing just the sanitizer and broth without bacteria was used as negative control and positive control, respectively. Finally, the tubes were incubated for 18–24 hours and visible growth (turbidity) was assessed. When compared with the controls, the concentration of the sanitizers at which no visible growth was regarded as MIC.

### Minimum Bactericidal Concentration (MBC) Determination

During this study MBC (the lowest concentration of a specific antimicrobial substance that can kill 99.9% of a given bacterial strain<sup>16</sup>) was determined from the MIC tests that showed no visible growth by taking a loopful of inoculum living test organisms from the MIC tubes by streaked on fresh Mueller Hinton agar. The streaked

**Table 2** The Susceptibility Pattern of the Selected Test Organisms to the Hand Sanitizers in the Agar Diffusion Test

Test Organism	Zones of Inhibition (mm) of Hand Sanitizers and Antimicrobial Disk Against Test Organisms									
	MTU	Folium	Fine	Epharm	Harego	Taflen	Sheba	D.Wat	Van	Cip
<i>Staphylococcus aureus</i>	15	9	15	14	14	17	16	6	12	–
<i>Escherichia coli</i>	18	8	20	17	15	18	15	6	–	30
<i>Klebsiella pneumoniae</i>	20	6	20	15	13	15	15	6	–	28
<i>Pseudomonas aeruginosa</i>	16	9	13	11	12	16	27	6	–	25
<i>Salmonella typhi</i>	12	8	13	14	15	13	12	6	–	20
<i>Shigella boydii</i>	15	8	15	17	16	16	15	6	–	22

**Abbreviations:** D.wat, sterilized distilled water; Van, Vancomycin, Cip, Ciprofloxacin.

Mueller Hinton agar plates were incubated at 37°C for 24 hours and were observed for growth. Streaked Mueller Hinton agar plates that cannot show any growth indicates a 99.9% bactericidal effect of the sanitizer at that concentration or MBC.

## Data Analysis

Data analysis was done by descriptive methods.

## Quality Control

The reliability of the study findings was guaranteed by implementing strict quality control (QC) methods throughout the whole process of the laboratory work, such as pre-analytical (specimen collection from hospital laboratory room, transportation from hospital laboratory to testing laboratory, etc.), analytical (reagent preparation, sample inoculation, gram stain, bacterial isolation, biochemical test, MIC, MBC agar diffusion test, etc.) and post-analytical (reporting MIC, MBC result, etc.) quality control. During MBC and MIC only broth as a negative control to test the sterility of the media and equipment, and culture inoculated broth without antibiotics as a positive control to test the growth ability of the medium was also used.

All materials and equipment are controlled and Standard Operating Procedures (SOPs) were strictly followed.

## Results

### Hand Sanitizers' Agar Diffusion Test

In our study, efforts were made to assess the efficiency of seven hand sanitizers (MTU, Folium Fine, Epharm, Harego, Taflen and Sheba) by using agar disc inhibitory activity against the selected test organisms (*Staphylococcus aureus*, *Escherichia coli*, *klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Salmonella typhi* and *Shigella boydii*). The zones of inhibition of the hand sanitizers at their claim

concentration including the unknown (unlabeled) concentration Taflen products were ranged from no inhibition zone (Folium) to 27mm (Sheba). For instance, 9mm (Folium) to 17mm (MTU) on *Staphylococcus aureus*; 8mm (Folium) to 20mm (Fine) on *Escherichia coli*; 6mm (Folium) to 20mm (Fine) on *K. pneumoniae*; 6mm (Folium) to 27mm (Sheba) on *Ps. aeruginosa*; 8mm (Folium) to 15mm (Hargo) on *Salmonella typhi*; whereas on *Shigella boydii* it ranged from 8mm (Folium) to 17mm (Ephram) (Table 2).

### Minimum Inhibitory Concentration (MIC)

All hand sanitizers were further tested to determine their MIC and MBC values at 10%, 20%, 30%, 40%, 50%, 60%, 70% and 80% (at claim concentration). The minimum inhibitory concentration against all selected test organisms was observed at 45%, 55%, and 65% except Taflen which have had unknown concentration but it was bacteriostatic at undiluted form (Table 3).

### Minimum Bactericidal Concentration (MBC)

The content of the concentrations tubes of MIC was further plated out on sterile Mueller Hinton agar plates in order to determine the bactericidal effect of concentration. MTU (except against *E. coli*), Fine, Hargo, Ephra and Sheba hand sanitizers were showed growth on 10%, 20%, 30%, 40%, 50%, and 60% concentrations against the selected test organisms. The growth of selected test organisms was decreased in colony-forming unit across increasing the concentration gradient of different hand sanitizers (Table 4). But a Folium product was indicating that the product was only bacteriostatic against the selected test organisms at 65% but not bactericidal. Similarly, MTU product had a MIC value of 65% concentration against *E. coli* but was not bactericidal (Table 5).

**Table 3** Minimum Inhibitory Concentration (MIC) of Selected Test Organisms to MTU, Folium, Fine, Epharm, Harego, Taflen and Sheba Hand Sanitizers

Hand Sanitizer	MIC (%) of Tested Sanitizers Against Tested Organisms					
	<i>Staphylococcus aureus</i>	<i>Escherichia coli</i>	<i>Klebsiella pneumoniae</i>	<i>Pseudomonas aeruginosa</i>	<i>Salmonella typhi</i>	<i>Shigella boydii</i>
MTU	55	65	55	55	55	55
Folium	65	65	65	65	65	65
Fine	55	55	55	55	65	65
Epharm	55	55	55	55	55	55
Harego	55	55	65	55	55	55
Sheba	55	55	45	45	55	45
Taflen	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown

**Notes:** MIC = minimum inhibitory concentration; Unknown = the manufacturer claim concentration was unknown (unlabeled) but in undiluted form, it inhibits the growth.

**Table 4** Counting Colony Forming Unit of Selected Test Organisms at Increasing Concentration of Different Hand Sanitizers

		MTU	Folium	Fine	Epharm	Harego	Sheba
<i>Staphylococcus aureus</i>	<= 40%	TMCC	TMCC	TMCC	TMCC	TMCC	TMCC
	50%	46	150	123	125	63	136
	60%	15	78	No	No	5	No
	>=70%	No	No	No	No	No	No
<i>Escherichia coli</i>	<=40%	TMCC	TMCC	TMCC	TMCC	TMCC	TMCC
	50%	TMCC	TMCC	TMCC	TMCC	71	86
	60%	TMCC	TMCC	TMCC	124	No	7
	70%	152	125	187	40	No	No
	80%	5	NA	No	No	No	No
<i>Klebsiella pneumoniae</i>	<=30%	TMCC	TMCC	TMCC	TMCC	TMCC	TMCC
	40%	141	TMCC	TMCC	TMCC	TMCC	TMCC
	50%	32	TMCC	231	TMCC	46	52
	60%	No	TMCC	53	TMCC	1	3
	70%	No	187	2	No	No	No
	80%	No	NA	No	No	No	No
<i>Pseudomonas aeruginosa</i>	<=40%	TMCC	TMCC	TMCC	TMCC	TMCC	TMCC
	50%	TMCC	TMCC	141	TMCC	21	33
	60%	313	TMCC	No	101	No	No
	70%	No	59	No	No	No	No
	80%	No	NA	No	No	No	No
<i>Salmonella typhi</i>	<=30%	TMCC	TMCC	TMCC	TMCC	TMCC	TMCC
	40%	TMCC	TMCC	TMCC	TMCC	TMCC	TMCC
	50%	93	TMCC	TMCC	TMCC	TMCC	92
	60%	2	113	TMCC	61	54	15
	70%	No	15	120	No	No	No
	80%	No	NA	6	No	No	No
<i>Shigella boydii</i>	<=40%	TMCC	TMCC	TMCC	TMCC	TMCC	TMCC
	50%	72	TMCC	TMCC	TMCC	70	69
	60%	11	TMCC	TMCC	59	4	11
	70%	No	41	No	No	2	No
	80%	No	NA	No	No	No	No

**Abbreviations:** TMCC, to many colony count; NA, not applicable; No, no growth.



**Table 5** Minimum Bactericidal Concentration (MBC) of Selected Test Organisms to MTU, Folium, Fine, Epharm, Harego, Taflen and Sheba Hand Sanitizers

Hand Sanitizer	Minimum Bactericidal Concentration (MBC) of Tested Sanitizers Against Tested Organisms					
	<i>Staphylococcus aureus</i>	<i>Escherichia coli</i>	<i>Klebsiella pneumoniae</i>	<i>Pseudomonas aeruginosa</i>	<i>Salmonella typhi</i>	<i>Shigella boydii</i>
MTU	65	No	65	65	65	No
Folium	No	No	No	No	No	No
Fine	65	65	75	65	75	65
Epharm	65	75	65	65	65	65
Harego	65	65	65	65	75	75
Sheba	65	65	65	75	75	75
Taflen (stock)	Stock conc.	Stock conc.	Stock conc.	Stock conc.	Stock conc.	Stock conc.

**Notes:** No = there is no minimum bactericidal concentration (the product were not had MBC at any concentration); stock conc.= the manufacturer claim concentration in undiluted form.

## Discussion

Now the use of hand sanitizers in the community has mostly gained recognition in the world, including Ethiopia since the emergence of COVID-19. This has led to the development, production and importation of several hand sanitizers by various companies with the intend of profit-making as well as supporting the health-care system and community in preventing transmission of disease especially COVID-19. But there is no information about laboratory-based efficacy evaluation.

In this study, the efficiency of the claim concentration of the seven hand sanitizers was assessed. The seven hand sanitizers MTU, Folium, Fine, Ephra, Hargo, and Sheba with claimed concentration of 80% ethanol alcohol, and Taflen with unknown (unlabeled) concentration of ethanol alcohol as the main active ingredient were displayed bacteriostatic activity against all the selected test organisms at a concentration of 45%, 55%, and 65%. Moreover, all Fine, Harego, Epharm and Sheba hand sanitizers showed bactericidal activity against all selected test organisms, with MBC values of 65% and 75%. Generally, the growth of selected test organisms was decreased in colony-forming unit across increasing the concentration gradient of different hand sanitizers. This bacteriostatic/bactericidal activity more probably due to alcohol components of alcohol-based hand sanitizer which is the major active ingredients intended to exert disinfectant activity in bacteria by causing protein denaturation, disruption of tissue membranes and dissolution of several lipids.<sup>17</sup>

Folium which contained 70% denatured ethanol was not active against all selected test organisms in bactericidal activity with no to narrow rim of inhibition zone in disk diffusion agar. The lack of bactericidal activity and more or less no disk inhibition zone observed in this product

could be due to the relatively decrement of the concentration (70%) denatured alcohol compared with others since the efficacy of alcohol-based hand sanitizers is affected by several factors such as the type of alcohol used, concentration of alcohol or amount of alcohol used, the possible contact time<sup>18</sup> and absence of active ingredient in product (hydrogen peroxide) which may limit the cidal effect of the alcohol from attainment the bacterial cells.

MTU product was the other hand sanitizer product which was not active against *E. coli* test organisms in our study. This could be poor or prolonged storage of the products which could lead to increased temperature causing evaporation of the active ingredient and characteristic of the *E. coli* strain since the strain is bacteria, which can easily develop resistance.

## Limitation

There are limitations in our study; we could assess the efficacy of only seven locally made but not imported sanitizers. Only bacterial strain was used because of laboratory setup constraint for virus strain growth.

## Conclusion

We evaluated the antibacterial efficacy of the seven most popular brands of hand sanitizers sold in Mizan-Aman town. Of the seven the six sanitizers except MTU product for *E. coli* and Folium were the products that were effective in inhibited growth of all the selected test organisms in vitro, and they were had effective bactericidal activity on their claim concentration. But MTU products against *E. coli* and Folium against all selected test organisms were not had effective bactericidal activity.

## Recommendation

Even though alcohol-based hand sanitizers are easy to prepare, they must be used after proper testing of their efficacy at their producer claimed concentration to protect consumers from buying poor quality products. The company should have to follow strict standard operational procedures to produce standardized products. The regulatory authorities should enforce strict quality control measures during production, distribution and routine laboratory-based inspections to ensure the efficacy of products. Lastly, consumers should be aware of the availability of poor quality sanitizers on the shelves of some retail outlets.

## Abbreviations

CDC, the Center for Disease Control and Prevention; WHO, World Health Organization; MTU, Mizan-Tepi University; MIC, minimum inhibitory concentration; MBC, minimum bactericidal concentration.

## Data Sharing Statement

The data used to support the findings of this study are available from the corresponding author upon request.

## Ethics Approval and Consent to Participate

We conduct our study in compliance with recognized international standards and the principles of the Declaration of Helsinki. Ethical clearance was acquired from the Research and Ethical Review Committee of College of Medicine and Health Sciences, at Mizan Tepi University. The permission letter was taken from the clinical director of the Mizan-Tepi University referral hospital and laboratory head. Data were collected after full written consent had been obtained from each participant. For the purpose of the study, codes were used instead of any personal identifiers which were labeled by laboratory technologist on the specimen tubes during specimen collection. Specimens with abnormal test results were informed to physicians for further diagnosis and treatment accordingly.

## Acknowledgments

We would like to thank the College of Medicine and Health Science and Mizan-Tepi University for giving us this chance to conduct this research. Moreover, we would like to express our appreciation to Mizan-Tepi Teaching Hospital for willingness to give leftover patient specimens

and the staff of the outpatient department (OPD) unit for their support during data collection.

## Author Contributions

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed to submit to the current journal; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

## Funding

Mizan-Tepi University.

## Disclosure

The authors declare that they do not have any conflict of interests.

## References

1. Hassan AO, Hassan RO, Muhibi MA, Adebimpe WO. A survey of Enterobacteriaceae in hospital and community acquired infections among adults in a tertiary health institution in Southwestern Nigeria. *Afr J Microbiol Res*. 2012;6(24):5162–5167.
2. Jain A, Singh K. Recent advances in the management of nosocomial infections. *JK Sci*. 2007;9(1):3–8.
3. World Health Organisation. WHO guidelines in hand hygiene in health care. WHO/IER/PSP/2009/01. Geneva, Switzerland: World Health Organisation; 2009
4. Zerr DM, Garrison MM, Allpress AL, Heath J, Christakis DA. Infection control policies and hospital-associated infections among surgical patients: variability and associations in a multicenter pediatric setting. *Pediatrics*. 2005;115(4):387–392. doi:10.1542/peds.2004-2014
5. World Health Organization. 2020. Coronavirus disease 2019 (COVID-19): situation report 88. World Health Organization. Available from: <https://apps.who.int/iris/handle/10665/331851>.
6. Boyce JM, Pittet D. Guideline for hand hygiene in health-care settings: recommendations of the Health care Infection Control Practices Advisory Committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force. *Am J Infect Control*. 2002;30(8):S1–46.
7. Verma DK, Tesfu K, Getachew M, Workneh Y, Mekuriaw F, Tilahun M. Evaluation of antibacterial efficacy of different hand gel sanitizers in University of Gondar Students, North-West Ethiopia. *J Glob Biosci*. 2013;2(6):166–173.
8. Zaragoza M, Sallés M, Gomez J, Bayas JM, Trilla A. Hand washing with soap or alcoholic solutions? A randomized clinical trial of its effectiveness. *Am J Infect Control*. 1999;27(3):258–261. doi:10.1053/ic.1999.v27.a97622
9. Kramer A, Rudolph P, Kampf G, Pittet D. Limited efficacy of alcohol-based hand gels. *Lancet*. 2002;359(9316):1489–1490. doi:10.1016/S0140-6736(02)08426-X
10. Wolfe MK, Gallandat K, Daniels K, Desmarais AM, Scheinman P, Lantagne D. Handwashing and Ebola virus disease outbreaks: a randomized comparison of soap, hand sanitizer, and 0.05% chlorine solutions on the inactivation and removal of model organisms Phi6 and E. coli from hands and persistence in rinse water. *PLoS One*. 2017;12(2):e0172734. doi:10.1371/journal.pone.0172734

11. Kramer A, Rudolph P, Kampf G, Pittet D. Showing classic G to A mutation at nucleotide 1896. *Lancet*. 2002;359:1489–1490.
12. Rotter M. Hand washing and disinfection. In: Mayhall CG, editor. *Hospital Epidemiology and Infection Control*. 2nd ed. Baltimore: Williams and Wilkins; 1999:1339–1355.
13. Cheesbrough M. *District Laboratory Practice in Tropical Countries, Part 2*. Cambridge university press; March 2, 2006.
14. Vandepitte J, Engbaek K, Rohner P, Piot P, Heuck CC; World Health Organization. Basic laboratory procedures in clinical bacteriology/ J. Vandepitte ... [et al.]. In: *Basic Laboratory Procedures in Clinical Bacteriology/ J. Vandepitte ... [et al.]*; 2003
15. Valgas C, Souza SM, Smânia EF, Smânia JA. Screening methods to determine antibacterial activity of natural products. *Braz J Microbiol*. 2007;38(2):369–380. doi:10.1590/S1517-83822007000200034
16. Nester EW, Anderson DG, Roberts CE, Nester MT. *Microbiology: A Human Perspective, 6th Edition*. 2009.
17. Kar A. *Pharmaceutical Microbiology*. Delhi: New Age International (P) Ltd Publishers; 2008:216–217.
18. Harmanci H. Guideline for Hand Hygiene in Health-Care Setting Recommendations of the Healthcare Infection Control Practices Advisory Committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force. *Marmara Med J*. 2016;15(4):281–286.

## Infection and Drug Resistance

Dovepress

### Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>