### ORIGINAL RESEARCH

# The Relationship Between the Serum Anion Gap and All-Cause Mortality in Acute Pancreatitis: An Analysis of the MIMIC-III Database

This article was published in the following Dove Press journal: International Journal of General Medicine

Fang Gong<sup>1</sup> Quan Zhou<sup>2</sup> Chunmei Gui<sup>1</sup> Shaohua Huang<sup>1</sup> Zuoan Qin<sup>3</sup>

<sup>1</sup>Department of Intensive Care Unit, The First People's Hospital of Changde, Changde City, Hunan Province, 415000, People's Republic of China; <sup>2</sup>Department of Science and Education Section, The First People's Hospital of Changde, Changde City, Hunan Province, 415000, People's Republic of China; <sup>3</sup>Cardiovascular Medicine Department, The First People's Hospital of Changde, Changde City, Hunan Province, 415000, People's Republic of China

submit your manuscript | www.dovepress.com DovePress **Purpose:** We aimed to investigate the relationship between the serum anion gap (AG) and all-cause mortality in patients with acute pancreatitis (AP) in intensive care units (ICUs).

**Patients and Methods:** In this retrospective cohort analysis, data of patients with AP were extracted from the Medical Information Mart for Intensive Care database (version III). We collected the maximum serum AG value within the first 24 hours of ICU admission. The main outcome was 90-day all-cause mortality. A multivariate Cox proportional hazard regression model was used to examine the association between the serum AG and mortality. The restricted cubic spline curve was used to confirm a non-linear relationship between serum AG values and mortality.

**Results:** Of the 279 patients included in the study, 87 (31.18%) died. The serum AG value was positively associated with 90-day all-cause mortality (hazard ratio [HR] 1.08, 95% confidence interval [CI] 1.02–1.14), after adjusting for age, sex, alcohol consumption, congestive heart failure, diabetes mellitus, hypertension, eGFR, albumin, and the SOFA score. There was a non-linear relationship between serum AG values and mortality after adjusting for potential confounders. We used a two-piecewise regression model to obtain a threshold inflection point value of 13.8 mmol/L. The HR and the 95% CI on the left inflection point were 0. 82 (0.61–1.09; p = 0.1719), and on the right inflection point were 1.15 (1.08–1.23; p < 0.0001).

**Conclusion:** The relationship between all-cause mortality in patients with acute pancreatitis and serum AG values was non-linear. All-cause mortality and serum AG values were positively correlated when the serum AG value was >13.8 mmol/L.

Keywords: anion gap, all-cause mortality, acute pancreatitis, MIMIC III database

## Introduction

Acute pancreatitis (AP) is an acute inflammatory disease that can result in multiple system dysfunction.<sup>1</sup> Studies have reported mortality rates for mild AP to range from <1% to 5%.<sup>2,3</sup> However, 20% of patients with AP may develop severe pancreatitis, which has a reported mortality rate of 30% due to complications such as pancreatic necrosis and organ failure.<sup>2</sup> An acid-base balance is the basis for maintaining cell metabolism and physiological function.<sup>4</sup> The serum anion gap (AG) is a commonly used indicator to evaluate acid-base balance, which helps to identify the types of acidosis and the causes of acid-base imbalance.<sup>5</sup> At present, the serum AG value is usually calculated based on the concentration of sodium, chlorine, potassium, and bicarbonate in serum.<sup>6</sup> Albumin carries most of the

53 I

© 2021 Gong et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress.com/terms.php and incorporate the Creative Commons Attribution – Non Commercial (unported, v3.0). License (http://creativecommons.org/licenses/by-md/3.0). By accessing the work you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (https://www.dovepress.com/terms.php).

Correspondence: Fang Gong Department of Intensive Care Unit, The First People's Hospital of Changde, 818 Renmin Road, Changde City, Hunan Province, 415000, People's Republic of China Tel +86-0736-7788035 Email gficu2009@163.com

negative charge in the body, so hypoproteinemia can lead to a reduction in the serum AG.<sup>7,8</sup> The serum AG is commonly used in clinical practice as an important biochemical indicator for disease diagnosis or prognosis.<sup>9,10</sup> The initial serum AG in adult patients in the intensive care unit (ICU) has been reported to be a sensitive and specific tool for predicting prognosis or death.<sup>11,12</sup> Moreover, it has been used to predict mortality in children in the ICU.<sup>13</sup> However, the relationship between serum AG and allcause mortality in patients with AP has not been studied. In this study, we aimed to investigate the relationship between serum AG values and 90-day all-cause mortality in patients with AP in the ICU.

# **Patients and Methods**

## Data Source

This study was a retrospective cohort analysis. Data concerning patients with AP were retrieved from the Medical Information Mart for Intensive Care version III (MIMIC-III), which is a large, single-center, public database of critical care. This database includes >40,000 ICU inpatients who had been admitted to Beth Israel Deaconess Medical Center in Boston from 2001 to 2012.<sup>14</sup> To access the database, we applied the Protecting Human Research Participants test (no. 36,208,651). The International Classification of Diseases, Ninth Revision (ICD-9) code for the diagnosis of AP was 5770. Diagnostic criteria for AP are based primarily on conformance with the 2012 Atlanta Consensus.<sup>15</sup> Patients with recurrent AP were evaluated only at the first admission. We extracted clinical indicators such as demographic characteristics, ICD-9 codes, physiological indicators, and laboratory tests. Data were obtained through the Structured Query Language (SQL) performed in the MIMIC-III database. This study was approved by the Institutional Review Boards of Beth Israel Deaconess Medical Center (Boston, MA) and the Massachusetts Institute of Technology (Cambridge, MA), and the requirement for patient informed consent was waived due to the study design.

Baseline characteristics, collected within 24 hours of admission, included sex, age, medical history, the sequential organ failure assessment (SOFA) score, and the systemic inflammatory response system (SIRS) score for patients with AP who had been admitted to the ICU. Other clinical indicators included mean arterial pressure (BP), heart rate, body temperature, white blood cell count (WBC), and blood urea nitrogen (BUN), creatinine, pulse oxygen saturation (SpO<sub>2</sub>), and maximum values of serum AG within 24 hours of ICU admission. When the above indicators had multiple results within 24 hours, we took only the worst value. The study endpoint was all-cause mortality within 90 days from the date of ICU admission.

# Study Population

Data concerning 904 patients (aged  $\geq$  18 years) with AP according to the ICD-9 code 5770 were retrieved. Only patients at the first admission and with a first diagnosis of AP were included. Patients with an ICU length of stay <24 hours or >90 days; those with immune dysfunctions such as tumors, metastatic tumors, or acquired immune deficiency syndrome (AIDS); and those with incomplete (>10% of values missing), incorrect, or uninterpretable data were excluded.

# Statistical Analysis

In this study, data were expressed as mean (SD) or median (min, max; skewed distribution) for continuous variables and percentages (%) for categorical variables. A one-way analysis of variance (ANOVA) test (normal distribution) or Kruskal-Wallis H-test (skewed distribution) and chisquared test or Fisher's exact test (categorical variables) were used to detect the differences among different serum AG tertiles. We constructed three distinct models using Cox proportional hazards regression models to investigate the association between the serum AG value and mortality: the crude model, in which no covariates were adjusted; model 1, in which only age and sex were adjusted; and model 2, in which model-1 and other potential confounders were adjusted. We selected background variables based on existing literature and clinical judgment. We also adjusted for features such that if the influence of a confounder changed by >10%, it was incorporated into the adjusted model.<sup>16</sup> To test the robustness of our results, we performed sensitivity analysis. We converted the serum AG value into a categorical by tertiles, to verify the results of serum AG as a continuous variable and to study the possibility of nonlinearity. Because Cox proportional hazard regression model-based methods may not always be capable of dealing with non-linear models, non-linearity between the serum AG value and mortality was addressed using a Cox proportional hazard regression model with restricted cubic spline and smooth curve fitting. If nonlinearity was detected, we first calculated the threshold inflection point using a recursive algorithm. We then constructed a twopiecewise regression model on both sides of the inflection

point. Statistical software packages R (<u>http://www.</u> <u>R-project.org</u>, The R Foundation, Vienna, Austria) and Empower Stats (<u>http://www.empowerstats.com</u>, X&Y Solutions, Inc, Boston, MA, USA) were used for all statistical analyses in this study. A p-value <0.05 was considered statistically significant.

# Patient and Public Involvement

No patients actively participated in this study; however, data concerning patients with AP were retrieved from the Medical Information Mart for Intensive Care version III (MIMIC-III), which is a large, single-center, public database of critical care.

# Results

### Participant Selection

Data concerning 904 patients (aged  $\geq$  18 years) with AP were retrieved according to the ICD-9 code. We excluded patients whose first diagnosis was not AP (n = 556) and those who lacked ICU admission information (n = 3). Patients who had been admitted to the ICU for >90 days (n = 3) or <24 hours (n = 7) were excluded. Patients with solid tumors (n = 8), metastatic tumors (n = 3), or AIDS (n = 6) were excluded. We also excluded patients with missing AG values (n = 2) and those with AG values >40 mmol/L (n = 1). In total, 279 patients were included in the study (Figure 1).

# Population and Baseline Characteristics

Participants were stratified into tertiles (T1, T2, and T3) according the maximum serum AG value within the first 24 hours of ICU admission (Table 1). Of 279 eligible patients with AP whose data were extracted from the MIMIC-III database (women, n = 125; men, n = 154), 87 (31.18%) participants died. Patients with high serum AG values (>=17.0 mmol/L) were more likely to have accompanying renal failure. The SOFA scores along with WBC, creatinine, BUN, glucose, and lactic acid values were significantly higher in those with high serum AG values than in those with low serum AG values.



Figure I Flowchart of subject screening.

Abbreviations: AP, acute pancreatitis; MIMIC-III, Medical Information Mart for Intensive Care version III; ICU, intensive care unit.

### Table I Participants' Baseline Characteristics

Characteristics	AG Tertiles (mmol/L)			
	T1: <13.00 (n = 65)	T2: 14.00 -16.00 (n = 102)	T3: ≥ 17.00 (n = 112)	
Demographics				
Age, years, (SD)	58.66 (16.76)	61.77 (17.01)	58.88 (15.62)	0.345
Sex, n (%)				0.060
Male	30 (46.15)	53 (51.96)	71 (63.39)	
Female	35 (53.85)	49 (48.04)	41 (36.61)	
Clinical features, n (%)	•	•		
Congestive heart failure				0.464
No	54 (83.08)	77 (75.49)	90 (80.36)	
Yes	( 6.92)	25 (24.51)	22 (19.64)	
Hypertension				0.128
No	36 (55.38)	42 (41,18)	46 (41.07)	
Yes	29 (44.62)	60 (58.82)	66 (58.93)	
				0.070
COPD	E2 (80.00)	00 (0( )7)		0.069
140	52 (00.00) 13 (20.00)	00 (00.27)	9 (9 04)	
103	13 (20.00)		7 (0.04)	
Diabetes mellitus				0.084
No	52 (80.00)	/3 (/1.5/)	/2 (64.29)	
Yes	13 (20.00)	29 (28.43)	40 (35.71)	
Renal failure				<0.001
No	64 (98.46)	95 (93.14)	91 (81.25)	
Yes	l (l.54)	7 (6.86)	21 (18.75)	
Liver disease				0.446
No	53 (81.54)	87 (85.29)	88 (78.57)	
Yes	12 (18.46)	15 (14.71)	24 (21.43)	
Obesity				0.184
No	57 (87.69)	96 (94.12)	106 (94.64)	
Yes	8 (12.31)	6 (5.88)	6 (5.36)	
Alcohol consumption				0.109
No	50 (76.92)	75 (73.53)	71 (63.39)	
Yes	15 (23.08)	27 (26.47)	41 (36.61)	
Biomarkers mean (SD)/median (min-max)				
Albumin, g/L	2.65 (0.56)	2.85 (0.58)	3.05 (0.69)	0.002
Lactic acid, mmol/L	1.53 (0.76)	1.94 (1.31)	2.78 (1.82)	<0.001
Glucose, mg/dL	158.48 (59.54)	166.45 (73.65)	201.91 (102.56)	<0.001
WBC, 10 <sup>9</sup> /L	13.89 (5.58)	16.77 (7.21)	15.28 (7.02)	0.028
SpO <sub>2</sub> , %	91.64 (4.39)	91.16 (3.51)	92.26 (3.04)	0.084
Mean BP, mmHg	64.20 (14.33)	68.60 (16.72)	66.67 (15.04)	0.209
BUN, mmol/L	15.00 (3.00-64.00)	19.00 (5.00-91.00)	30.00 (3.00-128.00)	<0.001
Creatinine, mg/dL	0.80 (0.20-2.10)	0.90 (0.40-3.40)	1.60 (0.50-7.00)	<0.001
eGFR, mL/min/1.73 m <sup>2</sup>	67.19 (3.04–199.12)	70.55 (6.59–200.64)	58.17 (5.44–227.57)	0.954
SIRS score	3.23 (0.75)	3.14 (0.91)	3.16 (0.81)	0.774

(Continued)

#### Table I (Continued).

Characteristics	AG Tertiles (mmol/L)			
	T1: <13.00 (n = 65)	T2: 14.00 -16.00 (n = 102)	T3: ≥ 17.00 (n = 112)	
SOFA score	3.00 (0.00-11.00)	3.00 (0.00-11.00)	4.00 (0.00-18.00)	<0.001
Death, n (%) No Yes	47 (72.31) 18 (27.69)	77 (75.49) 25 (24.51)	68 (60.71) 44 (39.29)	0.052

Abbreviations: SD, standard deviation; BP, blood pressure (arterial); BUN, blood urea nitrogen; COPD, chronic obstructive pulmonary disease; eGFR, estimated glomerular filtration rate; max, maximum; min, minimum; SIRS, systemic inflammatory response system; SOFA, sequential organ failure assessment; SpO<sub>2</sub>, pulse oxygen saturation; WBC, white blood cell count.

# Univariate Analysis

Table 2 shows the results of univariate analysis that indicated congestive heart failure (p = 0.0443), renal failure (p = 0.0019), BUN (p < 0.0001), serum AG (p = 0.0017), and creatinine (p < 0.0001) values, age (p < 0.0001), and SOFA scores (p < 0.0001) positively correlated with mortality. Moreover, mean arterial blood pressure (p < 0.0001) negatively correlated with mortality (Table 2).

# Relationship Between Serum AG Values and All-Cause Mortality Using Different Models

A multivariate Cox proportional hazards regression model was used to evaluate the association between the serum AG value and 90-day all-cause mortality. Crude, minimally adjusted, and fully adjusted models are shown in Table 3. In the crude model, the serum AG was found to be positively correlated with mortality (hazard ratio [HR] 1.07, 95% confidence interval [CI] 1.02–1.11, p = 0.0017). In model 1, which was adjusted for age and sex, the serum AG also showed a positive correlation with mortality (HR 1.08, 95% CI 1.03–1.12, p = 0.0004). In model 2, which was adjusted for age, sex, alcohol consumption, congestive heart failure, diabetes mellitus, hypertension, eGFR, albumin, and the SOFA score, no obvious changes were observed (HR 1.08, 95% CI 1.02-1.14, p = 0.0091). Next, tests for trend were conducted with multivariate proportional hazards regression models by entering the median value of each serum AG tertile as a continuous variable in the models. Patients with a lower serum AG value (<13 mmol/L) were included in the reference group. In model 1, high serum tertile (AG  $\geq$ 17 mmol/L) was associated with an increased risk of all-cause mortality (HR 1.95, 95% CI 1.11-3.42) after adjusting for age and sex. In model 2, after adjusting for age, sex, alcohol consumption, congestive heart failure, diabetes mellitus, hypertension, eGFR, albumin, and the SOFA score, high serum AG value ( $\geq$ 17 mmol/L) remained positively associated with 90-day all-cause mortality, compared with a low serum AG tertile (HR 1.88, 95% CI 1.04–3.41); the p-for-trend was 0.0182.

# Analyses of Non-Linear Relationships

We found a non-linear relationship between the maximum serum AG within 24 hours of ICU admission and all-cause mortality after adjusting for age, sex, alcohol consumption, congestive heart failure, diabetes mellitus, hypertension, eGFR, albumin, and the SOFA score (Figure 2). We calculated that the inflection point was 13. 8 mmol/L using a two-piecewise regression model (Table 4). There was no significant correlation to the left of the inflection point (HR [95% CI] 0.82 ([0.61–1.09]), p = 0.1719; however, there was a positive correlation between the serum AG value and all-cause mortality to the right of the threshold inflection point (HR 1.15, 95% CI 1.08–1.23, p < 0.0001).

# Discussion

In this study, we investigated the association between serum AG value and all-cause mortality in critically ill patients with AP. First, the serum AG value was associated with mortality in patients with AP after adjusting for age, sex, alcohol abuse, congestive heart failure, diabetes mellitus, hypertension, eGFR, albumin, and the SOFA score. Second, there was a non-linear relationship between serum AG values and death from all causes in patients with AP, and we calculated a threshold inflection point of 13.8mmol/L, using a two-segment linear regression model. To the right of the threshold, the number of all-

Table 2 Factors Correlated	d to	Mortality	in	Acute	Pancreatitis
Using Univariate Analysis					

Variables	Statistics	OR (95% CI)	p-value
Demographics			
Age, years	59.62 ± 16.54	1.04 (1.02, 1.06)	<0.0001
Sex n (%) Male Female	154 (55.20) 125 (44.80)	1.0 0.98 (0.64–1.50)	0.9345
Clinical features, n (%)			
Congestive heart failure No Yes	221 (79.2) 58 (20.8)	1.0 1.61 (1.01–2.56)	0.0443
Hypertension No Yes	124 (44.44) 155 (55.56)	1.0 1.12 (0.73–1.71)	0.6177
COPD No Yes	243 (87.1) 36 (12.9)	1.0 1.51 (0.86–2.64)	0.1470
Diabetes mellitus No Yes	197 (70.61) 82 (29.39)	1.0 0.83 (0.50–1.39)	0.4820
Renal failure No Yes	250 (81.72) 29 (18.28)	1.0 2.59 (1.42–4.74)	0.0019
Liver disease No Yes	228 (92.83) 51 (7.17)	1.0 1.17 (0.68–2.19)	0.6179
Obesity No Yes	259 (92.83) 20 (7.17)	1.0 0.46 (0.11–1.86)	0.2761
Alcohol consumption No Yes	196 (70.25) 83 (29.75)	1.0 0.58 (0.33–1.04)	0.0590
Biomarkers mean (SD)/median (min- max) Albumin, g/L	2.90 ± 0.64	0.84 (0.58–1.20)	0.3322
Lactic acid, mmol/L WBC, 10 <sup>9</sup> /L SpO <sub>2</sub> vital, % BUN, mmol/L	$2.34 \pm 2.14$ $15.50 \pm 6.84$ $91.73 \pm 3.59$ $21$ $(12.75-32.25)$	1.04 (0.93–1.16) 1.03 (1.00–1.06) 0.97 (0.92–1.03) 1.02 (1.01–1.03)	0.5003 0.0915 0.3263 <0.0001

(Continued)

Variables	Statistics	OR (95% CI)	p-value
Serum AG, mmol/L	16.39 ± 4.47	1.07 (1.02–1.11)	0.0017
eGFR, mL/min/ 1.73 m <sup>2</sup>	71.33 ± 44.47	1.00 (0.99–1.00)	0.2687
Creatinine, mg/	l (0.7–1.7)	1.35 (1.18–1.56)	<0.0001
dL			
Glucose, mg/dL	178.71 ± 85.48	1.00 (1.00–1.00)	0.4734
Mean BP, mmHg	66.84 ± 15.53	0.97 (0.95-0.98)	<0.0001
SIRS score SOFA score	3.15 ± 0.87 4 (2–6)	1.05 (0.83–1.33) 1.18 (1.12–1.25)	0.6810 <0.0001

Abbreviations: AG, anion gap; OR, odds ratio; CI, confidence interval; BP, blood pressure (arterial); BUN, blood urea nitrogen; WBC, white blood cell count; COPD, chronic obstructive pulmonary disease; eGFR, estimated glomerular filtration rate; min, minute; SD, standard deviation; SIRS, systemic inflammatory response system; SOFA, sequential organ failure assessment; SpO<sub>2</sub>, pulse oxygen saturation.

cause deaths increased with an increase in the serum AG value, which was positively correlated. However, the trend was the opposite to the left of the threshold, although it was not significant. This may have something to do with our insufficient sample size.

The serum AG has been reported to be associated with death and complications in a variety of diseases. Cheng et al,<sup>17</sup> found that high serum AG values in patients with acute kidney injury had higher all-cause mortality at 30 days, 90 days, and 365 days, HR (95% CI) were: 1.54 (1.33-1.75), 1.55, (1.38-1.73), and 1.46 (1.31-1.60) respectively. Abramowitz et al.<sup>18</sup> found that the serum AG value was associated with mortality, the Relative Hazard (95% CI) was 1.20 (1.03-1.41), and higher levels of serum AG were present in patients with impaired renal function. A retrospective analysis of 440 participants data suggested that adjusted serum anion gap is an independent risk factor for all -cause mortality in advanced CKD patients, with HR (95% CI) 2.968 (1.143-7.708).<sup>19</sup> The relationship between serum AG and mortality has been reported not only in adults but also in children. Kim and his team<sup>20</sup> reported that corrected AG in children at ICU admission was associated with mortality, with odds ratio (95% CI) 1.110 (1.06-1.17). Cheng et al,<sup>17</sup> and Tang et al,<sup>21</sup> confirmed a U-shaped relationship between serum AG levels and all-cause mortality in critically ill patients with acute kidney injury and patients with congestive heart failure, respectively. Our findings are consistent with previous studies. We found a non-linear relationship between the serum AG and mortality, after adjusting for

Variable	Crude Model HR (95% CI)	p-value	Model   HR (95% CI)	p-value	Model 2 HR (95% CI)	p-value
Serum AG	1.07 (1.02–1.11)	0.0017	1.08 (1.03–1.12)	0.0004	1.08 (1.02–1.14)	0.0091
Serum AG ter	tiles					
ті	I.0 (Ref)		I.0 (Ref)		I.0 (Ref)	
T2	0.91 (0.50–1.69)	0.7731	0.79 (0.42–1.48)	0.4542	0.72 (0.38-1.38)	0.2564
Т3	1.82 (1.05–3.17)	0.0332	1.95 (1.11–3.42)	0.0207	1.88 (1.04–3.41)	0.0370
p-for-trend		0.0131		0.0069		0.0182

Table 3 The Relationship Between the Serum AG and All-Cause Mortality in Patients with AP in Different Models

Notes: Crude model: No covariates were adjusted. Model 1: adjusted for age and sex. Model 2: adjusted for age, sex, alcohol abuse, congestive heart failure, diabetes mellitus, hypertension, eGFR, albumin, and the SOFA score.

Abbreviations: AG, anion gap; AP, acute pancreatitis; CI, confidence interval; eGFR, estimated glomerular filtration rate; HR, hazards ratio; SOFA, sequential organ failure assessment; Ref, reference.

serum albumin level, eGFR and other confounding factors, and we further identified a threshold inflection point of the impact of AG on all-cause mortality in AP. As reported, the main component of the serum AG is the sum of the anionic charges on circulating proteins,<sup>22</sup> changes in serum albumin concentration levels will cause changes in the serum AG. After adjusting for covariates including serum albumin, the serum AG was still associated with all-cause mortality in critical patients with AP. The serum AG value, which is a low-cost and easily available biomarker, may facilitate clinical assessment of the mortality risk for patients with AP.



**Figure 2** The relationship between serum AG values and all-cause mortality in patients with AP. There was a nonlinear relationship between serum AG values and all-cause mortality in patients with AP after adjusting for age, sex, alcohol abuse, congestive heart failure, diabetes, hypertension, eGFR, albumin, and SOFA score. Solid rad line indicates the cubic spline functions between variables. Imaginary lines represent the 95% of confidence interval from the fit. **Abbreviations:** AG, anion gap; AP, acute pancreatitis.

# **Study Strengths and Limitations**

Our study had the following strengths. First, both a Cox proportional hazard regression model and the restricted cubic spline curve were used to study the association between serum AG and 90-day all-cause mortality in patients with AP. Second, this was an observational study. Potential confounders were unavoidable; therefore, we combined existing literature, clinical judgment, and statistical adjustments to minimize the effect of confounders.

Our study also had some limitations. First, although we adjusted for certain factors, our results may have been influenced by other unknown factors. Second, only adults were included in our study; therefore, the relationship between serum AG levels and outcomes in children with AP could not be determined.

## Conclusions

The relationship between the serum AG and all-cause mortality in patients with AP was non-linear after adjustment for potential confounders. Large prospective studies are needed to further confirm our results.

**Table 4** Threshold Effect Analysis of the Relationship Betweenthe Serum AG and Death Using a Two-Piecewise RegressionModel

Serum AG Inflection Point	HR (95% CIs)	p-value
<13.8 ≥13.8	0.82 (0.61, 1.09) 1.15 (1.08, 1.23)	0.1719 <0.0001
P for log likelihood ratio test		0.046

Note: Adjusted for age, sex, alcohol abuse, congestive heart failure, diabetes mellitus, hypertension, eGFR, albumin, and the SOFA score.

Abbreviations: AG, anion gap; Cl, confidence interval; eGFR, estimated glomerular filtration rate; HR, hazard ratio; SOFA, sequential organ failure assessment.

# **Data Sharing Statement**

The clinical data of this study were derived from MIMIC-III (v. 1.4). This is an open and free database, and researchers must complete an online course at the National Institutes of Health, known as Protecting Human Research Participants, before they can apply for permission to access it.

# Acknowledgments

We wish to thank Dr. Jie Liu of the Department of Vascular and Endovascular Surgery, Chinese PLA General Hospital, for his contribution to the study design, and for his advice and comments regarding the manuscript. We would also like to thank Editage (www.editage.cn) for English language editing.

# Funding

This study was supported by the Scientific Research Foundation of Changde (grant number: 2017so317).

# Disclosure

The authors declare that there are no conflicts of interest regarding the publication of this paper.

# References

- 1. Agarwal N, Pitchumoni CS. Acute pancreatitis: a multisystem disease. *Gastroenterologist.* 1993;1(2):115–128.
- Wu BU, Johannes RS, Sun X, Tabak Y, Conwell DL, Banks PA. The early prediction of mortality in acute pancreatitis: a large population-based study. *GUT*. 2008;57(12):1698–1703. doi:10.1136/ gut.2008.152702
- Dervenis C, Johnson CD, Bassi C, et al. Diagnosis, objective assessment of severity, and management of acute pancreatitis. Santorini consensus conference. *Int J Pancreatol.* 1999;25(3):195–210. doi:10.1007/BF02925968
- Rosner MH, Perazella MA, Choi MJ. American Society of Nephrology quiz and questionnaire 2014: acid-base and electrolyte disorders. *Clin* J Am Soc Nephrol. 2015;10(3):530–539. doi:10.2215/CJN.10911114
- Kraut JA, Madias NE. Serum anion gap: its uses and limitations in clinical medicine. *Clin J Am Society Nephrol.* 2007;2(1):162–174. doi:10.2215/CJN.03020906
- Co I, Gunnerson K. Emergency department management of acute kidney injury, electrolyte abnormalities, and renal replacement therapy in the critically ill. *Emerg Med Clin North Am.* 2019;37(3):459–471. doi:10.1016/j.emc.2019.04.006

#### **International Journal of General Medicine**

Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies

- Figge J, Jabor A, Kazda A, Fencl V. Anion gap and hypoalbuminemia. *Crit Care Med.* 1998;26(11):1807–1810. doi:10.1097/00003246-199811000-00019
- Nanji AA, Campbell DJ, Pudek MR. Decreased anion gap associated with hypoalbuminemia and polyclonal gammopathy. *JAMA*. 1981;246(8):859–860. doi:10.1001/jama.1981.03320080045027
- Park M, Jung SJ, Yoon S, Yun JM, Yoon HJ. Association between the markers of metabolic acid load and higher all-cause and cardiovascular mortality in a general population with preserved renal function. *Hypertension Research*. 2015;38(6):433–438. doi:10.1038/hr.2015.23
- Abramowitz MK, Hostetter TH, Melamed ML. Lower serum bicarbonate and a higher anion gap are associated with lower cardiorespiratory fitness in young adults. *Kidney Int.* 2012;81(10):1033–1042. doi:10.1038/ki.2011.479
- Imran MN, Leng PH, Yang S, Kurup A, Eng P. Early predictors of mortality in pneumococcal bacteraemia. *Ann Academy Med Singapore*. 2005;34(7):426–431.
- Sahu A, Cooper HA, Panza JA. The initial anion gap is a predictor of mortality in acute myocardial infarction. *Coron Artery Dis.* 2006;17 (5):409–412. doi:10.1097/00019501-200608000-00002
- 13. Kim MJ, Kim YH, Sol IS, et al. Serum anion gap at admission as a predictor of mortality in the pediatric intensive care unit. *Sci Rep.* 2017;7(1):1456. doi:10.1038/s41598-017-01681-9
- Johnson AE, Pollard TJ, Shen L, et al. MIMIC-III, a freely accessible critical care database. *Scientific Data*. 2016;3:160035. doi:10.1038/ sdata.2016.35
- Banks PA, Bollen TL, Dervenis C, et al. Classification of acute pancreatitis–2012: revision of the Atlanta classification and definitions by international consensus. *GUT*. 2013;62(1):102–111. doi:10.1136/gutjnl-2012-302779
- 16. Agoritsas T, Merglen A, Shah ND, O'Donnell M, Guyatt GH. Adjusted analyses in studies addressing therapy and harm: users' guides to the medical literature. *JAMA*. 2017;317(7):748–759. doi:10.1001/jama.2016.20029
- Cheng B, Li D, Gong Y, Ying B, Wang B. Serum anion gap predicts All-Cause mortality in critically ill patients with acute kidney injury: analysis of the MIMIC-III database. *Dis Markers*. 2020;2020:1–10.
- Abramowitz MK, Hostetter TH, Melamed ML. The serum anion gap is altered in early kidney disease and associates with mortality. *Kidney Int.* 2012;82(6):701–709. doi:10.1038/ki.2012.196
- 19. Lee SW, Kim S, Na KY, et al. Serum anion gap predicts All-Cause mortality in patients with advanced chronic kidney disease: a retrospective analysis of a randomized controlled study. *PLoS One.* 2016;11(6):e0156381. doi:10.1371/journal.pone.0156381
- 20. Kim MJ, Kim YH, Sol IS, et al. Serum anion gap at admission as a predictor of mortality in the pediatric intensive care unit. *Sci Rep.* 2017;7:1.
- Tang Y, Lin W, Zha L, et al. Serum anion gap is associated with All-Cause mortality among critically ill patients with congestive heart failure. *Dis Markers*. 2020;2020:1–10. doi:10.1155/2020/8833637
- Feldman M, Soni N, Dickson B. Influence of hypoalbuminemia or hyperalbuminemia on the serum anion gap. J Lab Clin Med. 2005;146(6):317–320. doi:10.1016/j.lab.2005.07.008

#### **Dove**press

across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/ testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/international-journal-of-general-medicine-journal