The Impact of Treatment Adherence on Quality of Life Among Type 2 Diabetes Mellitus Patients – Findings from a Cross-Sectional Study

This article was published in the following Dove Press journal:
Patient Preference and Adherence

Abdul Majeed¹
 Muhammad Rehman¹
 Iltaf Hussain¹
 Imran Imran²
 Muhammad Usman Saleem³
 Hamid Saeed⁴
 Furqan K Hashmi⁴
 Muqarrab Akbar⁵
 Muhammad Asad Abrar¹
 Basit Ramzan⁶
 Muhammad Omer Chaudhry⁷
 Muhammad Islam⁴
 Naveed Nisar⁸
 Muhammad Fawad Rasool¹

¹Department of Pharmacy Practice, Faculty of Pharmacy, Bahauddin Zakariya University, Multan, 60800, Pakistan; ²Department of Pharmacology, Faculty of Pharmacy, Bahauddin Zakariya University, Multan, 60800, Pakistan; ³Department of Biosciences, Faculty of Veterinary Sciences, Bahauddin Zakariya University, Multan, 60800, Pakistan; ⁴University College of Pharmacy, University of the Punjab, Allama Iqbal Campus, Lahore, 54000, Pakistan; ⁵Department of Political Science, Bahauddin Zakariya University, Multan, 60800, Pakistan; ⁶Al Shifa Pharmacy, Multan, Pakistan; ⁷School of Economics, Bahauddin Zakariya University, Multan, 60800, Pakistan; ⁸Department of Pharmaceutics, Faculty of Pharmacy, Bahauddin Zakariya University, Multan, 60800, Pakistan

Correspondence: Muhammad Fawad Rasool
 Department of Pharmacy Practice, Faculty of Pharmacy, Bahauddin Zakariya University
 Multan, Multan, Pakistan
 Tel +923008639046
 Email fawadrasool@bzu.edu.pk

Aim: The current study was conducted to assess the factors contributing to treatment adherence and its impact on the quality of life (QoL) in type 2 diabetes mellitus (T2DM) patients.

Methods: A cross-sectional study was conducted between January 2020 to March 2020 among T2DM patients. The data was collected from T2DM out-patient clinics. The participants were recruited by using a simple random sampling method. To assess the association of demographics with the level of adherence, binary logistics regression analysis was applied. Moreover, the Mann–Whitney *U*-test was used to evaluate the impact of adherence on QoL.

Results: A total of 384 patients participated in this study. Amongst them, 60.2% were male and 39.8% were female. Low adherence was seen in illiterate patients and patients older than 40 years. The results showed that good QoL in T2DM patients was significantly associated with treatment adherence ($p = 0.004$).

Conclusion: The finding of the current study showed that the non-adherence prevailed in illiterate strata of the study population and the medication adherence significantly affects the QoL in T2DM patients. These findings suggest that health regulatory agencies should focus on implementing disease-education interventions for improving the adherence to medications in patients with long-term conditions.

Keywords: type 2 diabetes, drug adherence, quality of life

Introduction

The prevalence of long-term conditions is increasing with every passing day, as 50% of the visits in primary care clinics are attributed to these conditions.¹ It has been reported that 70% of the total health budget is consumed by patients having long-term conditions.¹ Since patients diagnosed with long-term conditions have to receive treatment for a long period, therefore the adherence to treatment and the quality of life (QoL) have become important parameters for assessing the effectiveness of the disease management plan. The improvement of patients' QoL is very important in the management of long-term conditions, as the poor QoL may lead to the development of disease-associated complications.²

Adherence is the integral component for the implementation of pharmacological and non-pharmacological approaches, and it plays an essential role in the effective management of long-term diseases. However, medication non-adherence is a common problem in patients with long-term conditions.^{3,4} Studies have shown that adherence to medications is generally higher in patients with an acute illness, as

compared to patients with chronic diseases.^{3,5–7} Adherence to medications is around 50% in the developed world and it is expected to be lower in the developing world.⁸ The patient and medication-related factors associated with treatment non-adherence have been reported in the literature.⁹

QoL is a multidimensional parameter that encompassing the emotional, physical, material, and social well-being of the individual.^{10,11} The healthcare providers goal of therapy in long-term conditions is to maintain or improve the health-related or disease-related QoL.¹² The QoL is not only associated with the disease manifestations but it is also related to the patient's perception toward their illness.¹³ To improve patient's QoL in long-term conditions, the goal should be to provide personalized care by actively involving them in designing their treatment plans and considering their personal preferences.¹⁰

According to the world health organization (WHO), Pakistan, in 2019, ranked at fourth position with a diabetic population of 19.4 million.¹⁴ It is estimated that in 2030, this number will increase to 26.2 million, and in 2045, it will be ranked number third in the world with a projected diabetic population of 37.12.¹⁵ Most of the patients are suffering from type 2 diabetes mellitus (T2DM) which comprises 90% of all diabetes mellitus cases in the world. T2DM is spreading at an alarming rate in Pakistan, therefore it is the need of time to take drastic steps towards its management and improving medication adherence among the diabetic population.^{14,15}

In Pakistan, limited information is available regarding the assessment of drug adherence with the QoL of T2DM patients.¹⁶ Therefore, if a study is conducted that can assess the impact of treatment adherence with the QoL in T2DM patients, it can be of great clinical importance. The current study was conducted to assess the factors contributing to treatment adherence and its effect on the QoL of T2DM patients.

Method and Instruments

Study Design, Setting, and Participants

A cross-sectional study was conducted between January 2020 to March 2020. Study participants were T2DM patients. The study was conducted in the outpatient diabetic clinics of Multan, Pakistan. The patients having T2DM with age greater than 25 years, and a diabetic history of more than 1 year were included in this study. The patients were enrolled in the study after signing an informed consent form.

Sample Size and Sampling Method

According to the International Diabetes Federation (IDF), the prevalence of T2DM in Pakistan was 17.1% in 2019.¹⁷ Therefore, a prevalence-based sample size was calculated by using the Raosoft online sample size calculator (Raosoft, Inc. USA)¹⁸ A sample size of 384 was obtained with a margin of error of 5%, and a Confidence level of 95%. The simple random sampling technique was utilized for the selection of study participants.

Ethical Consideration

This study was performed in accordance with the ethical standard of the institution and the Declaration of Helsinki and was approved by the ethical committee of the department of pharmacy practice, Bahauddin Zakariya University, Multan, Pakistan (Reference No: Acad/PRAC/18-20/20). To ensure the privacy and anonymity of patients, their names were replaced by a sequence of allotted numbers. Moreover, the data in hard form were kept in the departmental laboratory locker.

Study Instrument

In the present study, two instruments were used ie drug attitude inventory- 10 (DAI-10) to assess the treatment adherence and EQ-5D-5L to assess the QoL. Both scales were used after getting the permission.

DAI-10

DAI-10 scale is a self-administered and pre-validated scale that comprises of 10 questions based on True/ False responses.^{19,20} In this scale, a 1-point score is awarded to a correct response, and the total score greater than 7 shows high adherence, and a score less than 7 shows low adherence.

QoL Scale

The QoL scale EQ-5D-5L²¹ consisted of five (5) domains that include mobility, self-care, usual activities, pain/discomfort, and anxiety/ depression. Each domain is comprised of five levels of severity, which include: no problem, slight problem, moderate problem, severe problem, and extreme problem. The EQ-5D preference weight for each health state was not available for the Pakistani population. Therefore, the states were derived from time trade-off (TTO) tariff of preference weights from the general population of the United Kingdom.²²

Data Collection Procedure

The participants were randomly selected from the outpatient diabetic clinics from selected hospitals of Multan during their visit to the health care providers. Informed consent was taken from each patient before being included in the study, ensuring confidentiality, and describing to them the objectives of the study. Incomplete questionnaires were not included in the final analysis. The numbers of excluded questionnaire were sixty-six. The process of data collection is shown in [Figure 1](#).

Statistical Analysis

The statistical package for Social Sciences (SPSS) v 25 (Inc., Chicago, USA) was used for statistical analysis. Descriptive statistics were used to summarize the data. The general linear multivariate analysis model (GLM) was used to predict the significant differences between the independent and dependent variables, whereas the Mann–Whitney-*U* test was used to assess the impact of adherence level on QoL.

Figure 1 The data collection process.

Results

Demographic Characteristics of the Participants

A total of 384 patients participated in this study. The age of most participants was greater than 40 years (77.1%). The majority of the participants were male (60.2%) and residing in urban areas (59.9%). Most of the participants were married (95.8%) and illiterate (34.1%). One twenty-one participants were jobless (31.5%). The majority of the participants were using oral medications (52.1%), followed by insulin (25.8%) and 22.1% were taking both insulin and oral medications. The detailed demographics are provided in [Table 1](#).

Medication Adherence

The majority of the participants gave a positive response regarding the good and beneficial use of medications

Table 1 Demographic Characteristics of the Participants

Characteristics of the Participants		Frequency	Percentage
Age	25–40	88	22.9
	>40	296	77.1
Gender	Man	231	60.2
	Woman	153	39.8
Marital status	Married	368	95.8
	Unmarried	16	4.2
Education level	Illiterate	131	34.1
	Primary	37	9.6
	Secondary	47	12.2
	Graduate	99	25.8
	Postgraduate	70	18.2
Occupation	Public-sector job	61	15.9
	Private-sector job	109	28.4
	Business	93	24.2
	Unemployed	121	31.5
Residence	Rural	154	40.1
	Urban	230	59.9
Treatment	Oral	200	52.1
	Insulin	99	25.8
	Oral + insulin	85	22.1

Table 2 The Participant Responses to DAI-10 Scale

Questions		Frequency	Percentage
Q1	True	298	77.6
Q2	True	173	44.5
Q3	True	146	38.0
Q4	True	282	73.4
Q5	True	148	38.5
Q6	True	113	29.4
Q7	True	293	76.3
Q8	True	250	65.1
Q9	True	101	26.3
Q10	True	251	65.4

(62.0%) and most of the participants showed fear while taking medications (44.5%). The majority of the participants were using medication of their choice (38%) and they believed that the medications make them calmer and more relaxed (73.4%). The majority of the participants complained of sluggish and lazy behavior due to medication use (38.5%) and most of the participants felt normal

when they use their medication (76.3%). The participant responses to the DAI-10 scale can be seen in [Table 2](#).

Impact of Demographics on QoL and Drug Adherence

In the current study, a significant difference was seen in age ($p < 0.001$) and residency ($p = 0.041$) of the participants to their QoL. Moreover, concerning drug adherence, there was a significant difference across marital status and level of education of the participants. The detail is given in [Table 3](#).

QoL

In terms of mobility, most of the participants were having no problem (44%) followed by a slight (29.2%) and moderate problem (21.4%). The majority of participants had a slight problem in performing self-care (34.1%) while 56.3% had a slight problem in carrying usual activities. Most of the participants felt pain/discomfort (49.5%) and depression (38%). The QoL of the participants can be seen in [Table 4](#).

A total of 46 different health states were reported by the study participants, whereby 13 (3.4%) showed no problems in all five domains while 124 (32.4%)

Table 3 The Association of Demographics with the Level of Adherence and Quality of Life by Using the Multivariate Analysis

Source	Dependent Variable	Mean Square	F-value	P-value
Age	QoL	1202.07	100.36.	<0.001*
	Drug Adherence	0.142	0.04	0.842
Gender	QoL	38.937	3.045	0.124
	Drug Adherence	0.222	0.082	0.783
Marital status	QoL	4.04	0.337	0.56
	Drug Adherence	31.11	8.71	0.003*
Educational level	QoL	7.632	0.597	0.677
	Drug Adherence	42.012	15.478	0.001*
Occupation	QoL	14.875	1.163	0.389
	Drug Adherence	0.557	0.205	0.890
Treatment	QoL	16.837	1.317	0.327
	Drug Adherence	3.464	1.276	0.337
Residence	QoL	80.080	6.263	0.041*
	Drug Adherence	0.051	0.019	0.895

Note: * $p < 0.05$.

Abbreviation: QoL, quality of life.

Table 4 The Quality of Life of the Participants

Level of Problems	QoL Domains				
	Mobility	Self-Care	Usual Activity	Pain/ Discomfort	Anxiety/ Depression
Level 1 (No Problem)	169(44%)	178(46.4%)	72(18.8%)	60(15.6%)	93(24.2%)
Level 2 (Slight Problem)	112(29.2%)	131(34.1%)	216(56.3%)	190(49.5%)	146(38%)
Level 3 (Moderate Problem)	82(21.4%)	62(16.1%)	60(15.6%)	66(17.2%)	70(18.2%)
Level 4 (Severe Problem)	21(5.5%)	13(3.4%)	30(7.8%)	29(7.6%)	38(9.9%)
Level 5 (Extreme problem)	0	0	6(1.6%)	39(10.2%)	37(9.6%)

respondents pointed out no problem in mobility and self-care domain while moderate problem in pain and anxiety/depression domain ([Supplementary Table S1](#)).

Impact of Level of Adherence on QoL

The independent-samples Mann Whitney *U*-test revealed that the good quality of life in T2DM patients was significantly associated with a high level of adherence ($p = 0.004$) ([Supplementary Figure S1](#)).

Discussion

In the present study, we have assessed the relationship between treatment adherence and QoL of T2DM patients. Our results showed that the marital status and level of literacy were significantly associated with treatment adherence. Moreover, it was observed that the good QoL in the T2DM patients was associated with higher medication adherence rates.

Adherence to the treatment is an important factor in the management of chronic conditions.²³ Poor medication adherence is a major barrier to the management of T2DM. Poor medication adherence is also associated with health expenditures.²⁴ The medication adherence rate identified in this study was 33.3%. This number was low as compared to a similar study from the United State of America (79%),²⁵ but it was consistent with a study reported from Indonesia (29.7%).²⁶ This comparison indicates that in developed countries, there is a well-established healthcare system where the healthcare team works together to deliver health and provide individualized treatment (if needed) to the patients. While in developing countries with a focus on Pakistan, due to poor healthcare system, poor quality of pharmaceutical care to the patients, and an increase in the cost of therapy may contribute to the low treatment adherence in the T2DM

population. The incorporation of pharmaceutical care in the healthcare system has significantly contributed to treatment adherence in patients with long-term conditions.²⁷

It has been reported that the participants with a higher lower literacy level, living with the disease for a long time had a negative effect on their QoL and treatment adherence.^{26,28} In the current study, it was seen that the literacy level of the participants was significantly associated with higher treatment adherence and that is consistent with the previous reports.^{29,30} This finding highlights the need for focused health literacy programs in the illiterate stratum of the community.

In the current study, most of the participants report slight to moderate problems in terms of QoL. This finding was similar to the reported studies from Pakistan¹⁶ and Indonesia.²⁶ The QoL in T2DM patients is an important outcome parameter. Therefore, it is the primary goal of the healthcare providers to formulate sustainable health goals for the betterment of QoL of diabetic patients.³¹

Patients with chronic conditions are more likely to have physical, psychological, and social health issues as compared to patients with acute illnesses.^{32–34} In this study, a positive relationship between QoL and treatment adherence was seen in T2DM patients. The patients with poor treatment adherence showed a lower QoL and those with good treatment adherence showed a good QoL. This result was consistent with the previous reports in which the DAI-10 scale was used with similar cutoff-points.^{33,34} Moreover, medication adherence can be improved by focusing on patient counseling, disease education, and by decreasing the complexity of the prescribed dosing regimen. According to WHO, improving adherence to treatment in chronic conditions can significantly produce health and economic benefits.³⁵

Treatment non-adherence leads to disease-related complications and can increase the direct and indirect

healthcare costs.³ It is reported that in adults, around 10% of hospital admissions can be attributed to treatment non-adherence.³⁶ Moreover, the 10% increase in treatment adherence in diabetic patients can decrease the annual healthcare costs by 8.6–28.9%.³⁵ Also, non-adherence to medication can develop an uncontrolled disease condition that can have a considerable effect on patients' QoL, as it may lead to physical, social, and financial problems.³⁷

Practice Implications and Future Recommendation

The present study can have many implications at the national and global levels. The poor QoL and treatment non-adherence in T2DM patients is alarming. The Physician and Pharmacist should emphasize the importance of treatment adherence in the management of chronic conditions to their patients. Moreover, the barriers to treatment adherence in patients with chronic conditions can be addressed by allowing their active participation in designing personalized disease management plans.

Conclusion

The current study showed that T2DM patients with a higher level of treatment adherence showed good QoL. Moreover, the literacy level of T2DM patients was significantly associated with their level of adherence. The treatment adherence can be improved by simplifying the prescribed dosing regimen and ensuring appropriate patient counseling. To increase treatment adherence in T2DM patients, there is a need for the initiation of diabetes education programs at the community levels.

Author Contributions

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed to submit to the current journal; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

1. UK department of health. Long Term Conditions Compendium of Information Third Edition; 2020. Available from: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/216528/dh_134486.pdf. Accessed 05 October 2020.

2. Adriaanse MC, Drewes HW, Van Der Heide I, Struijs JN, Baan CA. The impact of comorbid chronic conditions on quality of life in type 2 diabetes patients. *Quality Life Res.* 2016;25(1):175–182. doi:10.1007/s11136-015-1061-0
3. Osterberg L, Blaschke T. Adherence to medication. *N Engl J Med.* 2005;353(5):487–497. doi:10.1056/NEJMra050100
4. World Health Organization. Noncommunicable Diseases and Mental Health; 2020. Available from: <https://www.who.int/mip/2003/progress/en/nmhmp2003.pdf>. Accessed 5 November, 2020.
5. Jackevicius CA. Adherence with statin therapy in elderly patients with and without acute coronary syndromes. *JAMA.* 2002;288(4):462–467. doi:10.1001/jama.288.4.462
6. Caro JJ, Salas M, Speckman JL, Raggio G, Jackson JD. Persistence with treatment for hypertension in actual practice. *CMAJ.* 1999;160(1):31–37.
7. Haynes RB, McDonald HP, Garg AX. Helping patients follow prescribed treatment: clinical applications. *JAMA.* 2002;288(22):2880–2883. doi:10.1001/jama.288.22.2880
8. Khayat SM, Mohamed MMA, Khayat SMS, et al. Association between medication adherence and quality of life of patients with diabetes and hypertension attending primary care clinics: a cross-sectional survey. *Quality Life Res.* 2019;28(4):1053–1061. doi:10.1007/s11136-018-2060-8
9. Marcum ZA, Seivick MA, Handler SM. Medication nonadherence: a diagnosable and treatable medical condition. *JAMA.* 2013;309(20):2105–2106. doi:10.1001/jama.2013.4638
10. Choudry MI, Stewart K, Woodhead T. The Royal College of Physician's Quality Improvement Hub – how can it help physicians to improve patient care? *Future Hospital J.* 2016;3(3):211–216. doi:10.7861/futurehosp.3-3-211
11. Davis MV, Mahanna E, Joly B, et al. Creating quality improvement culture in public health agencies. *Am J Public Health.* 2014;104(1):e98–104. doi:10.2105/ajph.2013.301413
12. Porterfield DS, Marcial LH, Brown S, Throop C, Pina J. Evaluation of a Quality Improvement Resource for Public Health Practitioners. *Public Health Rep.* 2017;132(2):140–148. doi:10.1177/0033354916689609
13. Martin JC, Avant RF, Bowman MA, et al. The Future of Family Medicine: a collaborative project of the family medicine community. *Ann Fam Med.* 2004;2 Suppl 1(Suppl 1):S3–32. doi:10.1370/afm.130
14. World Health Organization. Diabetic country profile; 2020. Available from: https://www.who.int/diabetes/country-profiles/pak_en.pdf. Accessed 5 November, 2020.
15. Sherin A. National diabetes action plan of Pakistan: need and challenges. *Khyber Med Univ J.* 2015;7(1):1–2.
16. Nazir R, Hassali MA, Saleem F, Bashir S, Aljadhey H. Does adherence to the therapeutic regimen associate with health related quality of life: findings from an observational study of type 2 diabetes mellitus patients in Pakistan. *Pak J Pharm Sci.* 2017;30(6):2159–2165.
17. International Diabetic Federation. *IDF DIABETES ATLAS 9th Edition.* 9th ed. 2019.
18. Raosoft I. Sample size calculator; 2004. Available From: <http://www.raosoft.com/samplesize.html>. Accessed 15 February, 2021
19. Hogan TP, Awad AG, Eastwood R. A self-report scale predictive of drug compliance in schizophrenics: reliability and discriminative validity. *Psychol Med.* 1983;13(1):177–183. doi:10.1017/S0033291700050182
20. Saleem F, Hassali MA, Shafie AA, Awad AG, Bashir S. Association between knowledge and drug adherence in patients with hypertension in Quetta, Pakistan. *Tropical J Pharm Res.* 2011;10:2.
21. Euqator Group. EQ-5D-5L; 2020. Available from: <https://euroqol.org/eq-5d-instruments/eq-5d-5l-about/>. Accessed 05 November 2020.
22. Dolan P, Gudex C, Kind P, Williams A. *A Social Tariff for EuroQol: Results from a UK General Population Survey.* 1995.
23. Bassett SM, Schuette SA, O'Dwyer LC, Moskowitz JT. Positive affect and medication adherence in chronic conditions: a systematic review. *Health Psychol.* 2019;38(11):960. doi:10.1037/hea0000778

24. Polonsky WH, Henry RR. Poor medication adherence in type 2 diabetes: recognizing the scope of the problem and its key contributors. *Patient Preference Adherence*. 2016;10:1299. doi:10.2147/PPA.S106821
25. Boccuzzi SJ, Wogen J, Fox J, Sung JC, Shah AB, Kim J. Utilization of oral hypoglycemic agents in a drug-insured U.S. population. *Diabetes Care*. 2001;24(8):1411–1415. doi:10.2337/diacare.24.8.1411
26. Perwitasari DA, Urbayatun S. Treatment adherence and quality of life in diabetes mellitus patients in Indonesia. *Sage Open*. 2016;6(2):2158244016643748. doi:10.1177/2158244016643748
27. Shao H, Chen G, Zhu C, et al. Effect of pharmaceutical care on clinical outcomes of outpatients with type 2 diabetes mellitus. *Patient Preference Adherence*. 2017;11:897. doi:10.2147/PPA.S92533
28. Saleh F, Mumu SJ, Ara F, Hafez MA, Ali L. Non-adherence to self-care practices & medication and health related quality of life among patients with type 2 diabetes: a cross-sectional study. *BMC Public Health*. 2014;14(1):431. doi:10.1186/1471-2458-14-431
29. Ueno H, Ishikawa H, Suzuki R, et al. The association between health literacy levels and patient-reported outcomes in Japanese type 2 diabetic patients. *SAGE Open Med*. 2019;7:2050312119865647. doi:10.1177/2050312119865647
30. Rodrigues da Rocha M, Domingas dos Santos S, Rafaela de Moura K, de Sousa Carvalho L, Holanda de Moura I, Vilarouca da Silva AR. Health literacy and adherence to drug treatment of type 2 diabetes mellitus. *Escola Anna Nery*. 2019;23:2.
31. Amelia R, Lelo A, Lindarto D, Mutiara E. Quality of life and glycemic profile of type 2 diabetes mellitus patients of Indonesian: a descriptive study. *InIOP Conf Series*. 2018. doi:10.1088/1755-1315/125/1/012171
32. Lam CL, Lauder IJ. The impact of chronic diseases on the health-related quality of life (HRQOL) of Chinese patients in primary care. *Fam Pract*. 2000;17(2):159–166. doi:10.1093/fampra/17.2.159
33. Azar FF, Solhi M, Chabaksva F. Investigation of the quality of life of patients with hypertension in health centers. *J Educ Health Promot*. 2020;9(1):185. doi:10.4103/jehp.jehp_741_19
34. ElShazly HM, Hegazy NN. Socioeconomic determinants affecting the quality of life among diabetic and hypertensive patients in a rural area, Egypt. *J Family Med Prim Care*. 2017;6(1):141–145. doi:10.4103/jfmpe.jfmpe_31_17
35. Saleem F, Hassali MA, Shafie AA. A cross-sectional assessment of health-related quality of life (HRQoL) among hypertensive patients in Pakistan. *Health Expect*. 2014;17(3):388–395. doi:10.1111/j.1369-7625.2012.00765.x
36. Iqbal Q, Ul Haq N, Bashir S, Bashaar M. ul Haq N, Bashir S, Bashaar M. Profile and predictors of health related quality of life among type II diabetes mellitus patients in Quetta city, Pakistan. *Health Qual Life Outcomes*. 2017;15(1):142. doi:10.1186/s12955-017-0717-6
37. Brown MT, Bussell JK. Medication adherence: WHO cares? *Mayo Clinic Proc*. 2011;86(4):304–314. doi:10.4065/mcp.2010.0575

Patient Preference and Adherence

Dovepress

Publish your work in this journal

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focusing on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize clinical outcomes for existing disease

states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/patient-preference-and-adherence-journal>