

Seroprevalence and Associated Risk Factors of Canine *Leptospira* and *Brucella* Species Infection in West Shewa Zone, Central Ethiopia

This article was published in the following Dove Press journal:
Veterinary Medicine: Research and Reports

Lencho Megersa Marami¹
Endrias Zewdu
Gebremedhin²
Edilu Jorga Sarba²
Getachew Kebebew Tola²
Solomon Shiferaw Endalew¹
Asamenew Tesfaye Melkamsew³
Vincenzo Di Marco Lo Presti⁴
Maria Vitale⁴

¹Ambo University, College of Agriculture and Veterinary Sciences, Department of Veterinary Laboratory Technology, Ambo, Oromia, Ethiopia; ²Ambo University, College of Agriculture and Veterinary Sciences, Department of Veterinary Sciences, Ambo, Oromia, Ethiopia; ³National Animal Health Diagnostic and Investigation Center, Department of Immunology, Sebeta, Oromia, Ethiopia; ⁴Istituto Zooprofilattico Sperimentale di Sicily, Laboratory of Genetics of Microorganisms, Palermo, Italy

Purpose: Canine leptospirosis and brucellosis are significant zoonotic gram-negative bacterial diseases that affect humans and animal species. This study was aimed to estimate the seroprevalence and assess associated risk factors of canine *Leptospira* and *Brucella* species infections in Ambo, Bako, and Gojo Towns, West Shewa Zone, Ethiopia.

Methods: A total of 385 serum samples were collected from randomly selected dogs. Antibodies against *Brucella* and *Leptospira* species infection were tested using the Enzyme-Linked immunosorbent Assay technique (ELISA). A structured questionnaire survey was administered to each dog-owning household to gather information on potential risk factors. The association between independent and outcome variables was analyzed using the Chi-square test followed by univariable, and multivariable logistic regression analyses.

Results: The overall seroprevalence of *Brucella* and *Leptospira* species infections were found to be 15.06% (95% confidence interval (ci): 11.64–19.04%) and 4.16% (95% ci: 2.39–6.67%), respectively. Multivariable logistic regression analysis showed that the town was the significant risk factor of *Brucella* species infection seropositivity. Accordingly, dogs found in Gojo ((odds ratio (OR): 17.72 CI: 1.17–117.54, $p=0.036$) and Bako (OR: 7.99, 95% CI: 0.96–66.37), $p=0.054$ towns were more at risk to be infected with *Brucella* species than dogs living in Ambo.

Conclusion: The seroprevalence of *Leptospira* and *Brucella* species infections in dogs of the West Shewa Zone was high and moderate, respectively, suggesting the possible transmission of the disease between dogs and other animals. These infections might be an under-recognized threat to public health and animal welfare. Further research on the identification of the serovars of *Leptospira* and biotypes of *Brucella* circulating in dogs is encouraged. Finally, knowledge of the comprehensive epidemiology of these diseases is an invaluable input for veterinarians, healthcare professionals, and policy-makers to avoid or manage canine leptospirosis and brucellosis.

Keywords: canine, *Brucella*, Ethiopia, *Leptospira*, risk factor, seroprevalence

Introduction

The pathogenic bacteria, *Leptospira* and *Brucella* species cause leptospirosis and brucellosis respectively in many species of animals including cattle, sheep, camels, buffaloes, dogs, and horses around the globe. Leptospirosis and brucellosis are substantial causes of abortion in dogs.¹ In addition to causing economic losses in the dog breeding industry, these infectious diseases also pose a potential risk to human health, especially to dog breeders and children who come into close contact with dogs.^{2,3}

Correspondence: Endrias Zewdu
Gebremedhin
Ambo University, College of Agriculture and Veterinary Sciences, Department of Veterinary Sciences, P.O. BOX 19, Ambo, Oromia, Ethiopia
Tel +251911894802
Email endrias.zewdu@ambou.edu.et

Canine leptospirosis is one of the zoonotic bacterial diseases affecting dogs, other animals, and humans globally.^{4,5} According to the report of Costa et al,⁶ every year 1.03 million people become infected, and 58,900 die from leptospirosis. Asia, Latin America, and sub-Saharan Africa have witnessed this high rate of disease and death. The prevalence of leptospirosis is high in tropical and subtropical wetlands, due to the climatic and environmental conditions in these regions that support the presence of *Leptospira* spp attributes to high temperatures, rainfall, humidity, the presence of infected domestic and wild animals, and the poor drainage of wastes.^{7,8} In Sub-Saharan Africa, the disease is seen as prevalent, knowledge regarding its epidemiology is still lacking in most countries.⁹

Over 200 *Leptospira* serovars from 17 genospecies have been identified, and they are distinct from each other by geographical distribution, their host of infection, reservoir, and their virulence factors.^{4,10} Dogs are susceptible to most *Leptospira* serovars, including Canicola, Icterohaemorrhagiae, Pomona, Grippotyphosa, and other serovars.^{11,12} However, dogs are the main hosts for Canicola and Icterohaemorrhagiae serovars.¹³ Leptospirosis is recognized to cause late-term abortion in any pregnant animal, including people.⁴ In dogs, the common sign of this disease is septicemic, hepatic, and renal infection with a mortality rate of 10–20%.^{4,14}

Healthy animals or humans become infected when they come in contact with infected mucous membranes or skin or milk or vaginal fluid. Besides, humans acquire leptospirosis by direct contact with contaminated drinking water or soil. The main disease reservoirs are rodents, but a variety of wild and domestic animals, livestock, and insectivores are considered as reservoirs. *Leptospira* enters the body of the susceptible animal or human through mucous membranes, conjunctiva, small cuts, abrasions, and possibly wet skin.⁴

Brucellosis is the most common and serious global distributed zoonotic disease.¹⁵ It is an endemic disease in North and East Africa, including other developing countries, but has not been reported or less recognized.¹⁶ Globally, an estimated 500,000 new human cases of brucellosis are reported each year.^{17,18}

The genus *Brucella* has 10 known host-specific species and causes brucellosis in various domestic animals and humans.¹⁹ *B. melitensis*, *B. abortus*, *B. suis*, and *B. canis* are the most recognized species of *Brucella* known to cause brucellosis in people.^{16,18,20,21} Human brucellosis causes acute febrile illness (undulant fever) which may

progress to a more persistent form and can also produce serious complications in muscle-skeletal, cardiovascular, and central nervous systems.²² *B. canis* causes asymptomatic infection in humans and may last unconfirmed for a prolonged period, and thus the symptoms include headache, recurrent fever, weakness, fatigue, chills, sweats, and weight loss.^{23–25} The zoonotic risk is moderately high in people who handle rearing dogs in kennels and are exposed to reproductive tissues and fluids of infected dogs. However, chasers and proprietors of hunting dogs are at high risk for transmission of brucellosis from wild-life. Occasional transmission of *B. canis* from pet canine to their proprietors has been reported.^{21,25}

Brucella canis is one of the intracellular gram-negative bacteria that mainly causes reproductive failure in dogs. The primary sources of brucellosis in dogs are infected with vaginal discharge and urine of male dogs.^{26,27} The disease is mainly transmitted between dogs through genital, oronasal, conjunctiva, and infected placenta.²⁷ Late-term abortions in bitches, epididymitis in males, and infertility in both sexes are the most common symptoms of canine brucellosis.^{20,27} Dogs are also infected by *B. abortus*, *B. suis*, and *B. melitensis* whenever they share a common environment with cattle, sheep, goats, and pigs.^{26,27}

There is no recent data on the seroprevalence of leptospirosis of dogs in Ethiopia after the study done by Moch et al.²⁸ Similarly, there is no accessible study conducted on the seroprevalence of canine brucellosis in Ethiopia. However, there is a report of *Leptospira* seroprevalence in humans in 2004²⁹ and cart horses in 2016;³⁰ and *Brucella* seroprevalence in cattle in 2011,³¹ 2016,³² and 2020.³³ Knowledge about the status of these neglected diseases in dogs might help to design prevention and control strategies that limit the spread of *Leptospira* and *Brucella* spp infections to other animals and humans. Therefore, this research aims to estimate the seroprevalence of canine *Leptospira* and *Brucella* spp infections and to determine the associated risk factors in Ambo, Bako, and Gojo towns, West Shewa Zone, Oromia, Ethiopia.

Materials and Methods

Study Areas

The study was conducted in Ambo, Bako, and Gojo towns of West Shewa Zone of Oromia Regional State, Ethiopia. Gojo, Ambo, and Bako towns were purposively selected to

represent highland, midland, and lowland altitudes, respectively.

Ambo is the administrative center of the West Shewa Zone of the Oromia region is 114 Km located west of Addis Ababa. The town has a latitude and longitude of 8° 59'N 37°51'E and an elevation of 2101 meters. The average annual temperature and rainfall are 22° C and 900 mm respectively. According to Central Statistical Agency (CSA),³⁴ the total population of Ambo was 48,171, of which 24,634 were males and 23,537 females.

Bako town, the administrative center of the Bako-Tibe district, is located 260 Km West of Addis Ababa. The town has a longitude and latitude of 9°08'N 37°03'E with an elevation of 1743 meters. The average annual temperature is 19.7 °C while the rainfall is 1281 mm. According to the Central Statistical Agency, in 2005 Bako town has an estimated total human population of 18,641, of whom 9370 are men and 9271 women.³⁴

Gojo town is the administrative center of Jeldu district located 120 Km West of Addis Ababa. The town has a latitude and longitude of 9.26°N 38.09° E and an elevation of 2905 meters. The mean annual temperature for the town is 20°C; it receives an annual rainfall of 2500 mm. The human population of Jeldu district is 202,655 of which 102,796 are females and the remaining 99,859 males.³⁴ There were no recorded or known data on the dog population in the three towns.

Study Population

Dogs found in three towns were the study population. Only owned dogs were included and the stray dogs were excluded from the study. These owned dogs were confined to the home or outdoors. Dogs younger than 3 months of age were not included in the study due to the presence of maternal antibodies that exist at an early age.³⁵

Study Animals

Domestic owned dogs (*Canis familiaris*) found in selected Gotes ("Gote" is a subdivision of *Kebele* containing 20–30 households and "Kebele" refers to the smallest administrative unit in a district) were sampled during the house-to-house survey. Dogs in Ethiopia including in the study towns are primarily used for protection and companionship. Dogs are fed with whatever food is available such as household leftovers and animal products. Vaccination of dogs against brucellosis and leptospirosis was not practiced in the study areas.

Study Design

The cross-sectional study design was conducted from January 2015 to June 2017 to estimate the seroprevalence of *Brucella* and *Leptospira* species infections and to determine their associated risk factors in Ambo, Bako, and Gojo towns of West Shewa Zone, Oromia Regional State.

Sample Size and Sampling Technique

In Ethiopia, no study was previously done on canine leptospirosis and brucellosis seroprevalence. Thus, 8.3% expected prevalence,²⁸ 5% desired absolute precision, and a 95% confidence interval was used to calculate the required sample size. The sample size was calculated based on the Thrusfield³⁶ formula. $N = Z^2 P_{exp} (1 - P_{exp}) / D^2$. Where, $Z = 1.96$, N = sample size, P_{exp} = expected prevalence, and D = absolute precision. The calculated sample size (117) was inflated 3.2 times to account for clustering of prevalence during multistage sampling. Therefore, 384 was considered as sample size.

There was no information on the number of dogs in the three towns. Therefore, it is estimated that the number of dogs in the towns was evenly distributed. The number of dog owners living in the study towns was also taken into account. A multi-stage sampling procedure was used to choose households for this study. There are three, two, and one *Kebeles* in Ambo, Bako, and Gojo towns, respectively. From each "Kebeles," four "Gotes" were randomly selected using the list of *Gotes* in each *Kebeles* (sampling frame) provided by local administrators. The index household in a *Gote* was selected using a simple random sampling technique and subsequent households were surveyed door to door. Of the sample size ($n = 385$), 169 samples were collected from Ambo town, 148, and 68 samples were collected from Bako and Gojo towns, respectively.

Sampling and Transportation

The blood sample was obtained from the dogs' cephalic vein using a 10 mL plain vacutainer tubing. The blood specimens were then centrifuged at 3000 RPM for at least 10 minutes. The sera were harvested and transported to the Ambo University Veterinary Microbiology Laboratory in labeled sterile cryovials kept in an icebox with ice packs. The serum samples were transported to the National Animal Health Diagnostic Center (NAHDIC) at Sebeta, under the cold chain, and stored at -20 °C until serological testing.

Serological Tests

Leptospira Species

The serological tests were conducted at the National Animal Health and Diagnostic Center (NAHDIC), Sebeta, Ethiopia. Enzyme-linked immunosorbent assay (NovaTec VetLine *Leptospira* ELISA, Germany) was conducted according to the manufacturer's instruction to screen antibodies against *Leptospira* species infection. Microtiter strip wells that pre-coated with a *Leptospira* antigen were used. Firstly, 100 μ L controls and diluted samples were dispensed into their respective wells. Secondly, the assay was covered by a foil and incubated for 1-hour \pm 5 min at $37 \pm 1^\circ$. Thirdly, the content of each well was aspirated and washed three times with 300 μ L of Washing Solution. Fourthly, 100 μ L VetLine *Leptospira* Protein A/G Conjugate was dispensed into all wells except for the blank well. Fifthly, the assay was covered with foil and incubated for 30 min at room temperature. Again, the contents of each well were aspirated and washed. Next, 100 μ L Tetramethylbenzidine (TMB) Substrate Solution were dispensed and incubated for 15 minutes at room temperature in the dark. Finally, the reaction was stopped by adding 100 μ L of stopping solution. Using the ELISA micro-well plate reader, the optical density (OD) value at 450 nm was recorded and interpreted as the manufacturer's instruction.

Brucella Species

The commercial test kits of indirect enzyme-linked immunosorbent assay (ID Screen[®] Brucellosis Serum Indirect Multi-species, ID. Vet, France) designed to detect *B. abortus*, *B. melitensis*, and *B. suis* were used. The wells of the indirect *Brucella* ELISA kit were coated with *Brucella abortus* lipopolysaccharide (LPS). The method is performed according to the instructions given by the manufacturer. In brief, after the dilution buffer, negative control, and positive control were added to the wells. According to the instruction given, 10 μ L serum samples were added to the remaining wells. Then the test was incubated at room temperature for 45 minutes and the contents of the wells were aspirated. Immediately, empty wells were washed three times by 300 μ L of the wash solution. Next, 100 μ L 1x conjugate was added to each well and incubated at room temperature for 30 minutes. Again, it was washed three times by 300 μ L of the wash solution. Later, 100 μ L of the substrate was added to each well and incubated at 25 $^\circ$ C for 15 minutes in the dark. Finally, the reaction was stopped by adding 100 μ L of stopping

solution. Using the ELISA micro-well plate reader, the O.D. value at 450 nm was recorded and interpreted as the manufacturer's instruction.

Questionnaire Survey

A structured questionnaire survey was administered to the dog owners regarding information related to dog husbandry and others. The close-ended questions included were age, sex (male, female), breed (exotic, cross, indigenous), housing system (indoor, outdoor), type of dog feed (cooked animal products, raw animal products), the existence of other domestic animals in the household (yes, no), dog owner education level (illiterate, primary, secondary, tertiary), living area/residence (urban, peri-urban), marital status of dog-owning household (single, married, divorced), and presence of rodents in the house. The age of dogs was categorized as a juvenile (6 weeks to 6 months), adolescent (6 months to 18 months), adult (18 months to 7 years), and geriatric (greater than 7) years according to Kiflu et al³⁷ and based on owners' information.

Statistical Analysis

The data from the questionnaire survey and laboratory findings were filled and coded in Microsoft Excel and analyzed using STATA version 14.2.³⁸ Chi-square, univariable, and multivariable logistic regression were used to evaluate the relationship between independent (potential risk factors) and outcome variables (seroprevalence). To identify risk factors associated with infection of the *Brucella* species, a multivariable logistic regression model was used after checking collinearity, and variables with a p-value < 0.25 in the univariable logistic regression analysis were selected. An adjusted odds ratio with a 95% confidence interval (ci) was used to express the degree of association. The 95% confidence level for the subgroup and overall prevalence values were calculated using the exact binomial test. A p-value of < 0.05 was considered significant.

Ethical Issue

The purpose of the study was clarified to the owners of the dog and the members of the family, and then verbal informed consent was obtained from dog owners before blood sample collection. The verbal informed consent from the dog owners was approved by the Research and Ethical Committee of Ambo University. The research team administered the questionnaire face-to-face. The standard

sample collection protocols were followed to collect blood samples and dogs were handled with best practices of veterinary care. All dogs in the study towns were given an anti-rabies vaccine and anti-parasitic drugs free of charge and service cost.

Results

Overall Seroprevalence

The majority of the dogs involved in the study were adults (59.7%), males (76.1%), and had outdoor access (69.1%). Similarly, most of the dogs studied were indigenous (76.6%) and dog-owning households also had other domestic animals (53.3%). About 88.6% of the dogs sampled were from urban areas, while the remaining 11.4% were from peri-urban areas. The presence of rodents was reported from 46.8% of dog-owning households.

The overall seroprevalence of *Leptospira* and *Brucella* spp infection in dogs was 15.06% (95% CI: 11.64–19.04) and 4.16% (95% CI: 2.39–6.67), respectively. There was no significant difference in *Leptospira* spp seropositivity between study towns ($p>0.05$). However, *Brucella* species seropositivity was significantly different between study towns ($p<0.05$) (Table 1).

Risk Factors

Leptospira Species Infection

The results of univariable logistic regression analysis showed that the age of the dog, sex, housing system, breed, type of dog feed, the existence of other domestic animals, dog owner education level, towns, community type, and presence of rodents were not significantly associated with *Leptospira* seropositivity ($p>0.05$) (Table 2). A multivariable logistic regression model was not modeled due to univariable p-values greater than 0.25.

Brucella Species Infection

As shown in Table 3, the age of dogs, sex of the dog, housing system, the existence of other domestic animals, dog owner education level, community type, marital status of the head of the household were not significantly associated with *Brucella* seropositivity ($p>0.05$). Sex of the dog, type of dog feed, existence of other domestic animals, dog owner education level, and the marital status of the household were not fitted to the final model due to their univariable p-value were greater than 0.25. A multivariable logistic regression analysis indicated that the town

was an independent predictor of *Brucella* species infection seropositivity ($p<0.05$).

Discussion

In Ethiopia, there are no recent usable data on the seroprevalence of leptospirosis and brucellosis in dogs, so the current study used data from other countries to compare and contrast. This research report on the infection seroprevalence of the *Brucella* species in dogs was the first report from Ethiopia. The overall seroprevalence of *Leptospira* species infection was 15.06%. Our finding on seroprevalence of *Leptospira* infection was analogous to the seroprevalence in dogs (15.6%) in Zimbabwe.³⁹ Similarly, the seroprevalence of *Leptospira* infection in the present study was in agreement with the finding of Spangler et al⁴⁰ who reported 13.1% seroprevalence in southeastern Appalachia.

Compared to the present findings, the lower seroprevalence of leptospirosis was reported in Malaysia (3.1%), Thailand (4.4%), Brazil (7.1%), and Ethiopia.^{7,28,41,42} On the contrary, the higher seroprevalence of *Leptospira* infection was reported by Aslantaş et al,⁴³ Delaude et al⁴⁴ and Pratt et al⁴⁵ from Turkey, Switzerland, and the Caribbean Islands of Saint Kitts with the prevalence of 43.96% (51/116), 55.7% (210/377), and 73.2% (74/101), respectively. These variations in seroprevalence figures recorded by various researchers may be attributable to geographical variance (*Leptospira* lives longer and persists in hot, humid, and sub-tropical areas), distribution of wild animals, rodents,⁹ types of serological test, sample size, and season of sample collection. Similar to the findings of Meeyam et al⁴⁶ the present study did not come up with a significant association of canine *Leptospira* seroprevalence and investigated risk factors like sex and age ($p>0.05$).

In this study, the seroprevalence of the *Brucella* infection was comparable to the findings of Mosalanezhad et al,⁴⁷ who reported 4.9% from companion dogs in Ahvaz, Iran. The current finding is also consistent with the 7.5% seroprevalences of *B. canis* in shelter dogs in Turkey,⁴⁸ 4.9% in Mississippi,⁴⁹ and 7.9% in Japan.⁵⁰

The present study showed that there was a statistically significant difference in *Brucella* infection among the towns. The seroprevalence of *Brucella* infection was higher in Gojo and Bako than in Ambo town. In accord with the present findings, a significant difference in the seroprevalence of canine brucellosis concerning the location was also previously reported in southwestern Nigeria.²⁶ Although the presence of other domestic animals in the households was not significantly associated

Table 1 Overall Seroprevalence of *Leptospira* and *Brucella* Species Infection in Dogs in Ambo, Bako, and Gojo Towns

Towns	No. Tested	Leptospira Species*		Brucella Species**	
		No. Positive	Prevalence (95% CI)	No. Positive	Prevalence (95% CI)
Ambo	169	25	14.79 (9.80–21.06)	1	0.59 (0.01–3.25)
Gojo	68	12	17.65 (9.47–28.80)	7	10.29 (4.24–20.06)
Bako	148	21	14.19 (9.0–20.86)	8	5.41 (2.36–10.37)
Overall	385	58	15.06 (11.64–19.04)	16	4.16 (2.39–6.67)

Notes: *Chi-square: 0.45; p-value: 0.797. **Chi-square: 12.40; p-value: 0.002.

Abbreviations: No, number; CI, confidence interval.

with seropositivity, the variation in *Brucella* species seroprevalence between towns could be due to extensive breeding of cattle and sheep or abundance and uncontrolled movement of dogs in Gojo and Bako towns.

Thus, dogs might access *Brucella* spp infected tissues of these domestic animals. Therefore, the transmission of disease between dogs and other animals may occur. The spread of brucellosis between the vulnerable host occurs,

Table 2 Univariable Regression Analysis of Potential Risk Factors Associated with the Seropositivity of Dog *Leptospira* Species Infection in Ambo, Gojo, and Bako Towns

Variables	Categories	No. of Tested	Number of Positive (Prevalence)	Univariable	
				OR (95% CI)	p-value
Age	Geriatric	51	7 (13.73)	1.0	–
	Adult	230	34 (14.78)	1.09 (0.45–2.62)	0.847
	Juvenile	27	4 (14.81)	1.09 (0.29–4.12)	0.895
	Adolescent	77	13 (16.88)	1.28 (0.47–3.46)	0.631
Sex	Female	92	12 (13.04)	1.0	–
	Male	293	46 (15.70)	1.24 (0.63–2.46)	0.535
Housing system	Outdoor	266	38 (14.29)	1.0	–
	Indoor	119	20 (16.81)	1.21 (0.67–2.19)	0.523
Breed	Exotic	15	2 (13.33)	1.0	–
	Indigenous	296	41 (13.85)	1.05 (0.23–4.80)	0.955
	Cross	74	15 (20.27)	1.65 (0.34–8.12)	0.537
Type of dog feed	Raw meat	282	41 (14.54)	1.0	–
	Cooked meat	103	17 (16.50)	1.16 (0.63–2.15)	0.633
Existence of other domestic animals	No	181	24 (13.26)	1.0	–
	Yes	204	34 (16.67)	1.30 (0.74–2.30)	0.352
Dog owner education level	Secondary	125	15 (12.00)	1.0	–
	Illiterate	47	7 (14.89)	1.28 (0.49–3.38)	0.613
	Primary	98	16 (16.33)	1.43 (0.67–3.06)	0.356
	Tertiary	115	20 (17.39)	1.54 (0.75–3.18)	0.239
Towns	Bako	148	21 (14.19)	1.0	–
	Ambo	169	25 (14.79)	1.05 (0.56–1.97)	0.879
	Gojo	68	12 (17.65)	1.30 (0.60–2.82)	0.513
Community type	Urban	341	48 (14.08)	1.0	–
	Peri-urban	44	10 (22.73)	1.80 (0.83–3.87)	0.135
Presence of rodents	No	205	29 (14.15)	1.0	–
	Yes	180	29 (16.11)	1.17 (0.67–2.04)	0.591

Table 3 Univariable and Multivariable Logistic Regression Analysis of Potential Risk Factors Associated with the Seropositivity of Dog *Brucella* Species Infection in Ambo, Gojo, and Bako Towns

Variables	Categories	No. of Tested	Number of Positive (Prevalence)	Univariable		Multivariable	
				OR (95% CI)	p-value	OR (95% CI)	p-value
Age	Adolescent	77	1 (1.30)	1.0	–	1.0	–
	Juvenile	27	1 (3.70)	2.92 (0.18–48.42)	0.454	2.02 (0.12–35.50)	0.628
	Geriatric	51	2 (3.92)	3.10 (0.27–35.16)	0.361	2.27 (0.19–26.77)	0.516
	Adult	230	12 (5.22)	4.18 (0.53–32.71)	0.173	5.70 (0.69–47.23)	0.106
Sex	Male	293	11 (3.75)	1.0	–		
	Female	92	5 (5.43)	1.47 (0.50–4.36)	0.484		
Housing system	Indoor	119	2 (1.68)	1.0	–	1.0	–
	Outdoor	266	14 (5.26)	3.25 (0.73–14.53)	0.123	2.08 (0.44–9.91)	0.356
Type of dog feed	Cooked meat	103	2 (1.94)	1.0	–	1.0	–
	Raw meat	282	14 (4.96)	2.64 (0.59–11.81)	0.205	3.59 (0.74–17.29)	0.111
Existence of other domestic animals	No	181	6 (3.31)	1.0	–		
	Yes	204	10 (4.90)	1.50 (0.53–4.22)	0.439		
Dog owner education level	Illiterate	47	1 (2.13)	1.0	–		
	Primary	98	3 (3.06)	1.45 (0.15–14.35)	0.749		
	Secondary	125	5 (4.00)	1.92 (0.22–16.85)	0.557		
	Tertiary	115	7 (6.09)	2.98 (0.36–24.93)	0.313		
Towns	Ambo	169	1 (0.59)	1.0	–	1.0	–
	Bako	148	8 (5.41)	9.60 (1.19–77.69)	0.034	7.99 (0.96–66.37)	0.054
	Gojo	68	7 (10.29)	19.27 (2.32–159.92)	0.006	11.72 (1.17–117.54)	0.036
Community Type	Urban	341	10 (2.93)	1.0	–	1.0	–
	Peri-urban	44	6 (13.64)	5.22 (1.80–15.18)	0.002	2.56 (0.68–9.68)	0.165
The marital status of the household	Divorced	25	1 (4.00)	1.0	–		
	Married	343	14 (4.08)	1.02 (0.13–8.10)	0.984		
	Single	17	1 (5.88)	1.50 (0.08–25.75)	0.780		

mainly direct contact with an affected animal after abortion and sharing a home.⁵¹

In the current research, when other domestic animals were present in the household, the seropositivity of dogs to *Leptospira* and *Brucella* infections was moderately high as compared to the absence of other domestic animals in the household owning dogs.

The seroprevalence of *Brucella* and *Leptospira* infections was higher in those dogs fed raw meat as compared to dogs fed cooked meat. This distinction is possible because cooking animal tissues or meat could kill microorganisms.

The seropositivity of *Brucella* infection was relatively higher in female than in male dogs and peri-urban dogs than urban dogs. Similarly, the seroprevalence of *Brucella* species infection in adult and geriatric dogs was 4.18 and 3.10 times higher, respectively, than the seroprevalence in young dogs. Dogs held in the outdoor housing system

(5.26%) had a slightly higher seroprevalence of *Brucella* species infection than indoor kept dogs (1.68%). However, the statistical analysis revealed that there were no significant differences among categories of these variables concerning *Brucella* species infection.

Leptospirosis and brucellosis are still a serious public health problem worldwide, and a dog could serve as a source of infection for other animals and humans. Ethiopia is an agrarian country with huge untapped livestock resources dominated by traditional husbandry and owned by poor farmers. In Africa, Ethiopia ranks second only to Nigeria in the health burden of zoonotic diseases and stands in the first place as a hot spot of leptospirosis in the world.⁵²

The high humidity and rainfall in the studied towns coupled with the fact that infected dogs remain chronic carriers and shedders of *Leptospira* through their urine,⁴² it is likely that the environment of the towns is

contaminated, posing a risk of infection to other animals and humans.

The present study showed the status of *Brucella* and *Leptospira* infections and their associated factors in dogs in the study area. In Ethiopia, dog ownership is common, and humans live in close contact with dogs and other animals. In addition, there is no vaccination and control program for brucellosis and leptospirosis. *Brucella* and *Leptospira* circulate between dogs, other animals, and human beings. Both diseases reduce the productivity of livestock and induce medical problems in humans. Although Ethiopia mainly depends on livestock production, information that supports the prevention and control of these diseases is not well accessible. Therefore, the present findings will have an invaluable input to prevent and control brucellosis and leptospirosis in Ethiopia.

Despite the important findings, the main limitation of the current study was the failure to identify the serovars of *Leptospira* and species of *Brucella* due to the nature of the diagnostic kits used. The *Brucella* ELISA kit in the current study uses a multi-species conjugate, which recognizes mammalian antibodies and therefore should work on dog sera. However, the kit is not validated for canine sera. It does not detect also *Brucella canis* hence the results might be an underestimate. Furthermore, the association of some significant risk factors like abortion, stillbirth, neonatal death, retained placenta with *Brucella* seropositivity were not assessed. The association of rainfall with *Leptospira* seropositivity was not also investigated. Those households not willing to participate in the study might not be evenly distributed in all sampling areas and door-to-door sampling was done once the index case was identified. These might entail some bias. Such studies could have given more data into the epidemiology of brucellosis and leptospirosis in the studied dogs. Therefore, the findings of this study should be observed in light of these limitations.

Conclusions

In conclusion, a relatively high seroprevalence of *Leptospira* and moderate *Brucella* spp infections were found in dogs of the West Shewa Zone. Consequently, the risk of contracting these infections is higher among humans and other domestic animals. *Leptospira* and *Brucella* spp infections might constitute an under-recognized threat to public health and animal welfare. The town is the predictor of *Brucella* spp seropositivity. None of the considered variables were independent predictors of *Leptospira* spp seropositivity. Further research is

encouraged on the detection of *Leptospira* serovar and *Brucella* species circulating in dogs. Finally, comprehensive knowledge of the epidemiology of these infections is valuable input for veterinarians, health care professionals, and policy-makers to manage these infections.

Acknowledgments

We would like to thank Ambo University and the Italian National Reference Centre for Toxoplasmosis at the Istituto Zooprofilattico Sperimentale della Sicilia A. Mirri, Italy for their financial support. We would also like to thank the dog owners and veterinarians for their cooperation during the field study. The National Animal Health Diagnostic Center (NAHDIC) at Sebeta is highly acknowledged for allowing the laboratory facilities.

Author Contributions

All authors made a significant contribution to the work reported either in the conception of the study, execution, acquisition of data, analysis, and interpretation or in all these areas. Moreover, all authors took part in revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

This work was supported by Ambo University and the Italian National Reference Centre for Toxoplasmosis at the Istituto Zooprofilattico Sperimentale della Sicilia A. Mirri, Italy (grant No. RF 2007/RC).

Disclosure

The authors declare that there are no conflicts of interest regarding the publication of this paper.

References

1. Cutler S, Whatmore A, Commander N. Brucellosis—new aspects of an old disease. *J Appl Microbiol*. 2005;98(6):1270–1281. doi:10.1111/j.1365-2672.2005.02622.x
2. Heinemann MB, Garcia JF, Nunes CM, et al. Detection and differentiation of *Leptospira* spp. serovars in bovine semen by polymerase chain reaction and restriction fragment length polymorphism. *Vet Microbiol*. 2000;73(4):261–267. doi:10.1016/s0378-1135(00)00150-4
3. Ocholi R, Kwaga J, Ajogi I, Bale J. Abortion due to *Brucella abortus* in sheep in Nigeria. *Rev Sci Tech off Int Epiz*. 2005;24(3):973. doi:10.20506/rst.24.3.1627
4. McVey DS, Kennedy M, Chengappa MM. *Veterinary Microbiology*. 3rd ed. USA: John Wiley & Sons, Inc.; 2013.

5. Bharti AR, Nally JE, Ricaldi JN, et al. Leptospirosis: a zoonotic disease of global importance. *Lancet Infect Dis*. 2003;3(12):757–771. doi:10.1016/s1473-3099(03)00830-2
6. Costa F, Hagan JE, Calcagno J, et al. Global morbidity and mortality of leptospirosis: a systematic review. *PLoS Negl Trop Dis*. 2015;9(9):e0003898. doi:10.1371/journal.pntd.0003898
7. Lavinsky MO, Abou Said R, Strenzel GMR, Langoni H. Seroprevalence of anti-*Leptospira* spp. antibodies in dogs in Bahia, Brazil. *Prev Vet Med*. 2012;106(1):79–84. doi:10.1016/j.prevetmed.2012.03.015
8. Levett PN. Leptospirosis. *Clin Microbiol Rev*. 2001;14(2):296. doi:10.1128/CMR.14.2.296-326.2001
9. de Vries SG, Visser BJ, Nagel IM, Goris MGA, Hartskeerl RA, Grobusch MP. Leptospirosis in Sub-Saharan Africa: a systematic review. *Int J Infect Dis*. 2014;28:47–64. doi:10.1016/j.ijid.2014.06.013
10. Markey B, Leonard F, Archambault M, Cullinane A, Maguire D. *Clinical Veterinary Microbiology e-Book*. Elsevier Health Sciences; 2013.
11. López M, Vila A, Rodón J, Roura X. *Leptospira* seroprevalence in owned dogs from Spain. *Heliyon*. 2019;5(8):e02373. doi:10.1016/j.heliyon.2019.e02373
12. Schuller S, Arent Z, Gilmore C, Nally J. Prevalence of antileptospiral serum antibodies in dogs in Ireland. *Vet Rec*. 2015;177(5):126. doi:10.1136/vr.102916
13. Lunn KF Leptospirosis in dogs - generalized conditions - veterinary manual; 2020. Available from: <https://www.msdsvetmanual.com/generalized-conditions/leptospirosis/leptospirosis-in-dogs#:~:text=Dogs%20are%20the%20primary%20host,in%20tissues%2C%20blood%20or%20urine>. Accessed August 18, 2020.
14. Alton GD, Berke O, Reid-Smith R, Ojkic D, Prescott JF. Increase in seroprevalence of canine leptospirosis and its risk factors, Ontario 1998–2006. *Can J Vet Res*. 2009;73(3):167–175.
15. Ramamoorthy S, Woldemeskel M, Ligett A, Snider R, Cobb R, Rajeev S. *Brucella suis* infection in dogs, Georgia, USA. *Emerg Infect Dis*. 2011;17(12):2386–2387. doi:10.3201/eid1712.111127
16. Akhvediani T, Bautista CT, Garuchava N, et al. Epidemiological and clinical features of brucellosis in the country of Georgia. *PLoS One*. 2017;12(1):e0170376. doi:10.1371/journal.pone.0170376
17. Pappas G, Papadimitriou P, Akritidis N, Christou L, Tsianos E. The new global map of human brucellosis. *Lancet Infect Dis*. 2006;6:91–99. doi:10.1016/S1473-3099(06)70382-6
18. Seleem MN, Boyle SM, Sriranganathan N. Brucellosis: a re-emerging zoonosis. *Vet Microbiol*. 2010;140(3–4):392–398. doi:10.1016/j.vetmic.2009.06.021
19. Whatmore AM. Current understanding of the genetic diversity of *Brucella*, an expanding genus of zoonotic pathogens. *Infect Genet Evol*. 2009;9(6):1168–1184. doi:10.1016/j.meegid.2009.07.001
20. Hensel ME, Negron M, Arenas-Gamboa AM. Brucellosis in dogs and public health risk. *Emerg Infect Dis*. 2018;24(8):1401–1406. doi:10.3201/eid2408.171171
21. Nomura A, Imaoka K, Imanishi H, et al. Human *Brucella canis* infections diagnosed by blood culture. *Emerg Infect Dis*. 2010;16(7):1183–1185. doi:10.3201/eid1607.090209
22. Lopes LB, Nicolino R, Haddad JPA. Brucellosis-risk factors and prevalence: a review. *Open Vet J*. 2010;4(1):72–84. doi:10.2174/1874318801004010072
23. Pujol M, Castillo F, Alvarez C, et al. Variability in the response of canine and human dendritic cells stimulated with *Brucella canis*. *Vet Res*. 2017;48(1):72. doi:10.1186/s13567-017-0476-8
24. Sayan M, Erdenlig S, Stack J, et al. A serological diagnostic survey for *Brucella canis* infection in Turkish patients with Brucellosis-like symptoms. *Jpn J Infect Dis*. 2011;64(6):516–519.
25. Lucero N, Corazza R, Almuzara M, et al. Human *Brucella canis* outbreak linked to infection in dogs. *Epidemiol Infect*. 2010;138(2):280–285. doi:10.1017/S0950268809990525
26. Ayoola MC, Ogugua AJ, Akinseye VO, et al. Sero-epidemiological survey and risk factors associated with brucellosis in dogs in south-western Nigeria. *Pan Afr Med J*. 2016;23(1):1–8. doi:10.11604/pamj.2016.23.1.8442
27. Wanke M. Canine brucellosis. *Anim Reprod Sci*. 2004;82:195–207. doi:10.1016/j.anireprosci.2004.05.005
28. Moch RW, Ebner EE, Barsoum LS, Botros BA. Leptospirosis in Ethiopia: a serological survey in domestic and wild animals. *J Trop Med Hyg*. 1975;78(2):38–42.
29. Yimer E, Koopman S, Messele T, et al. Human leptospirosis, in Ethiopia: a pilot study in Wonji. *Ethiop J Health Dev*. 2004;18(1):01–06.
30. Tsegay K, Potts A, Aklilu N, Lötter C, Gummow B. Circulating serovars of *Leptospira* in cart horses of central and southern Ethiopia and associated risk factors. *Prev Vet Med*. 2016;125:106–115. doi:10.1016/j.prevetmed.2016.01.009
31. Megersa B, Biffa D, Niguse F, Rufael T, Asmare K, Skjerve E. Cattle brucellosis in traditional livestock husbandry practice in Southern and Eastern Ethiopia, and its zoonotic implication. *Acta Vet Scand*. 2011;53(1):24. doi:10.1186/1751-0147-53-24
32. Geresu MA, Ameni G, Tuli G, Arenas A, Kassa GM. Seropositivity and risk factors for *Brucella* in dairy cows in Asella and Bishoftu towns, Oromia Regional State, Ethiopia. *Afr J Microbiol Res*. 2016;10(7):203–213. doi:10.5897/AJMR2015.7707
33. Edao BM, Ameni G, Assefa Z, Berg S, Whatmore AM, Wood JL. Brucellosis in ruminants and pastoralists in Borena, Southern Ethiopia. *PLoS Negl Trop Dis*. 2020;14(7):e0008461. doi:10.1371/journal.pntd.0008461
34. CSA. Population and Housing Census of Ethiopia; 2007.
35. Chappuis G. Neonatal immunity and immunisation in early age: lessons from veterinary medicine. *Vaccine*. 1998;16(14–15):1468–1472. doi:10.1016/S0264-410X(98)00110-8
36. Thrusfield M. *Veterinary Epidemiology; Describing Disease Occurrence*. 3rd ed. Blackwell Publishing; 2005.
37. Kiflu B, Alemayehu H, Abdurahaman M, Negash Y, Eguale T. *Salmonella* serotypes and their antimicrobial susceptibility in apparently healthy dogs in Addis Ababa, Ethiopia. *BMC Vet Res*. 2017;13(1):134. doi:10.1186/s12917-017-1055-y
38. *Statistical Software* [computer program]. Version 14.2. USA: StataCorp; 2018.
39. Dhliwayo S, Matope G, Dutlow K, Pfukenyi DM, Marabini L. Seroprevalence of leptospirosis in dogs in urban Harare and selected rural communities in Zimbabwe. *Onderstepoort J Vet Res*. 2012;79(1):1–6. doi:10.4102/ojvr.v79i1.447
40. Spangler D, Kish D, Beigel B, et al. Leptospiral shedding and seropositivity in shelter dogs in the Cumberland Gap Region of Southeastern Appalachia. *PLoS One*. 2020;15(1):e0228038. doi:10.1371/journal.pone.0228038
41. Lau S, Wong J, Khor K, et al. Seroprevalence of leptospirosis in working dogs. *Top Companion Anim Med*. 2017;32(4):121–125. doi:10.1053/j.tcam.2017.12.001
42. Altheimer K, Jongwattapanis P, Luengyosuechakul S, et al. *Leptospira* infection and shedding in dogs in Thailand. *BMC Vet Res*. 2020;16(1):1–13. doi:10.1186/s12917-020-2230-0
43. Aslantaş Ö, Özdemir V, Kiliç S, Babür C. Seroepidemiology of leptospirosis, toxoplasmosis, and leishmaniasis among dogs in Ankara, Turkey. *Vet Parasitol*. 2005;129(3–4):187–191. doi:10.1016/j.vetpar.2004.11.037
44. Delaude A, Rodriguez-Campos S, Dreyfus A, et al. Canine leptospirosis in Switzerland-A prospective cross-sectional study examining seroprevalence, risk factors and urinary shedding of pathogenic leptospires. *Prev Vet Med*. 2017;141:48–60. doi:10.1016/j.prevetmed.2017.04.008
45. Pratt N, Conan A, Rajeev S. *Leptospira* seroprevalence in domestic dogs and cats on the Caribbean Island of Saint Kitts. *Vet Med Int*. 2017;2017:1–6. doi:10.1155/2017/5904757

46. Meeyam T, Tablerk P, Petchanok B, Pichpol D, Padungtod P. Seroprevalence and risk factors associated with leptospirosis in dogs. *Southeast Asian J Trop Med Public Health*. 2006;37(1):148.
47. Mosalanezhad B, Ghorbanpour NM, Avizeh R, Mohammadian N. A serological survey on *Brucella canis* in companion dogs in Ahvaz. *Iran J Vet Res*. 2009;10(4):383–386.
48. Öncel T, Akan M, Sareyyüpoğlu B, Tel OY, Çiftci A. Seroprevalence of *Brucella canis* infection of dogs in two provinces in Turkey. *Turk J Vet Anim Sci*. 2005;29(3):779–783.
49. Hubbard K, Wang M, Smith DR. Seroprevalence of brucellosis in Mississippi shelter dogs. *Prev Vet Med*. 2018;159:82–86. doi:10.1016/j.prevetmed.2018.09.002
50. Kei N, Shingo S, Hidenori K, Soichi M. Seroepidemiological survey of *Brucella canis* infection in dogs in Japan. *Jpn J Vet Res*. 2020;68(2):129–132. doi:10.14943/jjvr.68.2.129
51. Upadhyay A, Singh P, Nagpal A. Epidemiology of brucellosis in India: a review. *Pantnagar J Res*. 2019;17(3):199–205.
52. Grace D, Mutua F, Ochungo P, et al. Mapping of poverty and likely zoonoses hotspots. Report to the UK Department for International Development; 2012.

Veterinary Medicine: Research and Reports

Dovepress

Publish your work in this journal

Veterinary Medicine: Research and Reports is an international, peer-reviewed, open access journal publishing original research, case reports, editorials, reviews and commentaries on all areas of veterinary medicine. The manuscript management system is completely online

and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/veterinary-medicine-research-and-reports-journal>