Predictors of Hospitalization Among Older Adults with COVID-19 in Saudi Arabia: A Cross-Sectional Study of a Nationally Representative Sample

This article was published in the following Dove Press journal:
Risk Management and Healthcare Policy

Fahad Alamri^{1,*}
Yousef Alsafayan¹
Yazed AlRuthia^{1,2,3,*}
Ahmed Alahmari¹
Yasir Almuzaini¹
Fouad Abo Gazalah⁴
Faten Alradini⁵
Tareef Alaama^{1,6}
Anas Khan^{1,7}

¹Global Center for Mass Gatherings Medicine, Ministry of Health, Riyadh, Saudi Arabia; ²Department of Clinical Pharmacy, College of Pharmacy, King Saud University, Riyadh, Saudi Arabia; ³Pharmacoeconomics Research Unit, College of Pharmacy, King Saud University, Riyadh, Saudi Arabia; ⁴General Director of Health Centers Affairs, Ministry of Health, Riyadh, Saudi Arabia; ⁵Department of Family Medicine, Faculty of Medicine, Princess Nourah Bint Abdulrahman University, Riyadh, Saudi Arabia; ⁶Deputyship of Therapeutic Services, Ministry of Health, Riyadh, Saudi Arabia; ⁷Department of Emergency Medicine, College of Medicine, King Saud University, Riyadh, Saudi Arabia

*These authors contributed equally to this work

Background: The aim of this study was to identify the predictors of hospitalization in older (≥ 60 years) patients with coronavirus disease-19 (COVID-19) in Saudi Arabia.

Methods: Patients were randomly selected from a COVID-19 database maintained by the Ministry of Health, Saudi Arabia. All patients were aged ≥ 60 years, had reverse transcriptase-polymerase chain reaction (RT-PCR)-confirmed COVID-19, and were registered in the database during March 2020 to July 2020. Medical and sociodemographic characteristics were retrieved from the database. Additional data were collected by telephone interviews conducted by trained health professionals. Descriptive statistics and multiple logistic regression analyses were used to analyze the relationship between patient characteristics and the risk of hospitalization.

Results: Of the 613 included patients (51.1% females), more than half (57.3%) were between 60 to 69 years of age, and 53% (324/613) had been hospitalized. The independent predictors of hospitalization included age ≥ 65 years (OR = 2.35, 95% CI: 1.66–3.33, $P < 0.001$), having more than one comorbidity (OR = 1.55, 95% CI: 1.09–2.20, $P = 0.01$), diabetes mellitus (OR = 1.52, 95% CI: 1.09–2.11, $P = 0.01$), hypertension (OR = 1.40, 95% CI: 1.007–1.97, $P = 0.04$), chronic kidney disease (OR = 3.87, 95% CI: 1.41–10.58, $P = 0.008$), and history of hospital admission within the preceding year (OR = 1.69, 95% CI: 1.11–2.55, $P = 0.013$). Risk of hospitalization was lower in males (OR = 0.65, 95% CI: 0.43–0.90, $P = 0.01$) and in patients co-living with health care workers (OR = 0.64, 95% CI: 0.43–0.96, $P = 0.03$).

Conclusion: Factors associated with higher risk of COVID-19-associated hospitalization should be used in prioritizing older adults' admission. Future studies with more robust designs should be conducted to examine the risk of COVID-19-associated illness severity and mortality.

Keywords: COVID-19, older adults, hospitalization, diabetes, hypertension, chronic kidney disease, Saudi Arabia

Introduction

The coronavirus disease-2019 (COVID-19) pandemic has ravaged the world for more than a year and caused more than two million deaths.^{1,2} Multiple studies have shown that older adults with COVID-19 have high risk of mortality, mainly due to the presence of multiple comorbidities.^{3–7} In addition, many studies have shown that older patients are more likely to need intensive care unit (ICU) admission for management of respiratory complications.^{8–11} Among the elderly, heart failure, peripheral artery disease, C-reactive protein

Correspondence: Yazed AlRuthia
Tel +996 114677483
Fax +966 114677480
Email yazeed@ksu.edu.sa

Fahad Alamri
Email faabalamri@moh.gov.sa

level, need for oxygen support, respiratory rate, presence of lung crackles at presentation, shorter interval from onset of symptoms to hospitalization, length of hospital stay, and male sex have been shown to be associated with higher risk of mortality;¹² however, old age remains an independent risk factor for both hospitalization and mortality after controlling for a myriad of factors.^{13–16}

The Kingdom of Saudi Arabia reported its first case of COVID-19 on March 2, 2020, and by November 17, 2020, a total of 353,918 confirmed COVID-19 cases had been reported.^{17,18} As in other countries, it is highly likely that older Saudi patients with COVID-19 — especially those with comorbidities such as, diabetes, hypertension, chronic kidney disease (CKD), and bronchial asthma — have higher risk of hospitalization and death.^{12–16,19–21} However, no study has so far explored the predictors of hospitalization in this vulnerable population in Saudi Arabia. Therefore, in this study we aimed to determine the sociodemographic and clinical predictors of hospitalization in older adults with COVID-19 in Saudi Arabia. Better awareness regarding the risk factors for hospitalization will be useful for health policy planning and will enable more efficient allocation of limited resources.

Methods

Study Population and Data Collection

The study population was selected from the COVID-19 database maintained by the Ministry of Health (MOH) Saudi Arabia.¹⁸ This database, which is part of the Health Electronic Surveillance Network (HESN), contains the details of all reverse transcriptase polymerase chain reaction (RT-PCR)-confirmed cases of COVID-19 in Saudi Arabia. A total of 10,368 elderly (≥ 60 years) COVID-19 patients were registered in the database from March 2020 to June 2020. We performed systematic random sampling and selected every ninth patient for this cross-sectional study. The 1152 patients thus selected were contacted over the telephone and asked about their willingness to participate in the study. Only those who were willing to sign written informed consent and provide all requested information were included in the study.

Medical and sociodemographic data of the enrolled patients including proof of hospitalization were retrieved from the database. Further, seven trained

MOH employees (one physician, five nurses, and one administrative assistant) contacted each patient over the telephone to collect additional data regarding marital status, educational level, smoking history, whether co-living with a healthcare worker (HCW), whether hospitalized for COVID-19, number and type of comorbidities, history of hospital admission for other diseases in the preceding year, symptoms, and so on. Comorbidities were classified according to the International Classification of Diseases-10 (ICD-10).²² The data collection took place between July and September 2020. The collected information was systematically entered into an electronic database that was regularly reviewed by the first and third authors to ensure consistency.

The ethical principles of the Declaration of Helsinki were adhered to during collection, handling, and storage of data, and all care was taken to protect patient confidentiality. This study was approved by the Ethics Review Board

Figure 1 Flowchart showing patient recruitment.

Committee of the Central Ministry of Health, Riyadh, Saudi Arabia (No: 20–156M).

Statistical Analysis

Assuming a response rate of 50%, a minimum sample size of 371 would be required to detect a significant difference between groups with a margin of error of 5%.

However, to ensure sampling adequacy, a sample size of at least 415 patients was planned. Descriptive statistics were used to analyze baseline sociodemographic and clinical characteristics. Categorical variables were summarized as counts and percentages and compared between groups using the Chi-squared or Fisher's exact test, as appropriate. Numerical data were checked for

Table I Sociodemographic Characteristics of the COVID-19 Patients

Characteristics	Total N = 613	Hospitalized n = 324	Nonhospitalized n = 289	P
Age (years)				
60–69	351 (57.3%)	157 (48.5%)	194 (67.1%)	<0.001
70–79	180 (29.4%)	110 (34.0%)	70 (24.2%)	
80–89	62 (10.1%)	44 (13.6%)	18 (6.2%)	
≥90	20 (3.3%)	13 (4.0%)	7 (2.4%)	
Sex				
Male	300 (48.9%)	144 (44.4%)	156 (54.0%)	0.018
Female	313 (51.1%)	180 (55.6%)	133 (46.0%)	
Co-living with HCW				
Yes	130 (21.2%)	58 (17.9%)	72 (24.9%)	0.034
No	483 (78.7%)	266 (82.1%)	217 (75.1%)	
Smoking status				
Former smoker	94 (15.3%)	59 (18.2%)	35 (12.1%)	0.056
Current smoker	21 (3.4%)	7 (2.2%)	14 (4.8%)	
Never smoker	495 (80.8%)	257 (79.3%)	238 (82.4%)	
Missing	3 (0.5%)	1 (0.3%)	2 (0.7%)	
Region of residence				
Riyadh	209 (34.1%)	99 (30.6%)	110 (38.1%)	0.139
Eastern region	141 (23.0%)	66 (20.4%)	74 (25.6%)	
Makkah	160 (26.1)	94 (29.0)	66 (22.8)	
Madinah	41 (6.7%)	24 (7.4%)	17 (5.9%)	
Al-Qassim	12 (2.0%)	8 (2.5%)	4 (1.4%)	
Aseer	16 (2.61)	11 (3.4)	5 (1.73)	
Najran	8 (1.3%)	6 (1.9%)	2 (0.7%)	
Al-Baha	5 (0.8%)	4 (1.2%)	1 (0.3%)	
Tabuk	4 (0.7%)	2 (0.6%)	2 (0.7%)	
Hail	4 (0.7%)	3 (0.9%)	1 (0.3%)	
Northern borders	1 (0.2%)	-	1 (0.3%)	

Table 2 Clinical Characteristics of the Patients

Characteristics	Total N = 613	Hospitalized n = 324	Nonhospitalized n = 289	P
Comorbidity				
Any condition	506 (82.5%)	278 (85.8%)	228 (78.9%)	0.024
More than one condition	212 (34.6%)	131 (40.4%)	81 (28.0%)	0.001
Flu vaccination				
Yes	83 (13.5%)	40 (12.3%)	43 (14.9%)	0.537
No	439 (71.6%)	238 (73.5%)	201 (69.6%)	
Not sure	91 (14.8%)	46 (14.2%)	45 (15.6%)	
Hospital admission within the past year				
Yes	137 (22.3%)	91 (28.1%)	46 (15.9%)	<0.001
No	476 (77.7%)	233 (71.9%)	243 (84.1%)	
Symptoms				
Fever	426 (69.5%)	242 (74.7%)	184 (63.7%)	0.005
Cough	348 (56.8%)	205 (63.3%)	143 (49.5%)	0.002
Sore throat	232 (37.8%)	124 (38.3%)	108 (37.4%)	0.518
Runny nose	120 (19.6%)	61 (18.8%)	59 (20.4%)	0.450
Shortness of breath	269 (43.9%)	176 (54.3%)	93 (32.2%)	<0.001
Vomiting	96 (15.7%)	64 (19.8%)	32 (11.1%)	0.012
Nausea	146 (23.8%)	88 (27.2%)	58 (20.1%)	0.065
Dizziness	189 (30.8%)	123 (38.0%)	66 (22.8%)	<0.001
Headache	211 (34.4%)	103 (31.8%)	108 (37.4%)	0.147
Loss of smell or taste	272 (44.4%)	137 (42.3%)	135 (46.7%)	0.007
Change in taste	28 (4.6%)	14 (4.3%)	14 (4.8%)	0.757
Mouth ulcers	6 (1.0%)	1 (0.3%)	5 (1.7%)	0.105
Diarrhea	205 (33.4%)	121 (37.3%)	84 (29.1%)	0.056
Tiredness or general fatigue	478 (78.0%)	276 (85.2%)	202 (69.9%)	<0.001
Weight loss	145 (23.7%)	83 (25.6%)	62 (21.5%)	0.226
Weight gain	5 (0.8%)	3 (0.9%)	2 (0.7%)	>0.999
Eye redness/itching	21 (3.4%)	9 (2.8%)	12 (4.2%)	0.350
Back pain	101 (16.5%)	52 (16.0%)	49 (17.0%)	0.763
Leg pain	71 (11.6%)	39 (12.0%)	32 (11.1%)	0.710
Joint pain	158 (25.8%)	79 (24.4%)	79 (27.3%)	0.404
Abdominal pain	50 (8.2%)	31 (9.6%)	19 (6.6%)	0.176
Muscle spasms	2 (0.3%)	2 (0.6%)	–	0.501

(Continued)

Table 2 (Continued).

Characteristics	Total N = 613	Hospitalized n = 324	Nonhospitalized n = 289	P
Swollen ankles or feet	3 (0.5%)	1 (0.3%)	2 (0.7%)	0.604
Chills	15 (2.4%)	10 (3.1%)	5 (1.7%)	0.278
Loss of appetite	279 (45.5%)	172 (53.1%)	107 (37.0%)	<0.001
Hemoptysis	2 (0.3%)	2 (0.6%)	–	0.501
Altered smell of urine	3 (0.5%)	2 (0.6%)	1 (0.3%)	>0.999
Dark urine discoloration	4 (0.7%)	2 (0.6%)	2 (0.7%)	>0.999
Difficulty urinating	12 (2.0%)	7 (2.2%)	5 (1.7%)	0.701
Fainting	5 (0.8%)	4 (1.2%)	1 (0.3%)	0.377
Itching	5 (0.8%)	2 (0.6%)	3 (1.0%)	0.671
Rash	3 (0.5%)	1 (0.3%)	2 (0.6%)	0.604
Skin ulcers	2 (0.3%)	2 (0.6%)	–	0.501
Numbness or tingling	20 (3.3%)	10 (3.1%)	10 (3.5%)	0.795
Presence of symptoms				
Asymptomatic	50 (8.2%)	17 (5.2%)	33 (11.4%)	0.005
Symptomatic	563 (91.8%)	307 (94.8%)	256 (88.6%)	

normality using the Shapiro–Wilk test and summarized as medians (and interquartile ranges; IQR). For the median length of stay (LOS), the Mann–Whitney *U*-test and Kruskal–Wallis tests were used to compare differences between two and more than two independent variables, respectively. Simple and multiple logistic regression analyses were conducted to identify the independent predictors of hospitalization, such as, age groups, gender, co-living with an HCW, smoking status, presence of any health condition, number of comorbidities, specific chronic health conditions, such as diabetes, hypertension, CKD, heart disease, and history of admission within the past year; the odds ratios (ORs) and 95% confidence intervals (CIs) were calculated. The significance level was at $\alpha = 0.05$. Statistical analysis was performed using SPSS 24 (IBM Corp., Armonk, NY, USA).

Results

Sociodemographic Characteristics

Of the 1152 patients initially selected from the database, only 613 (53.21%) consented to participate and were

interviewed. These patients were divided into two groups: a hospitalized group (324/613, 52.85%) and a nonhospitalized group (289/613, 47.1%) (Figure 1). Table 1 summarizes the sociodemographic characteristics of the patients.

Most of the patients (531/613, 86.7%) were between 60 and 79 years of age. The proportion of patients aged ≥ 70 years was significantly higher in the hospitalized group than in the nonhospitalized group: 51.5% (167/324) vs 32.9% (95/289), $P < 0.001$. The proportion of female patients was significantly higher in the hospitalized group than in the nonhospitalized group (55.6% vs 46.0%, $P = 0.018$). Overall, 21% (130/613) of patients reported co-living with HCWs; the proportion was significantly lower in the hospitalized group than in the nonhospitalized group: 17.9% (58/324) vs 24.9% (72/289), $P = 0.034$.

Clinical Characteristics

The majority of patients hospitalized for COVID-19 (270/324, 83.33%) were admitted to the hospital wards; only 54/324 (16.67%) patients were admitted to ICUs. Among the patients admitted to ICUs, the

majority (45/54, 83.3%) were ≥ 65 years old. More than half of the patients admitted to ICUs (30/54, 55.6%) needed mechanical ventilation, and among these the majority (83.3%, 25/30) were ≥ 65 years. Table 2 lists the clinical characteristics of the patients.

Comorbidities were significantly more common in hospitalized patients, with at least one comorbidity being present in 85.8% (278/324) of the patients in the hospitalized group vs 78.9% (228/289) of the patients in the nonhospitalized group ($P = 0.024$); moreover,

Figure 2 Rates of chronic health conditions among hospitalized and nonhospitalized patients.

Figure 3 Rates of commonly reported symptoms in hospitalized and nonhospitalized patients.

a significantly larger proportion of hospitalized patients had more than one comorbidity: 40.4% of hospitalized patients vs 28% of nonhospitalized patients ($P = 0.001$). Diabetes mellitus, hypertension, heart disease, and CKD were all significantly more prevalent in hospitalized patients than in nonhospitalized patients (all $P < 0.05$; Figure 2). Only 13.5% (83/613) of patients reported receiving flu vaccination prior to COVID-19 infection; there was no significant difference between hospitalized and nonhospitalized patients. History of hospital admission within the past year was reported by 22.3% (137/613) of patients; the proportion was significantly higher in the hospitalized group than in the nonhospitalized group: 28.1% (91/324) vs 15.9% (46/289), $P < 0.001$.

COVID-19-related symptoms (such as fatigue, fever, cough, loss of appetite, shortness of breath, dizziness, and vomiting) were significantly more common in the hospitalized group than in the nonhospitalized group ($P < 0.05$; Figure 3); however, loss of smell/taste was more common among nonhospitalized patients ($P = 0.007$).

In the hospitalized group, median length of hospital stay was 10 days (IQR, 11.5 days). The median length of stay was significantly longer for those with heart disease than for those without heart disease (13.5 days vs 10.5 days, $P = 0.04$), and for patients with CKD than for those without CKD (19 days vs 10 days, $P = 0.01$). Table 3 shows the median length of hospital stay across some patients' characteristics. Herbal and food supplements were largely utilized among patients in the study. Honey, ginger, lemon, and vitamin C were used by at least 40% of the patients, as shown in Figure 4.

Predictors of Hospitalization

Table 4 and Figure 5 show how different sociodemographic and clinical variables were associated with higher risk of hospitalization. Patients aged ≥ 65 years had more than two times higher risk of hospitalization than patients aged < 65 years (OR = 2.35, 95% CI: 1.66–3.33, $P < 0.001$). Male patients had 34.7% lower risk of hospitalization than female patients (OR = 0.65, 95% CI: 0.47–0.90, $P = 0.01$). Patients living with HCWs had 35.5% lower risk of hospitalization than patients not living with HCWs (OR = 0.64, 95% CI: 0.43–0.96, $P = 0.03$). Patients with more than one comorbidity had 55.8% higher risk of hospitalization than patients with only one or no comorbidity (OR = 1.55, 95% CI: 1.09–2.20, $P = 0.01$). Patients

Table 3 Total Length of Stay in the Hospital (Ward/ICU) Stratified by Some Baseline Characteristics

Characteristics	Median (IQR)	P
Total	10 (11.5)	–
Age (years)		
<65	10 (12)	0.979
≥65	11 (11)	
Age-group		
60–69	10 (12.5)	0.957
70–79	12 (10.0)	
80–89	10 (9.0)	
≥90	11 (9.5)	
Sex		
Male	10 (14.0)	0.784
Female	11 (9.0)	
Co-living with HCW		
Yes	13 (12.0)	0.164
No	10 (10.0)	
Comorbidity		
Yes	11 (13.0)	0.157
No	9.0 (7.0)	
More than one comorbidity		
Yes	12.5 (12.0)	0.087
No	10 (11.25)	
Diabetes		
Yes	11 (13)	0.545
No	10 (8.0)	
Hypertension		
Yes	11 (12.0)	0.540
No	10 (8.5)	
Heart disease		
Yes	13.5 (10.5)	0.044
No	10 (12.0)	
Chronic kidney disease		
Yes	19 (17.5)	0.010
No	10 (10)	

Abbreviations: IQR, interquartile range; HCW, healthcare worker.

Figure 4 The rates of herbal and food supplements' utilization among the study sample.

with diabetes had 52.1% higher risk of hospitalization than patients without diabetes (OR = 1.52, 95% CI: 1.09–2.11, $P = 0.01$). Patients with hypertension had 40.9% higher risk of hospitalization than patients without hypertension (OR = 1.40, 95% CI: 1.007–1.97, $P = 0.04$). Patients with CKD had almost four times higher risk of hospitalization than patients without CKD (OR = 3.87, 95% CI: 1.41–10.58, $P = 0.008$). Patients with history of admission to hospital in the past year had 69.2% higher risk of hospitalization than patients without history of hospital admission within the past year (OR = 1.69, 95% CI: 1.11–2.55, $P = 0.013$). Patients with heart disease had 63.5% higher risk of hospitalization than patients without heart disease, but it was not statistically significant (OR = 1.45, 95% CI: 0.95–2.23, $P = 0.08$).

Discussion

This study aimed to identify the factors associated with risk of hospitalization among adults aged 60 years and above in Saudi Arabia. We found age ≥ 65 years to be an

independent predictor of hospitalization. This is consistent with several previous studies, including a recently published study from Denmark that found older age to be a significant predictor of hospitalization and mortality after controlling for sex and a number of comorbidities.¹⁴ Consistent with earlier studies, we found that the presence of multiple comorbidities — which is common in this age-group — increased the risk of hospitalization.^{5,14–16,19,20}

Specific health conditions that were associated with higher risk of hospitalization included diabetes, hypertension, and CKD. A recently published meta-analysis of 47 studies (with a total sample size of 13,268 COVID-19 patients) found diabetes to be significantly associated with a higher risk of severe disease and mortality.²³ This is very concerning, given the high prevalence of diabetes in the Saudi population in general and in older adults in particular, where the prevalence is as high as 20%.²⁴ Although hypertension has been found to be associated with worse clinical

Table 4 Unadjusted Odds Ratios (ORs) for the Risk of Hospitalization of Selected Patients' Characteristics

Characteristics	Unadjusted OR (95% CI)	P
Age (≥65 years)	2.287 (1.623–3.221)	<0.001
Age-group		
60–69 (ref)	-	-
70–79	1.922 (1.332–2.772)	<0.001
80–89	2.989 (1.661–5.380)	<0.001
≥90	2.271 (0.885–5.380)	0.088
Sex (male)	0.682 (0.496–0.938)	0.019
Co-living with HCW (Yes)	0.657 (0.445–0.970)	0.035
Smoking status		
Never (ref)	-	-
Current smoker	0.463 (0.184–1.167)	0.103
Former smoker	0.463 (0.042–5.140)	0.531
Comorbidity (Yes)	1.617 (1.061–2.463)	0.025
More than one comorbidity (Yes)	1.464 (1.156–1.855)	0.002
Diabetes mellitus (Yes)	1.631 (1.184–2.248)	0.003
Hypertension (Yes)	1.574 (1.140–2.175)	0.006
Heart disease (Yes)	1.635 (1.078–2.479)	0.021
Chronic kidney disease (Yes)	3.937 (1.465–10.580)	0.007
Hospital admission within the past year (Yes)	2.063 (1.386–3.070)	<0.001

Abbreviation: HCW, healthcare worker.

outcomes in COVID-19 patients, no study has so far shown hypertension to be an independent risk factor for severe COVID-19 illness or mortality.²⁵ In the present study, we found that hypertension and CKD were independent predictors of hospitalization and longer hospital stay. Multiple previous studies have reported significantly increased risk of mortality among COVID-19 patients with compromised renal function.^{14,26} Williamson et al analyzed a large database in England and found that patients with estimated glomerular filtration rate <30/min/1.73 m² and those on dialysis had significantly higher risk of mortality.²⁶ This is alarming because there is a high prevalence of CKD among older adults in Saudi Arabia.²⁷ Cardiovascular disease (including coronary artery disease and heart failure) has been previously shown to be associated

with higher risk of hospitalization and mortality in patients with COVID-19.^{14,26} In the present study, however, although heart disease was associated with high risk of hospitalization and prolonged hospital stay, it was not an independent predictor of these outcomes; the failure to demonstrate a significant association may have been because of the relatively small number of patients with heart disease in our sample.^{14,26} In a previous study, severe asthma (ie, with history of recent use of oral corticosteroid) was a risk factor for mortality in COVID-19 patients.²⁶ In our study, bronchial asthma was not significantly associated with risk of hospitalization; however, we did not stratify patients based on history of recent use of oral corticosteroids.

Interestingly, in the present study, male sex was associated with lower risk of hospitalization. While most previous studies have found that male patients have significantly higher risk of hospitalization, ICU admission, and death,^{14,26,28–30} a recently published study from Saudi Arabia found no significant difference in mortality between male and female patients hospitalized for COVID-19.³¹ Meanwhile, it should be emphasized that the lower risk of hospitalization that we found among older adult males does not necessarily mean lower mortality, as this study did not follow-up patients post-discharge.

Another interesting finding was the lower risk of hospitalization among patients co-living with HCWs. This is likely due to the better, and more informed care provided to family members by HCWs.³² In our sample, history of hospital admission within the past year was an independent predictor of hospitalization for COVID-19. This is in line with a previous Danish study on a large cohort of COVID-19 patients.¹⁴ Current smoking was not associated with risk of hospitalization, which is in accordance with the Williamson et al study.²⁶

COVID-19-related symptoms such as fatigue, fever, cough, sore throat, shortness of breath, vomiting, dizziness, and loss of appetite were common in all patients, though they were relatively more common among hospitalized patients. This finding is consistent with two previously published studies from Saudi Arabia.^{28,33}

This study has many limitations. First, the study only included older adults who survived COVID-19 infection since complete data were not available on patients who

Figure 5 Adjusted odds ratios (ORs) with 95% confidence intervals (CIs) of some patients' characteristics.

died. Second, the study findings are subject to information and documentation bias since the data were retrieved from an electronic database or via telephone - interviews.³⁴ Moreover, the cognitive strain on older adults asked multiple questions over the telephone may have resulted in an acquiescence bias.³⁵ Third, nonresponse bias cannot be excluded, since not all selected patients agreed to participate in the study, which can influence the results significantly.³⁶ Finally, this was a cross-sectional study and so no causal relationship between any of the examined clinical/sociodemographic variables and the risk of hospitalization can be established.

Conclusion

The factors associated with high risk of hospitalization for COVID-19 in Saudi Arabia include older age; presence of multiple comorbidities; presence of diabetes, hypertension, or CKD; female sex; and hospital admission within the previous year. Further well-designed studies are necessary to confirm our findings. Future studies should examine the factors associated with higher risk of hospitalization and mortality among elderly patients with COVID-19 in Saudi Arabia using more robust study designs, which will help in efficient allocation of health resources.

Abbreviations

CI, confidence interval; CKD, chronic kidney disease; COVID-19, coronavirus disease-19; HESN, Health Electronic Surveillance Network; HCW, healthcare worker; ICD-10, international classification of diseases-10; ICU, intensive care unit; IQR, interquartile range; MOH, Ministry of Health; OR, odds ratio; RT-PCR, reverse transcriptase polymerase chain reaction.

Data Sharing Statement

Study data are de-identified and can be made available upon request from the corresponding authors.

Acknowledgments

The authors acknowledge the help and support of data collectors at the Saudi Ministry of Health.

The first and third authors contributed equally to this study and are both corresponding authors for this paper.

Disclosure

The authors report no conflicts of interest in this work.

References

1. World Health Organization. WHO Coronavirus Disease (COVID-19) Dashboard. Available from: <https://covid19.who.int/>. Accessed 24 October, 2020.

2. WHO. Director-General's opening remarks at the media briefing on COVID-19; 2020. Available from: <https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19—11-march-2020>. accessed 25 October 2020.
3. Liu K, Chen Y, Lin R, Han K. Clinical features of COVID-19 in elderly patients: a comparison with young and middle-aged patients. *J Infect.* 2020;80(6):e14–e18. doi:10.1016/j.jinf.2020.03.005
4. Song J, Hu W, Yu Y, et al. A Comparison of clinical characteristics and outcomes in elderly and younger patients with COVID-19. *Med Sci Monit.* 2020;26:e925047. doi:10.12659/MSM.925047
5. Li P, Chen L, Liu Z, et al. Clinical features and short-term outcomes of elderly patients with COVID-19. *Int J Infect Dis.* 2020;97:245–250. doi:10.1016/j.ijid.2020.05.107
6. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet.* 2020;395(10229):1054–1062. doi:10.1016/S0140-6736
7. Grasselli G, Zangrillo A, Zanella A, et al. Baseline characteristics and outcomes of 1591 patients infected with SARS-CoV-2 admitted to ICUs of the Lombardy region, Italy. *JAMA.* 2020;323(16):1574–1581. doi:10.1001/jama.2020.5394
8. Niu S, Tian S, Lou J, et al. Clinical characteristics of older patients infected with COVID-19: a descriptive study. *Arch Gerontol Geriatr.* 2020;89:104058. doi:10.1016/j.archger.2020.104058
9. Wu C, Chen X, Cai Y, et al. Risk factors associated with acute respiratory distress syndrome and death in patients with coronavirus disease 2019 pneumonia in Wuhan, China. *JAMA Intern Med.* 2020;180(7):934–943. doi:10.1001/jamainternmed.2020.0994
10. Zou L, Dai L, Zhang Y, et al. Clinical characteristics and risk factors for disease severity and death in patients with coronavirus disease 2019 in Wuhan, China. *Front Med.* 2020;7:532. doi:10.3389/fmed.2020.00532
11. Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet.* 2020;395(10223):507–513. doi:10.1016/S0140-6736(20)30211-7
12. Mendes A, Serratrice C, Herrmann FR, et al. Predictors of in-hospital mortality in older patients with COVID-19: the COVIDAge study. *J Am Med Dir Assoc.* 2020;21(11):1546–1554. doi:10.1016/j.jamda.2020.09.014
13. Bialek S, Boundy E, Bowen V, COVID TC, Team R. Severe Outcomes Among Patients with Coronavirus Disease 2019 (COVID-19) — United States, February 12–March 16, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69(12):343–346. doi:10.15585/mmwr.mm6912e2
14. Reilev M, Kristensen KB, Pottegård A, et al. Characteristics and predictors of hospitalization and death in the first 11 122 cases with a positive RT-PCR test for SARS-CoV-2 in Denmark: a nationwide cohort. *Int J Epidemiol.* 2020;49(5):1468–1481. doi:10.1093/ije/dyaa140
15. Tehrani S, Killander A, Astrand P, Jakobsson J, Gille-Johnson P. Risk factors for death in adult COVID-19 patients: frailty predicts fatal outcome in older patients. *Int J Infect Dis.* 2021;102:415–421. doi:10.1016/j.ijid.2020.10.071
16. Palmieri L, Vanacore N, Donfrancesco C, et al. Clinical characteristics of hospitalized individuals dying with COVID-19 by age group in Italy. *J Gerontology.* 2020;75(9):1796–1800. doi:10.1093/geronola/glaa146
17. MOH News M.O.H. Reports First Case of Coronavirus Infection; 2020. Available from: <https://www.moh.gov.sa/en/Ministry/MediaCenter/News/Pages/News-2020-03-02-002.aspx>. Accessed 5 April 2020.
18. COVID-19 Daily Updates. Saudi Arabia: ministry of Health; 2020. Available from: <https://covid19.moh.gov.sa/>. accessed 14 Jun 2020.
19. Guo T, Shen Q, Guo W, et al. Clinical Characteristics of Elderly Patients with COVID-19 in Hunan Province, China: a Multicenter, Retrospective Study. *Gerontology.* 2020;29:1–9.
20. Medetalibeyoglu A, Senkal N, Kose M, et al. Older adults hospitalized with Covid-19: clinical characteristics and early outcomes from a single center in Istanbul, Turkey. *J Nutr Health Aging.* 2020;1.
21. Khoja AT, Aljawadi MH, Al-Shammari SA, et al. The health of Saudi older adults; results from the Saudi National Survey for elderly health (SNSEH) 2006–2015. *Saudi Pharm J.* 2018;26(2):292–300. doi:10.1016/j.jsps.2017.11.008
22. ICD-10 for Mortality and Morbidity Statistics. World Health Organization; 2019. Available from: <https://icd.who.int/browse10/2019/en>. Accessed 5 Sep 2020.
23. Varikasuvu SR, Dutt N, Thangappazham B, Varshney S. Diabetes and COVID-19: a pooled analysis related to disease severity and mortality. *Prim Care Diabetes.* 2021;15(1):24–27. doi:10.1016/j.pcd.2020.08.015
24. Whiting DR, Guariguata L, Weil C, Shaw J. IDF diabetes atlas: global estimates of the prevalence of diabetes for 2011 and 2030. *Diabetes Res Clin Pract.* 2011;94(3):311–321. doi:10.1016/j.diabres.2011.10.029
25. Huang S, Wang J, Liu F, et al. COVID-19 patients with hypertension have more severe disease: a multicenter retrospective observational study. *Hypertension Res.* 2020;1–8.
26. Williamson EJ, Walker AJ, Bhaskaran K, et al. Factors associated with COVID-19-related death using OpenSAFELY. *Nature.* 2020;584(7821):430–436. doi:10.1038/s41586-020-2521-4
27. Alsuwaidia AO, Farag YMK, Al Sayyari AA, et al. Epidemiology of chronic kidney disease in the Kingdom of Saudi Arabia (SEEK-Saudi investigators) - a pilot study. *Saudi J Kidney Dis Transplant.* 2010;21(6):1066.
28. Khan AA, Althunayyan SM, Alsafayan YM, Alotaibi RN, Mubarak AM. Risk factors associated with worse outcomes in COVID-19: a retrospective study in Saudi Arabia. *East Mediterr Health J.* 2020;26(11):1371–1380. doi:10.26719/emhj.20.130
29. Gebhard C, Regitz-Zagrosek V, Neuhauser HK, Morgan R, Klein SL. Impact of sex and gender on COVID-19 outcomes in Europe. *Biol Sex Differ.* 2020;11(1):1–3. doi:10.1186/s13293-020-00304-9
30. Mallapaty S. The coronavirus is most deadly if you are older and male-new data reveal the risks. *Nature.* 2020;16–17.
31. Khan AA, AlRuthia Y, Balkhi B, et al. Survival and estimation of direct medical costs of hospitalized COVID-19 patients in the kingdom of Saudi Arabia (Short Title: COVID-19 Survival and Cost in Saudi Arabia). *Int J Environ Res Public Health.* 2020;17(20):7458. doi:10.3390/ijerph17207458
32. Bianchetti A, Bellelli G, Guerini F, et al. Improving the care of older patients during the COVID-19 pandemic. *Aging Clin Exp Res.* 2020;32(9):1883–1888.
33. Alsafayan Y, Althunayyan S, Khan A, Hakawi A, Assiri A. Clinical characteristic of COVID-19 in Saudi Arabia: a national retrospective study. *J Infect Public Health.* 2020;13(7):920–925. doi:10.1016/j.jiph.2020.05.026
34. Matt V, Matthew H. The retrospective chart review: important methodological considerations. *J Educ Eval Health Prof.* 2013;10.
35. Bowling A. Mode of questionnaire administration can have serious effects on data quality. *J Public Health (Bangkok).* 2005;27(3):281–291.
36. Cheung KL, Peter M, Smit C, de Vries H, Pieterse ME. The impact of non-response bias due to sampling in public health studies: a comparison of voluntary versus mandatory recruitment in a Dutch national survey on adolescent health. *BMC Public Health.* 2017;17(1):1–10.

Risk Management and Healthcare Policy

Dovepress

Publish your work in this journal

Risk Management and Healthcare Policy is an international, peer-reviewed, open access journal focusing on all aspects of public health, policy, and preventative measures to promote good health and improve morbidity and mortality in the population. The journal welcomes submitted papers covering original research, basic science, clinical & epidemiological studies, reviews and evaluations,

guidelines, expert opinion and commentary, case reports and extended reports. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/risk-management-and-healthcare-policy-journal>