Co-Infection Pneumonia with *Mycobacterium abscessus* and *Pneumocystis jirovecii* in a Patient without HIV Infection Diagnosed by Metagenomic Next-Generation Sequencing

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Dan Xie *

Ying Xian*

Jingya You

Wen Xu

Min Fan

Xiaogang Bi

Kouxing Zhang

Department of General Intensive Care
Unit, Lingnan Hospital, The Third
Affiliated Hospital of Sun Yat-Sen
University, Guangzhou, People's Republic
of China

*These authors contributed equally to
this work

Introduction: Co-infection pneumonia with *Mycobacterium abscessus* (*M. abscessus*) and *Pneumocystis jirovecii* (*P. jirovecii*) is rarely reported in previously healthy patients without HIV infection. The diagnosis of pneumonia of *M. abscessus* and *P. jirovecii* remains challenging due to its nonspecific clinical presentation and the inadequate performance of conventional diagnostic methods.

Case Report: We report the case of a 44-year-old previously healthy male transferred to our hospital in February 2020 with a 4-month history of productive cough and one month of intermittent fever. At local hospital, the metagenomic next-generation sequencing (mNGS) detected *P. jirovecii* sequences in blood; with the antifungal therapy (Caspofungin, trimethoprim-sulfamethoxazole [TMP-SMX] and methylprednisolone [MP]), the patient still had hypoxemia, cough and fever. Then he was transferred to our hospital, the mNGS of bronchoalveolar lavage fluid (BALF) detected the sequences of *M. abscessus* and *P. jirovecii*. CD4+ T-lymphocytopenia in the peripheral blood cells was presented and HIV serology was negative. Caspofungin, TMP-SMX, clindamycin and MP were used to treat *P. jirovecii* pneumonia (PJP). Moxifloxacin, imipenem cilastatin and linezolid were used to treat *M. abscessus* infection. Clinical progress was satisfactory following antifungal combined with anti-*M. abscessus* therapy.

Conclusion: Co-infection pneumonia with *M. abscessus* and *P. jirovecii* as reported here is exceptionally rare. mNGS is a powerful tool for pathogen detection. *M. abscessus* infection could be a risk factor for *P. jirovecii* infection. This case report supports the value of mNGS in diagnosing of *M. abscessus* and *P. jirovecii*, and highlights the inadequacies of conventional diagnostic methods.

Keywords: *Pneumocystis jirovecii*, *Mycobacterium abscessus*, mNGS, co-infection, HIV negative

Introduction

Pneumocystis jirovecii (*P. jirovecii*) is a common opportunistic fungus causing *Pneumocystis jirovecii* pneumonia (PJP) in HIV infected patients. It is also a major opportunistic infection in HIV-negative immunocompromised patients, such as malignancy, stem cell transplantation (SCT), solid organ transplantation, rheumatologic disease, and respiratory disease.¹ *Mycobacterium abscessus* (*M. abscessus*) is a rapidly growing mycobacteria (RGM) of non-tuberculosis mycobacterium (NTM), which is an opportunistic pathogen of pulmonary infection

Correspondence: Kouxing Zhang
Department of General Intensive Care
Unit, Lingnan Hospital, The Third
Affiliated Hospital of Sun Yat-Sen
University, 2693 Kaichuang Avenue,
Guangzhou, 510530, People's Republic of
China
Email zhkoux@mail.sysu.edu.cn

in HIV infected patient. In HIV-negative people, *M. abscessus* infects both patients without underlying risk factors,^{2,3} and patients with a history of chronic lung diseases, such as cystic fibrosis and bronchiectasis, which led it to being the first RGM to be isolated in these contexts.^{2,4} In HIV-negative patients, there were a few cases of co-infection of *P. jiroveci* and *Mycobacterium tuberculosis*.⁵⁻⁷ But the co-infection of *M. abscessus* and *P. jiroveci* had not been reported before by search of Pubmed with the key words “‘*Mycobacterium abscessus*’ AND ‘*Pneumocystis*’”.

The pneumonia caused by *M. abscessus* and *P. jirovecii* present similar clinical features including fever and cough, which are nonspecific.^{2,5} Traditional diagnostic techniques in the microbiology laboratory for *P. jirovecii* and *M. abscessus* have some limits: *P. jirovecii* still cannot be reliably grown in vitro and *P. jirovecii* organisms in respiratory specimens following immunofluorescence staining suffer from low sensitivity;⁸ when the direct microscopic examination is positive for acid-fast bacilli, it is necessary to exclude the diagnosis of tuberculosis and other NTM organisms; *M. abscessus* cultures are not pigmented (neither scoto- nor photochromogen) and are similar to many other RGMs, such as *M. chelonae*;⁹ and the PCR of *P. jirovecii* and *M. abscessus* is not available in clinical settings in most hospitals in China including our hospital.

Metagenomic next generation sequencing (mNGS) is a powerful tool for detecting pathogens that can be performed directly on clinical specimens.^{10,11} It had been used for detection of *P. jirovecii*,^{8,10} and *M. abscessus*.¹² In this report, we describe a previously healthy HIV-negative male coinfecting with *M. abscessus* and *P. jirovecii* detected by mNGS.

Case Report

A 44-year-old previously healthy male was transferred to our hospital in February 2020 with a 4-month history of productive cough and one month of intermittent fever. He had no relevant transplantation and no significant medical history (particularly no corticosteroid and no immunosuppressive drug history), and never had any episode of tuberculosis or chronic pulmonary disorders. He was a never-smoker and no alcoholism. He had no family history of chronic lung disease. He had no recent travel history of Wuhan. Four months prior, the patient developed cough with white phlegm, accompanied with chest tightness and shortness of breath when he coughed

severely. Sometimes he had blood-streak sputum. He was suspected as upper respiratory tract infection and given symptomatic treatment, but the cough was not relieved. One month later, chest x-ray showed pulmonary infection. With the treatment of clarithromycin, ribavirin and ampicillin for 5 days, cough was relieved. However, one month prior, he developed low-grade fever. 22 days prior, he had a fever with maximum body temperature of 39.4°C. He was admitted to local hospital. Blood tests showed a white blood cell (WBC) count of $8.32 \times 10^9/L$ and C-reactive protein (CRP) of 128.06mg/L. Chest computed tomography (CT) scan showed bilateral infiltrations. Piperacillin tazobactam, oseltamivir and doxycycline were initiated after admission to treat pulmonary infection. One day later, chest CT showed the aggravated infection of both lungs. He developed to be hypoxia, septic shock. Acute respiratory distress syndrome (ARDS) was diagnosed. Then he was treated with meropenem, voriconazole, doxycycline. Reexamination of chest CT showed the infection was relieved. Five tests of nucleic acid of Corona Virus Disease 2019 (COVID-19) were all negative, which were carried out by regional Centers for Disease Control and Prevention (CDC). Eight days prior, the metagenomic next-generation sequencing (mNGS) of blood detected the sequences of *P. jirovecii* and *Cytomegalovirus* (CMV). Caspofungin, trimethoprim-sulfamethoxazole (TMP-SMX) and methylprednisolone (MP) were used for antifungal therapy. One day prior, the oxygen saturation decreased again. Then he was transferred to our hospital and admitted to our department.

Chest CT on admission showed bilateral infiltrations, ground glass opacity, crazy paving pattern (Figure 1A1 and A2). Blood tests performed on admission showed WBC count of $9.38 \times 10^9/L$, absolute value of lymphocyte (LYMP) of $0.65 \times 10^9/L$, hemoglobin (Hb) level of 126g/L, platelet count of $291 \times 10^9/L$, aspartate aminotransferase (AST) of 37 IU/L, alanine aminotransferase (ALT) of 40 IU/L, blood urea nitrogen (BUN) of 5.6mmol/L, and serum creatinine of 68umol/L. Hypersensitive C-reactive protein (hsCRP) was 43.2mg/L, and procalcitonin (PCT) was 0.025ng/mL. The physical examination showed blood pressure of 125/69mmHg, heart rate of 94 beats/minute, respiratory rate of 25 rates/minute, and temperature of 37.7°C, oxygen saturation of 99% with therapy of high-flow nasal cannula (HFNC) with fraction of inspiration (FiO2) 0.55). Moist rales were audible over bilateral lungs, without lower-extremity edema. Physical examination of the heart and abdomen

Figure 1 Chest CT performed on admission (**A1, A2**), on day 9 (**B1, B2**), on day 17 (**C1, C2**), on day 26 (**D1, D2**), and five months after discharge (**E1, E2**).

were normal. The arterial blood gas analysis revealed a pH of 7.45, the partial pressure of oxygen (PO₂) was 124mmHg, the partial pressure of carbon dioxide (PCO₂) was 45mmHg, PaO₂/FiO₂ was 113mmHg, alveolar-arterial oxygen gradient ((A-a) DO₂) was 212mmHg.

Intravenous moxifloxacin (400mg/day), ganciclovir (250mg, every 12 hours), caspofungin (50mg/day), clindamycin (600mg, every 8 hours) and methylprednisolone (40mg/day), and oral trimethoprim-sulfamethoxazole (TMP-SMX) (3 tablets (240/1200mg) every 6 hours, with

each tablet containing 80mg trimethoprim and 400mg sulfamethoxazole) were prescribed. Human immunoglobulin for intravenous injection (IVIG) was given with the dosage of 10g for 5 days. Thymalfasin was used with the dosage of 1.6mg for 12 days. Hepatitis B and C viruses, HIV, and Syphilis were all tested negative by serology. The sputum analysis of acid-fast bacilli smear and mycobacterial culture was negative. Mycobacterium tuberculosis complex was negative based on blood test with the GeneXpert MTB/RIF assay. The immunofluorescence staining of sputum smear did not find Pneumocystis organisms. Serum 1,3-beta-D-glucan (BDG) level was normal (<3.836pg/mL). In addition, bronchoalveolar lavage fluid (BALF) culture revealed no growth of bacteria or fungi. The DNA copies of CMV was 9E2 copies/mL. The lactate dehydrogenases level was 429 U/L. Antinuclear antibody (ANA), anti-double stranded DNA antibody(dsDNA), anti-neutrophil cytoplasmic antibody (ANCA) series and extractable nuclear antigen (ENA) series were all negative. Flow cytometry of peripheral blood cells showed CD3+T cells/lymphocyte was 86% (normal range 50%-84%), CD3+CD4+T cells/lymphocyte was 17%, CD3+CD8+T cells/lymphocyte was 55%, CD3+CD4+T cells/CD3+CD8+T cells was 0.31, B cells/lymphocyte was 12%, NK cells (CD3-CD16+ or CD3-CD56+)/lymphocyte was 2%. The thyroid stimulating hormone (TSH) level was normal. Peripheral blood and bronchoalveolar lavage fluid (BALF) samples collected on day 3 were sent for mNGS test, which was performed by Vision Medicals CO. Ltd (Guangzhou, China) using Illumina NextSeq 550 (USA). On day 4, the mNGS detected *M. abscessus* with 1315 high-confidence sequence reads, CMV with 649 high-confidence sequence reads and *P. jirovecii* with 44 high-confidence sequence reads in BALF sample. mNGS detected CMV with 74 high-confidence sequence reads and *P. jirovecii* with 368 high-confidence sequence reads in blood (Table 1). The *M. abscessus* sequence reads mapped to a *M. abscessus* reference genome (NC_010397.1), the *P. jirovecii* sequence reads assembled to a *P. jirovecii* reference genome (NW_017264779.1), and the sequence reads for CMV allowed assembly of a CMV reference genome (NC_006273.2) (Figure 2).

Meanwhile, the oxygen saturation decreased to about 90%, and the partial pressure of oxygen dropped to 54 mmHg. Moxifloxacin (400 mg/day) was switched to imipenem cilastatin (1.0g every 8 hours) for the treatment of *M. abscessus* infection. Linezolid (600mg every 12 hours) was synergistically used on day 6 to anti-*M. abscessus*.

Table 1 The Pathogen and Sequence Reads Detected by mNGS in BALF and Blood Samples

Sample	Pathogen	Sequence Reads
BALF	<i>M. abscessus</i>	1315
BALF	<i>P. jirovecii</i>	44
BALF	CMV	649
Blood	<i>M. abscessus</i>	0
Blood	<i>P. jirovecii</i>	368
Blood	CMV	44

Abbreviations: *M. abscessus*, *Mycobacterium abscessus*; *P. jirovecii*, *Pneumocystis jirovecii*; CMV, Cytomegalovirus.

On day 9, The patient discontinued HFNC, and be treated with nasal catheter for oxygen inhalation, and chest CT was performed and revealed that worsening infiltration in lower lungs and absorbing of other lungs (Figure 1B1 and B2). With the treatment, the patient still had intermittent low-grade fever, with the highest temperature of 38.2 °C, and the heat peak was gradually decreased, the body temperature returned to normal on day 10. The symptom of cough and sputum was also significantly relieved, the hsCRP decreased to 4.8mg/L. The PCT was 0.022ng/mL. On day 12, the patient was transferred to General Medical Wards. Moxifloxacin (400 mg/day), imipenem cilastatin (1.0g every 8 hours) and linezolid (600mg every 12 hours) were used to treat *M. abscessus* infection. Caspofungin (50mg/day), TMP-SMX (240/1200mg, every 6 hours), clindamycin (600mg, every 8 hours) and MP (20mg/day) were used to treat PJP. The copies of CMV were below 500 copies/mL tested on Day 11, so the ganciclovir was discontinued. On day 17, MP was switched to oral taken with 16mg/day for 7 days, then it was reduced to 12mg/day for 3 days. Chest CT performed on day 17 revealed absorbing infiltrations (Figure 1C1 and C2). On day 18, the count of WBC dropped to 3×10E9/L, recombinant human granulocyte stimulating factor injection was given. On day 20, caspofungin was discontinued, and clindamycin was changed from intravenous to oral administration with the dosage of 450mg/day (150mg every 8hours), moxifloxacin was also switched to oral taken (400mg/day). On day 23, linezolid was discontinued for the plantlet dropping to 89×10E9/L. On day 26, chest CT showed pulmonary infection was significantly absorbed (Figure 1D1 and D2). The level of LYMP elevated to 1.1×10E9/L, with the WBC of 3.97×10E9/L. The levels of AST, ALT, BUN and creatinine was normal. On day 27, he was discharged with 7-day oral

Figure 2 A, B, C detected from BALF; D and E detected from Blood. **(A)** The sequence reads for CMV allowed assembly of a CMV reference genome (NC_006273.2); **(B)** The *P. jirovecii* sequence reads assembled to a *P. jirovecii* reference genome (NW_017264779.1); **(C)** The sequence *M. abscessus* reads mapped to *M. abscessus* reference genome (NC_010397.1). **(D)** The sequence reads for CMV allowed assembly of a CMV reference genome (NC_006273.2); **(E)** The *P. jirovecii* sequence reads assembled to a *P. jirovecii* reference genome (NW_017264779.1).

medicines (SMZ 240/1200mg every 6hours, clindamycin 150mg every 8hours, MP 8mg/day, clarithromycin 500mg every 12hours, levofloxacin 0.5g/day). The important laboratory results on admission and after treatment are summarized in Table 2. The details of antibiotic and main process of disease are summarized in Figure 3. Five months after

discharge, he returned to our hospital for reexamination of chest CT which showed the inflammation in both lungs was basically completely absorbed (Figure 1E1 and E2). During this period, the patient had no symptoms such as cough and fever. Unfortunately, the patient refused blood tests including flow cytometry test.

Table 2 The Important Laboratory Results on Admission and After Treatment

Laboratory Parameters	On Admission	After Treatment	Normal Range
WBC count	9.38	3.97	3.5–9.5×10E9/L
Platelet count	291	89	100–350×10E9/L
LYMP	0.65	1.01	1.1–3.2×10E9/L
Hemoglobin	126	100	130–175g/L
ALT	40	18	3–35IU/L
AST	37	15	13–35IU/L
BUN	5.6	4.27	2.4–8.2mmol/L
Creatinine	68	88	31.8–116umol/L
hsCRP	43.2	4.8	0–10mg/L
PCT	0.025	0.022	0–0.05ng/mL
HIV	Negative	NA	Negative
Syphilis	Negative	NA	Negative
Hepatitis B and C	Negative	NA	Negative
Syphilis	Negative	NA	Negative
BDG	<3.836	NA	0–10 pg/mL
ANCA	Negative	NA	Negative
ENA	Negative	NA	Negative
Anti-dsDNA	Negative	NA	Negative
ANA	Negative	NA	Negative
TSH	0.7793	NA	0.35–4.94uIU/mL
DNA copies of CMV	9E2	<500	<500 copies/mL
CD4+T cells/lymphocyte	17%	NA	27%–51%
NK cells/lymphocyte	2%	NA	7%–40%
Acid-fast bacilli smear	Negative	NA	Negative
Culture of sputum	Negative	NA	Negative
GeneXpert MTB/RIF assay	Negative	NA	Negative
Sputum immunofluorescence staining for Pneumocystis organisms	Negative	NA	Negative

Abbreviations: WBC, white blood cell; LYMP, absolute value of lymphocyte; AST, aspartate aminotransferase; ALT, alanine aminotransferase; BUN, blood urea nitrogen; hsCRP, hypersensitive C-reactive protein; PCT, procalcitonin; BDG, 1,3-beta-D-glucan; ANA, antinuclear antibody; anti-dsDNA, anti-double stranded DNA antibody; ANCA, antineutrophil cytoplasmic antibody; ENA, extractable nuclear antigen; TSH, thyroid stimulating hormone; N/A, not available.

Figure 3 The details of antibiotic and main process of disease.

To further confirming *P. jirovecii*, the BALF and blood samples detected by mNGS were sent for PCR. The commercial PneumoGenius® real-time PCR (PathoNostics, Maastricht, The Netherlands) performed by Shanghai GeneDx Biotech Co., Ltd, according to the manufacturers' instructions using an LC480 II real-time system, detected *Pneumocystis* mitochondrial large subunit of ribosomal RNA (*mtLSU*) gene and wild-type sequence at the dihydropyrimidine synthase (*DHPS*) locus both in BALF and blood samples. The cycle threshold (Ct) value for *mtLSU* of BALF and blood were 32.5 and 36.31, respectively (Figure 4).

Discussion

To the best of our knowledge, this is the first report of pulmonary co-infection with *M. abscessus* and *P. jirovecii* responsible for respiratory distress in an HIV-negative previously healthy man without any known underlying illness.

P. jirovecii is a ubiquitous organism that commonly causes pneumonia in an AIDS-defining condition, it also caused pneumonia in HIV-negative immunocompromised patients. Predisposing factors of PJP in HIV-negative patients include corticosteroids, cirrhosis, CD4⁺T-lymphocytopenia, chemotherapy plus corticosteroids, alcoholism and malnutrition, visceral leishmaniasis, and so on.^{1,6,13–15} In our case, the patient was previously healthy (he had no significant medical history (especially no chemotherapy or corticosteroids), no

chronic pulmonary disorders, and no alcoholism). Also, the serology of Hepatitis B and C viruses, HIV, and Syphilis were negative, the autoantibodies (ANA, anti-dsDNA, ANCA, ENA) were all negative. There was no underlying disease in this patient. However, we observed CD4⁺T-lymphocytopenia in this patient (flow cytometry of peripheral blood showed CD3⁺CD4⁺T cells/lymphocyte was 17%). Therefore CD4⁺T-lymphocytopenia may be the predisposing factor of PJP in this patient.

M. abscessus is an emerging pathogen of increasing medical and microbiological concern. This bacterium is responsible for lung disease, most often in predisposed patients such as those with cystic fibrosis, chronic obstructive pulmonary disease and bronchiectasis, but this bacterium affects healthy patients in one-third of cases.¹⁶ Chronic infection is the most frequent presentation. Symptoms are variable and nonspecific, and a majority of patients suffer from a recurrent or chronic cough.^{2,16} In our case, prior to developed fever, the patient had three-month productive cough and hemoptysis, x-ray revealed pulmonary infection, with 5-day clarithromycin treatment, the cough relieved. Then he developed fever, hypoxia, and respiratory distress, *P. jirovecii* sequences were detected in blood by mNGS. But just given the antifungal therapy, the patient still had hypoxemia, cough and fever. Until mNGS detected *M. abscessus* sequences in BALF, *M. abscessus* pulmonary infection was considered. Following the

Figure 4 The cycle threshold (Ct) value for mtLSU of BALF and blood detected by PneumoGenius® real-time PCR were 32.5 and 36.31, respectively.
Abbreviations: PC, positive control; NC, negative control; Ct, cycle threshold.

combination of anti-*M. abscessus* therapy, the clinical symptom was significantly improved. So, it was confirmed that the patient was infected with *M. abscessus* firstly.

M. abscessus and *M. tuberculosis* shared many characteristics. *M. abscessus* is an intracellular bacterium and replicates in macrophages. Little is known about the immune response to *M. abscessus*. The control of *M. abscessus* infection relies, as for *M. tuberculosis*, on interferon gamma (IFN- γ) and tumor necrosis factor (TNF)- α production by T lymphocytes. Activation of the toll-like receptor2 (TLR2) signaling cascade leads to the activation of nuclear factor kappa B, resulting in a pro-inflammatory cytokine response.^{17,18} *M. abscessus* MAB2560 enhanced dendritic cell (DC) maturation via TLR4 and activated mitogen-activated protein kinases' (MAPKs') production of IL-12.¹⁹ The secretion of IL-12 will then activate and polarize naive CD4⁺ T cells towards Th1 cells to produce IFN- γ . Macrophages and natural killer (NK) cells can both release IL-12 and IFN- γ to guide T cells to the Th1 type phenotype.²⁰ Persistent untreated *M. abscessus* infection may lead to the depletion of CD4⁺ T cells and NK cells, which presented as decreasing number of CD4⁺ T cells and NK cells in our case. Wang H et al reported a case of disseminated cutaneous infection with *M. abscessus* with low level of CD4⁺ T cells in the peripheral blood cells and with the treatment of rifampin, isoniazid, ofloxacin, clarithromycin and thymosin for 6 months, the skin lesions were greatly improved and the level of CD4⁺ T cells in the peripheral blood cells became normal.²¹ Also, some cases reported the HIV-negative patients of co-infection *P. jiroveci* and *M. tuberculosis* had transiently CD4⁺ T-lymphocytopenia and completely resolved after anti-TB treatment.^{6,22} The complex interaction between innate immune cells, such as alveolar macrophage (AM)s and DCs with CD4⁺T-cells is important for an effective host adaptive response, essential for *Pneumocystis* clearance.²³ Therefore, the CD4⁺ T-lymphocytopenia leaves the host vulnerable to opportunistic fungal infection such as *Pneumocystis*. So, our patient had developed pulmonary infection caused by *M. abscessus* firstly which then induced a transient immunosuppression leading to PJP.

Traditional diagnostic techniques in the microbiology laboratory include specimen smear staining, growth and isolation of microorganisms in culture, detection of pathogen-specific antibodies (serology) or antigens and molecular identification of microbial nucleic acids (DNA or RNA), most commonly via PCR. In our case, the acid-fast bacilli staining and culture for *M. abscessus* was negative; the immunofluorescence staining for *P. jirovecii* was negative. The PCR of NTM and *P. jirovecii* was not available in our hospital.

Recently, mNGS has rapidly emerged as a promising single, culture-independent pathogen detection tool that can be performed directly on clinical specimens. mNGS targets all DNA or RNA present in a sample, allowing detection of the entire microbiome as well as the human host genome or transcriptome in patient samples.²⁴ Fortunately, in our case, mNGS directly detected the sequences of *P. jirovecii* in blood, and the sequences of *M. abscessus* and *P. jirovecii* in BALF samples, allowing confirmation of the diagnosis of PJP and the *M. abscessus* pneumonia, and allowing timely adjustment of treatment regimens. The clinical progress was satisfactory following antifungal therapy combined with anti-*M. abscessus* therapy. Later, the PneumoGenius[®] real-time PCR detected the *Pneumocystis mtLSU* gene both in BALF and blood samples, which was consistent with the results of mNGS. However, PCR can only detect target DNA or RNA in the sample, while mNGS can detect all DNA or RNA present in a sample.

Unfortunately, the patient refused blood test and mNGS of BALF at the follow-up visit. The lymphocytes count and CD4⁺ T-cells ratio and the sequence reads of *M. abscessus* and *P. jiroveci* of this patient after cured is not known, which is the major limitation of this study.

Conclusion

Co-infection pneumonia with *M. abscessus* and *P. jirovecii* as reported here is exceptionally rare. *M. abscessus* infection could be a risk factor for *P. jirovecii* infection.

mNGS is a powerful tool for pathogen detection. This case report supports the value of mNGS in diagnosing of *M. abscessus* and *P. jirovecii*, and highlights the inadequacies of conventional diagnostic methods.

Statement of Ethics

This research complies with the guidelines for human studies and is in accordance with the Declaration of Helsinki. This work was approved by the medical ethics committee of the Third Affiliated Hospital of Sun Yat-sen University, China (No. [2020]-02-246-01). Consent was obtained from the patient for participation in this study and the publication of associated data including radiological images. The authors confirmed that personal identity information of the patient data was unidentifiable from this report.

Acknowledgments

We thank Vision Medicals Co. Ltd (Guangzhou, China) and Shanghai GeneDx Biotech Co., Ltd (Shanghai, China) for providing the technical assistance.

We thank Ziyang Wang (Hangzhou Matridx Biotechnology Co., Ltd.) for providing the technical assistance.

Funding

This work was supported by the Major Science and Technology Project of Guangdong Province, China (No. 2013B020224002).

Disclosure

The authors declare no conflicts of interest in this work.

References

- Young N, McBride S, Morpeth S, Bryce A, Siddiqui A, Bhally H. Pneumocystis pneumonia in HIV-negative adults: missed opportunities for prevention. *N Z Med J*. 2020;133(1520):27–34.
- Griffith DE, Aksamit T, Brown-Elliott BA, et al. An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. *Am J Respir Crit Care Med*. 2007;175(4):367–416. doi:10.1164/rccm.200604-571ST
- Varghese B, Shajan SE, Al MO, Al-Hajj SA. First case report of chronic pulmonary lung disease caused by Mycobacterium abscessus in two immunocompetent patients in Saudi Arabia. *Ann Saudi Med*. 2012;32(3):312.
- Floto RA, Olivier KN, Saiman L, et al. US Cystic Fibrosis Foundation and European Cystic Fibrosis Society consensus recommendations for the management of non-tuberculous mycobacteria in individuals with cystic fibrosis: executive summary. *Thorax*. 2016;71(1):88–90.
- Ben-Mustapha I, Belkhouja K, Kheder S, et al. Coinfection with Mycobacterium tuberculosis and Pneumocystis jirovecii in immunocompetent young woman. *Arch Inst Pasteur Tunis*. 2013;90(1–4):55–60.
- Mongardon N, Bruneel F, Henry-Lagarigue M, Legriel S, Azarian R, Bedos JP. Pneumonia involving Mycobacterium tuberculosis and Pneumocystis jirovecii in HIV-seronegative patients. *Eur J Intern Med*. 2008;19(7):e70.
- Filoch P, Adoun M, Caron F, Godet C, Robert R, Meurice JC. Coinfection with Mycobacterium tuberculosis and Pneumocystis jirovecii in a patient without HIV infection. *Rev Mal Respir*. 2006;23(1 Pt 1):83–87.
- Chen J, He T, Li X, Wang X, Peng L, Ma L. Metagenomic Next-Generation Sequencing in Diagnosis of a Case of Pneumocystis jirovecii Pneumonia in a Kidney Transplant Recipient and Literature Review. *Infect Drug Resist*. 2020;13:2829–2836.
- Mougari F, Guglielmetti L, Raskine L, Sermet-Gaudelus I, Veziris N, Cambau E. Infections caused by Mycobacterium abscessus: epidemiology, diagnostic tools and treatment. *Exp Rev Anti Infect Ther*. 2016;14(12):1139–1154.
- Wang J, Han Y, Feng J. Metagenomic next-generation sequencing for mixed pulmonary infection diagnosis. *BMC Pulm Med*. 2019;19(1):252. doi:10.1186/s12890-019-1022-4
- Huang J, Jiang E, Yang D, et al. Metagenomic Next-Generation Sequencing versus Traditional Pathogen Detection in the Diagnosis of Peripheral Pulmonary Infectious Lesions. *Infect Drug Resist*. 2020;13:567–576. doi:10.2147/IDR.S235182
- Leo S, Gaia N, Ruppé E, et al. Detection of Bacterial Pathogens from Broncho-Alveolar Lavage by Next-Generation Sequencing. *Int J Mol Sci*. 2017;18(9):2011. doi:10.3390/ijms18092011
- Koffi N, Ngom A, Aka-Danguy E. [Association of pneumocystosis and pulmonary tuberculosis in an HIV-negative patient]. *Rev Mal Respir*. 1997;14(5):399–400.
- Toledo Jr. AC, de Castro MRD. Pneumocystis carinii pneumonia, pulmonary tuberculosis and visceral leishmaniasis in an adult HIV negative patient. *Braz J Infect Dis*. 2001;5(3):154–157. doi:10.1590/S1413-86702001000300008
- Onorati P, Carfagna P, Palange P, Venditti M, Serra P. CD4+ T-lymphocytopenia and Pneumocystis carinii pneumonia in a patient with miliary tuberculosis. *Eur J Intern Med*. 2001;12(2):134–136.
- Griffith DE, Girard WM, Wallace RJ. Clinical Features of Pulmonary Disease Caused by Rapidly Growing Mycobacteria: an Analysis of 154 Patients. *Am Rev Respir Dis*. 1993;147(5):1271–1278. doi:10.1164/ajrccm/147.5.1271
- Aulicino A, Dinan AM, Miranda-CasoLuengo AA, et al. High-throughput transcriptomics reveals common and strain-specific responses of human macrophages to infection with Mycobacterium abscessus smooth and rough variants. *BMC Genomics*. 2015;16(1):1046. doi:10.1186/s12864-015-2246-1
- Davidson LB, Nessar R, Kempaiah P, Perkins DJ, Byrd TF, Fessler MB. Mycobacterium abscessus Glycopeptidolipid Prevents Respiratory Epithelial TLR2 Signaling as Measured by HβD2 Gene Expression and IL-8 Release. *PLoS One*. 2011;6(12):e29148. doi:10.1371/journal.pone.0029148
- Lee SJ, Shin SJ, Lee SJ, et al. Mycobacterium abscessus MAB2560 induces maturation of dendritic cells via Toll-like receptor 4 and drives Th1 immune response. *BMB Reports*. 2014;47(9):512–517. doi:10.5483/BMBRep.2014.47.9.001
- Tomioka H, Tatano Y, Maw WW, Sano C, Kanehiro Y, Shimizu T. Characteristics of suppressor macrophages induced by mycobacterial and protozoal infections in relation to alternatively activated M2 macrophages. *Clin Dev Immunol*. 2012;2012:635451.
- Wang H, Jin P, Wu Q. Disseminated cutaneous infection with Mycobacterium abscessus in a patient with a low CD4+ T cell count. *Eur J Dermatol*. 2008;18(3):337–340. doi:10.1684/ejd.2008.0400
- Onorati P, Carfagna P, Palange P, Venditti M, Serra P. CD4+ T-lymphocytopenia and Pneumocystis carinii pneumonia in a patient with miliary tuberculosis. *Eur J Intern Med*. 2001;12(2):134–136. doi:10.1016/S0953-6205(01)00113-3
- Kelly MN, Shellito JE. Current understanding of Pneumocystis immunology. *Future Microbiology*. 2001;12(2):43–65. doi:10.2217/fmb.09.116
- Chiu CY, Miller SA. Clinical metagenomics. *Nat Rev Genet*. 2019;20(6):341–355.

Infection and Drug Resistance

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>

Dovepress