

A Retrospective Analysis of the Initial Effect of COVID-19 on German Prehospital Care During Lockdown in Germany

This article was published in the following Dove Press journal:
Open Access Emergency Medicine

Marcus Örgel¹
Manfred Gogol¹
Tilman Graulich¹
Mohamed Omar¹
Alexander Ranker²
Christian Böttcher¹
Emmanouil Liodakis¹
Christian Krettek¹
Christian Macke¹

¹Trauma Department, Hannover Medical School, Hanover, Germany; ²Department of Rehabilitation Medicine, Hannover Medical School, Hanover, Germany

Introduction: This retrospective cohort analysis examines the impact of the COVID-19 pandemic in the prehospital setting in Germany. The data of two emergency physician response units of a northern German region with 1.2 million citizens was analyzed retrospectively.

Materials and Methods: We analyzed the period March 16 to April 16 for the year 2020 when the lockdown took place in Germany and compare the results for the same period for the year 2019 and 2018. 1004 patients were included. Demographic data, the type of rescue missions, the number of missions per day, the National Advisory Committee for Aeronautics Score (NACA-score), the frequency of respiratory emergencies (COVID-19 and non-COVID-19 associated), as well as the number of deaths were documented.

Results: Mean age was 62.3±24.8 years and 576 (56.5%) were male. Number of missions were 397, 403 and 333 in 2018, 2019 and 2020 respectively. The control room registered a 22% reduction of rescue missions for the year 2020. Even the amount of emergency calls via the emergency number 112 was reduced by 17.4% between 2018 and 2020. 150 (14.9%) missions were due to respiratory emergencies. In 2020 10 missions (28.6% of respiratory emergencies) were COVID-19 related. In 2020 the NACA score increased significantly.

Conclusion: We found a decreasing effect of the COVID-19 pandemic in Germany on the number of emergency calls as well as missions, and an increase of the severity of cases in preclinical care for a northern German region with 1.2 million citizens. The effect of these findings caused by COVID-19 on the health care system remains to be seen.

Keywords: emergency medicine, SARS-CoV-2, COVID-19, prehospital care, emergency physician response units, pandemic

Background

Since the COVID-19 pandemic¹⁻⁴ started in China in December 2019, Europe has become a center of this new disease by number of cases and deaths. Germany is less affected with a case fatality proportion^{5,6} (CFP) of 0.041 compared to Italy (0.137),⁷ Spain (0.102)⁷ or France (0.148).⁷ Germany is, with its absolute number of cases (n=165,745) ranked 6 in Europe after Russia, Spain, UK, Italy, and France. Some authors report during the pandemic of less patient treatments due to heart disease or stroke,^{8,9} and that vaccination rates were lower than in former years (Morbidity and Mortality Weekly Report 2020 [MMWR]).¹⁰ The authors also noticed fewer medical emergencies in the emergency room of our hospital as well as a lower number of emergency calls for the hospital-based physician

Correspondence: Marcus Örgel
Department for Traumatology, Hannover Medical School (MHH), Carl – Neuberg – Straße 1, Hannover, 30625, Germany
Tel +49 511 532 2099
Email Oergel.Marcus@mh-hannover.de

response unit (PRU) and air ambulance (HEMS [helicopter emergency medical service]).

Objectives

This study compares during lockdown in Germany while the COVID-19 pandemic six parameters (1–6) in 2020 with the two previous years of 2018 and 2019 to identify the influence of the COVID-19 pandemic of the prehospital emergency system: (1) to quantify for each year number of emergency calls and (2) subsequent rescue missions, (3) number of respiratory diseases and (4) COVID-19-associated diseases, as well as (5) the severity of the diseases and (6) number of deaths. We suggest that the COVID-19 pandemic has an impact on all of these aspects in the prehospital emergency system.

Methods

Study Design and Setting

This retrospective single center cohort analysis included all rescue missions of a physician response unit (PRU, 24/7 on service) and air ambulance (HEMS, 07:00 a.m./sunrise to sunset on service) that were conducted from March 16 to April 16, 2020, the time when the lockdown took place in Germany and during the same period in 2018 and 2019. The PRU and HEMS are located at the Trauma Department at Hannover Medical School (MHH), a level 1 trauma center, which is a university hospital in Lower Saxony, Germany, with approximately 1600 inpatient beds. Other PRUs and HEMS in lower Saxony were not included.

Furthermore, a survey of all emergency calls and rescue missions carried out in the city and region of

Hanover, an area with 1.2 million citizens, was conducted by Hanover Fire Department. The control room, run by Hanover Fire Department, coordinates overall one primary HEMS (air ambulance [helicopter emergency medical service]), five PRUs (physician response unit) for the city and seven PRUs for the greater area of Hannover. In addition, the control room coordinates a second HEMS too, whose primary task is the physician assisted airborne secondary transport of critically ill patients between hospitals.

A potential confounder of this study could be the varying number of emergency physicians assigned during this time period with different levels of medical education and individual experience in emergency medicine.

Participants and Data Assessment

For the above-mentioned period, age, gender and emergency characteristics (traumatology, internal medicine, neurology, psychiatry, pediatrics or gynecology, [Table 1](#)) of all patients as well as the number of emergency calls per day were obtained. We looked at disease severity with the National Advisory Committee for Aeronautics (NACA) score (0–7, 0= no injury or illness, 7=death, detailed information is shown in [supplementary material](#)), respiratory emergencies and the likelihood of association with COVID-19, and deaths. The NACA score was prospectively documented in the medical reports. Data assessment was performed anonymously by two of the authors.

Additionally, we documented the number of emergency calls received by Hanover Fire Department. Considering the total number of infections and deaths

Table 1 Percentage Distribution of the Emergency Physician Missions

Type of Rescue Mission	PRU*				HEMS**				PRU + HEMS n (%)
	2018 n (%)	2019 n (%)	2020 n (%)	Total n (%)	2018 n (%)	2019 n (%)	2020 n (%)	Total n (%)	
Traumatology	19 (8.2)	28 (11.9)	26 (12.9)	73 (10.9)	53 (41.1)	45 (39.1)	46 (48.7)	144 (42.6)	217 (22.6)
Internal Medicine	169 (72.8)	162 (68.6)	148 (73.6)	479 (71.6)	62 (51.8)	56 (47.1)	41 (46.0)	159 (48.5)	638 (63.1)
Neurology	30 (12.9)	36 (15.3)	22 (10.9)	88 (13.2)	8 (6.4)	15 (12.3)	3 (2.7)	26 (7.4)	114 (11.0)
Psychiatric	6 (2.6)	5 (2.1)	2 (1.0)	13 (1.3)	0 (0.0)	0 (0.0)	1 (1.8)	1 (1.0)	14 (1.2)
Pediatric	4 (1.7)	3 (1.3)	2 (1.0)	9 (1.9)	1 (0.7)	1 (0.7)	2 (0.9)	4 (0.3)	13 (1.3)
Gynecology	4 (1.7)	2 (0.8)	1 (0.5)	7 (1.0)	0 (0.0)	1 (0.7)	0 (0.0)	1 (0.3)	8 (0.8)

Abbreviations: *PRU, physician response unit; **HEMS, air ambulance (helicopter emergency medical service).

Table 2 COVID-19 Suspected Patients Treated by the Emergency Services PRU* and HEMS**

	Sex	Age	COVID-19- Infection Identified Prehospitally	Initial Symptoms	COVID Confirmed	Treatment	Death	Emergency Services
Patient 1	♂	54	No	Dyspnea, oxygen saturation <90%, cough, fever >38.5°C	Positive	Oxygen and volume supply, admitted to ICU	No	PRU*
Patient 2	♀	53	No	Dyspnea, oxygen saturation <90%, cough, fever >38.5°C	Negative	Oxygen and volume supply, admitted to ICU	No	PRU*
Patient 3	♀	89	No	Dyspnea, oxygen saturation >90%, cough, no fever	Negative	Oxygen and volume supply, admitted to a normal ward	Yes	PRU*
Patient 4	♀	83	No	Dyspnea, oxygen saturation <90%, cough, fever >38.5°C	Positive	NIV, volume supply, admitted to ICU	Yes	PRU*
Patient 5	♂	58	No	Dyspnea, oxygen saturation <90%, cough, fever >38.5°C, cardiovascular arrest	Negative	Resuscitation, admitted to ICU	Yes	PRU*
Patient 6	♂	83	Yes	Dyspnea, oxygen saturation <90%, cough, fever >38.5°C	Positive	Oxygen and volume supply, admitted to ICU	Yes	PRU*
Patient 7	♂	80	No	Dyspnea, oxygen saturation <90%, cough, fever >38.5°C	Positive	Oxygen and volume supply, admitted to ICU	No	PRU*
Patient 8	♂	45	No	Dyspnea, oxygen saturation <90%, cough, no fever	Positive	Oxygen and volume supply, admitted to ICU	No	PRU*
Patient 9	♀	81	No	No Dyspnea, oxygen saturation >90%, fever >38.5°C	Positive	Oxygen and volume supply, admitted to a normal ward	No	PRU*
Patient 10	♀	93	No	Dyspnea, oxygen saturation <90%, cough, no fever <38°C	Positive	Oxygen and volume supply, admitted to a normal ward	Yes	PRU*
Patient 11	♂	96	No	Bradycardia and cardiovascular arrest	Negative	Resuscitation	Yes	HEMS**
Patient 12	♂	95	No	Dyspnea, oxygen saturation <90%, cough, fever >38.5°C	Negative	Oxygen and volume supply, admitted to a normal ward	No	HEMS**

Notes: ♀= female, ♂= male.

Abbreviations: ICU, intensive care unit; *PRU, physician response unit; **HEMS, air ambulance (helicopter emergency medical service).

related to COVID-19 the case fatality proportion^{5,6} was calculated for Germany, its selected states and counties as well as cities (Tables 2 and 3) and compared to the other countries (USA, Spain, Italy and France).

COVID-19 Disease in Hannover

We assessed publicly available data from local authorities, and from the Robert Koch-Institut (RKI)¹¹ in Hamburg. For comparison of CFP (case fatality proportion) values of European

Table 3 Demographic Figures Related to COVID-19 Infections and Deaths for the International Comparison Based on Data from the Johns Hopkins University¹ (as of March 3, 2020)

Countries	Death (n)	Infections (n)	CFP* ^{2,3}
USA	68,957	1,192,906	0.058
Italy	28,884	210,717	0.137
Spain	25,428	248,301	0.102
France	24,895	168,693	0.148
Germany	6,866	165,745	0.041

Abbreviation: *CFP, case fatality proportion.

countries and the USA, data from Johns Hopkins University & Medicine Coronavirus Resource Center¹² was used.

Statistical Methods

Statistical analysis was performed using SPSS 26 (IBM, SPSS Inc., Chicago, IL). For dichotomous variables, the Fisher's exact test was used. For mean variables, the Mann-Whitney *U*-test was used after checking for normal distribution. Significance value was set to $p < 0.05$.

Results

Participants

From March 16 to April 16 in 2018, 2019 and 2020 $n=1133$ rescue missions were recorded. 1004 patients were included for evaluation of patient data (Figure 1). The only exclusion criterion for patient data was a mission being canceled before arrival at the scene. All included patients had complete data sets. Figure 1 shows a flow chart of the study design.




Figure 1 Flow chart of the inclusion and exclusion criteria for the cohort. *The proportion of 10% cancelled missions is comparable throughout the years (data not shown).

Main Results

The average age of the total cohort was 62.3 ± 24.8 years. Patients treated by HEMS (air ambulance [helicopter emergency medical service]) were on average 3.5 years younger (HEMS: 59.9 ± 25.5 vs PRU: 63.6 ± 24.3 , $p=0.027$). Comparing 2018 to 2019 and 2020 respectively, patients treated by HEMS became also significantly younger over the years (2018: 63.4 ± 25.8 ; 2019: 60.1 ± 25.9 , $p=0.131$; 2020: 54.9 ± 23.9 , $p=0.011$; both compared to 2018). The age of patients treated by PRUs (physician response unit) increased significantly compared to 2018 and 2019 (2018: 61.5 ± 25.3 ; 2019: 63 ± 24.9 , $p=0.118$; 2020: 66.6 ± 22.1 , $p=0.029$; both compared to 2018).

Five hundred and seventy-six (56.5%) of the 1004 patients were male. The subgroup analysis of HEMS and PRU showed that significantly more men than women were treated by HEMS (228 (65.3%) vs 121 (34.7%), $p < 0.001$). The gender distribution within the subgroups did not vary significantly (PRU 2018–2020: men: 46.4–56.2%, women: 43.8–53.6%; $p=0.09$; HEMS 2018–2020: men: 61.7–72.2%, women: 27.8–38.3%; $p=0.237$). Figure 2 shows the gender distribution within PRUs and HEMS cumulated for 2018 to 2020.

In the study period of 2018 $n=21,008$, 2019 $n=18,946$ and 2020 $n=17,360$ emergency calls (objective 1) were registered, respectively. The assessment of the total number of rescue missions in the City and Region of Hanover showed $n=12,564$ missions in the analyzed period for 2018, $n=13,070$ for 2019 and $n=9,973$ for 2020. This amounts to a reduction of 22%.

The number of emergency missions (objective 2) in 2020 ($n=333$) decreased by about 17% compared to 2018 ($n=403$) and 2019 ($n=397$). The amount of missions




Figure 2 Bar chart for graphical representation of gender distribution, separated into the subgroups PRU and HEMS.

per day in 2020 declined significantly compared to previous years to 3.53 ± 1.30 (2018: 4.41 ± 1.60 ; $p=0.019$; 2019: 4.31 ± 1.71 ; $p=0.044$) for HEMS and to 6.88 ± 1.66 (2018: 8.19 ± 1.66 ; $p=0.002$; 2019: 8.09 ± 2.67 ; $p=0.032$) for PRUs. However, medical reasons for emergency physician missions did not change significantly over time (Table 1).

A total of $n=150$ (14.9%) missions were due to respiratory emergencies (objective 3), with HEMS ($n=22$, 5.6%) treating significantly fewer respiratory emergencies than PRU ($n=128$, 19.1%, $p<0.001$). However, there was no difference in the number of respiratory emergencies in 2020 compared to previous years (2020: HEMS $n=3$ (3.1%), PRU $n=35$ (17.5%)).

In 2020 $n=10$ patients (28.6%) with COVID-19 (objective 4) associated complaints were treated by the PRU of Hannover Medical School. None of the COVID-associated cases died in the prehospital phase but in three cases severe respiratory failure had to be managed by two times endotracheal intubation and one times non-invasive ventilation (NIV). When NIV was used for respiratory support in the preclinical phase, a COVID-19 infection of the attending physician occurred. In this case there was a relative indication for an intubation. It was up to his/her experience and evaluation of the situation. He/she made the decision for NIV. The clinical findings of these ten patients are shown in Table 2.

The average NACA (National Advisory Committee for Aeronautics) score (objective 5) of the total cohort was 3.52 ± 1.36 . The comparison of the complete NACA score (PRU and HEMS) for both rescue services for the years 2018 and 2019 showed no significant difference (2018: 3.39 ± 1.28 , 2019: 3.44 ± 1.32 , $p=0.625$). However, for the period under investigation in 2020, a significant increase in the NACA score was found compared to the same periods in 2018 and 2019 (3.76 ± 1.47 , $p=0.004$ compared to 2019, $p=0.001$ compared to 2018).

The subgroup analysis showed that the NACA score was significantly higher for HEMS than in the PRU group (3.64 ± 1.43 vs 3.46 ± 1.32 ; $p=0.048$).

While in the PRU group the COVID-19 pandemic did not have a significant impact on the NACA score (Figure 3, 2018: 3.45 ± 1.24 ; 2019: 3.39 ± 1.32 , $p=0.632$; 2020: 3.54 ± 1.40 ; $p=0.450$; each compared to 2018), the NACA score for HEMS increased significantly in 2020 (2018: 3.30 ± 1.35 ; 2019: 3.54 ± 1.34 , $p=0.155$; 2020: 4.17 ± 1.50 , $p<0.001$; each compared to 2018).

In total, prehospital deaths (objective 6) were recorded in 78 (7.8%) patients. Mortality did not differ significantly between years (2018 $n=22$, 6.2%; 2019 $n=32$, 9.1%; 2020




Figure 3 Graphical display of the NACA Score for PRU and HEMS as well as for the whole cohort; $^{*}p=0.001$ 2020 vs 2018; $^{§}p=0.004$ 2020 vs 2019; $^{*}p<0.001$ 2020 vs 2018, $^{§}p=0.001$ 2020 vs 2019.

$n=24$, 8.2%; $p=0.332$). There were also no differences in rescue missions related to mortality for 2020, neither between the years nor between HEMS ($n=8$, 8.6%) and PRU ($n=16$, 8.0%, $p=0.88$). Data clarifying death by COVID-19 was not available for these patients.

To assess the severity of disease in the districts and states as well as for international comparison, the CFP values are shown in detail in Tables 3 and 4.

Discussion

Key Results

In summary, our study shows a significant reduction in the number of PRU and HEMS missions per day and a significant increase in the severity of the disease in HEMS cases. The number of emergency missions handled by the control room for the period under investigation in 2020 compared to the same period in 2018 and 2019 decreased by 17%. We observed no excessive increase of interventions due to respiratory or COVID-19 associated complaints. The number of respiratory non-COVID-19 associated emergencies did not change significantly compared to 2018 and 2019. The same applies to the number of deaths.

Strengths and Limitations

For the data assessment of the study, we decided to analyse the period when the lockdown took place in Germany. At this time we expected the strongest impact of COVID-19 on people. In addition, the quarantine measures were carried out for this period and thus external disturbance factors as sources of infection, such as an increased risk of infection by returning holidaymakers or foreigners

Table 4 Demographic Figures Related to COVID-19 Infections and Deaths in the Individual Federal States and Districts as Well as for Germany Based on Data from the Robert Koch Institute¹ (as of 03.05.2020)

Location	Infections (n)	Deaths (n)	CFP ^{2,3}	Number of COVID-19 Cases by Age (n)											
				0–4 (n)		5–14 (n)		15–34 (n)		35–59 (n)		60–79 (n)		80+ (n)	
				♀	♂	♀	♂	♀	♂	♀	♂	♀	♂	♀	♂
Germany	165,745	6,866	0.041	647	757	1,656	1,755	20,579	19,321	35,627	32,402	14,666	16,237	11,575	6,648
Bavaria	42,792	1,910	0.045	215	234	550	558	5,425	5,114	9,271	8,047	3,908	4,333	3,155	1,815
District of Starnberg*	504	9	0.017	3	1	10	11	70	64	122	110	40	44	12	16
North Rhine Westphalia	33,428	1,287	0.039	125	136	306	311	4,270	3,972	7,334	7,238	2,884	3,228	2,314	1,261
District of Heinsberg**	1,756	62	0.035	5	5	6	9	191	230	341	408	203	191	78	88
Lower Saxony	10,283	453	0.044	27	36	94	122	1,276	1,193	2,229	1,972	1,006	1,042	787	430
District of Hanover	1,946	71	0.036	5	7	30	20	231	278	392	412	170	134	74	144

Notes: *First reported patient with a COVID-19 infection; ♀ = female, ♂ = male; **First major outbreak reported in Germany on February 25, 2020 after carnival party in Heinsberg.

entering the country as well as increased social contacts, were reduced.

The used cohort (n=1004) was made as a single center study up from two of six physician staffed services of Hanover city. Therefore, a selection bias could be possible. The study size was derived from the calls processed in the control room run by Hanover Fire Department for the above mentioned period. In this context, it should be mentioned that only certain keywords provide the indication for the deployment of a PRU (physician response unit) or HEMS (air ambulance [helicopter emergency medical service]). These are for example severe chest pain, serious injuries, eg, in traffic accidents, but also acute respiratory distress, eg, in COPD, as well as symptoms of a potential COVID-19 infection.

Nevertheless, as far as the selection of the data origin is concerned, the only limiting factor is the affiliation of the rescue services to the authors' hospital. The control room of Hanover Fire department is completely independent, therefore the data are reliable and objective. Thus, neither the reason for alerting nor the frequency of deployment or the decision which emergency service was sent out could be influenced. Within the scope of our investigation, the detailed clinical pictures of the rescue operations could not be analyzed separately. This is due to the fact that pre-hospital diagnoses are made based on telephone information only. It could neither be revised on scene or in hospital, and could not be assessed by us later. Therefore, the missions are divided into six categories: traumatology, internal medicine, neurology, psychiatric, pediatric or gynecological, without further explanation of clinical symptoms (Table 1).

For a better classification and objectivity of our results, we calculated the CFP. CFP is evaluated in the literature as a valid epidemiological parameter,^{5,6} and is a simple indicator to assess the severity of the pandemic. The higher this parameter is, the more severe is the impact of the disease on the population. Thus, countries, states and regions as well as cities can be easily compared.

In order to present data which are as objective as possible, the selection of these three states as a comparative cohort is justified by the fact that the western (North Rhine-Westphalia with a major outbreak after a carnival party in Heinsberg on February 27, 2020) and southeastern parts of Germany (Bavaria with the first COVID-19 patient in Germany on January 28, 2020) showed a higher incidence and mortality compared to the nationwide infection numbers. In addition, there is

a geographical proximity between Bavaria and Northern Italy, which had been badly affected by the COVID-19 pandemic. For international comparison, the infection and death figures due to COVID-19 of the USA, Italy, Spain and France were used. At the time of the data collection these countries were seriously affected by the COVID-19 pandemic, although, with the exception of the USA, in these European neighbouring countries, nearly comparable health services and socio-economic standards as in Germany, existed. Nevertheless, due to less ICU beds and ventilators per 100,000 citizen the comparability could be relatively limited and this has indeed an impact on the CFP and leads to more worse CFP rates in Italy, Spain and France than in Germany.

Interpretation

COVID-19 takes on a special role, as it not only acts as a final diagnosis in the sense of an upper respiratory tract infection, but also as a secondary powerful contributor. In addition to the primary effect of the disease, it can also have a secondary effect, intensified by quarantine measures, on the avoidance behavior of patients in the sense of fear of an increased risk of infection with COVID-19, eg, through close contacts, especially in hospitals. The closure of schools, pre-schools, day-care centers and sports clubs has led to a reduction in everyday activities and thus to a lower risk of acquiring infections at school, during sports activities and on the daily commute. Nevertheless, our data showed even a reduction across all types of rescue missions, so that we could not confirm this effect. Despite this, an increase in the severity of illness of the NACA (National Advisory Committee for Aeronautics) score could be shown for HEMS (air ambulance [helicopter emergency medical service]). This could be due to the fact that the patients with minor illnesses did not call the emergency services out of fear and thus a shift to higher NACA values occurred. However, it could also be an expression of the time delay until the emergency call is made since patients wait and only call when the symptoms are more severe. These circumstances could have led to a reduction in the use of emergency medical services as well as greatly delayed decision making until the emergency consultation occurred. These facts could also explain the reduction in the number of interventions. This is also confirmed by the reduced number of emergency calls. Nevertheless, the quarantine situation may have led to a reduced stress situation in addition to

avoidance behavior, which could also explain the reduction of ambulance requests.

Whereas Marijon et al showed a significant increase in out-of-hospital cardiac arrest incidence, coupled with a reduction in survival during the specified time period of the pandemic when compared with the equivalent time period in previous years with no pandemic,¹³ Trabattoni et al reported that there has been a demonstrable reduction of about 25% in the treatment of heart attacks.⁸ This would be in line with our results, as we were also able to prove a reduction in the number of treatments for the above-mentioned period. But this is contradicted by a UK (United Kingdom) study, which reports no changes in the number of heart attacks and strokes.¹⁴ Also, Baldi et al showed a cumulative incidence of diagnosed Covid-19 and cases of out-of-hospital cardiac arrests in Italy. Furthermore, the authors showed that the cumulative incidence of out-of-hospital cardiac arrests in 2020 was strongly associated with the cumulative incidence of COVID-19. The number of cases of out-of-hospital cardiac arrests compared to 2019 increased also.¹⁵ These findings may be explained with different CFPs (case fatality proportion) in these regions. A region with comparably low CFPs like ours may have a problem with avoidance behavior of the general public, thus reducing the utilization of medical services. Whereas regions with higher CFPs as an expression of more severe courses of the pandemic itself have to deal with COVID linked diseases and deaths, clouding the reduction in utilization of medical services and resulting in equivalent or even higher disease and death rates.¹⁶

Generalizability

In assessing external validity, it should be taken in mind that our cohort was analyzed under special conditions (emergency situations). The objectives 1, 2, 3, 4, 5 and 6 were confirmed. This suggests that the pandemic triggered by the SARS-CoV-2 virus had and still has an impact on prehospital care in the German health care system. Equally, COVID-19 also has had an indirect (collateral) impact on the health of citizens in other countries.^{8,13,15}

Based on the calculated CFP, a representative classification of our results for the infection and death figures available in Germany at the time of our study was possible. It was noticeable that 17% fewer emergency medical interventions were carried out in the period under review compared with the same periods in 2018 and 2019. Nevertheless, the NACA score of the emergency medical

interventions we analyzed increased, so that an effect of the COVID-19 pandemic can until now only be shown with regard to the number of rescue missions and emergency calls.

Conclusion

We found a decreasing effect of the COVID-19 pandemic in Germany on the number of emergency calls as well as missions, and an increase of the severity of cases in pre-clinical care for a northern German region with 1.2 million citizens. The effect of these findings caused by COVID-19 on the health care system remains to be seen.

Abbreviations

CFP, case fatality proportion; Control room, Control center, guidance office for the emergency teams; HEMS, air ambulance, helicopter emergency medical service; MHH, Hannover Medical School; MMWR, Morbidity and Mortality Weekly Report; NACA, National Advisory Committee for Aeronautics; NIV, non-invasive ventilation; PRU, physician response unit; RKI, Robert Koch-Institut; UK, United Kingdom; USA, United States of America.

Transparency

Availability of data and material: The data sets used and analyzed during the current study are available from the corresponding author on request.

Ethics Approval and Consent to Participate

Each author certifies that all investigations were conducted in conformity with ethical principles of research. Due to the retrospective data collection ethical approval was given as a waiver (No. 9099_BO_K_2020) and consent was granted by the Ethics Committee of Hannover Medical School.

Acknowledgments

We thank Hanover Fire Department for offering their help in data collecting.

Funding

We acknowledge support by the German Research Foundation (DFG) and the Open Access Publication Fund of Hannover Medical School (MHH).

Disclosure

The authors declare that they have no competing interests.

References

1. Chung M, Bernheim A, Mei X, et al. CT imaging features of 2019 Novel Coronavirus (2019-nCoV). *Radiology*. 2020;295:202–207. doi:10.1148/radiol.202000230
2. Fu L, Wang B, Yuan T, et al. Clinical characteristics of coronavirus disease 2019 (COVID-19) in China: a systematic review and meta-analysis. *J Infect*. 2020;80:656–665. doi:10.1016/j.jinf.2020.03.041
3. Linton NM, Kobayashi T, Yang Y, et al. Incubation period and other epidemiological characteristics of 2019 Novel Coronavirus infections with right truncation: a statistical analysis of publicly available case data. *J Clin Med*. 2020;9:538. doi:10.3390/jcm9020538
4. World Health Organization: Coronavirus disease 2019. (COVID-19) situation report – 87; 2020. Available from: www.who.int/docs/default-source/coronaviruse/situation-reports/20200416-sitrep-87-covid-19.pdf?sfvrsn=9523115a_2. Accessed April 20, 2020.
5. Rothman KJG, Sander L, Timothy L. *Modern Epidemiology*. Lippincott Williams & Wilkins (LWW); 2008.
6. Gianicolo E, Blettner M, Karch A, Karch A. Epidemiological measures in the context of the COVID-19 pandemic. *Dtsch Arztebl Int*. 2020;117:336–342. doi:10.3238/arztebl.2020.0336
7. John Hopkins University of Medicine. Coronavirus Resource Center. Available from: <https://coronavirus.jhu.edu/map.html>. Accessed February 15, 2021.
8. Trabattoni D, Montorsi P, Merlino L. Late STEMI and NSTEMI patients' emergency calling in CoVID-19 outbreak. *Can J Cardiol*. 2020;36:1161.e7–1161.e8. doi:10.1016/j.cjca.2020.05.003
9. Mehra MR, Desai SS, Kuy S, Henry TD, Patel AN. Cardiovascular disease, drug therapy, and mortality in Covid-19. *N Engl J Med*. 2020;382:2582. doi:10.1056/NEJMc2021225
10. Bramer CA, Kimmins LM, Swanson R, et al. Decline in child vaccination coverage during the COVID-19 pandemic – Michigan care improvement registry, May 2016–May 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69:630–631. doi:10.15585/mmwr.mm6920e1
11. Robert Koch Institut. Available from: https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Fallzahlen.html. Accessed February 15, 2021.
12. Robert Koch-Institut: COVID-19-Dashboard Available from: https://experience.arcgis.com/experience/478220a4c454480e823b17327b2bf1d4/page/page_1/. Accessed February 15, 2021.
13. Marijon E, Karam N, Jost D, et al. Out-of-hospital cardiac arrest during the COVID-19 pandemic in Paris, France: a population-based, observational study. *Lancet Public Health*. 2020;5:e437–e443. doi:10.1016/S2468-2667(20)30117-1
14. Holmes JL, Brake S, Docherty M, Lilford R, Watson S. Emergency ambulance services for heart attack and stroke during UK's COVID-19 lockdown. *Lancet*. 2020;395:e93–e94. doi:10.1016/S0140-6736(20)31031-X
15. Baldi E, Sechi GM, Mare C, et al. Out-of-hospital cardiac arrest during the Covid-19 outbreak in Italy. *N Engl J Med*. 2020;383:496–498. doi:10.1056/NEJMc2010418
16. Statista. Coronavirus (COVID-19) deaths in Italy as of February 21, 2021, by region. Available from: <https://www.statista.com/statistics/1099389/coronavirus-deaths-by-region-in-italy/>. Accessed February 15, 2021.

Open Access Emergency Medicine**Dovepress****Publish your work in this journal**

The Open Access Emergency Medicine is an international, peer-reviewed, open access journal publishing original research, reports, editorials, reviews and commentaries on all aspects of emergency medicine. The manuscript management system is completely online

and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/open-access-emergency-medicine-journal>