

Diagnosing MonoMAC Syndrome in GATA2 Germline Mutated Myelodysplastic Syndrome via Next-Generation Sequencing in a Patient with Refractory and Complex Infection: Case Report and Literature Review

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Yingying Shen¹
Yuzhu Li²
Hangchao Li²
Qi Liu²
Huijie Dong²
Bo Wang¹
Baodong Ye¹
Shenyun Lin¹
Yiping Shen¹
Dijiong Wu¹

¹Department of Hematology, First Affiliated Hospital of Zhejiang Chinese Medical University, Zhejiang Provincial Hospital of TCM, Hangzhou, Zhejiang Province, People's Republic of China;
²First Clinical Medical College, Zhejiang Chinese Medical University, Zhejiang Provincial Hospital of TCM, Hangzhou, Zhejiang Province, People's Republic of China

Correspondence: Dijiong Wu; Yiping Shen
Department of Hematology, First Affiliated Hospital of Zhejiang Chinese Medical University, No. 54 Youdian Road, Hangzhou, Zhejiang Province, People's Republic of China
Email wudijiong@zcmu.edu.cn; shenyp_@126.com

Abstract: Monocytopenia and mycobacterial infection (MonoMAC) syndrome is a rare disease. Herein, we reported a 65-year-old Asian woman, previously diagnosed with myelodysplastic syndrome (MDS), suffering from recurrent pneumonia, intermittent fever, fatigue, and chest tightness lasting for five months. She was ultimately diagnosed with MonoMAC syndrome with *Mycobacterium kansasii* (*M. kansasii*) infection and GATA2 mutation through metagenomic generation sequencing (mNGS) of peripheral blood specimen, for which she was given anti-NTM therapy. Her situation significantly improved within 2 weeks of therapy. We discussed the clinical features, genetic characteristic, and prognosis of this disorder, aiming to further elucidate this rare syndrome. For MDS/AML patient with recurrent mixed infection and pancytopenia (especially with monocyte absence), MonoMAC syndrome should be highly suspected, and germline mutation and pathogen sequencing should be performed.

Keywords: monocytopenia and mycobacterial infection syndrome, MonoMAC, *M. kansasii*, GATA-2 germline mutation, nontuberculous mycobacteria, next-generation sequencing

Introduction

Monocytopenia and mycobacterial infection (MonoMAC) syndrome is a rare autosomal dominant syndrome associated with monocytopenia, deficiency of B and natural killer (NK) cells, as well as mycobacterial, viral, fungal, and bacterial opportunistic infections. More than 50% of the patients with MonoMAC are diagnosed with myelodysplastic syndrome (MDS) or acute myeloid leukemia (AML).^{1,2} Patients with MonoMAC syndrome typically present with severe or recurrent infection of nontuberculous mycobacterial (NTM) or opportunistic fungal infection;¹ the mortality rate in those patients is 28%.¹

The allogeneic hematopoietic stem cell transplantation (HSCT) has been shown to be an effective treatment to reconstitute the depleted hematopoietic systems and reverse the clinical phenotype seen in affected patients.² However, if the diagnosis is performed too late, the treatment may not be effective. Thus, early diagnosis is crucial. Herein, we diagnosed a MonoMAC syndrome in a MDS patient with GATA2 mutation by next-generation sequencing of the blood specimen, and the mutation was

further confirmed as germline mutation with oral mucosal specimen from her son. To better understand the sensitivity and specificity of a blood specimen by NTM detection, we further reviewed the available reported MonoMAC cases and analyzed the disease's diagnosing process, which may hopefully help clinicians perform earlier diagnoses of this kind of rare disease.

Case Report

A 65-year-old Asian woman was admitted to our clinic in August 2019 due to fever (38.3–39 °C), chest tightness, cough producing sputum, and shortness of breath for one day. Her clinical history could be defined by long-term leukopenia. She was diagnosed with MDS in 2014, after which she received supportive treatments. In 2019, she was admitted to the hospital with increased bone marrow blasts (10% vs

4%) and decreased megakaryocytes with small round separated nuclei (Figure 1A–E, Table 1). She received four courses of amifostine, but her symptoms did not improve. During hospitalization, she was infected with recurrent pneumonia, which could be relieved by multiple antibiotic treatments (Figure 2A and B). She received voriconazole after the last course in July 2019. Two months later, the symptoms of fever and pneumonia re-appeared again (Figure 2C).

On admission, she presented with fever and chest pain, and blood cell count at that time revealed lymphocytopenia and monocytopenia. The blood routine test showed a white blood cell (WBC) count of 1800 cells/mm^3 , a low monocyte count of $0\text{--}0.1 \text{ cells/mm}^3$; a hemoglobin level of 7.6 g/dL ; and a platelet count of $17 \times 10^9/\text{L}$. The absolute count of B cell ($\text{CD}19^+$) was $6 \times 10^6/\text{L}$ ($50\text{--}670 \times 10^6$), NK cell ($\text{CD}3/\text{CD}16+56+$) count was $34 \times 10^6/\text{L}$ (40--




Figure 1 Bone marrow images at the first admission. Pathological change with multi-lineage myelodysplasia, including erythropathy (megaloblastic changes, (A)), granulocytopenia (rod thickening of neutrophils, (B); binuclear, (C)), megakaryocytopenia (binuclear, (D)) and blasts (E).

Table I Changes in Peripheral Blood and Bone Marrow During the Disease Process

Date	Hemogram	Absolute Value of Monocyte (0/dl)	NK Cell Count (CD3-/CD16+56+) ($40\text{--}1000\times 10^6/\text{L}$)	B Cell Count (CD19+) ($50\text{--}670\times 10^6/\text{L}$)	Bone Marrow Blasts%	Myelodysplasia
24 July 2014	WBC 3,000/dl; HB 10.3g/dl PLT 86,000/dl	0	Unknow	Unknow	3	Dysplasia granulocyte and erythrocyte
19 July 2017	WBC 1,500/dl; HB 9.8g/dl PLT 50,000/dl	0	Unknow	Unknow	2.5	Trilineage Dysplasia
11 April 2019	WBC 1,700/dl; HB 8.8g/dl PLT 45,000/dl	0	$34\times 10^6/\text{L}$	$6\times 10^6/\text{L}$	10	Trilineage Dysplasia
4 June 2019	WBC 2,000/dl; HB 8.1g/dl PLT 26,000/dl	0	Not detected	Not detected	5.5	Dysplasia granulocyte and erythrocyte
24 July 2019	WBC 2,300/dl; HB 9.2g/dl PLT 31,000/dl (Platelet transfusion dependent)	0	$19\times 10^6/\text{L}$	$11\times 10^6/\text{L}$	5	Dysplasia granulocyte and erythrocyte

$1000\times 10^6/\text{L}$). The CD4^+ T-cell count was 141 cells/ mm^3 and the CD4/CD8 ratio was 0.99. She also had splenomegaly (4.1 cm in size) and iron overload (ferritin: 2361.4ng/mL). The results of Epstein-Barr virus (EBV)-DNA and cytomegalovirus (CMV)-DNA were negative. Next, the patient was treated with tigecycline, ceftazidime, and voriconazole according to the manifestation of lung computed tomography (CT) scan (Figure 2D). Furthermore, *Klebsiella pneumoniae* infection was confirmed by sputum culture analysis, after which she

received therapy with caspofungin, posaconazole, cefoperazone, linezolid, ganciclovir, sulfamethoxazole, amphotericin B, and polymyxin subsequently. However, no significant improvement was observed with reference to fever and pneumonia. During this time, she developed chest tightness and dyspnea which required continuous high-flow oxygen inhalation. She was not able to undergo a bronchoscopy. Complicated with severe infection, the patient also developed acute heart failure. Eventually, we sequenced the patient's peripheral blood specimens despite




Figure 2 Computed tomography (CT) manifestation of recurrent pulmonary infection. The recurrent pneumonia was observed during the hospitalization (A), and could be relieved by multiple antibiotic treatments (B). (C) showed exacerbation of infection before the detection of mycobacterium kansasii (with air bronchogram and pulmonary consolidation), and the situation did not get improved after multiple treatments (D).

her repeated negative sputum culture, after which the *M. kansasii* infection was confirmed. Consequently, she was given antibiotic therapy, consisting of ethambutol, rifabutin, and clarithromycin as recommended.³ Her situation significantly improved within 2 weeks of therapy (without fever). C-reactive protein (CRP) level remarkably decreased from 250.87 mg/L to an almost normal level. According to the above-mentioned signs and symptoms, the patient was suspected of having MonoMAC syndrome. The next-generation sequencing (NGS, peripheral blood) found a heterozygous mutation in *GATA2* (exon6: c.1126_1128del:p.K376del) along with U2AF1 (exon2:c. C101T:p.S34F), SETBP1 (exon4, cG2602A:pD868N, exon4:cG2608A:pG870S), ASXL1 (exon12:c.G2548T:p. E850X) as well as KMT2D, BRAF, EPPK1, and ETV6. Her son was further confirmed with the same *GATA2* mutation with oral mucosal specimen. Unfortunately, the patient died of carbapenem-resistant *Klebsiella pneumoniae* bloodstream infection on October 2019, 41 days after the MonoMAC diagnosing.

Discussion

MonoMAC syndrome was firstly reported in 2010. Approximately 50% of MonoMAC patients reported so far presented with MDS or AML. Studies have found that MonoMAC syndrome is closely related to *GATA2* germline mutation. Heterozygous mutation in *GATA2* causes the loss of gene function that regulates many aspects of development from hematopoiesis to lymphatic, leading to immunodeficiency and bone marrow failure.⁴ Among individuals with *GATA2* deficiency progressing to MDS/AML, acquired secondary mutations in the ASXL1 that encode chromatin-binding protein ASXL1 have been detected in approximately 30% of cases;⁵ this was also a case of our patient. Up to date, at least 30 cases of MonoMAC in MDS patients have been reported, some of whom were confirmed based on family *GATA2* mutation. Various *GATA* mutation site have been reported, mostly in exon 6 and exon 7, and some in exon 4 and 5^{6–15} (Figure 3). In our case, a *GATA2* mutation (p.K376 del), located in exon6 was found. The results showed that there might not be a close relationship between mutation site and the prognosis. Although HSCT has been shown to be an effective treatment for MonoMAC, the overall survival rate of 57% at 36 months.¹⁶

Early diagnosis improves prognosis and treatment outcomes. Unlike other microbial infections, culturing nontuberculous mycobacterial (NTM) is challenging and requires a long time for analysis (7 days to several weeks),¹⁷ while the rate of




Figure 3 Distribution of *GATA2* germline mutation sites in MonoMAC syndrome. Various site of *GATA* germline mutation had been reported, mostly in exon 6 (45%) and exon 7 (32%), and some in exon 4 (14%) and exon 5 (9%).

positive sputum test is mostly under 35%.¹⁸ In order to increase the specificity and sensitivity of detection, NGS assays are very commonly used for gene screening in those with predisposition syndrome. Usually, bronchoalveolar lavage fluid (BALF) is an ideal specimen for mNGS detection. However, patients with hematological disorders, especially those with severe thrombocytopenia, cannot tolerate bronchoscopy.¹⁹ In our case, we preferred the peripheral blood specimen; other reported specimen including secretion and tissue fluid, were optional samples for mNGS (Table 2).

In our case, the NTM was finally confirmed as *M. kansasii*, which was firstly isolated in 1953.¹⁹ *M. kansasii* is ubiquitous in the environment as a group I non-tuberculous mycobacterium (NTM).²⁰ The *M. kansasii* extrapulmonary disease is rare.²¹ In patients with poor immunity, *M. kansasii* infection may lead to underlying systemic disease or abscess formation,²² with a 1-year mortality rate of 43% and median survival of 71 days.²³ The current therapies for *M. kansasii* infection are similar but not the same. The British Thoracic Society (BTS) recommends daily two-drug therapy with rifampicin (450–600mg/day) and ethambutol (15mg/kg) for 9 months.²⁴ The American Thoracic Society (ATS) and the Infectious Diseases Society of America (IDSA) guidelines recommend daily therapy with isoniazid (5mg/kg/day), rifampicin (10mg/kg/day), ethambutol (15mg/kg/day) and pyridoxine (50mg/d) until sputum culture is negative for at least 12 months.²⁵ The standard treatment regimen is usually effective, but is always time-consuming and not suitable for all the patients.²⁶ Alternative therapies include clarithromycin (500–1000mg/day) plus rifampicin (10mg/kg/day) and ethambutol (15–25mg/kg/day).²⁷ Rifampicin is the main drug of the therapy; ethambutol can prevent acute resistance to rifampicin. If rifampicin resistance occurs, varonones, macrolides,

Table 2 Mutation Sites of the Related Literature in MDS Patients with MonoMAC Syndrome

Case	Study	Age (Years)/ Gender	Pathogenic Specimens Type	Genotyping Specimens	Mutation and Site	Prognosis
1	Portich et al,2020 ⁷	16/F	Unreported	Unreported	Exon6: Chr3:128200759C>T (or alternatively c.1.046G>A), p.Cys349Tyr	Unreported
2	Mendes-de-Almeida,et al,2019 ⁸	43/M	(NTM)Blood culture and arthrocentesis	PB	Exon6: c.1061 C > T; p.T354 M	Died(unknow)
3	Simonis,et al,2018 ⁹	17/F	(<i>Mycobacterium</i>) Skin secretions	BM	Exon6: 1045T>G C349G	Alive(HSCT)
4	Damian,et al,2018 ¹⁰	19/M	(NTM)clinical diagnosis	PB	Exon6: p.C349R	Alive(HSCT)
5	Sologuren, et al,2018 ¹¹	54/M	(H1N1 and <i>Klebsiella pneumoniae</i>)blood test	Unknow	Exon7:R396L	Died(infection)
6	Yamamoto,et al,2018 ¹²	18/M	Unreported	BM and control cells (buccal swab)	Exon4: p.R230Hfs*44	Alive(HSCT)
7	Eguchi,et al,2017 ¹³	14/M	(<i>Mycobacterium kansasii</i>) BM and peripheral blood cultures	Unknow	Exon6: c.1077_1078insA	Alive(HSCT)
8	Vila,et al,2016 ¹⁴	24/F	(NTM)Blood,sputum, BAL cultures	PB	Exon5: (c.1009C>T, p.R337X)	Died(infection)
9	Ganapathi, et al,2016 ¹⁵	39/M	(MAC) unreported	PB & BM	Exon7: c.1192C>T	Unknow
10	Ganapathi, et al,2016 ¹⁵	44/F	(MAC) unreported	PB & BM	Exon6: c.1061C>T	Unknow
11	Ganapathi, et al,2016 ¹⁵	33/M	(MAC) unreported	PB & BM	Exon4: c.243_244delAinsGC	Unknow
12	Ganapathi, et al,2016 ¹⁵	22/M	(<i>M. fortuitum</i>) unreported	PB & BM	Exon7: c.1192C>T	Unknow
13	Ganapathi, et al,2016 ¹⁵	14/F	(MAC) unreported	PB & BM	Exon7: c.1186C>T	Unknow
14	Ganapathi, et al,2016 ¹⁵	32/M	(MAC) unreported	PB & BM	Exon6: c.1061C>T	Unknow
15	Ganapathi, et al,2016 ¹⁵	15/M	(MAC) unreported	PB & BM	Exon4: c.769_778dup	Unknow
16	Ganapathi, et al,2016 ¹⁵	33/M	(MAC) unreported	PB & BM	Exon7: c.1192C>T	Unknow
17	Ganapathi, et al,2016 ¹⁵	45/F	(MAC) unreported	PB & BM	Exon6: c.1018-1G>A	Unknow
18	Ganapathi, et al,2016 ¹⁵	24/M	(MAC) unreported	PB & BM	Exon7: c.1186C>T	Unknow
19	Ganapathi, et al,2016 ¹⁵	17/F	(MAC) unreported	PB & BM	Exon6: c.1099insG	Unknow
20	Ganapathi, et al,2016 ¹⁵	27/M	(<i>M. kansasii</i>) unreported	PB & BM	Exon6: c.1083_1094del12	Unknow
21	Ganapathi, et al,2016 ¹⁵	26/F	(<i>M. kansasii</i>) unreported	PB & BM	Exon5: c.941_951dup	Unknow
22	Ishida, et al,2012 ⁶	19/F	(Severe varicella and/or <i>Salmonella</i> infection) unreported	BM	Exon7: 1187 G > A	Unknow
23	Our case	65/F	(<i>M. kansasii</i>) peripheral blood	BM	Exon6: c.1126> T; p.k376del	Died(infection)

Abbreviations: MAC, *M. avium* intracellulare complex; BM, bone marrow; PB, peripheral blood.

nitroimidazoles and clofazimine can be used as therapeutic options. Physicians should make the therapeutic schedule individually considering the hepatorenal function and bone marrow hematopoiesis during the practice. In our case, Rifampicin, ethambutol and clarithromycin were selected, and the fever, respiratory, constitutional symptoms were improved within 2 weeks.

In conclusion, we reported a case of a rare MDS patient with refractory infection and immune deficiency ascribe to MonoMAC syndrome. The sequencing of NTM and germline GATA2 mutation should be performed as early as possible.

Data Sharing Statement

The data used and/or analyzed during the current study are available from the corresponding author upon a reasonable request.

Ethics Approval and Consent to Participate

This study was approved by the ethical committee of First Affiliated Hospital of Zhejiang Chinese Medical University.

Patient Consent for Publication

Written informed consent was obtained from the patient's son for the publication of clinical results.

Author Contributions

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed to submit to the current journal; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Funding

This study was supported by National Natural Science Foundation of China (NO. 81603573), Zhejiang Traditional Chinese Medicine Scientific Research Foundation (NO. 2020ZB085) and Science and Technology Department of Zhejiang Research Foundation (NO.2019C03047).

Disclosure

The authors declare that they have no competing interests.

References

- Vinh DC, Patel SY, Uzel G, et al. Autosomal dominant and sporadic monocytopenia with susceptibility to mycobacteria, fungi, papilloma-viruses, and myelodysplasia. *Blood*. 2010;115(8):1519–1529. doi:10.1182/blood-2009-03-208629
- Bigley V, Haniffa M, Doulatov S, et al. The human syndrome of dendritic cell, monocyte, B and NK lymphoid deficiency. *J Exp Med*. 2011;208(2):227–234. doi:10.1084/jem.20101459
- Destefano MS, Shoen CM, Cynamon MH. Therapy for Mycobacterium kansasii infection: beyond 2018. *Front Microbiol*. 2018;9:2271. doi:10.3389/fmicb.2018.02271
- Kazenwadel J, Secker GA, Liu YJ, et al. Loss-of-function germline GATA2 mutations in patients with MDS/AML or MonoMAC syndrome and primary lymphedema reveal a key role for GATA2 in the lymphatic vasculature. *Blood*. 2012;119(5):1283–1291. doi:10.1182/blood-2011-08-374363
- Bödör C, Renneville A, Smith M, et al. Germ-line GATA2 p. THR354MET mutation in familial myelodysplastic syndrome with acquired monosomy 7 and ASXL1 mutation demonstrating rapid onset and poor survival. *Haematologica*. 2012;97(6):890–894. doi:10.3324/haematol.2011.054361
- Ishida H, Imai K, Honma K, et al. GATA-2 anomaly and clinical phenotype of a sporadic case of lymphedema, dendritic cell, monocyte, B- and NK-cell (DCML) deficiency, and myelodysplasia. *Eur J Pediatr*. 2012;171(8):1273–1276. doi:10.1007/s00431-012-1715-7
- Portich JP, CONDINO NETO A, Faulhaber GAM. Humoral deficiency in a novel GATA2 mutation: a new clinical presentation successfully treated with hematopoietic stem cell transplantation. *Pediatr Blood Cancer*. 2020;67(9):e28374. doi:10.1002/pbc.28374
- Mendes-de-almeida DP, Andrade FG, Borges G, et al. GATA2 mutation in long stand Mycobacterium kansasii infection, myelodysplasia and MonoMAC syndrome: a case-report. *BMC Med Genet*. 2019;20(1):64. doi:10.1186/s12881-019-0799-6
- Simonis A, Fux M, Nair G, et al. Allogeneic hematopoietic cell transplantation in patients with GATA2 deficiency-a case report and comprehensive review of the literature. *Ann Hematol*. 2018;97(10):1961–1973. doi:10.1007/s00277-018-3388-4
- Damian L, Sauvetre G, Marguet F, et al. Pseudo-sarcoidosis revealing MonoMAC syndrome. *J Clin Immunol*. 2018;38(7):739–741. doi:10.1007/s10875-018-0551-6
- Sologuren I, Martinez-saavedra MT, Sole-violan J, et al. Lethal influenza in two related adults with inherited GATA2 deficiency. *J Clin Immunol*. 2018;38(4):513–526. doi:10.1007/s10875-018-0512-0
- Yamamoto H, Hattori H, Takagi E, et al. [MonoMAC syndrome patient developing myelodysplastic syndrome following persistent EBV infection]. *Rinsho Ketsueki*. 2018;59(3):315–322. doi:10.11406/rinketsu.59.315.Japanese.
- Eguchi K, Ishimura M, Sonoda M, et al. Nontuberculous mycobacteria-associated hemophagocytic lymphohistiocytosis in MonoMAC syndrome. *Pediatr Blood Cancer*. 2018;65(7):e27017. doi:10.1002/pbc.27017
- Vila A, Dapas JI, Rivero CV, et al. Multiple opportunistic infections in a woman with GATA2 mutation. *Int J Infect Dis*. 2017;54:89–91. doi:10.1016/j.ijid.2016.11.408
- Ganapathi KA, Townsley DM, Hsu AP, et al. GATA2 deficiency-associated bone marrow disorder differs from idiopathic aplastic anemia. *Blood*. 2015;125(1):56–70. doi:10.1182/blood-2014-06-580340
- Chu VH, Curry JL, Elghetany MT, et al. MonoMAC versus idiopathic CD4+ lymphocytopenia. Comment to Haematologica. 2011;96(8):1221–5. *Haematologica*. 2012;97(4):e9–11; author reply e2. doi:10.3324/haematol.2011.058669
- Sharma SK, Upadhyay V. Epidemiology, diagnosis & treatment of non-tuberculous mycobacterial diseases. *Indian J Med Res*. 2020;152(3):185–226. doi:10.4103/ijmr.IJMR_902_20

18. Hatta M, Sultan AR, Tandirogang N, et al. Detection and identification of mycobacteria in sputum from suspected tuberculosis patients. *BMC Res Notes*. 2010;3:1–6. doi:10.1186/1756-0500-3-72
19. Sugihara E, Hirota N, Niizeki T, et al. Usefulness of bronchial lavage for the diagnosis of pulmonary disease caused by *Mycobacterium avium-intracellulare* complex (MAC) infection. *J Infect Chemother*. 2003;9(4):328–332. doi:10.1007/s10156-003-0267-1
20. Buhler VB, Pollak A. Human infection with atypical acid-fast organisms; report of two cases with pathologic findings. *Am J Clin Pathol*. 1953;23(4):363–374. doi:10.1093/ajcp/23.4.363
21. Kaustova J, Chmelik M, Ettlova D, et al. Disease due to *Mycobacterium kansasii* in the Czech Republic: 1984–89. *Tuber Lung Dis*. 1995;76(3):205–209. doi:10.1016/S0962-8479(05)80006-1
22. Smith MB, Molina CP, Schnadig VJ, et al. Pathologic features of *Mycobacterium kansasii* infection in patients with acquired immunodeficiency syndrome. *Arch Pathol Lab Med*. 2003;127(5):554–560. doi:10.5858/2003-127-0554-PFOMKI
23. Liu CJ, Huang HL, Cheng MH, et al. Outcome of patients with and poor prognostic factors for *Mycobacterium kansasii*-pulmonary disease. *Respir Med*. 2019;151:19–26. doi:10.1016/j.rmed.2019.03.015
24. Subcommittee of the Joint Tuberculosis Committee of the British Thoracic Society. Management of opportunist mycobacterial infections: Joint Tuberculosis Committee Guidelines 1999. *Thorax*. 2000;55(3):210–218. doi:10.1136/thorax.55.3.210
25. Griffith DE, Aksamit T, Brown-elliott BA, et al. An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. *Am J Respir Crit Care Med*. 2007;175(4):367–416.
26. Santin M, Dorca J, Alcaide F, et al. Long-term relapses after 12-month treatment for *Mycobacterium kansasii* lung disease. *Eur Respir J*. 2009;33(1):148–152. doi:10.1183/09031936.00024008
27. Shitrit D, Priess R, Peled N, et al. Differentiation of *Mycobacterium kansasii* infection from *Mycobacterium tuberculosis* infection: comparison of clinical features, radiological appearance, and outcome. *Eur J Clin Microbiol Infect Dis*. 2007;26(10):679–684. doi:10.1007/s10096-007-0331-3

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>