Psychometric Properties of the Chinese Version of the Cognitions About Body and Health Questionnaire

Shih-Cheng Liao^{1,2}
Wei-Lieh Huang²⁻⁴

¹Department of Psychiatry, National Taiwan University Hospital, Taipei, Taiwan; ²Department of Psychiatry, College of Medicine, National Taiwan University, Taipei, Taiwan; ³Department of Psychiatry, National Taiwan University Hospital Yunlin Branch, Yunlin, Taiwan; ⁴Graduate Institute of Clinical Medicine, College of Medicine, National Taiwan University, Taipei, Taiwan

Introduction: The Cognitions About Body and Health Questionnaire (CABAH) is a self-rating questionnaire measuring cognitions about health. The psychometric properties of its Chinese version and its performance on evaluating somatic symptom disorder (SSD) were examined in this study.

Methods: After translating the CABAH into Chinese, we collected data of the CABAH and another four questionnaires (Patient Health Questionnaire-15, Health Anxiety Questionnaire, Beck Depression Inventory-II, Beck Anxiety Inventory) in 208 SSD patients and 197 healthy individuals. We used exploratory factor analysis (EFA) to explore the structure of the Chinese CABAH. Internal consistency and criterion-related validity were analyzed. An independent *t*-test and receiver operating characteristic (ROC) curve analysis were used to evaluate the performance of the CABAH for evaluating SSD.

Results: Six factors were suggested by EFA. Five (bodily weakness, health habits, catastrophizing interpretation of specific bodily complaints, somatosensory amplification, catastrophizing interpretation of general bodily complaints) are conceptually associated with the original CABAH. The sixth factor (reverse, with three items) is different from the original construct. Cronbach's alpha for the CABAH was 0.885. The CABAH score was moderately correlated with scores of the other four questionnaires. Scores for the whole CABAH and scores of the six factors were all significantly higher in SSD patients than in healthy individuals. The results of ROC curve analysis were as follows: area under the curve=0.700; suggested cutoff=58/59; Youden's *J*=0.295.

Conclusion: The reliability and validity of the Chinese CABAH were fair, although the three items in the reverse factor should be interpreted cautiously.

Keywords: Cognitions About Body and Health Questionnaire, psychometric properties, bodily weakness, somatic symptom disorder

Introduction

Somatoform disorders are defined in the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) for the individuals with medically unexplained physical symptoms. In the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5), the name of this diagnostic category was changed to somatic symptom and related disorders, in which the psychological features (such as catastrophizing cognitive style and health anxiety) are emphasized to replace the medically unexplained criteria.^{1,2} Somatic symptom disorder (SSD), the most common diagnosis of this category, has a criterion (criterion B) for the

Correspondence: Wei-Lieh Huang
Department of Psychiatry, National
Taiwan University Hospital Yunlin Branch
Email weiliehuang@gmail.com

cognitive, affective or behavioral features of body and health.³ Although there have been questionnaires developed for measuring such psychological features (such as Somatic Symptom Disorder–B Criteria Scale, SSD-12)⁴ or questionnaires that examine associations with SSD criterion B (such as Health Anxiety Questionnaire, HAQ),^{2,5} these tools do not focus on the cognitive domain about health. The cognitive aspect of the somatic symptoms has gained more and more attention in the current diagnostic system. For example, the catastrophizing cognitive style is listed as one feature of SSD, whereas excessive attention to the body is described in bodily distress disorder, a similar construct to SSD in the International Classification of Diseases 11th Revision.⁶ Several other cognitive features have been reported to be associated with somatic symptoms, including memory bias, the feature of expectation, and health attitude.⁷ Perceptual features, such as somatosensory amplification, are also associated with cognition.⁷ Therefore, a quantitative tool focusing on the cognitive domain should be helpful for clarifying the relations between the above features and the contemporary somatic diagnoses. To the best of our knowledge, the Cognitions About Body and Health Questionnaire (CABAH) is one of the most comprehensive tools for measuring cognition about health.⁸

The CABAH was developed by Rief et al;⁸ it has different versions with 31 and 39 items, and the 31-item version was originated from the previous 68-item version. A four-point Likert scale (0–3) was used in the CABAH, with a higher value indicating higher severity.⁸ The 31-item CABAH includes catastrophizing interpretation of bodily complaints (14 items), autonomic sensations (4 items), bodily weakness (6 items), intolerance of bodily complaints (4 items) and health habits (3 items). The 39-item version of the CABAH additionally incorporates the Somatosensory Amplification Scale (10 items, two overlaps with autonomic sensations) developed by Barsky et al.^{8,9} Two items in the CABAH were scored in reverse. The scoring principle is to sum the scores of all items; the scores of the different factors can be estimated separately.^{8,10} According to Rief et al.'s research,⁸ the internal consistency for the whole CABAH was 0.90 and 0.67–0.88 for the different factors. The correlations between several factors and the Whiteley Index (WI, a measurement of hypochondriasis) score were 0.04–0.45. These psychometric properties indicate CABAH to be a useful tool, but there was no Chinese version of CABAH in the past.

The CABAH has been widely applied in the psychosomatic field. Comparing patients with somatoform disorder, patients with hypochondriasis had significantly higher CABAH scores.¹⁰ After receiving cognitive-behavioral therapy (CBT), CABAH scores of individuals with health anxiety significantly improved.¹¹ For patients with hypochondriasis, responders to CBT had significantly lower CABAH scores than non-responders.¹² CABAH scores were associated with functioning and quality of life in SSD patients.¹³ Apart from the above findings, we are also interested in whether the cognitive feature in the diagnostic criteria of SSD can be presented via the CABAH score, and whether the CABAH can be used for screening, diagnosing or evaluating SSD. This issue can be explored by comparing CABAH scores in individuals with and without SSD. A similar approach has been adopted for investigating other psychosomatic questionnaires, such as the Patient Health Questionnaire-15 (PHQ-15), WI-7 and HAQ.^{5,14,15}

There are two major aims in the present study. The first is to explore the psychometric properties of the Chinese version of the CABAH. The second is to investigate whether the CABAH is suitable for differentiating between SSD and healthy individuals and whether it can be applied to evaluate SSD.

Methods

Translation of the CABAH

We contacted the author of the CABAH to obtain permission for translating its 39-item version (which includes the Somatosensory Amplification Scale) into Chinese.⁸ The English version of the CABAH was initially translated into Chinese (forward-translation) by one author, then two authors discussed the translation and made necessary changes. Three psychosomatic patients were invited to read the questionnaire in order to limit any possible misunderstanding. We then requested a bi-lingual (English and Chinese) expert to perform a backward-translation (Chinese to English). When the two authors both considered that the meaning of sentences after backward-translation changed, we assumed that the sentences of forward-translation were not understood correctly; the Chinese sentences were revised under this circumstance. At the end of this process, we attained the Traditional Chinese version (based on the vocabulary in Taiwan) of the CABAH, which was used in the following procedure. This material originally appeared in English as Cognitive

aspects of hypochondriasis and the somatization syndrome. Copyright © 1998 by American Psychological Association. Translated and reproduced with permission. American Psychological Association is not responsible for the quality or accuracy of this translation. This translation cannot be reproduced or distributed further without prior written permission.

Procedure of the Psychometric Analysis

The CABAHA data from two studies were collected and analyzed. The Institutional Review Board of National Taiwan University Hospital approved the execution of this analysis (approval number: 201808047RINB). The aims of the two studies were both to investigate the psychological and biological features of SSD patients, some results of these studies have been published.^{13,16} All participants completed informed consent before gathering the data; this analysis was conducted in accordance with the declaration of Helsinki. Enrollment was from 2016 to 2020 at the National Taiwan University Hospital Yunlin Branch (NTUHYL). The inclusion criteria of the two studies are as follows: (1) patients with SSD, anxiety, or depression; (2) healthy individuals. The exclusion criteria of them are as follows: (1) age lower than 20 or higher than 70 years old; (2) having psychotic symptoms; (3) having difficulty completing questionnaires. To examine whether the CABAHA is suitable for evaluating SSD, two populations were included in this analysis: SSD patients who visited psychiatric clinics at the NTUHYL and healthy individuals living in communities near the NTUHYL. Because there is not yet a structuralized tool for performing DSM-5-based diagnostic interview in Taiwan (such as The Structured Clinical Interview for DSM-5), we developed some semi-structured questions for diagnosing SSD; the content has been described in our another study.¹⁷ We found 410 eligible subjects at first but five did not complete the questionnaire, leaving a final sample size of 405 for statistical analysis (208 SSD patients and 197 healthy individuals; mean age 44.16 ± 12.42 years, 139 male). Their detailed demographic data are shown in [Supplementary Table 1](#). All included subjects completed the CABAHA and also another four questionnaires commonly used in the psychosomatic field in Taiwan: the PHQ-15, HAQ, Beck Depression Inventory-II (BDI-II) and Beck Anxiety Inventory (BAI). PHQ-15 and HAQ have been applied for screening SSD in a previous study.² The data of all questionnaires from each subject were gathered in one single day.

Other Questionnaires

Scores of the PHQ-15, HAQ, BDI-II and BAI were gathered for examining the criterion-related validity (concurrent validity) of CABAHA.

The PHQ-15, developed by Kroenke et al.,¹⁸ measures the types and severity of somatic symptoms. It has 15 items; the score for each item is 0–2 (higher score indicates more severe symptoms). The items can be separated into three major factors: cardiopulmonary, pain–fatigue and gastrointestinal.¹⁴ The internal consistency (Cronbach's alpha) of the Chinese PHQ-15 was 0.861.

The authors of the HAQ are Lucock and Morley.¹⁹ HAQ is a 21-item questionnaire aimed at measuring health anxiety and hypochondriacal ideation, with scoring on a four-point Likert scale (0–3). There were three major factors in the Chinese version of the HAQ: excessive worry about health and illness; the extent to which symptoms interfere with a person's life; and reassurance-seeking behavior. Its Cronbach's alpha was 0.943.⁵

The BDI-II was developed by Beck.²⁰ It measures the level of depression, with coverage of cognitive/affective and somatic/vegetative dimensions. There are 21 items (score 0–3 for each) and Cronbach's alpha of the Chinese BDI-II was 0.94.²¹

The BAI was also developed by Beck.²² BAI emphasizes the somatic features of anxiety. It has 21 items, with scoring on a four-point Likert scale (0–3). Cronbach's alpha of the Chinese BAI was 0.95.²³

Statistical Analysis

We firstly used confirmatory factor analysis to examine the structure reported by Rief et al.,⁸ but the model fit was not satisfying and did not obviously improve under minor modification (it was described in detail in Results). Therefore, exploratory factor analysis (EFA) was used to clarify the structure of the Chinese version of the CABAHA. Principal component analysis with the Kaiser criterion was adopted for determining the number of factors to extract (the scree plot was shown in [Supplementary Figure 1](#)). We then used principal axis factoring with direct oblimin rotation to generate a pattern matrix. For comparison with the original CABAHA, all items were classified into one factor according to the highest loadings (even when the highest loadings were lower than 0.4). We also performed item analysis for clarifying the features of each item. We then estimated internal consistency with Cronbach's alpha for the whole CABAHA and for each

factor; above 0.6–0.7 and above 0.8 are usually considered as the acceptable and good values of Cronbach's alpha, respectively.²⁴ With regard to criterion-related validity, we used Pearson's correlation analysis to calculate the association between the scores of the whole CABAHA/six factors and four questionnaires (PHQ-15, HAQ, BDI-II, BAI). Moreover, the difference in the scores of the whole CABAHA/six factors between SSD patients and healthy individuals was compared using an independent *t*-test. Finally, receiver operating characteristic (ROC) curve analysis was adopted to examine whether the CABAHA was suitable for diagnosing SSD; Youden's J (sensitivity + specificity – 1) was calculated for determining the cutoff (the score with the highest Youden's J was considered as the optimal cutoff). Two-sided analysis with an alpha value of 0.05 was set for the *t*-test and correlation analysis. We used SPSS 25 and AMOS 25 (IBM, USA) for performing these analyses.

Results

Confirmatory Factor Analysis According to the Original Structure

The model fit statistics of confirmatory factor analysis according to the structure of the original CABAHA were: $X^2=1827.138$, $X^2/df=2.667$, $NFI=0.646$, $RFI=0.597$, $IFI=0.745$, $CFI=0.740$, $TLI=0.704$. These model fits were unsatisfactory, and EFA was warranted.

Exploratory Factor Analysis

The results of EFA revealed that there were six factors in the Chinese CABAHA (Table 1). According to their features and similarity with the original CABAHA, we named them: bodily weakness (factor 1; items 1, 4, 5, 11, 14, 17, 23, 25, 30), health habits (factor 2; items 6, 9, 13, 18, 19, 26), catastrophizing interpretation of specific bodily complaints (factor 3; items 2, 10, 15, 20, 21, 24, 27, 28, 32), somatosensory amplification (factor 4; items 33, 34, 35, 36, 37, 38, 39), catastrophizing interpretation of general bodily complaints (factor 5; items 7, 8, 12, 16, 29) and reverse (factor 6; items 3, 22, 31). The detailed content of each factor and item (including a comparison of the structure of the Chinese and original CABAHA versions) is shown in Supplementary Material.

Internal Consistency

Cronbach's alpha for the whole questionnaire (CABAHA total) was good (0.885) and for factors 1–5 it was 0.643–

0.826, which is acceptable. However, Cronbach's alpha for factor 6 (reverse) was quite low (–0.023). The results are shown in Table 2.

Item Analysis

Supplementary Table 2 gives the results of item analysis. In this step, we estimated the following values for all items: mean, variance, coefficient of skewness, correlation coefficient with CABAHA total score and Cronbach's alpha of CABAHA total after excluding this item. There was obvious skewness for item 3. All items except item 3 and item 22 were significantly correlated with CABAHA total score. With regard to Cronbach's alpha of CABAHA total after excluding an item, several items were related with elevating internal consistency (items 3, 19, 22, 31); these items were highly overlapping with the factor 6 (reverse).

Criterion-Related Validity

The correlations between the scores of CABAHA total, the six factors and other questionnaires are shown in Table 3. The score of CABAHA total was moderately correlated with the scores of the four questionnaires ($r = 0.416$ – 0.535). Among the six factors, factor 1 (bodily weakness) score revealed the highest correlations with the four questionnaires ($r = 0.591$ – 0.637) scores. Factor 3 (catastrophizing interpretation of specific bodily complaints), factor 4 (somatosensory amplification) and factor 5 (catastrophizing interpretation of general bodily complaints) scores were significantly correlated with the four questionnaires ($r = 0.2$ – 0.4) scores. Factor 2 (health habits) score was only significantly correlated with the HAQ score ($r = 0.230$). Although factor 6 (reverse) score was significantly correlated with the HAQ and BAI scores, the correlation coefficients were low (0.106 and 0.124, respectively).

Comparison of CABAHA Scores in SSD Patients and Healthy Individuals

Table 4 and Supplementary Table 3 show the results from the comparison of CABAHA total/six factors and all item scores, respectively. SSD patients showed significantly higher scores for CABAHA total and all six factors than healthy individuals, especially factor 1 (bodily weakness, $t = 12.595$). For most items, SSD patients had significantly higher scores, but there were exceptions: the scores on items 19 and 33 were significantly higher in healthy individuals;

Table 1 Exploratory Factor Analysis of the Chinese Version of CABAQ

	Factor 1	Factor 2	Factor 3	Factor 4	Factor 5	Factor 6
Factor 1 (bodily weakness)						
CABAQ25	0.643	-0.050	0.041	0.183	0.035	0.046
CABAQ11	0.592	-0.083	0.015	0.120	0.189	-0.048
CABAQ14	0.561	-0.067	0.151	0.021	0.110	0.072
CABAQ01	0.549	0.085	-0.065	0.192	0.129	-0.182
CABAQ17	0.545	0.127	0.286	-0.019	-0.207	0.077
CABAQ04	0.423	-0.001	0.107	0.067	0.263	0.114
CABAQ05	0.376	0.344	-0.049	0.021	0.291	-0.040
CABAQ30	0.318	-0.101	0.302	0.262	0.062	0.055
CABAQ23	0.264	-0.201	0.155	0.210	-0.092	0.163
Factor 2 (health habits)						
CABAQ19	-0.108	0.655	-0.009	0.043	-0.165	0.198
CABAQ13	-0.082	0.584	0.082	0.136	0.141	-0.108
CABAQ09	0.079	0.443	0.002	0.101	0.341	-0.234
CABAQ26	0.036	0.416	0.054	0.073	0.094	0.034
CABAQ06	0.169	0.322	0.181	0.001	0.303	-0.307
CABAQ18	0.076	0.258	0.243	0.119	0.213	-0.156
Factor 3 (catastrophizing interpretation of bodily complaints, specific diseases)						
CABAQ10	-0.142	0.020	0.590	0.031	0.211	-0.181
CABAQ15	0.209	0.012	0.533	-0.014	0.022	-0.079
CABAQ02	0.135	-0.007	0.525	0.020	-0.017	-0.170
CABAQ27	0.033	0.007	0.510	-0.060	0.244	-0.060
CABAQ28	0.235	-0.035	0.509	0.134	-0.132	0.052
CABAQ32	-0.097	0.009	0.460	0.048	-0.073	0.066
CABAQ21	0.239	0.254	0.363	-0.022	-0.200	-0.036
CABAQ24	0.030	0.122	0.354	0.093	0.055	0.153
CABAQ20	0.107	0.120	0.332	0.078	0.254	0.088
Factor 4 (somatosensory amplification)						
CABAQ35	0.045	0.245	0.062	0.589	-0.038	0.035
CABAQ38	0.028	0.017	0.205	0.571	-0.209	-0.196
CABAQ36	0.130	-0.069	0.019	0.548	0.089	0.025
CABAQ39	0.110	0.036	0.091	0.477	-0.090	-0.178
CABAQ34	0.071	0.141	-0.183	0.451	0.210	0.117
CABAQ37	0.131	-0.018	0.084	0.427	0.058	-0.013
CABAQ33	-0.178	0.078	-0.122	0.348	0.094	0.056
Factor 5 (catastrophizing interpretation of bodily complaints, general)						
CABAQ08	0.057	0.010	0.072	0.134	0.597	0.014
CABAQ07	0.159	0.032	0.017	-0.052	0.548	0.047
CABAQ12	-0.068	0.190	0.012	0.061	0.493	0.011
CABAQ16	-0.027	0.056	0.258	0.237	0.335	0.005
CABAQ29	0.066	-0.180	0.125	0.217	0.226	0.125
Factor 6 (reverse)						
CABAQ31	0.303	-0.141	-0.154	0.058	-0.046	-0.316
CABAQ22	-0.153	-0.033	0.018	0.059	0.024	-0.296
CABAQ03	-0.050	-0.022	-0.051	0.037	0.054	0.251

Abbreviation: CABAQ, Cognitions About Body and Health Questionnaire.

Table 2 Internal Consistency of the Chinese Version of CABAH, Including the Whole Questionnaire and 6 Factors

	Internal Consistency (Cronbach's α)
CABAH total	0.885
Factor 1 (bodily weakness)	0.826
Factor 2 (health habits)	0.741
Factor 3 (catastrophizing interpretation of bodily complaints, specific diseases)	0.774
Factor 4 (somatosensory amplification)	0.728
Factor 5 (catastrophizing interpretation of bodily complaints, general)	0.643
Factor 6 (reverse)	−0.023

Abbreviation: CABAH, Cognitions About Body and Health Questionnaire.

and the scores on items 3, 12, 13, 16, 18, 21, 22, 32 and 35 did not reveal the significant inter-group difference.

ROC Curve Analysis

The results of the ROC curve analysis are shown in Figure 1. The area under the curve was 0.700, which is acceptable, and the suggested cutoff was 58/59; however, Youden's J equaled 0.295 (sensitivity 0.615, specificity

Figure 1 ROC curve analysis for the Chinese version of CABAH on detecting SSD. Area under curve=0.700, suggested cutoff 58/59, Youden's J =0.295.

Abbreviations: ROC, receiver operating characteristic; CABAH, Cognitions About Body and Health Questionnaire; SSD, somatic symptom disorder.

0.680), which means that using the CABAH alone for diagnosing SSD is unsatisfactory.

Discussion

Compared with the structure of the original CABAH, the Chinese version of CABAH revealed high similarity in the following three factors: bodily weakness, health habits and

Table 3 Criterion-Related Validity of the Chinese Version of CABAH

	Factor 1	Factor 2	Factor 3	Factor 4	Factor 5	Factor 6	PHQ-15	HAQ	BDI-II	BAI
CABAH total	0.815***	0.661***	0.772***	0.736***	0.691***	0.103*	0.442***	0.535***	0.416***	0.493***
Factor 1	I	0.321***	0.559***	0.491***	0.457***	0.012	0.606***	0.611***	0.591***	0.637***
Factor 2		I	0.437***	0.417***	0.474***	−0.054	0.028	0.230***	−0.004	0.081
Factor 3			I	0.416***	0.392***	−0.054	0.274***	0.295***	0.262***	0.314***
Factor 4				I	0.455***	0.029	0.299***	0.379***	0.279***	0.325***
Factor 5					I	0.028	0.257***	0.359***	0.256***	0.293***
Factor 6						I	0.081	0.106*	0.056	0.124*

Notes: * $p < 0.05$, *** $p < 0.001$.

Abbreviations: CABAH, Cognitions About Body and Health Questionnaire; PHQ-15, Patient Health Questionnaire; HAQ, Health Anxiety Questionnaire; BDI-II, Beck Depression Inventory-II; BAI, Beck Anxiety Inventory.

Table 4 Comparison of the Chinese Version of CABAH Scores in Patients with SSD and Healthy Individuals

	SSD (n=208)		Healthy (n=197)		Statistics	
	Mean	SD	Mean	SD	t	p value
CABAH total	62.51	13.06	52.25	14.03	7.624	<0.001***
Factor 1 (bodily weakness)	15.09	4.60	9.47	4.34	12.595	<0.001***
Factor 2 (health habits)	10.25	3.16	9.57	3.46	2.069	0.039*
Factor 3 (catastrophizing interpretation of bodily complaints, specific diseases)	9.61	4.20	8.20	4.30	3.323	0.001**
Factor 4 (somatosensory amplification)	11.89	3.69	10.50	3.45	3.915	<0.001***
Factor 5 (catastrophizing interpretation of bodily complaints, general)	10.50	2.46	9.65	2.63	3.367	0.001**
Factor 6 (reverse)	5.17	2.04	4.81	1.20	2.189	0.029*

Notes: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Abbreviations: CABAH, Cognitions About Body and Health Questionnaire; SSD, somatic symptom disorder.

somatosensory amplification.⁸ The factor of catastrophizing interpretation of bodily complaints, which has the most items in the original CABAHA, can be separated into two factors in our sample: catastrophizing about specific and general bodily complaints. The other two factors in the original CABAHA, autonomic sensations and intolerance of bodily complaints, are not so distinct in our study. In this analysis, we found another factor consisting of three items, two of which were reversely scored and showed negative loadings in the pattern matrix. We have carefully checked the coding and confirm that the negative loadings did not originate from the coding error. We named this factor “reverse”; its three items showed problems also in other analyses besides EFA. Moreover, low loadings (<0.3) were found in several items, such as in item 23 (I can sometimes hear my pulse or my heartbeat throbbing in my ear) of factor 1 (bodily weakness), in item 18 (If I do not observe my body often, I could become seriously ill without noticing it) of factor 2 (health habits) and in item 29 (I hate to be too hot or too cold) of factor 5 (catastrophizing interpretation of general bodily complaints). These items had relatively low convergent validity and did not belong to the similar factors of the original CABAHA; therefore, it is rational that they were less connected with other items of the same factors.⁸ Items with low factor loadings are sometimes deleted during the development of questionnaires; however, because the full version of CABAHA was used in most published articles, we chose to preserve the full CABAHA in these analyses. If performing an additional confirmatory factor analysis after excluding items with low factor loadings, the model fit would improve ($X^2=278.353$, $X^2/df=2.554$, $NFI=0.848$, $RFI=0.786$, $IFI=0.901$, $CFI=0.899$, $TLI=0.858$).

The internal consistency of 0.855 for the whole CABAHA is a good level and similar to that of the original version (0.9).⁸ Separating the different factors, factor 1 (bodily weakness) has the highest internal consistency of 0.826, followed by 0.6–0.8 for factors 2–5; these values are similar to those for the original CABAHA.⁸ However, the internal consistency of factor 6 (reverse) was quite low (-0.23); this may be explained by the one positive (item 3) and two negative loadings (items 22, 31). Removing these three items from analysis would lead to higher internal consistency; these items also showed lower correlations with CABAHA total scores or high levels of skewness. Reverse item performance in item analysis is often not as good as other items in psychometric studies, so the results are not surprising.²⁵ With regard to item 3, we suppose the

finding to be associated with the translated sentence. The original content of item 3 is “I am healthy when I do not have any bodily sensations”; this means that the healthy cognition exists “only when” the subjects did not feel any discomfort; however, we translated this sentence directly and did not additionally emphasize the concept “only when”, so some subjects might have misunderstood this statement. In summary, items 3, 22 and 31 of the Chinese CABAHA should be used and interpreted cautiously.

The CABAHA total scores were moderately correlated with the scores of four commonly used questionnaires in the psychosomatic field. This implies that the constructs in the CABAHA cannot be replaced by other quantitative measurements. Due to the coexisting tendency of somatic distress, depression, anxiety and hypochondriacal ideation, it is rational that the correlations were not very low.^{5,15} Factor 1 (bodily weakness) scores showed the highest correlations with all four questionnaires scores. A similar finding was reported in the original CABAHA; the bodily weakness factor score showed the highest correlation with the WI score than the other factors.⁸ This implies that “weakness” may be an overlapping concept with both somatic distress and negative emotions. On the other hand, factor 2 (health habits) seemed quite different from all the constructs of the other questionnaires; only the HAQ score revealed a significant (but low) correlation with it. An explanation for this is that health habits are features that transit from cognition to behavior, which are not measured in the four questionnaires. Whether a measurement of illness behavior (such as the Scale for the Assessment of Illness Behavior) shows a higher association with the health habits factor warrants further exploration.²⁶

Scores of the CABAHA total and of the six factors were significantly higher in SSD patients than in healthy individuals, which is quite rational. Score of factor 6 (reverse) was also significantly higher in SSD patients; this phenomenon indicates that although the reverse items had a different presentation approach, not all subjects misunderstood their meanings. The inter-group difference was highest for factor 1 (bodily weakness), implying that the cognition about “weakness” is the clearest feature of SSD patients in Taiwan; thus, “weakness” questions may be helpful for the clinical purpose of SSD screening. “Neurasthenia” is still a commonly used term in Taiwan and in China; its original meaning is about the weakness.²⁷ Therefore, the emphasis on bodily weakness may have a cultural meaning, especially in Chinese society.

Besides, the cognition about weakness may be considered as an extended criterion of SSD or similar diagnoses. From ROC curve analysis, the low Youden's *J* indicates that the sensitivity and specificity when applying CABAHA alone for diagnosing SSD are not satisfactory. It is rational because only part of the SSD criteria (the "cognitive" criterion B1) can be assessed by CABAHA. Some studies reported that combining different questionnaires (such as one for SSD criterion A and others for criterion B) may generate better performance in ROC curve analysis.^{2,28,29} We also considered this approach and found when combining the scores of CABAHA, PHQ-15 and HAQ, the area under the curve increased from 0.700 to 0.859 (Supplementary Figure 2).

Several limitations of this study need to be discussed. First, we did not collect data on test–retest reliability, therefore we cannot establish whether the Chinese CABAHA is temporally stable. Second, we did not include data of non-SSD patients in the psychiatric clinic or psychosomatic patients in consultation–liaison situations, therefore, the suitability of applying the Chinese CABAHA for these populations is not clear.

In sum, there are several implications of the present study. First, the structures of CABAHA may be different in Taiwan and in Europe. Among the dimensions of somatic symptoms, PHQ-15, a questionnaire measuring the level of somatic distress, was found to have distinct structures in Chinese and European samples.³⁰ Second, using only CABAHA is not suitable for diagnosing SSD; the questionnaire SSD-12 should be the better option for this purpose.³¹ Different from the previous construct somatoform disorders, the concept "medically unexplained" has been removed in SSD; therefore, SSD may co-exist in patients with medical comorbidity.^{32,33} SSD-12 has been found to effectively diagnose SSD in a population with medical comorbidity.³¹ Third, though diagnosing SSD is not the strength of CABAHA, its abundant cognitive measurements may still have clinical values. A study revealed that SSD had high complexity and was more complex than some other diseases, such as multiple sclerosis and rheumatoid arthritis.³⁴ It indicates further classification or detailed evaluation of SSD to be needed; CABAHA can be an option for these aims. Another study showed that when patients with somatic symptom and related disorders had comorbid depression, they had poor neurocognitive performance; it may explain why some patients easily dropped out from CBT.³⁵ Besides, the different diagnoses in somatic symptoms and related disorders may have

distinct neurocognitive performance. For example, patients with conversion disorder were found to have relatively poor processing speed.³⁶ The above features may be explored by using CABAHA.

This study is a first attempt at clarifying the psychometric features of the Chinese version of the CABAHA. Besides providing information on the structure of the Chinese CABAHA and the suitability of items, we also investigated the possibility of applying the CABAHA to the evaluation of SSD. In general, the Chinese CABAHA has fair reliability and validity, although three items (items 3, 22, 31) in factor 6 (reverse) should be used and interpreted cautiously. We expect that the CABAHA can be used more extensively in Chinese society; apart from clinical purposes, it may be beneficial for showing the potential cultural difference between Eastern and Western populations.

Data Accessibility

Data of this study are not shared.

Acknowledgments

We thank Professor Winfried Rief for providing us permission to translate the Cognitions About Body and Health Questionnaire into the Chinese version. We also thank Professor Arthur Joseph Barsky III for providing us permission to translate the Somatosensory Amplification Scale into the Chinese version.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

This work was supported by the National Taiwan University Hospital Yunlin Branch (grant numbers NTUHYL105.X002, NTUHYL108.X001).

Disclosure

The authors report no conflicts of interest in this work.

References

- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5th ed. Arlington, VA: American Psychiatric Association; 2013.
- Huang WL, Chen IM, Chang FC, Liao SC. Somatic symptom disorder and undifferentiated somatoform disorder, which is broader? Response to "Clinical value of DSM IV and DSM 5 criteria for diagnosing the most prevalent somatoform disorders in patients with medically unexplained physical symptoms (MUPS)". *J Psychosom Res*. 2016;89:114–115. doi:10.1016/j.jpsychores.2016.07.014
- Rief W, Martin A. How to use the new DSM-5 somatic symptom disorder diagnosis in research and practice: a critical evaluation and a proposal for modifications. *Annu Rev Clin Psychol*. 2014;10:339–367. doi:10.1146/annurev-clinpsy-032813-153745
- Toussaint A, Murray AM, Voigt K, et al. Development and validation of the somatic symptom disorder-B Criteria Scale (SSD-12). *Psychosom Med*. 2016;78(1):5–12. doi:10.1097/PSY.0000000000000240
- Chen IM, Huang WL, Lee MB, Liao SC. Chinese Version of Health Anxiety Questionnaire: psychometric properties and its association with DSM-IV/DSM-5 systems. *Formosan J Med*. 2019;23:21–30.
- Gureje O, Reed GM. Bodily distress disorder in ICD-11: problems and prospects. *World Psychiatry*. 2016;15(3):291–292. doi:10.1002/wps.20353
- Rief W, Broadbent E. Explaining medically unexplained symptoms-models and mechanisms. *Clin Psychol Rev*. 2007;27(7):821–841. doi:10.1016/j.cpr.2007.07.005
- Rief W, Hiller W, Margraf J. Cognitive aspects of hypochondriasis and the somatization syndrome. *J Abnorm Psychol*. 1998;107(4):587–595. doi:10.1037/0021-843X.107.4.587
- Barsky AJ, Wyshak G, Klerman GL. The somatosensory amplification scale and its relationship to hypochondriasis. *J Psychiatr Res*. 1990;24(4):323–334. doi:10.1016/0022-3956(90)90004-A
- Hiller W, Rief W, Fichter MM. Dimensional and categorical approaches to hypochondriasis. *Psychol Med*. 2002;32(4):707–718. doi:10.1017/S0033291702005524
- Newby JM, Mahoney AEJ, Mason EC, Smith J, Uppal S, Andrews G. Pilot trial of a therapist-supported internet-delivered cognitive behavioural therapy program for health anxiety. *Int Interventions*. 2016;6:71–79. doi:10.1016/j.invent.2016.09.007
- Hiller W, Leibbrand R, Rief W, Fichter MM. Predictors of course and outcome in hypochondriasis after cognitive-behavioral treatment. *Psychother Psychosom*. 2002;71(6):318–325. doi:10.1159/000065990
- Liao SC, Ma HM, Lin YL, Huang WL. Functioning and quality of life in patients with somatic symptom disorder: the association with comorbid depression. *Compr Psychiatry*. 2019;90:88–94. doi:10.1016/j.comppsych.2019.02.004
- Liao SC, Huang WL, Ma HM, et al. The relation between the patient health questionnaire-15 and DSM somatic diagnoses. *BMC Psychiatry*. 2016;16(1):351. doi:10.1186/s12888-016-1068-2
- Tu CY, Liao SC, Liu CY, et al. Application of the Chinese Version of the Whiteley Index-7 for Detecting DSM-5 Somatic symptom and related disorders. *Psychosomatics*. 2016;57(3):283–291. doi:10.1016/j.psym.2015.12.010
- Huang WL, Liao SC. Subgrouping somatic symptom disorder: an analysis based on the diagnostic criteria for psychosomatic research. *Psychother Psychosom*. 2018;87(5):316–318. doi:10.1159/000492227
- Tu CY, Liu WS, Chen YF, Huang WL. Is severity a clinically meaningful specifier of somatic symptom disorder? *J Psychosom Res*. 2020;133:110108. doi:10.1016/j.jpsychores.2020.110108
- Kroenke K, Spitzer RL, Williams JB. The PHQ-15: validity of a new measure for evaluating the severity of somatic symptoms. *Psychosom Med*. 2002;64(2):258–266. doi:10.1097/00006842-200203000-00008
- Luccock MP, Morley S. The health anxiety questionnaire. *Br J Health Psychol*. 1996;1(2):137–150. doi:10.1111/j.2044-8287.1996.tb00498.x
- Beck AT, Steer RA, Ball R, Ranieri W. Comparison of Beck Depression Inventories-IA and -II in psychiatric outpatients. *J Pers Assess*. 1996;67(3):588–597. doi:10.1207/s15327752jpa6703_13
- Lu ML, Che HH, Chang SW, Shen WW. Reliability and validity of the Chinese version of the Beck Depression Inventory-II. *Taiwanese J Psychiatry*. 2002;16(4):301–310.
- Beck AT, Epstein N, Brown G, Steer RA. An inventory for measuring clinical anxiety: psychometric properties. *J Consult Clin Psychol*. 1988;56(6):893–897. doi:10.1037/0022-006X.56.6.893
- Che HH, Lu ML, Chen HC, Chang SW, Li YC. Validation of the Chinese Version of the Beck Anxiety Inventory. *Formosan J Med*. 2006;4:447–454.
- Ursachi G, Horodnic IA, Zait A. How reliable are measurement scales? External factors with indirect influence on reliability estimators. *Procedia Economics Finance*. 2015;20:679–686. doi:10.1016/S2212-5671(15)00123-9
- Swain SD, Weathers D, Niedrich RW. Assessing three sources of misresponse to reversed Likert items. *J Marketing Res*. 2008;45(1):116–131. doi:10.1509/jmkr.45.1.116
- Rief W, Ihle D, Pilger F. A new approach to assess illness behaviour. *J Psychosom Res*. 2003;54(5):405–414. doi:10.1016/S0022-3999(02)00401-4
- Schwartz PY. Why is neurasthenia important in Asian cultures? *Western J Med*. 2002;176(4):257–258.
- Laferton JAC, Stenzel NM, Rief W, Klaus K, Brahler E, Mewes R. Screening for DSM-5 Somatic Symptom Disorder: diagnostic Accuracy of Self-Report Measures Within a Population Sample. *Psychosom Med*. 2017;79(9):974–981. doi:10.1097/PSY.0000000000000530
- Toussaint A, Husing P, Kohlmann S, Lowe B. Detecting DSM-5 somatic symptom disorder: criterion validity of the Patient Health Questionnaire-15 (PHQ-15) and the somatic symptom scale-8 (SSS-8) in combination with the Somatic Symptom Disorder-B Criteria Scale (SSD-12). *Psychol Med*. 2020;50(2):324–333. doi:10.1017/S003329171900014X
- Leonhart R, de Vroeghe L, Zhang L, et al. Comparison of the factor structure of the patient health questionnaire for somatic symptoms (PHQ-15) in Germany, the Netherlands, and China. A Transcultural Structural Equation Modeling (SEM) Study. *Front Psychiatry*. 2018;9:240. doi:10.3389/fpsy.2018.00240
- Kop WJ, Toussaint A, Mols F, Lowe B. Somatic symptom disorder in the general population: associations with medical status and health care utilization using the SSD-12. *Gen Hosp Psychiatry*. 2019;56:36–41. doi:10.1016/j.genhosppsych.2018.10.004
- van Eck van der Sluijs J, Ten Have M, Rijnders C, van Marwijk H, de Graaf R, van der Feltz-cornelis C. Medically unexplained and explained physical symptoms in the general population: association with prevalent and incident mental disorders. *PLoS One*. 2015;10(4):e0123274. doi:10.1371/journal.pone.0123274
- van Eck van der Sluijs JF, Ten Have M, de Graaf R, Rijnders CAT, van Marwijk HWJ, van der Feltz-cornelis CM. Predictors of persistent medically unexplained physical symptoms: findings from a general population study. *Front Psychiatry*. 2018;9:613. doi:10.3389/fpsy.2018.00613
- van Eck van der Sluijs JF, de Vroeghe L, van Manen AS, Rijnders CAT, van der Feltz-cornelis CM. Complexity Assessed by the INTERMED in patients with somatic symptom disorder visiting a specialized outpatient mental health care setting: a cross-sectional study. *Psychosomatics*. 2017;58(4):427–436. doi:10.1016/j.psym.2017.02.008

35. de Vroege L, Timmermans A, Kop WJ, van der Feltz-cornelis CM. Neurocognitive dysfunctioning and the impact of comorbid depression and anxiety in patients with somatic symptom and related disorders: a cross-sectional clinical study. *Psychol Med*. 2018;48(11):1803–1813. doi:10.1017/S0033291717003300
36. de Vroege L, Koppenol I, Kop WJ, Riem MME, van der Feltz-cornelis CM. Neurocognitive functioning in patients with conversion disorder/functional neurological disorder. *J Neuropsychol*. 2020;15(1):69–87. doi:10.1111/jnp.12206

Neuropsychiatric Disease and Treatment

Dovepress

Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS, and

is the official journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>