Safety, Tolerability, Pharmacokinetics, and Pharmacodynamics of the MK2 Inhibitor ATI-450 in Healthy Subjects: A Placebo-Controlled, Randomized Phase I Study

David Gordon¹
Edward T Hellriegel¹
Heidi Rath Hope²
David Burt¹
Joseph B Monahan²

¹Research and Development, Aclaris Therapeutics, Inc., Wayne, PA, USA;

²Research and Development, Aclaris Therapeutics, Inc., and Confluence Discovery Technologies, Inc., St. Louis, MO, USA

Purpose: ATI-450 is an oral, small-molecule inhibitor of the p38 α mitogen-activated protein kinase (MAPK)/MAPK-activated protein kinase 2 (MK2) inflammatory signaling pathway. This phase 1, single and multiple ascending dose (SAD, MAD) study evaluated ATI-450 safety, tolerability, pharmacokinetics, and pharmacodynamics.

Patients and Methods: Healthy adults were randomly assigned to SAD (10, 30, 50, 100 mg; n=24) and MAD (10, 30, 50 mg twice daily [BID] for 7 days; n=24) cohorts of ATI-450 or placebo (n=14). Safety and tolerability were evaluated through clinical and laboratory assessments. Pharmacokinetic parameters were evaluated in plasma samples; pharmacodynamic assessments included quantification of cytokine levels (tumor necrosis factor α [TNF- α], interleukin [IL]-1 β , IL-6, IL-8) and phosphorylation of the MK2 downstream substrate, heat shock protein 27 (p-HSP27).

Results: The most common adverse events were headache (10/48, 20.8%), dizziness (6/48, 12.5%), upper respiratory tract infection (3/48, 6.3%), and constipation (3/48, 6.3%). Pharmacokinetics were dose-proportional, with a terminal half-life of 9–12 hours in the MAD cohorts on day 7. Dose- and concentration-dependent inhibition of ex vivo stimulated cytokines and target biomarker was observed. On day 7, patients in the 50 mg BID dose cohort recorded mean trough drug levels that were 1.4, 2.2, 2.3, and 2.4 times greater than the IC₈₀ for TNF- α , IL-1 β , IL-8, and p-HSP27, respectively. Mean C_{max} was 3.5, 5.4, 5.6, and 6.0 times greater than the IC₈₀ for TNF- α , IL-1 β , IL-8, and p-HSP27, respectively. IL-6 inhibition >50% was noted for part of the dosing interval.

Conclusion: ATI-450 was well tolerated at the doses investigated, exhibited dose- and time-independent (ie, linear) pharmacokinetics, and dose-related pharmacodynamic effects. These results support further study of ATI-450 in immunoinflammatory diseases in phase 2 trials.

Keywords: immunologic diseases, rheumatoid arthritis, serine-threonine kinases, protein kinase inhibitors, pharmacokinetics/pharmacodynamics

Introduction

The p38 mitogen-activated protein kinase (p38MAPK) signaling pathway has been a target for therapeutic intervention for inflammatory diseases because of its involvement in the regulation and expression of multiple cytokines (eg, tumor necrosis factor α [TNF- α], interleukin [IL] 1 β , and IL-6) and other inflammatory signals.^{1,2} MAPK-activated protein kinase 2 (MK2) is a direct downstream substrate of p38MAPK, which

Correspondence: David Gordon
Chief Medical Officer, Research and Development, Aclaris Therapeutics, Inc.,
640 Lee Road, Suite 200, Wayne, PA,
19087, USA
Tel +1 484-540-6273
Email dgordon@aclaristx.com

has been recognized as a key driver of inflammation.^{3,4} Inflammatory cytokines activated through the p38MAPK/MK2 pathway include TNF- α , IL-1 β , IL-6, and IL-8.⁴⁻⁹

Small-molecule inhibitors of p38MAPK have been evaluated in clinical trials for the treatment of inflammatory diseases; however, clinical development of these compounds has been limited by toxicity and lack of sustained efficacy.^{4,5,10,11} The clinical efficacy achieved with p38MAPK inhibitors has been generally disappointing. Of note, there is evidence in certain autoimmune diseases that p38MAPK inhibitors exhibit tachyphylaxis after multiple treatments, making sustained inhibition of the inflammatory response difficult to achieve in these therapeutic indications, such as rheumatoid arthritis.^{12,13} Tachyphylaxis may be explained by inhibition of the negative feedback and/or anti-inflammatory target substrates described above. Therefore, the approach of targeting key downstream kinase substrates of p38MAPK that specifically regulate inflammatory cytokines to potentially achieve greater specificity for blockade of pathological inflammation has gained interest.

ATI-450 (also known as CDD-450), a recently developed MK2 inhibitor, has a novel mechanism of action by which it targets the modified p38MAPK ATP binding pocket and juxtaposed MK2 formed upon the interaction between p38MAPK and MK2 ([Supplementary Figure S1](#)).⁵ The p38MAPK binding site on MK2 is located within the C terminus; thermodynamic studies have shown that the 2 proteins in the complex bind tightly, and deletion of the binding sequence in MK2 abrogates p38-dependent phosphorylation and activation of MK2.¹⁴ Following the formation of the p38MAPK-MK2 biomolecular complex, ATI-450 binds to the interface of the complex with significantly higher affinity than to either kinase alone, thereby selectively inhibiting p38MAPK phosphorylation of MK2 and locking MK2 in an inactive conformation.⁵ MK2 regulates inflammatory cytokine mRNA stability and translation through phosphorylation of downstream effectors including adenylate-uridylate-rich element-binding proteins (eg, tristetraprolin).¹⁵⁻¹⁷ Therefore, inhibition of p38MAPK-MK2 blocks downstream MK2-mediated inflammatory actions.

Preclinical animal disease models have been used to predict the potential of ATI-450 (ie, CDD-450) as a treatment in several immune-mediated inflammatory diseases. In a rat streptococcal cell wall arthritis model for rheumatoid arthritis, ATI-450 showed joint protective effects and preserved bone mineral density.⁵ ATI-450 also was studied in a murine model of neonatal-onset

multisystem inflammatory disease (NOMID), which is a severe form of a cryopyrin-associated periodic syndrome; NOMID and other cryopyrin-associated periodic syndromes comprise a spectrum of rare hereditary autoinflammatory disorders in which IL-1 β and IL-18 are overproduced. A dramatic attenuation of disease was observed in this transgenic animal model, including a reduction in bone marrow levels of IL-1 β in mice treated with ATI-450 relative to those treated with placebo.⁵ These preclinical results support clinical development of ATI-450 in humans for autoimmune and autoinflammatory diseases driven by the cytokines TNF- α , IL-1 β , IL-6, and IL-8.

A first-in-human, phase 1, randomized, observer-blind, placebo-controlled study was conducted to evaluate the safety, tolerability, pharmacokinetics (PK), and pharmacodynamics (PD) of ATI-450 in single and multiple ascending dose (SAD and MAD) cohorts in healthy subjects. In addition, the study included separate cohorts that were used to evaluate the PK of fed versus fasting administration and of co-administration with methotrexate; these findings will be reported separately because of content size limitations. Here, we report data from the SAD and MAD cohorts.

Methods

The study's clinical protocol, informed consent forms, and amendments to all documents were reviewed and approved by Midlands Independent Review Board of Overland Park, Kansas. The protocol ATI-450-PKPD-101 was first approved on June 7, 2019 and there were 4 amendments, with the final protocol approved on November 6, 2019. Informed written consent was obtained from all subjects before any study-related procedures began. The study was conducted by PRA Health Sciences – Early Development Services (Lenexa, KS) in accordance with the principles of the Declaration of Helsinki and in compliance with the International Council for Harmonization E6 Guideline for Good Clinical Practice, and any applicable national and local laws and regulations. Consistent with guidelines, this phase 1 study was not submitted to a clinical trial register. The conduct of the study included the ability of subjects to opt out and subject deidentification and anonymization. No personal information was collected.

Study Design

PRA Health Sciences led the study at a clinical site in Lenexa, Kansas, and prepared the randomization scheme for all cohorts while maintaining blinding of observers

associated with the study for the cohorts described here. PRA assigned randomization codes sequentially as healthy subjects became eligible for randomization after enrolling in the study. For each SAD dose cohort, subjects were randomly assigned to ATI-450 or placebo in an overall 6:2 ratio. For each MAD dose cohort, subjects were randomly assigned to ATI-450 or placebo in an overall 8:2 ratio.

Single Ascending Dose (SAD) Cohorts

Thirty-two subjects were randomly assigned to 1 of 4 cohorts to receive ATI-450 doses of 10, 30, 50, or 100 mg ($n=6$ per cohort) or placebo ($n=2$ per cohort). Each subject received a single oral dose of ATI-450 or placebo in the morning on an empty stomach. All subjects were admitted to the clinical study facility on day -1 and stayed until 48 hours after ATI-450 or placebo administration.

Multiple Ascending Dose (MAD) Cohorts

Thirty subjects were randomly assigned to 1 of 3 cohorts to receive ATI-450 doses of 10, 30, or 50 mg twice daily (BID) ($n=8$ per cohort) or placebo ($n=2$ per cohort). Subjects received each dose of ATI-450 or placebo on an empty stomach for 7 days, in the morning and the evening, with the final dose administered on the morning of day 7. Subjects were admitted to the clinical facility on day -1 and remained at the site until at least 72 hours after the morning dose on day 7.

Subjects

Subjects were healthy males or females between 18 and 55 years of age, inclusive; had a body mass index between 18 and 32 kg/m², inclusive, with a minimum body weight of 50 kg; and tested negative for HIV-1 and HIV-2 antibodies, hepatitis B surface antigen, and hepatitis C virus antibody at screening. Female subjects were not pregnant or nursing and agreed to use 2 effective methods of contraception if heterosexually active and of childbearing potential.

Subjects were excluded from the study if they had a current acute or chronic disease or used medications that may have affected their safety or other study assessments within 2 weeks of admission.

Safety and Tolerability Assessments

Adverse events, clinical laboratory tests, vital signs, an electrocardiogram, and physical examination results were evaluated for each subject in both parts of the study. Holter monitoring data were collected for future analysis.

Blood Sampling and Bioanalysis

Two-milliliter blood samples were collected for PK analysis in plasma samples. For the SAD cohorts, blood was collected pre-dose and 0.5, 1, 2, 4, 6, 8, 12, 24, 36, and 48 hours post-dose. For the MAD cohorts, blood was collected on day 1 prior to the morning dose and 0.5, 1, 2, 4, 6, 8, and 12 hours post-dose (before the evening dose); before the morning dose on days 2 through 6; and on day 7, before the morning dose and 0.5, 1, 2, 4, 6, 8, 12, 13, 14, 24, 36, 48, and 72 hours after the morning dose.

Blood samples of 10 mL were collected for PD analysis. For the SAD cohorts, blood was collected pre-dose and 1, 12, and 24 hours post-dose. For the MAD cohorts, on day 1 blood was collected before the morning dose and 4 and 12 hours after the morning dose (prior to the evening dose); on day 7, blood was collected before the last dose and 4 and 12 hours after the last dose.

ATI-450 concentrations in plasma samples were determined by PRA Health Sciences Bioanalytical Laboratory (Lenexa, Kansas) using a validated ultra-performance liquid chromatography with tandem mass spectrometry method (details in [Supplementary Information](#)). The lower limit of quantification was 0.500 ng/mL.

PD bioanalysis (performed by Confluence Discovery Technologies, Inc. [St. Louis, Missouri]) of phosphorylated heat shock protein 27 (p-HSP27), IL-1 β , IL-6, IL-8, and TNF- α levels in blood samples were analyzed under 3 conditions: (1) unstimulated, (2) ex vivo lipopolysaccharide-stimulated, and (3) ex vivo lipopolysaccharide-stimulated in the presence of exogenously added 10 μ M ATI-450. Historical data show the percentage of maximum inhibition by ATI-450 achieved for a given analyte is consistent and therefore useful for normalization across samples. Additional details of the assays for analyzing p-HSP27 and cytokines are included in the [Supplementary Information](#).

SAD cohort data were calculated for each subject set (pre-dose, 1 hour post-dose, and 12 hours post-dose on day 1, and 24 hours post-dose on day 2) as % day 1 pre-dose using the ex vivo lipopolysaccharide stimulation as the maximum signal and normalizing across the sample set using the day 1 ex vivo lipopolysaccharide plus 10 μ M ATI-450 as maximum inhibition.

MAD cohort data were calculated for each subject set (pre-dose, 4 hours post-dose, and 12 hours post-dose on days 1 and 7) as % day 1 pre-dose using the ex vivo lipopolysaccharide stimulation as the maximum signal and normalizing across the sample set using the day 1 ex

vivo lipopolysaccharide plus 10 μ M ATI-450 as maximum inhibition.

Statistics

Pharmacokinetics

ATI-450 plasma concentrations and PK parameters were determined for all SAD and MAD cohorts. Noncompartmental analysis with Phoenix WinNonlin™, version 8.2 (Certara, Princeton, NJ) was used to determine the following PK parameters from the plasma concentration–time data for ATI-450: maximum observed plasma concentration ($C_{\max}$); time to maximum observed plasma concentration ($t_{\max}$); area under the plasma concentration–time curve from time 0 through time t (AUC_{0-t}), where t is the time of the last quantifiable concentration; apparent terminal elimination rate constant (λ_z); and associated half-life ($t_{1/2}$). For the SAD cohorts, additional PK parameters determined were area under the plasma concentration–time curve from time 0 to infinity (AUC_{0-inf}), apparent clearance (CL/F), and apparent volume of distribution (V_z/F). For the MAD cohorts, several additional PK parameters were determined, including area under the plasma concentration–time curve over the 12-hour dosing interval, tau (AUC_{τ}); minimum observed plasma concentration in the dosing interval after multiple dosing, on day 7 ($C_{\min}$); average plasma concentration after multiple dosing, on day 7 (C_{avg}); measured concentration at the end of a dosing interval (C_{trough}); accumulation ratio for $C_{\max}$ and AUC_{τ} (R_{ac} , $C_{\max}$ and R_{ac} , AUC_{τ}); apparent clearance at steady state (CL_{ss}/F); and apparent volume of distribution at steady state (V_{ss}/F).

Pharmacodynamics

PD data analysis was conducted using Microsoft Excel 2010 (Microsoft, Redmond, WA), Meso Scale Discovery Workbench® analysis software (Rockville, MD), and Phoenix WinNonlin, version 8.1. For the PK/PD analysis in Phoenix WinNonlin, the ATI-450 plasma concentrations and the percentages of inhibition of production of lipopolysaccharide-stimulated p-HSP27, TNF- α , IL-1 β , IL-6 and IL-8 levels in human whole blood from all SAD/MAD cohorts were simultaneously fit using an inhibitory $E_{\max}$ model (WinNonlin, model 104). The parameters estimated by the model for each biomarker were the baseline effect at zero concentration (E_0), the maximum percent inhibition ($I_{\max}$), and the ATI-450 plasma concentration at half maximal inhibition (IC_{50}). The ATI-450 concentrations at 80% inhibition (IC_{80}) were subsequently calculated from the IC_{50} estimates using the following equation: $IC_x = (x/100-x)*IC_{50}$, where x is the percent inhibition (ie, 80%). Additionally, multiples

of the peak and trough concentrations of ATI-450 50 mg BID on day 7 to the IC_{50} and IC_{80} estimates for p-HSP27 and each cytokine were determined.

Sample Size Calculations

No prospective calculations of statistical power were made for sample size. Sample sizes of 32 subjects and 30 subjects for the SAD and MAD cohorts, respectively, were selected and are typical for a first-in-human study.

Results

Study Subjects

The first screening for subjects began on July 8, 2019, and the last follow-up visit was on December 10, 2019.

In the SAD cohorts, 32 subjects were enrolled, randomized, and included in the safety set. A total of 31 subjects completed the study; 1 subject in the 30 mg ATI-450 cohort withdrew from the study for non-safety reasons after receiving a single dose on day 1 and thus did not complete all study assessments. PK and PD analyses included the 24 subjects who received ATI-450. Most subjects in the SAD cohorts were female (84%; Table 1) and ages ranged from 18 to 51 years.

In the MAD cohorts, 30 subjects were enrolled, randomized, and completed the study, and all subjects were included in the safety set; the 24 subjects who received ATI-450 were included in the PK and PD sets. These cohorts were more evenly split between female and male subjects (57% female, 43% male) and ages ranged from 21 to 53 years.

Safety and Tolerability

No clinically significant findings in vital signs, electrocardiograms, or clinical laboratory values were observed. The most common treatment-emergent adverse events (TEAEs) among all study participants are shown in Table 2. In the SAD cohorts, 15 TEAEs were reported by 7 of 24 (29%) subjects who received ATI-450 and 6 TEAEs were reported by 4 of 8 (50%) subjects who received placebo. No deaths or serious adverse events (SAEs) and no discontinuations due to TEAEs occurred. All TEAEs were mild, transient, and resolved without sequelae by the end of the study. Four of 24 (17%) subjects who received ATI-450 reported TEAEs that were considered by the investigator to be related to the study drug (0 subjects in the 10 mg cohort; 2 [33%] subjects in the 30 mg cohort [1 event of nausea and 2 events each of headache and dizziness]; 1 [17%] subject in the 50 mg cohort [1 event of headache]; and 1 subject in the 100 mg cohort [1 event of dizziness]).

Table 1 Summary of Demographic Characteristics

	Single Ascending Dose Group						Multiple Ascending Dose Group				
	Placebo (n=8)	ATI-450				Overall (N=32)	Placebo (n=6)	ATI-450			Overall (N=30)
		10 mg (n=6)	30 mg (n=6)	50 mg (n=6)	100 mg (n=6)			10 mg BID (n=8)	30 mg BID (n=8)	50 mg BID (n=8)	
Sex, n (%)											
Female	6 (75.0)	5 (83.3)	6 (100.0)	5 (83.3)	5 (83.3)	27 (84.4)	3 (50.0)	5 (62.5)	4 (50.0)	5 (62.5)	17 (56.7)
Male	2 (25.0)	1 (16.7)	0	1 (16.7)	1 (16.7)	5 (15.6)	3 (50.0)	3 (37.5)	4 (50.0)	3 (37.5)	13 (43.3)
Race, n (%)											
White	3 (37.5)	2 (33.3)	3 (50.0)	4 (66.7)	3 (50.0)	15 (46.9)	4 (66.7)	3 (37.5)	6 (75.0)	2 (25.0)	15 (50.0)
Black	4 (50.0)	3 (50.0)	3 (50.0)	1 (16.7)	3 (50.0)	14 (43.8)	2 (33.3)	4 (50.0)	2 (25.0)	5 (62.5)	13 (43.3)
Other	1 (12.5)	1 (16.7)	0	1 (16.7)	0	3 (9.4)	0	1 (12.5)	0	1 (12.5)	2 (6.7)
Ethnicity, n (%)											
Hispanic or Latino	0	1 (16.7)	0	2 (33.3)	1 (16.7)	4 (12.5)	0	0	2 (25.0)	0	2 (6.7)
Not Hispanic or Latino	8 (100.0)	5 (83.3)	6 (100.0)	4 (66.7)	5 (83.3)	28 (87.5)	6 (100.0)	8 (100.0)	6 (75.0)	8 (100.0)	28 (93.3)
Age, y, mean $\pm$ SD	37.0 $\pm$ 7.5	32.2 $\pm$ 13.7	28.7 $\pm$ 8.8	25.2 $\pm$ 4.0	31.2 $\pm$ 3.7	31.2 $\pm$ 8.8	35.2 $\pm$ 6.9	36.3 $\pm$ 7.8	37.9 $\pm$ 8.0	37.9 $\pm$ 9.0	36.9 $\pm$ 7.7
Weight, kg, mean $\pm$ SD	76.4 $\pm$ 9.9	75.2 $\pm$ 16.8	72.8 $\pm$ 8.4	71.3 $\pm$ 8.2	61.5 $\pm$ 6.2	71.8 $\pm$ 11.2	79.7 $\pm$ 17.4	72.7 $\pm$ 10.7	77.2 $\pm$ 14.4	75.9 $\pm$ 9.1	76.1 $\pm$ 12.5
Height, cm, mean $\pm$ SD	166.1 $\pm$ 5.9	172.5 $\pm$ 10.5	159.5 $\pm$ 10.0	163.8 $\pm$ 6.9	165.0 $\pm$ 8.6	165.4 $\pm$ 8.9	170.7 $\pm$ 7.2	166.6 $\pm$ 7.1	168.1 $\pm$ 9.6	170.3 $\pm$ 10.5	168.8 $\pm$ 8.5
BMI, kg/m ² , mean $\pm$ SD	27.6 $\pm$ 2.6	25.1 $\pm$ 4.2	28.6 $\pm$ 2.1	26.6 $\pm$ 2.3	22.6 $\pm$ 1.1	26.2 $\pm$ 3.3	27.1 $\pm$ 4.4	26.2 $\pm$ 3.9	27.1 $\pm$ 2.5	26.3 $\pm$ 3.7	26.7 $\pm$ 3.5

Abbreviations: BID, twice daily; BMI, body mass index.

Table 2 Summary of Treatment-Emergent Adverse Events (TEAEs), Safety Set

Summary of Adverse Events in the SAD and MAD Cohorts				
Safety Parameter	SAD Cohort		MAD Cohort	
	ATI-450 (n=24)	Placebo (n=8)	ATI-450 (n=24)	Placebo (n=6)
Number of TEAEs	15	6	24	3
Patients reporting TEAEs, n (%)	7 (29.2)	4 (50.0)	12 (50.0)	3 (50.0)
Patients with treatment-related TEAEs, n (%)	4 (16.7)	0	9 (37.5)	2 (33.3)
Serious TEAEs	0	0	0	0
Discontinuations due to TEAEs	0	0	0	0
Deaths	0	0	0	0
Most Common TEAEs Occurring in 2 or More Subjects				
TEAE	ATI-450, n (%) (n=48)		Placebo, n (%) (n=14)	
Dizziness	6 (12.5)		0	
Headache	10 (20.8)		2 (14.3)	
Upper respiratory tract infection	3 (6.3)		1 (7.1)	
Constipation	3 (6.3)		1 (7.1)	
Nausea	2 (4.2)		1 (7.1)	
Abdominal pain	2 (4.2)		0	
Vomiting	0		2 (14.3)	

Abbreviations: MAD, multiple ascending dose; SAD, single ascending dose.

In the MAD cohorts, there were 24 TEAEs reported by 12 of 24 (50%) subjects who received ATI-450 and 3 TEAEs reported by 3 of 6 (50%) subjects who received placebo. As in the SAD cohorts, there were no deaths or SAEs and no subjects discontinued the study due to TEAEs; all TEAEs were mild, transient, and resolved without sequelae by the next visit. Nine of 24 (38%) subjects who received ATI-450 reported TEAEs that were considered by the investigator to be related to the study drug (1 of 8 [13%] subjects in the 10 mg BID cohort [1 event of headache], 5 of 8 [63%] subjects in the 30 mg BID cohort [1 event each of blurred vision, constipation, abdominal pain, diarrhea, arthralgia, dizziness, and oropharyngeal pain], and 3 of 8 [38%] subjects in the 50 mg BID cohort [1 event of headache and 2 events of dizziness]). In the placebo group, 2 of 6 (33%) subjects had TEAEs that were considered by the investigator to be related to the study drug (1 event each of vomiting and headache).

Laboratory changes were generally unremarkable. Dose-dependent reductions in neutrophils were observed at day 2 and day 5 with the maximum effect in the MAD cohort. The mean reduction at day 7 in the 50 mg BID

group was 24% (3318.4 cells/ μ L at baseline versus 2514.4 cells/ μ L at day 7).

Pharmacokinetics

SAD Cohorts

Administration of ATI-450 at single doses of 10 mg, 30 mg, 50 mg, and 100 mg under fasting conditions resulted in ATI-450 post-dose mean plasma concentrations above the lower limit of quantification through 24 hours post-dose at 10 mg and through 48 hours post-dose at all other dose levels. ATI-450 mean plasma concentrations increased with increasing doses (Figure 1). Systemic exposure increased in a dose-proportional manner across the ATI-450 dose range from 10 to 100 mg, with approximately 10.8-, 12.6-, and 12.8-fold increases in C_{max} , AUC_{0-t} , and AUC_{0-inf} , respectively (Table 3). Statistical analysis for dose proportionality using the power model is shown in Supplementary Table S1, where systemic exposure (C_{max} , AUC_{0-t} , and AUC_{0-inf}) increased in proportion to ATI-450 dose after single dosing as indicated by the slope estimates near unity and the 90% CIs including unity. ATI-450 concentrations appeared rapidly in plasma, with median t_{max} ranging from 2 to 4 hours across the dose

Figure 1 Mean (SD) plasma concentration–time profiles of ATI-450 after a single dose under fasted conditions, semi-log scale. Hour 0 is the pre-dose value.

Table 3 ATI-450 Pharmacokinetic Parameters

Parameter	Single Ascending Dose Group				Multiple Ascending Dose Group		
	ATI-450 10 mg (n=6)	ATI-450 30 mg (n=6) ^a	ATI-450 50 mg (n=6)	ATI-450 100 mg (n=6)	ATI-450 10 mg BID (n=8)	ATI-450 30 mg BID (n=8)	ATI-450 50 mg BID (n=8)
C_{max} , ng/mL	39.4 ± 10.4	122.0 ± 33.4	160.7 ± 20.4	426.0 ± 110.6	51.8 ± 15.8	146.5 ± 33.6	219.0 ± 77.8
t_{max} , h ^b	2.0 (2.0–4.0)	4.0 (2.0–4.1)	3.0 (2.0–4.0)	2.0 (2.0–4.0)	2.0 (2.0–2.0)	2.0 (1.0–4.0)	3.0 (1.0–4.0)
AUC_{0-t} , h*ng/mL	276.3 ± 77.8	1074.0 ± 243.5 ^a	1430.0 ± 254.0	3489.8 ± 475.7	364.6 ± 110.7	1204.6 ± 309.1	2260.3 ± 1074.7
AUC_{0-inf} , h*ng/mL	285.3 ± 78.4	1102.4 ± 247.4 ^a	1464.7 ± 269.5	3654.8 ± 522.9	N/A	N/A	N/A
AUC_{tau} , h*ng/mL	N/A	N/A	N/A	N/A	287.8 ± 81.5	908.9 ± 163.1	1507.8 ± 659.9
$t_{1/2}$, h	8.5 ± 3.2	10.7 ± 3.3 ^a	9.1 ± 2.4	11.2 ± 5.8	9.6 ± 2.1	10.3 ± 3.2	11.6 ± 3.7
CL/F or CL_{ss}/F , ^c L/h	37.0 ± 8.8	28.4 ± 7.1 ^a	35.1 ± 6.3	27.9 ± 4.3	37.4 ± 10.9	33.9 ± 5.9	37.3 ± 11.7
V_z/F or V_{ss}/F , ^c L	430.2 ± 124.0	451.4 ± 201.5 ^a	445.0 ± 94.7	439.8 ± 189.7	517.4 ± 203.2	483.1 ± 85.1	586.4 ± 163.2
C_{trough} , ng/mL ^d							
Day 2	N/A	N/A	N/A	N/A	10.1 ± 4.7	37.5 ± 18.8	84.0 ± 40.6
Day 3	N/A	N/A	N/A	N/A	9.8 ± 4.5	44.1 ± 29.3	80.6 ± 44.2
Day 4	N/A	N/A	N/A	N/A	9.2 ± 5.0	36.8 ± 14.9	81.5 ± 49.6
Day 5	N/A	N/A	N/A	N/A	9.6 ± 4.7	34.9 ± 13.8	81.5 ± 43.9
Day 6	N/A	N/A	N/A	N/A	10.7 ± 4.9	38.9 ± 16.2	76.4 ± 38.6
C_{min} , ng/mL	N/A	N/A	N/A	N/A	8.1 ± 3.1	28.8 ± 10.7	58.7 ± 32.8
C_{avg} , ng/mL	N/A	N/A	N/A	N/A	24.0 ± 6.8	75.6 ± 13.4	125.6 ± 55.0
R_{ac} , C_{max}	N/A	N/A	N/A	N/A	1.6 ± 0.6	1.2 ± 0.3	1.3 ± 0.5
R_{ac} , AUC_{tau}	N/A	N/A	N/A	N/A	1.4 ± 0.3	1.4 ± 0.3	1.3 ± 0.4

Notes: Data presented are means ± SD unless otherwise indicated. ^aOne subject was excluded for AUC_{0-t} , AUC_{0-inf} , $t_{1/2}$, CL/F, and V_z/F measurements due to withdrawn consent. No samples were collected past 12 hours. ^bData presented are medians (ranges). ^cValues for CL/F and V_z/F are for single doses and values for CL_{ss}/F and V_{ss}/F are for multiple doses. ^d C_{trough} values are presented for day 2 through day 6 for the MAD cohorts.

Abbreviations: AUC_{0-inf} , area under the plasma concentration–time curve from time 0 to time infinity; AUC_{0-t} , area under the plasma concentration–time curve from time 0 to time t; AUC_{tau} , area under the plasma concentration–time curve over the 12-hour dosing interval, tau; BID, twice daily; CL/F, clearance; CL_{ss}/F , clearance at steady state; C_{avg} , average plasma concentration after multiple dosing, on day 7; C_{max} , maximum plasma concentration; C_{min} , minimum plasma concentration; C_{trough} , measured concentration at the end of a dosing interval; N/A, not applicable; R_{ac} , AUC_{tau} , accumulation ratio for AUC_{tau} ; R_{ac} , C_{max} , accumulation ratio for C_{max} ; $t_{1/2}$, terminal elimination half-life; t_{max} , time to maximum plasma concentration; V_{ss}/F , volume of distribution at steady state; V_z/F , volume of distribution.

cohorts. Mean $t_{1/2}$ of ATI-450 in plasma ranged from 8.5 to 11.2 hours. CL/F, t_{max} , and $t_{1/2}$ values showed no clear relation with dose over the dose range of 10 to 100 mg. Across the dose cohorts, low inter-subject variability in C_{max} , AUC_{0-inf} , and AUC_{0-t} was observed with coefficients of variation ranging from 12.7% to 28.1%.

MAD Cohorts

Figure 2 shows the mean (SD) plasma concentration–time profiles of ATI-450 from 0 to 72 hours after the last dose

on day 7, representing dose-proportional PK for the MAD cohorts, and Table 3 shows ATI-450 PK parameters on day 7 after multiple doses. The results obtained after the first dose on day 1 in the MAD cohorts were consistent with those obtained after single doses in the SAD cohorts (data not shown). T_{max} appeared to be independent of dose, with median values ranging from 2 to 3 hours following the final dose administration on day 7. Mean $t_{1/2}$ was similar across doses following repeated dose administration (from 9.6 to 11.6 hours). AUC_{tau} and C_{max} increased with

Figure 2 Mean (SD) plasma concentration–time profiles of ATI-450 after 7 days of BID dosing under fasted conditions, semi-log scale. Hour 0 is the pre-dose value.
Abbreviation: BID, twice daily.

escalating doses and there was slight accumulation after multiple doses. Over the dose range of 10 to 50 mg BID, C_{max} and AUC_{tau} increased 5.4- and 5.9-fold on day 1, respectively, and 4.2- and 5.2-fold on day 7, respectively. Steady state appeared to be reached by day 2 of administration at all dose levels based on the similarity in the mean C_{trough} of ATI-450 on days 2 through 6. Mean accumulation of ATI-450 after 7 days of BID dosing ranged from 1.2- to 1.6-fold for C_{max} and 1.3- to 1.4-fold for AUC_{tau} . The mean values of $t_{1/2}$, CL, and apparent volume of distribution after multiple doses were comparable to those after a single dose, indicating time-independent PK of ATI-450 after 7 days of BID administration. Statistical analysis of dose proportionality on day 7 using the power model is shown in [Supplementary Table S2](#), where systemic exposure (C_{max} , AUC_{tau}) following BID administration of ATI-450 increased in proportion to dose over the 5-fold dose range as indicated by the slope estimates near unity and the 90% CIs including unity.

Pharmacodynamics

A relationship between concentrations of each analyte and ATI-450 plasma levels was developed and the data were then fit to a nonlinear inhibitory E_{max} model, from which IC_{50} and IC_{80} concentrations were established. The p-HSP27 and cytokine inhibition parameters generated from the model are shown in [Supplementary Table S3](#). Target modulation potency (inhibition of HSP27 phosphorylation) for ATI-450 was similar to potencies for inhibition of production of 3 of the 4 cytokines, TNF- α , IL-1 β , and IL-8, but the potency for inhibition of IL-6 production was lower (less potent). I_{max} was greatest for p-HSP27, TNF- α , and IL-6, with greater than 96% inhibition, and lower for IL-1 β (74%) and IL-8 (57%).

For the 50 mg BID dose MAD cohort, concentrations at C_{trough} were in excess of the relative IC_{80} concentrations (1.4- to 2.4-fold) for the target biomarker p-HSP27 and for 3 of the 4 cytokines (TNF- α , IL-1 β , and IL-8), but not for IL-6 ([Table 4](#)). TNF- α and IL-1 β were inhibited by up to 92.7% and

Table 4 Cytokine and Biomarker IC₈₀ Values and Multiples Across the Dosing Interval in the 50 mg BID Dose Cohort

Biomarker	IC ₈₀ ^a (ng/mL)	C _{trough} ^b (Multiple of IC ₈₀)	C _{max} ^b (Multiple of IC ₈₀)
p-HSP27	36.7	2.4x	6.0x
TNF- α	62.6	1.4x	3.5x
IL-1 β	40.8	2.2x	5.4x
IL-6	747.8	0.1x	0.3x
IL-8	38.8	2.3x	5.6x

Notes: ^aIC₈₀ values generated from combined SAD and MAD ATI-450 concentrations and ex vivo inhibition data using the WinNonlin inhibitory E_{max} model 104. ^bValues are from 50 mg BID MAD cohort on day 7; C_{trough} = 87.9 ng/mL and C_{max} = 219.0 ng/mL.

Abbreviations: BID, twice daily; C_{max}, maximum observed plasma concentration; C_{trough}, measured concentration at the end of a dosing interval; IC₈₀, concentration for 80% of maximal inhibition; IL, interleukin; MAD, multiple ascending dose; p-HSP27, phosphorylated heat shock protein 27; SAD, single ascending dose; TNF- α , tumor necrosis factor α .

83.0%, respectively, versus baseline. Plasma levels were greater than the IC₅₀ for IL-6 for at least part of the dosing interval. The effects of ATI-450 dose on the relative

concentration of each cytokine analyte and the biomarker p-HSP27 in the ex vivo stimulated assay, expressed as a percentage of pre-dose analyte levels (set to 100%) for the 10 mg, 30 mg, and 50 mg BID MAD cohorts on day 7, 4 hours post-dose and 12 hours post-dose, are shown in Figure 3. The 4-hour post-dose day 7 samples reflected approximate steady-state C_{max} ATI-450 concentrations while the 12-hour post-dose day 7 samples reflected steady-state C_{trough} concentrations of the drug. A marked dose-dependent reduction in concentration was observed for all 4 cytokines at the 4-hour time point and persisted through 12 hours. TNF- α , IL-1 β , IL-8, and p-HSP27 all demonstrated a reduction in concentration that persisted for the entire dosing interval. Concentration-dependent and dose-dependent modulation of p-HSP27 and inhibition of the 4 cytokines analyzed were demonstrated.

Discussion

A key challenge with global inhibition of p38MAPK is its ubiquitous expression and the broad effects it exerts on

Figure 3 Mean ($\pm$ SEM) cytokine and biomarker levels for (A) TNF- α , (B) IL-1 β , (C) IL-6, (D) IL-8, and (E) p-HSP27 in the BID dosing cohorts comparing day 1 pre-dose values (set to 100%) with day 7 values 4 hours post-dose (approximate C_{max}) and 12 hours post-dose (C_{trough}).

Abbreviations: BID, twice daily; C_{max}, maximum observed plasma concentration; C_{trough}, measured concentration at the end of a dosing interval; IL, interleukin; p-HSP27, phosphorylated heat shock protein 27; TNF, tumor necrosis factor.

cellular physiology as a consequence of the more than 60 substrates that it regulates.⁴ Through phosphorylation of these substrates, p38MAPK regulates negative feedback loops and anti-inflammatory pathways, which are both involved in downregulating inflammation;^{4,18} thus, global blockade of p38MAPK pathways may inhibit both proinflammatory and anti-inflammatory pathways, limiting its potential efficacy in some disease states. Furthermore, several p38MAPK substrates are involved in the regulation of cellular function and inhibition of these proteins may result in toxicity.⁴

The novel approach of selectively inhibiting the p38MAPK/MK2 biomolecular complex instead of p38MAPK alone results in effective blockade of the proinflammatory axis while sparing the anti-inflammatory pathways, negative feedback substrates, and proteins that regulate general cellular function. This approach has the potential to generate safer compounds with greater and more sustained anti-inflammatory efficacy compared with global p38MAPK inhibitors,⁵ as direct targeting of MK2 has been largely unsuccessful.⁴

ATI-450 is an oral, small-molecule inhibitor of the p38 α MAPK/MK2 inflammatory signaling pathway. In this study, ATI-450 was generally safe and well tolerated by healthy subjects over a range of single doses up to 100 mg and multiple doses up to 50 mg BID for 7 days. All TEAEs resolved without sequelae and there were no discontinuations due to adverse events. Dizziness occurred more frequently in the active arms but there was no clear dose response; the events were generally transient in nature and almost all resolved while on study drug. Dose-dependent reductions in neutrophils were observed and were consistent with the mechanism of action of ATI-450. PK parameters demonstrated that plasma concentrations of ATI-450 increased in a dose-proportional manner after administration of single or multiple ascending doses. Across all SAD and MAD cohorts, median t_{max} ranged from 2 to 4 hours and mean $t_{1/2}$ ranged from approximately 9 to 12 hours with no clear relationship to dose. ATI-450 demonstrated concentration-dependent and dose-dependent modulation of the target biomarker p-HSP27 and inhibition of the production of TNF- α , IL-1 β , IL-6, and IL-8.

ATI-450 presents a novel mechanism by which to inhibit MK2 and the proinflammatory cytokines activated by the p38MAPK pathway. As an orally administered drug, ATI-450 provides a potential alternative to injectable biologic medications that treat immune-mediated

inflammatory diseases. ATI-450 has demonstrated efficacy in animal models for various immune-mediated inflammatory diseases and some types of cancer.^{5,19} Inhibition of TNF- α and IL-1 β observed in healthy subjects in this study align with inhibition of the same cytokine biomarkers demonstrated in animal models for rheumatoid arthritis and cryopyrin-associated periodic syndromes.⁵

The PD responses in this study have been expressed as ratios of plasma concentrations of ATI-450 to the IC₈₀ for each cytokine. This measurement assists in identification of the key biomarkers that are modulated in response to ATI-450 and allows for adjustment of ATI-450 dosing to maintain plasma concentrations that are above the IC₈₀ for those biomarkers. ATI-450 demonstrated potent inhibition of 4 of the 5 biomarkers (p-HSP27, TNF- α , IL-1 β , and IL-8). This inhibition was observed even at trough levels, as demonstrated in the 50 mg BID cohort, in which systemic drug concentrations in excess of the IC₈₀ were achieved for p-HSP27, TNF- α , IL-1 β , and IL-8 at C_{max} (3.5x to 6.0x) and C_{trough} (1.4x to 2.4x). The more modest inhibition by ATI-450 of IL-6 induction in healthy subjects observed in this study may not be indicative of response in patients with rheumatoid arthritis because it does not take into account the impact of inhibiting IL-1 β on downstream IL-6 release. Studies evaluating the potency of ATI-450 in human whole blood stimulated with IL-1 β demonstrated potency for inhibiting IL-6 that was comparable to that for TNF- α and IL-8 (data not shown).

Limitations of this study include the small number of participants who were healthy volunteers; further study in patients who have an inflammatory disease is required to adequately quantify clinical efficacy and safety. However, this is a limitation that is present in most phase 1 studies of new compounds and is well accepted as the first step to determining whether a new compound is worth investigating further.

ATI-450 was well tolerated at the doses investigated, exhibited dose- and time-independent (ie, linear) PK, and dose-related PD effects. Results of this study support the progression of ATI-450 into phase 2 development for the treatment of rheumatoid arthritis and other immune-mediated inflammatory diseases.

Data Sharing Statement

All data requests should be submitted to the corresponding author for consideration. Access to de-identified data may be granted following review.

Acknowledgments

Writing and editorial assistance was provided to the authors by Peloton Advantage, LLC, an OPEN Health company, Parsippany, NJ, and was funded by Aclaris Therapeutics, Inc. The authors would like to thank the volunteers; clinical personnel; Aclaris CMC, Laboratory and Operational Team; and Data Management and Biostatistical staff. Also, the authors thank Daniel Dickerson, MD, the principal study investigator, and Marco Cardillo, CCRA, Madison Bangs, BA, Emma Huff, BS, Aparna Kaul, PhD, Anne Hildebrand, PhD, and Barry Burnette, PhD for their contributions to the conduct of the study and analysis.

Funding

This study was sponsored by Aclaris Therapeutics, Inc., Wayne, Pennsylvania, USA. Neither honoraria nor other form of payment was made for authorship. Writing and editorial assistance was provided to the authors by Peloton Advantage, LLC, an OPEN Health company, Parsippany, NJ, and was funded by Aclaris Therapeutics, Inc.

Disclosure

David Gordon, Joseph B Monahan, Edward T Hellriegel, Heidi Rath Hope and David Burt are employees and shareholders of Aclaris Therapeutics. Dr David Gordon has patents 63076689, 63126173, 63136080, 63136967, 63138672, 63128523, and 63140116 pending. Mr Edward T Hellriegel has patents 62986954, 62986955, 63076689, 63126173, 63136080, 63136967, 63138672, and 63140116 pending to Aclaris Therapeutics. Dr Joseph B Monahan has patents 8563558, 8507499, 9359300, 9365546, 9365547, 9056110, 9115089, 9636333, and 63136967 pending to Aclaris Therapeutics; patents 63138672, 63140116, 63136080, 63126173, 63076689, 62980955, 62986954, 63128523, and 63024160 issued to Aclaris Therapeutics.

References

- Saklatvala J. The p38 MAP kinase pathway as a therapeutic target in inflammatory disease. *Curr Opin Pharmacol*. 2004;4(4):372–377. doi:10.1016/j.coph.2004.03.009
- Schett G, Zwerina J, Firestein G. The p38 mitogen-activated protein kinase (MAPK) pathway in rheumatoid arthritis. *Ann Rheum Dis*. 2008;67(7):909–916. doi:10.1136/ard.2007.074278
- Kotlyarov A, Neininger A, Schubert C, et al. MAPKAP kinase 2 is essential for LPS-induced TNF- α biosynthesis. *Nat Cell Biol*. 1999;1(2):94–97. doi:10.1038/10061
- Singh RK, Najmi AK, Dastidar SG. Biological functions and role of mitogen-activated protein kinase activated protein kinase 2 (MK2) in inflammatory diseases. *Pharmacol Rep*. 2017;69(4):746–756. doi:10.1016/j.pharep.2017.03.023
- Wang C, Hockerman S, Jacobsen EJ, et al. Selective inhibition of the p38 α MAPK-MK2 axis inhibits inflammatory cues including inflammasome priming signals. *J Exp Med*. 2018;215(5):1315–1325. doi:10.1084/jem.20172063
- Muniyappa H, Das KC. Activation of c-Jun N-terminal kinase (JNK) by widely used specific p38 MAPK inhibitors SB202190 and SB203580: a MLK-3-MKK7-dependent mechanism. *Cell Signal*. 2008;20(4):675–683. doi:10.1016/j.cellsig.2007.12.003
- Ma W, Lim W, Gee K, et al. The p38 mitogen-activated kinase pathway regulates the human interleukin-10 promoter via the activation of Sp1 transcription factor in lipopolysaccharide-stimulated human macrophages. *J Biol Chem*. 2001;276(17):13664–13674. doi:10.1074/jbc.M011157200
- Wagh DJ, Wilson C. The interleukin-8 pathway in cancer. *Clin Cancer Res*. 2008;14(21):6735–6741. doi:10.1158/1078-0432.CCR-07-4843
- Murphy C, McGurk M, Pettigrew J, et al. Nonapical and cytoplasmic expression of interleukin-8, CXCR1, and CXCR2 correlates with cell proliferation and microvessel density in prostate cancer. *Clin Cancer Res*. 2005;11(11):4117–4127. doi:10.1158/1078-0432.CCR-04-1518
- Hammaker D, Firestein GS. “Go upstream, young man”: lessons learned from the p38 saga. *Ann Rheum Dis*. 2010;69(Suppl1):i77–82. doi:10.1136/ard.2009.119479
- Rokosz LL, Beasley JR, Carroll CD, et al. Kinase inhibitors as drugs for chronic inflammatory and immunological diseases: progress and challenges. *Expert Opin Ther Targets*. 2008;12(7):883–903. doi:10.1517/14728222.12.7.883
- De Buck S, Hueber W, Vitaliti A, et al. Population PK-PD model for tolerance evaluation to the p38 MAP kinase inhibitor BCT197. *Pharmacometr Syst Pharmacol*. 2015;4(12):691–700. doi:10.1002/psp4.12037
- Damjanov N, Kauffman RS, Spencer-Green GT. Efficacy, pharmacodynamics, and safety of VX-702, a novel p38 MAPK inhibitor, in rheumatoid arthritis: results of two randomized, double-blind, placebo-controlled clinical studies. *Arthritis Rheum*. 2009;60(5):1232–1241. doi:10.1002/art.24485
- Fiore M, Forli S, Manetti F. Targeting mitogen-activated protein kinase-activated protein kinase 2 (MAPKAPK2, MK2): medicinal chemistry efforts to lead small molecule inhibitors to clinical trials. *J Med Chem*. 2016;59(8):3609–3634. doi:10.1021/acs.jmedchem.5b01457
- Hitti E, Iakovleva T, Brook M, et al. Mitogen-activated protein kinase-activated protein kinase 2 regulates tumor necrosis factor mRNA stability and translation mainly by altering tristetraprolin expression, stability, and binding to adenine/uridine-rich element. *Mol Cell Biol*. 2006;26(6):2399–2407. doi:10.1128/MCB.26.6.2399-2407.2006
- Mahtani KR, Brook M, Dean JL, Sully G, Saklatvala J, Clark AR. Mitogen-activated protein kinase p38 controls the expression and posttranslational modification of tristetraprolin, a regulator of tumor necrosis factor α mRNA stability. *Mol Cell Biol*. 2001;21(19):6461–6469. doi:10.1128/MCB.21.9.6461-6469.2001
- Chrestensen CA, Schroeder MJ, Shabanowitz J, et al. MAPKAP kinase 2 phosphorylates tristetraprolin on in vivo sites including Ser178, a site required for 14-3-3 binding. *J Biol Chem*. 2004;279(11):10176–10184. doi:10.1074/jbc.M310486200
- Jones DS, Jenney AP, Joughin BA, Sorger PK, Lauffenburger DA. Inflammatory but not mitogenic contexts prime synovial fibroblasts for compensatory signaling responses to p38 inhibition. *Sci Signal*. 2018;11:520. doi:10.1126/scisignal.aal1601
- Murali B, Ren Q, Luo X, et al. Inhibition of the stromal p38MAPK/MK2 pathway limits breast cancer metastases and chemotherapy-induced bone loss. *Cancer Res*. 2018;78(19):5618–5630. doi:10.1158/0008-5472.CAN-18-0234

Clinical Pharmacology: Advances and Applications**Dovepress****Publish your work in this journal**

Clinical Pharmacology: Advances and Applications is an international, peer-reviewed, open access journal publishing original research, reports, reviews and commentaries on all areas of drug experience in humans. The manuscript management system is completely online and

includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-pharmacology-advances-and-applications-journal>