Is Prenatal Diagnosis Necessary for Fetal Isolated Nasal Bone Absence or Hypoplasia?

Feng Zhang¹
 Wei Long¹
 Qin Zhou¹
 Jing Wang¹
 Ye Shi¹
 Jianbing Liu^{1,*}
 Qiuwei Wang^{2,*}

¹Department of Medical Genetics, Changzhou Maternity and Child Health Care Hospital Affiliated to Nanjing Medical University, Changzhou, Jiangsu Province, People's Republic of China;

²Clinical Laboratory, Changzhou Children's Hospital Affiliated to Nantong Medical University, Changzhou, Jiangsu Province, People's Republic of China

*These authors contributed equally to this work

Purpose: This study aimed to explore the value of chromosomal microarray analysis (CMA) and whole exome sequencing (WES) in the prenatal diagnosis of fetal isolated nasal bone absence (INBA) or isolated nasal bone hypoplasia (INBH). We hope to provide additional relevant information for clinical counseling.

Patients and Methods: From November 1, 2018, to March 1, 2020, 55 pregnant women with isolated nasal bone dysplasia were admitted to the Changzhou Maternity and Child Health Care Hospital. Based on the degree of abnormality, the patients were divided into two groups: INBA and INBH. CMA was performed on all patients. The clinical data and prenatal genetic diagnoses of the two groups were retrospectively analyzed. According to the requirements of WES for samples, 12 cases with negative CMA results were selected for the WES test.

Results: A total of 55 cases with INBA or INBH met the inclusion criteria. In 35 INBA fetuses, there was one case of trisomy 21 and one case of 10q11.22 deletion (5.7Mb), and the abnormality rate was 5.71% (2/35). Compared with INBA fetuses, the abnormality rate was increased in the fetuses with INBH [15.00% (3/20)] (15.00% vs 5.71%); there was one case of 1q21.1 duplication (1.3Mb), one case of Xp22.31 duplication (1.67Mb), and one case of 4p deletion (7.6Mb). In a later retrospective study, two pathogenic variants were identified in two cases after the WES test; the abnormality rate was 16.67% (2/12), which involved *RUNX2* and *CDH4* genes, respectively.

Conclusion: A preliminary study confirmed that molecular prenatal diagnosis should be performed in fetuses with INBA or INBH. CMA followed by WES is an effective method.

Keywords: chromosomal microarray, whole exome sequencing, isolated nasal bone absence, isolated nasal bone hypoplasia

Introduction

Nasal bone abnormalities, including nasal bone absence and hypoplasia, have become the focus of many studies. Previous studies showed that the incidence of fetal nasal bone absence or hypoplasia was 0.5–4.5% among fetuses with a normal karyotype.^{1–3} Upon prenatal ultrasound examination, it is noted that nasal bone abnormalities are often accompanied by other structural abnormalities or soft markers, such as endocardial cushion defect, increased nuchal fold thickness (≥ 6 mm), short femur or humerus, ventriculomegaly (≥ 10 mm), intestinal hyperechogenicity, and echogenic intracardiac focus among others.¹ It is widely known that nasal bone absence or hypoplasia is associated with fetal chromosomal abnormalities, such as trisomy 13, 18, 21, and Turner syndrome. In particular, it is closely related to trisomy 21, with a likelihood ratio of 11.6 to 50.5.^{1–5} Rare

Correspondence: Jianbing Liu
 Department of Medical Genetics,
 Changzhou Maternity and Child Health
 Care Hospital Affiliated to Nanjing
 Medical University, Changzhou, Jiangsu
 Province, People's Republic of China
 Tel +86 0519-88581983
 Email liujb2222@126.com

Qiuwei Wang
 Clinical Laboratory, Changzhou
 Children's Hospital Affiliated to Nantong
 Medical University, Changzhou, Jiangsu
 Province, People's Republic of China
 Tel +86 0519-69808280
 Email qiuwei_wang@ntu.edu.cn

conditions have also been reported to be associated with nasal bone absence or hypoplasia, such as Cri du chat (5p-) syndrome, Wolf-Hirschhorn syndrome (4p-), and Fryns Syndrome.^{6–8} Ultrasound measurement of nasal bone length has become a routine part of prenatal care during the first or second trimester, and is recommended for all pregnant women. However, it is difficult to measure fetal nasal bone length during early pregnancy because of unclear imaging, but image clarity improves as the fetus grows. In the second trimester, the nasal bone is clearly visible and can be more easily measured and evaluated.

In the recent years, Europe, the United States, and China have issued statements recommending chromosome microarray analysis (CMA) as the preferred diagnostic method for fetal genetic detection of prenatal ultrasound abnormalities.^{9–11} In clinical practice, pregnant women with nasal bone absence or hypoplasia in association with other structural abnormalities are generally recommended to undergo prenatal karyotype analysis or CMA tests after clinical counseling.

However, even when the karyotype or CMA is normal, fetal genetic abnormalities may still occur.^{12,13} It is necessary to develop new efficient screening technologies to improve the detection rate of chromosomal abnormalities. Whole exome sequencing (WES) is used not only to detect known disease-related genes, but also to discover new disease-related genes, and diagnoses nearly 20% of rare diseases.¹⁴ A meta-analysis by Clark et al reported that the diagnostic and clinical utility of WGS/WES was greater than that of CMA in children with suspected genetic diseases.¹² During prenatal diagnosis, even when karyotype or CMA is normal, fetal genetic abnormalities may still occur, while WES may improve the identification of genetic disorders in fetuses with structural abnormalities.^{15,16} Currently, studies have focused on fetal ultrasound structural abnormalities and/or soft marker abnormalities. Few studies have examined the prenatal diagnosis of single soft marker abnormalities using CMA or WES. Whether prenatal diagnosis of fetal isolated nasal bone absence or hypoplasia is necessary and which technology should be used for diagnosis remains unclear. This makes clinical consultations complicated.

However, whether pregnant women with isolated nasal bone absence (INBA) or isolated nasal bone hypoplasia (INBH) should be advised to undergo prenatal diagnosis remains to be determined. The extent of the risk of genetic variation in these fetuses remains unclear. The aim of this study was to review the prenatal diagnosis and outcomes

of pregnancies with INBA or INBH in the second trimester, in order to explore the value of CMA and WES in the diagnosis of INBA or INBH. We hope to provide additional relevant information for clinical counseling.

Patients and Methods

Patients and Design

The study design and protocol were reviewed and approved by the ethics committee of the Changzhou Maternity and Child Health Care Hospital affiliated with Nanjing Medical University (No. 2017003). All participants received genetic counseling and provided informed consent prior to testing according to the guidelines of the Declaration of Helsinki.

From November 2018 to March 2020, 55 pregnant women were reported to have fetal nasal bone absence/hypoplasia after ultrasound examination in the second trimester, including 35 cases of INBA and 20 cases of INBH. Patients were 25–37 years old, and 23 weeks (and 5 days) to 29 weeks (and 2 days) pregnant, respectively. The characteristics of all women are shown in Table 1. They were recalled for genetic counseling, and prenatal diagnosis was accepted. All patients underwent traditional karyotype analysis and the CMA test. Moreover, with the development and wide application of WES, 12 cases with negative CMA results agreed to undergo further testing using WES.

Nasal Bone Measurement

Routine ultrasound examination of the abdomen was performed to show the median sagittal section of the fetal head, so that the fetal head was facing the probe. The nasal bone was measured at the level of synostosis using the method and reference range as previously described by Sonek et al.¹⁷ The nasal bone was considered absent if it was not visualized on any appropriate view. Cases with sagittal sections of the fetuses showed that one or both

Table 1 The Characteristics of INBA and INBH

	INBA (n=35)	INBH (n=20)
Maternal age(y)	28.06 ± 4.39	27.10 ± 4.50
Gestational age at ultrasound(w)	23.11 ± 3.62	23.14 ± 3.83
Gestational age at CMA(w)	24.58 ± 1.99	24.30 ± 2.18
Singleton	35	20
High risk of prenatal screening	1	0
Low risk of prenatal screening	32	19
No screening	2	1

Abbreviations: INBA, isolated nasal bone absence; INBH, isolated nasal bone hypoplasia.

sides of the nasal bone length below the 2.5th percentile were considered hypoplastic.¹⁸

Chromosomal Microarray Analysis for Prenatal Diagnosis

Amniotic fluid (10 mL) was collected with informed consent, and genomic DNA was extracted using a QIAamp DNA Mini Kit (Qiagen, Valencia, CA, USA). Using an Affymetrix CytoScan 750K Array chip, DNA (250 ng) was amplified, labeled, and hybridized according to the manufacturer's protocol. The data were analyzed using the Chromosome Analysis Suite v3.2 (ChAs) software package, and the pathogenicity was analyzed according to OMIM, UCSC, DECIPHER, ISCA, and DGV databases.

Whole Exome Sequencing for Prenatal Diagnosis

After the genomic DNA was processed by the M220 breaker (American Covaris), the SureSelect hereditary disease gene detection (2742 genes) kit (American Agilent) was used to build the library, and the HiSeq 2500 System (American Illumina company) was used for high-throughput sequencing. The data were read by NextGENe software and uploaded to Ingenuity Variant Analysis TM software for bioinformatic analysis.

Statistical Analyses

The parameters of age and gestational age were expressed as median (M), 2.5th percentile (P2.5), and 97.5th percentile (P97.5). The chi-square test or Fisher's exact test was used to compare differences between the two groups. All significance tests were two-sided and a p-value < 0.05 was considered statistically significant.

Results

In this study, 55 pregnant women with INBA or INBH underwent CMA prenatal diagnosis. WES was performed in 12 cases with negative CMA results. The characteristics of INBA and INBH are listed in Table 1.

After the prenatal CMA test, five women with fetal INBA or INBH were confirmed to have chromosomal abnormalities; the abnormality rate was 9.10% (5/55) (Table 2). Of the 55 participants, 15.00% (3/20) had INBH and 5.71% (2/35) had INBA. Although the abnormality rate of INBH was slightly higher, there was no significant difference between the two groups. One case of trisomy 21 with INBA was found, and the fetus had no

Table 2 The Results of CMA Prenatal Diagnosis of INBA and INBH

	INBA (n =35)	INBH (n =20)	p
Aneuploidies, n (%)	1 (2.86%)	0 (0%)	0.10
Micro-duplication/deletion, n (%)	1 (2.86%)	3 (15.00%)	
Total abnormalities, n (%)	2 (5.71%)	3 (15.00%)	

Abbreviations: INBA, isolated nasal bone absence; INBH, isolated nasal bone hypoplasia.

other ultrasound abnormalities. The mother's prenatal screening results did not suggest any abnormalities. Four cases were reported as micro-duplication/deletion, involving chromosomes 1, 4, 10, and X, and the fragment sizes ranged from 1.1Mb to 7.6Mb. According to the literature and database,^{14,19–25} they were identified as three cases with pathogenic copy number variants (CNVs) and one with unclear clinical significance. After genetic counseling, the mothers chose to undergo clinical intervention combined with the clinical phenotype and penetrance of CNV, as shown in Table 3. Notably, case 3 was confirmed as arr[hg19]1q21.1q21.2 (146, 586, 249–147, 844, 778) x3, which showed 1.3Mb duplication in the 1q21.1q21.2 region, containing 10 OMIM genes. Many studies have reported that 1q21.1 duplication syndrome complicates clinical manifestations and leads to incomplete penetrance.^{20,21} The mother decided to continue the pregnancy and successfully delivered a boy. After 6 months, the infant showed no signs of intellectual disability or other manifestations.

Twelve patients underwent a prenatal WES test after obtaining negative results of CMA. As a result, two cases were confirmed as pathogenic CNVs, and the additional abnormal rate of WES was 16.67% (2/12) (Table 4).

Case 5 was detected as INBA by ultrasound at 24 weeks gestation. After genetic counseling, the mother agreed to undergo the CMA test after amniocentesis, and the result was negative. The mother then underwent the WES test, and the result showed a splice-site mutation (c.1021+1G>T) involving the *RUNX2* gene, which was reported to cause cleidocranial dysplasia (CCD; MIM119600). CCD is a congenital autosomal dominant genetic disease with high penetrance and variable expressivity.²⁶ The most common clinical manifestations are delayed closure of the fontanel, widening of the cranial suture, dysplasia of the clavicle and teeth, and short stature.²⁷ The runt-related transcription factor 2 (*RUNX2*; MIM 600211) gene is known as the main

Table 3 4 Cases with Micro-Duplication/Deletion After CMA Test

Case	MA (y)	GA (w)	Ultrasonic Diagnosis	Chromosome Region	CMA Test				Pregnancy Outcome
					Variation Section	Fragment Size	Variant Type	Classification	
1	30	24.5	INBA	10q11.22	46,225,349–51,904,377	5.7Mb	Deletion	Pathogenic	ToP
2	25	26.4	INBH	1q21.1q21.2	146,586,249–147,844,778	1.3Mb	Duplication	Pathogenic	LB,2750g,37+5 weeks GA Normal newborn examination
3	27	25.5	INBH	4p16.3p16.1	1,124,732–8,721,580	7.6Mb	Deletion	Pathogenic	ToP
4	27	25	INBH	Xp22.31	6,455,151–8,127,579	1.1Mb	Duplication	VUS	LB,3140g,40 +5weeks GA Normal newborn examination

Abbreviations: MA, maternal age; GA, gestational age; INBA, isolated nasal bone absence; INBH, isolated nasal bone hypoplasia; CMA, chromosomal microarray; VUS, variants of uncertain significance; ToP, termination of pregnancy; LB, live birth.

Table 4 2 Cases with Monogenic Diseases After WES

Case	MA(y)	Ultrasonic Diagnosis	WES Results	Inheritance	Disease Association
5	29	INBA	RUNX2 c.1021+1G>T chr6-45480145	AD	Cleidocranial Dysplasia
6	31	INBA	CDH4c.4819het_delG, p.V1607Sfs	AD De novo	Sifrim-Hitz-Weiss syndrome

Abbreviations: MA, maternal age; INBA, isolated nasal bone absence; AD, autosomal dominant inheritance.

pathogenic gene of CCD. It was found that this mutation had not been previously reported. The mother gave birth to a boy, who was followed up for half a year, and showed no signs of intellectual disability or other manifestations.

Case 6 was a fetus with an ultrasonic diagnosis of INBA to our hospital for genetic counseling at 25 weeks 'gestation. WES was performed in the case of a negative CMA result. WES result demonstrated a frameshift mutation (c.581-4_581-1delGCAG), involved the *CDH4* gene, which was reported to cause Sifrim-Hitz-Weiss syndrome (SIHIWES; MIM617159). Sifrim-Hitz-Weiss syndrome is an autosomal dominant intellectual developmental disorder with variable congenital defects affecting other systems, including cardiac, skeletal, and urogenital disorders. Patients may have short stature, enlarged head circumference, hearing loss, and nonspecific dysmorphic facial features.²⁸ The mutation was identified as a novel frameshift mutation after parental verification (Figure 1). During the follow-up of the pregnancy outcome, it was found that the patient gave birth to a girl at term, and the the infant's nasal bone appeared normal. The infant was followed up

for ten months, and showed no signs of intellectual disability or other manifestations.

Discussion

In addition to its known association with trisomy 21, a hypoplastic nasal bone may be an objective marker of facial dysmorphism associated with clinically relevant CNVs.⁴⁻⁶ Compared to traditional G-band karyotyping, CMA is a high-resolution genomic technology that can reveal submicroscopic imbalances or CNVs.^{16,29} Numerous studies have shown that CMA provides additional information in 6–7% of pregnancies with abnormal ultrasound findings, and as such is now the recommended first-tier genetic test. In the study, four cases (7.27 (4/55)) were detected with clinically relevant CNVs, which may go undetected by standard karyotype analysis, and the incidence was consistent with the results reported in previous studies.^{30,31} Two patients chose to terminate their pregnancy voluntarily, while the other two opted to continue the pregnancy after clinical counseling. The later flow-up showed no signs of intellectual disability or other manifestations.

Figure 1 The result of WES in case b due to parental verification.

Notes: Sequencing results for patient (C), patient's father (B), and mother (A). A heterozygous deletion (c.4819het_delG) in *CDH4* was found in the fetus and was not detected in her parents.

WES can detect and indels in exon regions, as well as CNVs and gene structure variation. It can be used not only to detect known disease-related genes, but also to discover new disease-related genes; it diagnoses nearly 20% of rare diseases.³² During the study, two cases of pathogenic variants were found after WES; the incidence was 16.67% (2/12), and the abnormal detection rate improved after CMA detection.

However, onelimitation of WES in INBA or INBH in this study was that the test was carried out for the purpose of the research. Although the sequencing results suggested that there were pathogenic mutations in two cases, the pregnant women had to continue the pregnancy because the prenatal CMA results were normal. Another limitation was that a number of variants would not be found to lack phenotypes during pregnancy or even after birth. We will continue to follow up these two cases, and they may show clinical phenotypes consistent with the disease-causing genes as they age.

In conclusion, this preliminary study suggested that the abnormality rate was 7.27% after prenatal CMA diagnosis in fetal isolated nasal bone absence or hypoplasia. An additional abnormality rate of 16.67% was found in further WES tests. The present results confirm that molecular prenatal diagnosis should be performed in fetal isolated nasal bone absence or hypoplasia, as it may be useful in improving the clinical management of pregnancies and in improving the reproductive decisions of affected families.

Acknowledgments

We thank all of the participants for their contribution to this study.

Funding

This work was supported by Changzhou Key Laboratory of High-tech Research (CM20193009), Project funding for the training of high-level health professionals in Changzhou (2016CZLJ013), and key discipline of

maternal and child health care in Jiangsu Province (FXK201754).

Disclosure

The authors have no conflicts of interest to disclose.

References

- Du Y, Ren Y, Yan Y, Cao L. Absent fetal nasal bone in the second trimester and risk of abnormal karyotype in a prescreened population of Chinese women. *Acta ObstetGynecol Scand*. 2018;97(2):180–186. doi:10.1111/aogs.13263
- Agathokleous M, Chaveeva P, Poon LC, Kosinski P, Nicolaides KH. Meta-analysis of second-trimester markers for trisomy 21. *Ultrasound Obstet Gynecol*. 2013;41(3):247–261. doi:10.1002/uog.12364
- Manegold-Brauer G, Maymon R, Shor S, et al. Down's syndrome screening at 11–14 weeks' gestation using prenatal thickness and nasal bone length. *Arch Gynecol Obstet*. 2019;299(4):939–945. doi:10.1007/s00404-019-05083-2
- Moreno-Cid M, Rubio-Lorente A, Rodríguez MJ, et al. Systematic review and meta-analysis of performance of second-trimester nasal bone assessment in detection of fetuses with Down syndrome. *Ultrasound Obstet Gynecol*. 2014;43(3):247–253. doi:10.1002/uog.13228
- Gu YZ, Nisbet DL, Reidy KL, Palma-Dias R. Hypoplastic nasal bone: a potential marker for facial dysmorphism associated with pathogenic copy number variants on microarray. *Prenat Diagn*. 2019;39(2):116–123. doi:10.1002/pd.5410
- Dukhovny S, Wilkins-Haug L, Shipp TD, Benson CB, Kaimal AJ, Reiss R. Absent fetal nasal bone: what does it mean for the euploid fetus? *J Ultrasound Med*. 2013;32(12):2131–2134. doi:10.7863/ultra.32.12.2131
- Sherer DM, Eugene P, Dalloul M, et al. Second-trimester diagnosis of cri du chat (5p-) syndrome following sonographic depiction of an absent fetal nasal bone. *J Ultrasound Med*. 2006;25(3):387–388. doi:10.7863/jum.2006.25.3.387
- Xing Y, Holder JL Jr, Liu Y, et al. Prenatal diagnosis of Wolf-Hirschhorn syndrome: from ultrasound findings, diagnostic technology to genetic counseling. *Arch Gynecol Obstet*. 2018;298(2):289–295. doi:10.1007/s00404-018-4798-1
- On Genetics, Gynecologists Committee. Committee Opinion No. 581: the use of chromosomal microarray analysis in prenatal diagnosis. *ObstetGynecol*. 2013;122(6):1374–1377.
- American College of Obstetricians and Gynecologists and others. ACOG Committee Opinion No. 446: array comparative genomic hybridization in prenatal diagnosis. *Obstet Gynecol*. 2009;114(5):1161–1163. doi:10.1097/AOG.0b013e3181c33cad
- Miller DT, Adam MP, Aradhya S, et al. Consensus statement: chromosomal microarray is a first-tier clinical diagnostic test for individuals with developmental disabilities or congenital anomalies. *Am J Hum Genet*. 2010;86(5):749–764. doi:10.1016/j.ajhg.2010.04.006
- Clark MM, Stark Z, Farnaes L, et al. Meta-analysis of the diagnostic and clinical utility of genome and exome sequencing and chromosomal microarray in children with suspected genetic diseases. *NPJ GenomMed*. 2018;3:16.
- Yang X, Huang LY, Pan M, et al. Exome sequencing improves genetic diagnosis of fetal increased nuchal translucency. *PrenatDiag*. 2020;40(11):1426–1431.
- Stankiewicz P, Kulkarni S, Dharmadhikari AV, et al. Recurrent deletions and reciprocal duplications of 10q11.21q11.23 including CHAT and SLC18A3 are likely mediated by complex low-copy repeats. *Hum Mutat*. 2012;33(1):165–179. doi:10.1002/humu.21614
- Petrovski S, Aggarwal V, Giordano JL, et al. Whole-exome sequencing in the evaluation of fetal structural anomalies: a prospective cohort study. *Lancet*. 2019;393(10173):758–767. doi:10.1016/S0140-6736(18)32042-7
- Levy B, Wapner R. Prenatal diagnosis by chromosomal microarray analysis. *FertilSteril*. 2018;109(2):201–212.
- Sonek JD, McKenna D, Webb D, Croom C, Nicolaides K. Nasal bone length throughout gestation: normal ranges based on 3537 fetal ultrasound measurements. *Ultrasound Obstet Gynecol*. 2003;21(2):152–155. doi:10.1002/uog.41
- Hung JH, Fu CY, Chen CY, Chao KC, Hung J. Fetal nasal bone length and Down syndrome during the second trimester in a Chinese population. *J Obstet Gynaecol Res*. 2008;34(4):518–523.
- Ghai SJ, Shago M, Shroff M, Yoon G. Cockayne syndrome caused by paternally inherited 5 Mb deletion of 10q11.2 and a frameshift mutation of ERCC6. *Eur J Med Genet*. 2011;54(3):272–276. doi:10.1016/j.jmg.2011.02.008
- Brunetti-Pierri N, Berg JS, Scaglia F, et al. Recurrent reciprocal 1q21.1 deletions and duplications associated with microcephaly or macrocephaly and developmental and behavioral abnormalities. *Nat Genet*. 2008;40(12):1466–1471. doi:10.1038/ng.279
- Gourari I, Schubert R, Prasad A. 1q21.1 Duplication syndrome and epilepsy: case report and review. *Neurol Genet*. 2018;4(1):e219. doi:10.1212/NXG.0000000000000219
- Guy C, Wang X, Lu X, Lu J, Li S. Clinical report: variable phenotypic expression in a large sibling cohort with a deletion of 4p16.1. *Clin Case Rep*. 2016;4(10):913–918. doi:10.1002/ccr3.638
- Battaglia A, Carey JC, South ST. Wolf-Hirschhorn syndrome: a review and update. *Am J MedGenet SeminMedGenet*. 2015;169(3):216–223.
- Furrow A, Theisen A, Velsher L, et al. Duplication of the STS region in males is a benign copy-number variant. *Am J MedGenet SeminMedGenet*. 2011;155a(8):1972–1975.
- Li F, Shen Y, Köhler U, et al. Interstitial microduplication of Xp22.31: causative of intellectual disability or benign copy number variant? *Eur J MedGenet*. 2010;53(2):93–99.
- Guo YW, Chiu CY, Liu CL, Jap TS, Lin LY. Novel mutation of RUNX2 gene in a patient with cleidocranial dysplasia. *Int J Clin ExpPathol*. 2015;8(1):1057–1062.
- Muzio LL, Tetè S, Mastrangelo F, et al. A novel mutation of gene CBFA1/RUNX2 in cleidocranial dysplasia. *Ann ClinLabSci*. 2007;37(2):115–120.
- Weiss K, Terhal PA, Cohen L, et al. De novo mutations in CHD4, an ATP-dependent chromatin remodeler gene, cause an intellectual disability syndrome with distinctive dysmorphisms. *Am J HumGenet*. 2016;99(4):934–941. doi:10.1016/j.ajhg.2016.08.001
- Ganapathi M, Nahum O, Levy B. Prenatal diagnosis using chromosomal SNP microarrays. *Methods Mol Biol*. 2019;1885:187–205.
- Callaway JL, Shaffer LG, Chitty LS, Rosenfeld JA, Crolla JA. The clinical utility of microarray technologies applied to prenatal cytogenetics in the presence of a normal conventional karyotype: a review of the literature. *PrenatDiag*. 2013;33(12):1119–1123.
- Wapner RJ, Martin CL, Levy B, et al. Chromosomal microarray versus karyotyping for prenatal diagnosis. *N Engl J Med*. 2012;367(23):2175–2184. doi:10.1056/NEJMoa1203382
- Gahl WA, Markello TC, Toro C, et al. The national institutes of health undiagnosed diseases program: insights into rare diseases. *Gene Med*. 2012;14(1):51–59. doi:10.1038/gim.0b013e318232a005

International Journal of General Medicine

Dovepress

Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies

across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-general-medicine-journal>