#### ORIGINAL RESEARCH

Cost-Effectiveness of Apixaban versus Other Direct Oral Anticoagulants and Warfarin in the Prevention of Thromboembolic Complications Among Finnish Patients with Non-Valvular Atrial Fibrillation

Taru Hallinen<sup>1</sup> Erkki Soini <sup>1</sup> Christian Asseburg <sup>1</sup> Miika Linna<sup>2</sup> Pia Eloranta<sup>3</sup> Sari Sintonen<sup>3</sup> Mikko Kosunen<sup>3</sup>

<sup>1</sup>ESiOR Oy, Kuopio, Finland; <sup>2</sup>Aalto University, Department of Industrial Engineering and Management, Espoo, Finland; <sup>3</sup>Pfizer Oy, Helsinki, Finland

Correspondence: Taru Hallinen ESiOR Oy, Tulliportinkatu 2 LT4, Kuopio, FI-70100, Finland Tel +358 50 568 1894 Email taru.hallinen@esior.fi **Purpose:** Direct oral anticoagulant (DOAC) use for the prevention of thromboembolic complications in patients with non-valvular atrial fibrillation (AF) has increased steadily in Finland. DOACs have been shown to be cost-effective in comparison to warfarin, but published evidence of relative cost-effectiveness between DOACs is still scarce and mostly based on indirect comparisons of clinical trial evidence. The aim of this study was to compare the cost-effectiveness of apixaban to dabigatran, rivaroxaban and warfarin in a Finnish setting using real-life evidence where available.

**Patients and Methods:** A lifetime Markov simulation model used previously in a published Finnish assessment comparing apixaban and warfarin was modified and updated with the relative effectiveness and safety data available from the real-world NAXOS-study and representative Finnish input data for patient characteristics, event risks, mortality, resource use, costs, and quality of life. Apixaban's cost-effectiveness was assessed from health care payer perspective (using 3% per year discount rate) based on incremental cost-effectiveness ratio (ICER, cost per quality-adjusted life year [QALY] gained), probability of cost-effectiveness (at willingness-to-pay [WTP] of 35,000 euros/QALY), and net monetary benefit (NMB).

**Results:** Apixaban increased the average modelled quality-adjusted life-expectancy and reduced the average total health care costs of AF patients when compared to warfarin (+0.14 QALYs, -3691 euros), dabigatran (+0.11 QALYs, -404 euros), and rivaroxaban (+0.03 QALYs, -43 euros). The resulting NMB of apixaban versus warfarin, dabigatran and rivaroxaban was 8723, 4168, and 1129 euros, respectively. The respective probabilities of apixaban being cost-effective against each comparator were 100%, 92.7%, and 64.0%.

**Conclusion:** In this modelling study, apixaban dominated other anticoagulants in the Finnish real-life setting.

Keywords: apixaban, cost-utility, dabigatran, economic evaluation, rivaroxaban, warfarin

## Introduction

Atrial Fibrillation (AF) is the most common form of cardiac arrhythmia with increasing prevalence in the aging western populations. Currently lifetime risk for AF is 1 in 3 individuals of European ancestry at index age of 55.<sup>1</sup> It causes a significant burden to both primary and specialized healthcare. Approximately 30% of patients with AF have at least one hospitalization annually,<sup>1</sup> and in Finland over one third of arrhythmia related hospitalizations are caused by AF.<sup>2</sup> If

© 2021 Hallinen et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress.com/terms work you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission form Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial uses of this work, please see paragraphs A 21 and 5 of our Terms (https://www.dovepress.com/twws.dovepress.com/two hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission form Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs A 21 and 5 of our Terms (https://www.dovepress.com/terms.php).

Received: 22 April 2021 Accepted: 19 July 2021 Published: 13 August 2021

ClinicoEconomics and Outcomes Research 2021:13 745-755

untreated, AF predisposes patients to stroke, thromboembolic complications, heart failure and death, as well as worsens their quality of life.<sup>1,2</sup> The single most important prognostic factor for AF patients is well-managed anticoagulation treatment.<sup>2</sup>

Warfarin was the treatment of choice for anticoagulation treatment of AF patients before the advent of direct oral anticoagulants (DOACs) such as apixaban, dabigatran, rivaroxaban and edoxaban. DOACs have been shown to be superior or non-inferior to warfarin in the prevention of stroke and systemic embolism with similar or lower risk of major bleeds. Especially risk of intracranial bleeds is lower with DOACs.<sup>3–6</sup> Accumulating reallife evidence support these findings.<sup>7–9</sup>

DOACs have been shown to be cost-effective in comparison to warfarin in numerous treatment settings,<sup>10–16</sup> including Finland.<sup>17,18</sup> As a result, the use of DOACs as the first treatment alternative for the prevention of thromboembolic complications in patients with non-valvular atrial fibrillation (AF) has steadily increased over time.<sup>19,20</sup> In Finland, the use of warfarin has drastically decreased, with an almost 50% reduction in the number of users since 2016,<sup>21</sup> despite the aging of the population. At the same time, the use of DOACs has more than tripled with almost half of the DOAC-treated patients currently using apixaban.<sup>21</sup> Currently available published evidence of relative cost-effectiveness of DOACs is mostly based on indirect comparative evidence from the clinical trials.<sup>22-25</sup> Since clinical trials often enroll more homogeneous patient populations differing from those being treated in the real-life setting, it is of interest to investigate the relative cost-effectiveness of DOACs based on the comparative effectiveness findings from real-world studies. Such evidence is useful in assessing how the societal resources can be allocated in a cost-effective manner. Here, we report results from a cost-effectiveness assessment comparing apixaban to two other DOACs (dabigatran, rivaroxaban) and warfarin in the Finnish real-world setting.

## **Materials and Methods**

The cost-effectiveness assessment was performed using a lifetime Markov state transition model with 6-week cycles. The model has been previously adapted and modified for cost-effectiveness assessments in numerous countries including Finland,<sup>10,11,18,26</sup> and its structure and validation has been described in detail previously.<sup>10,11</sup> Here, the model structure is therefore only briefly summarized.

In the current assessment apixaban was compared to rivaroxaban, dabigatran and warfarin based on the relative effectiveness data available from the real-world NAXOS study (Evaluation of Apixaban in Stroke and Systemic Embolism Prevention in Patients With Nonvalvular Atrial Fibrillation),<sup>7</sup> and representative Finnish input data for event risks, mortality, resource use, costs, and quality of life. NAXOS-study was chosen to provide relative effectiveness estimates for the comparison, because it is one of the largest real-world studies conducted on DOACs and thus far the largest in Europe. Because the NAXOS-study compared apixaban separately to each of the comparators using propensity score matching,<sup>7</sup> the relative costeffectiveness was likewise assessed pair-wise as apixaban versus comparator, and comparisons between the other three anticoagulants could not be assessed in this setting. As NAXOS did not include comparison between apixaban and edoxaban (due to lack of reimbursement for the latter), edoxaban was excluded from the current analysis.

The primary outcome measures for this analysis were the incremental cost-effectiveness ratio (ICER), given as cost per quality-adjusted life year (QALY) gained and probability of cost-effectiveness based on the societal willingness-to-pay (WTP). In addition, net monetary benefit (NMB =  $\Delta$ QALY \* WTP threshold –  $\Delta$ costs) was estimated to assess the value of apixaban versus its comparators in monetary terms. In addition, a cost-effectiveness acceptability curve for each comparison was drawn to depict the probability of cost–effectiveness of apixaban versus each comparator at different values of WTP per QALY gained.

In Finland, Health Technology Assessments including cost-effectiveness analyses for new innovative treatments or significant extensions of a therapeutic indication are appraised by separate governmental appraisal committees depending on where the treatment is administered. New hospital-only medicinal products are appraised by the Finnish Medicines Agency (Fimea) whereas selfadministered new medicinal products that are used in an outpatient setting are appraised by the Finnish Pharmaceuticals Pricing Board. As for now, there are no publicly announced WTP thresholds in Finland that could be used to determine whether a treatment, new or existing, is considered as cost-effective. When evaluating possible price discount levels for new hospital only products, Fimea has used WTP thresholds ranging from 50,000 €/QALY to 100,000 €/OALY in its recent appraisals.<sup>27,28</sup> Since no official Finnish threshold value for cost-effectiveness exists for established outpatient products, we applied the value of  $35,000 \notin QALY$  gained as the threshold value, which has been used in Finnish cost-effectiveness assessments in chronic diseases.<sup>29</sup> This threshold is in line with the ICER-threshold (20,000–30,000  $\pounds QALY$ ) applied by NICE.<sup>30</sup>

#### Model

The modelled cohort consists of 50.6% male and have an average age of 76.3 years at baseline reflecting the current Finnish AF population using anticoagulants.<sup>19</sup> The model (Figure 1) captures the health and cost outcomes of the AF cohort starting anticoagulation treatment (AF health state) through transitions to mutually exclusive health states as the result of the following events: ischemic stroke (IS), myocardial infarction (MI), systemic embolism (SE), intracranial hemorrhage (ICH; further classified as either hemorrhagic stroke (HS) or other ICH), other major bleeds (further classified as gastrointestinal [GI] or non-GI bleed) and clinically relevant non-major bleeds. IS, HS, MI and SE events are modelled as permanent health states where the patients reside until death. The remaining health states are modelled as transient states where patients reside for one model cycle.

In the analysis, apixaban served as the reference product to which the other products were compared. The baseline risk of the modelled events for apixaban treated patients in Finnish real-world setting were derived from the rates reported for warfarin treated AF patients in Finnish real-world studies<sup>31-33</sup> and relevant clinical trial evidence where Finnish data is currently lacking (Table 1). The risk of event for treatment comparators were then derived from apixaban rates through the hazard ratios detailed in Table 1. To maintain comparability between the previously conducted clinical trial-based costeffectiveness analysis for warfarin and apixaban in the Finnish setting, the modelling assumptions and inputs (eg, age-associated increase in the risk of events, severity of IS and HS events, proportion of GI-bleeds among other major bleeds, proportion of HS among ICH, event fatality and post-event mortality risk adjustment factors) were mostly implemented as reported previously by Hallinen et al.<sup>18</sup> (Supplementary file, Tables S1–S3).

For simplicity, the assessment considers only the first line antithrombotic treatment. Patients experiencing HS were assumed to discontinue the first line anticoagulation treatment permanently whereas patients with other ICHs



Figure I Schematic presentation of the model. \*Mild, moderate or severe, #transient state (ie, the patients transit back to the previous health state after one cycle and either continue or discontinue anticoagulation treatment). Reproduced from Hallinen T, Soini EJ, Linna M, Saarni SI. Cost-effectiveness of apixaban and warfarin in the prevention of thromboembolic complications among atrial fibrillation patients. *Springerplus*. 2016;5(1):1354, under the terms of the creative commons attribution 4.0 international license (http://creativecommons.org/licenses/by/4.0/).<sup>18</sup>

	Rate per 100 Pa	tient Years	Hazard Ratio (95% CI) vs	Apixaban	
	Apixaban	No Treatment <sup>26</sup>	Warfarin	Dabigatran	Rivaroxaban
Stroke <sup>a</sup>	1.359 <sup>b</sup>	4.186	1.67 (1.54–1.79) <sup>7</sup>	1.08 (0.90–1.28) <sup>7</sup>	0.95 (0.87–1.03) <sup>7</sup>
Intracranial hemorrhage	0.249 <sup>b</sup>	0	2.38 (2.08–2.70) <sup>7</sup>	1.02 (0.62–1.68) <sup>7</sup>	1.15 (0.99–1.33) <sup>7</sup>
Other major bleed	1.603 <sup>c</sup>	0	2.33 (2.17–2.50) <sup>7</sup>	1.08 (0.93–1.24) <sup>7</sup>	1.49 (1.39–1.59) <sup>7</sup>
Clinically relevant non-major bleed	2.090 <sup>d</sup>	0	l.433 (l.241–1.655) <sup>d</sup>	1.00 (0.900–1.100) <sup>34</sup>	1.52 (1.28–1.80) <sup>34</sup>
Myocardial infarction	1.654 <sup>e</sup>	1.003	1.149 <sup>3</sup>	1.460 (0.960–2.200) <sup>34</sup>	1.060 (0.730–1.520) <sup>34</sup>
Systemic embolism <sup>a</sup>	0.060 <sup>f</sup>	0.956	1.67 (1.54–1.79) <sup>7</sup>	1.08 (0.90–1.28) <sup>7</sup>	0.95 (0.87–1.03) <sup>7</sup>
Other CV hospitalizations <sup>g</sup>	10.46	16.506	1.00	1.00	00.1
Treatment discontinuation for non-event related reasons	13.107 <sup>d</sup>		1.10 (1.0 <del>4</del> –1.16) <sup>d</sup>	1.50 (1.36–1.67) <sup>34</sup>	1.18 (1.08–1.61) <sup>34</sup>
<b>Notes:</b> "Assumed equal to stroke and systemic embolism. <sup>b</sup> Total number vere converted to a rate per 100 patient-years: 2.27 and 0.59, respectiv ubtracting the ICH rate from total bleeds, the rate for other maior bleed	of events among 16,741 F vely. <sup>c</sup> Total number of blev s was estimated at 3.73 pe	innish AF patients using warfarin eding events in 125,261 Finnish r 100 patient-vears. <sup>d</sup> Secondary a	were 375 for stroke and 99 for IC patients using warfarin was 5412, <sup>32</sup> analysis of ARISTOTLE-trial. °1.900 p	H. <sup>31</sup> Assuming a full follow-up of on and the corresponding rate was 4, ser 100 patient-vears for warfarin <sup>33</sup>	e year for each patient, these 32 per 100 patient-years. By 601 per 100 patient-years for

were converted to a rate per 100 patient-years: 2.27 and 0.59, respectively. <sup>c</sup>Total number of bleeding events in 125,261 Finnish patients using warfarin was 5412,<sup>32</sup> and the corresponding rate was 4.32 per 100 patient-years. <sup>ast</sup> for the rate for other major bleeds was estimated at 3.73 per 100 patient-years. <sup>d</sup>Secondary analysis of ARISTOTLE-trial. <sup>e</sup>1.900 per 100 patient-years for warfarin.<sup>33 f</sup>0.1 per 100 patient-years for the rate for other major bleeds was estimated at 3.73 per 100 patient-years. <sup>g</sup>Secondary analysis of AVERROES-trial, all treatments assumed equal. warfarin.<sup>8</sup>

and other major bleeds were assumed to discontinue treatment either permanently (56% and 25% of patients, respectively) or temporarily for 6 weeks (44% and 75%, respectively).<sup>10,18</sup> Treatment discontinuations that were not related to modelled events were modelled in accordance with the pivotal clinical trials for each product. The subsequent risk of modelled events for patients residing in AF state at the time of treatment discontinuation are detailed in Table 1. Apart for the cost of treatmentrelated dyspepsia, other adverse events were not considered in the model.

**Dove**press

# Treatment Costs and Quality of Life **Estimates**

The health-related quality of life and cost inputs were modelled in line with the analysis by Hallinen et al<sup>18</sup> apart for the health state associated cost inputs which were updated to year 2019 real values (Table 2). The assessment was performed from health care payer perspective with all costs and outcomes discounted at an annual rate of 3% in line with the national guidelines.<sup>40</sup>

According to Finnish treatment recommendations,<sup>39</sup> the routine monitoring of patients using anticoagulants should take place 1-4 times per year. Therefore, the cost of an annual general physician visit and monitoring of basic blood count (incl. thrombocytes) as well as renal and liver function were implemented as routine monitoring costs for all compared treatments. In addition, monitoring of international normalized ratio (INR) was modelled to take place for warfarin treated patients on average 18.5 times a year.<sup>41</sup> Drug costs were estimated at Finnish retail prices (excluding value added tax) in February 2021.<sup>37</sup> The DOAC costs were estimated based on the reimbursed price of the pack with approximately one-month supply for each product. Similar pack sizes for DOACs were used due to the modelling approach where drug wastage associated with treatment discontinuations are not considered due to the complexity of such modelling.

# Sensitivity Analyses

We tested the impact of discounting, modelling timeframe, and warfarin monitoring costs as deterministic sensitivity analyses. In addition, we performed the analyses using the wholesale and retail prices of the largest available pack size for DOACs. These scenarios were conducted to show the impact of the Finnish pharmaceutical pricing scheme on the results of the analyses. In Finland, the retail prices

#### Table 2 Costs and Quality of Life Inputs

Drug cost per day <sup>3</sup> Apixaban 2.5 or 5 mg, twice a day Dabigatran 110 or 150 mg, twice a day Rivaroxaban 20 mg, once a day Warfarin 5 mg2.53 2.58 2.59 0.104.58 2.59 2.59Routine care & Monitoring General physician visit INR-monitoring, warfarin only (Nurse visit Laboratory tests'119.97 <sup>35</sup> 4.85 <sup>35.46</sup> 4.58 3.20 <sup>36</sup> Health states/Events <sup>18</sup> 3.20 <sup>36</sup> 0.7430.743Schemic stroke Mild Acute Care Long-term Maintenance Severe Acute Care Lator7.719.71 967.57 4543.12 00.087Hemorrhagic Stroke Mild Acute Care Long-term Maintenance Severe Acute Care Long-term Maintenance Se		Cost <sup>b</sup>	Utility
Apixaban 2.5 or 5 ng, twice a day   2.53     Dabigatran 110 or 150 mg, twice a day   2.59     Warfarin 5 mg   0.10     Routine care & Monitoring   119.97 <sup>35</sup> General physician visit   119.97 <sup>35</sup> INR-monitoring, warfarin only (Nurse visit   13.20 <sup>36</sup> Health states/Events <sup>18</sup> 1     AF   0.00     Actue Care   4543.12     Long-term Maintenance   0     Moderate   7719.71     Actue Care   7719.71     Long-term Maintenance   967.57     Severe   7725.75     Long-term Maintenance   967.57     Severe   7725.75     Long-term Maintenance   967.57     Severe   7725.75     Long-term Maintenance   2696.51     Fata   543.61     Fata   5475.69     Long-term Maintenance   2696.51     Long-term Maintenance   2696.51     Long-term Maintenance   2183.41     Severe   9455.08     Long-term Maintenance   2183.41     Severe   9455.08     Long-term Maint	Drug cost per day <sup>a</sup>		
Dabigatran 110 or 150 mg, twice a day Rivaroxaban 20 mg, once a day Warfarin 5 mg2.59 0.10Routine care & Monitoring General physician visit INR-monitoring, warfarin only (Nurse visit and INR-test) Laboratory tests'119.9735 6.4.8535.36Health states/Events'73.03**Health states/Events'0.10Act0.743Ischemic stroke Mild Acute Care Long-term Maintenance Severe Long-term Maintenance Long-term Maintenance Long-term Maintenance Long-term Maintenance Long-term Maintenance Severe Long-term Maintenance Long-term Maintenance Severe Acute Care Long-term Maintenance Long-term Maintenance0.0081 Long-term Maintenance Long-term Maintenance Long-term Maintenance Long-term Maintenance0.0081 Long-term Maintenance Long-term Maintenance Long-term Maintenance0.0081 Long-term Maintenance Long-term Maintenance Long-term Maintenance0.0081 Long-term Maintenance Long-term Maintenance Long-term Ma	Apixaban 2.5 or 5 mg, twice a day	2.53	
Rivaroxaban 20 mg, once a day Warfarin 5 mg2.59 0.101Routine care & Monitoring General physician visit INR-monitoring, warfarin only (Nurse visit and INR-test) Laboratory tests'119.9735 64.8535.361INR-monitoring, warfarin only (Nurse visit and INR-test) Laboratory tests'13.20361Health states/Events1810.743AF00.743Ischemic stroke Midd Acute Care Long-term Maintenance Moderate4543.12 967.57-0.087 -0.198Acute Care Long-term Maintenance Moderate7719.71 967.57-0.087 -0.198Acute Care Long-term Maintenance Moderate7719.71 967.57-0.087 -0.644Acute Care Long-term Maintenance Moderate7719.71 967.57-0.071 -0.644Acute Care Long-term Maintenance Moderate2696.51 2696.51 209.508-0.071 -0.352Hemorrhagic Stroke Midd Moderate2183.41 2696.51 209.51-0.352 -0.352Acute Care Long-term Maintenance Moderate2183.41 2696.51 209.51-0.352 -0.578Acute Care Long-term Maintenance Stasio-0.084 2125.68 106.76-0.168 -0.168 -0.168 -0.168 -0.168 -0.168 -0.168 -0.168 -0.168 -0.168 -0.168-0.0821Systemic Embolism Acute Care Long-term Maintenance3537.48 -0.168 -0.1	Dabigatran 110 or 150 mg, twice a day	2.58	
Warfarin 5 mg0.10100Routine care & Monitoring General physician visit INR-monitoring, warfarin only (Nurse visit and INR-test) Laboratory tests*119.9735 64.8535.36119.9735 64.8535.36Health states/Events18IIIAFI0.743Ischemic stroke Mild Acute Care Long-term Maintenance Moderate4543.12 967.570.087 4543.12 0Moderate Acute Care Long-term Maintenance Moderate7719.71 967.57-0.087 -0.198Hemorrhagic Stroke Mild Acute Care Long-term Maintenance Moderate7725.75 4403.61 5475.59-0.071 -0.071 -0.198Hemorrhagic Stroke Mild Acute Care Long-term Maintenance Poffer Acute Care Long-term Maintenance Severe Acute Care Long-term Maintenance Site Site Site Site Site Site Site Site	Rivaroxaban 20 mg, once a day	2.59	
Routine care & Monitoring General physician visit INR-monitoring, warfarin only (Nurse visit and INR-test) Laboratory tests <sup>6</sup> I 19.97 <sup>35</sup> 64.85 <sup>35,36</sup> IINR-monitoring, warfarin only (Nurse visit and INR-test) Laboratory tests <sup>6</sup> IIIHealth states/Events <sup>18</sup> I0.743IAF00.743IIIschemic stroke Mild Acute Care Long-term Maintenance Severe4543.12 00-0.087Acute Care Long-term Maintenance7719.71 97.57-0.087-0.198Acute Care Long-term Maintenance7719.71 97.57-0.644Acute Care Long-term Maintenance7725.75-0.644Acute Care Long-term Maintenance2696.51 269.51-0.071Acute Care Long-term Maintenance2696.51 269.51-0.071Acute Care Long-term Maintenance2696.51 269.51-0.578Acute Care Long-term Maintenance2183.41 3818.62-0.578Acute Care Long-term Maintenance2125.68 381.62-0.084Care Care Long-term Maintenance2125.68 366.50-0.168 -0.168Gib Bieds Non-ICH and Non-GI Related Major Bleeds353.748 353.748-0.081MI Acute Care Long-term Maintenance353.748-0.0822CRNM Bleeds2058.10-0.0582MI Acute Care Long-term Maintenance353.748-0.0822CRNM Bleeds2058.10-0.0582MI Acute Care Long-term Maintenance353.748-0.00582MI Acute Care	Warfarin 5 mg	0.10	
Instruct care of Former only (Nurse visit and INR-test) Laboratory tests <sup>c</sup> 119.97 <sup>35</sup> 64.85 <sup>35,36</sup> INR-monitoring, warfarin only (Nurse visit and INR-test) Laboratory tests <sup>c</sup> 0Health states/Events <sup>18</sup> 0AF0Ischemic stroke Mild Acute Care Long-term Maintenance0Moderate Acute Care Long-term Maintenance0Noderate Acute Care Long-term Maintenance967.57 4004 4004Hemorrhagic Stroke Mild Acute Care Long-term Maintenance0Hemorrhagic Stroke Mild Acute Care Long-term Maintenance0Hemorrhagic Stroke Mild Acute Care Long-term Maintenance0Hemorrhagic Stroke Mild Acute Care Long-term Maintenance0Moderate Acute Care Long-term Maintenance0Moderate Acute Care Long-term Maintenance0Moderate Acute Care Long-term Maintenance0Systemic Embolism Acute Care Long-term Maintenance0Systemic Embolism Acute Care Long-term Maintenance0Systemic Embolism Acute Care Long-term Maintenance0Systemic Embolism Acute Care Long-term Maintenance0Suppleeds Other ICH (excluding hemorrhagic stroke) Other Major Bleeds (excluding ICH) GI Bleeds Non-ICH and Non-GI Related Major Bleeds0MI Acute Care Long-term Maintenance0-0.0812CRNM Bleeds2058.10-0.0582MI Acute Care Long-term Maintenance0Supple Other ICH (excluding hemorrhagic stroke) Acute Care L	Routine care & Monitoring		
INR-monitoring, warfarin only (Nurse visit and INR-test) Laboratory tests <sup>c</sup> Intermination (ALS)Intermination (ALS)Health states/Events <sup>18</sup> IIAFI0.743Ischemic stroke Mild Acute Care Long-term Maintenance Moderate4543.12 0 1.00870.7087 1.0087Acute Care Long-term Maintenance Moderate7719.71 967.57-0.087 1.0087Acute Care Long-term Maintenance Moderate7719.71 967.57-0.644 1.0087Acute Care Long-term Maintenance Moderate7725.75 4403.61 1.5475.59-0.071 1.0087Hemorrhagic Stroke Mild Acute Care Long-term Maintenance Moderate1.0087 1.0087-0.071 1.0097Acute Care Long-term Maintenance Moderate2696.51 2.096.51 2.009.51-0.051 1.009.51Acute Care Long-term Maintenance Severe945.08 2.018.41 2.009.51-0.0582Acute Care Long-term Maintenance Severe945.08 2.018.41-0.084 2.018.41Severe Acute Care Long-term Maintenance Cong-term Maintenance Long-term Maintenance-0.084 2.125.68 1.06.76-0.084 3.06.76Systemic Embolism Acute Care Long-term Maintenance3537.48 3.537.48-0.081 3.537.48CRNM Bleeds2058.10-0.0582Mi Acute Care Long-term Maintenance2058.10-0.0582Mi Acute Care Long-term Maintenance2574.02-0.0582Mi Acute Care Long-term Maintenance2585.01-0.0582Mi Acute Care Long-term Maintena	General physician visit	119.97 <sup>35</sup>	
and INR-test) Laboratory tests <sup>c</sup> 13.20 <sup>36</sup> Health states/Events <sup>18</sup> IAF0.743Ischemic stroke Mild Acute Care Long-term Maintenance Moderate4543.12 0 00.783Acute Care Long-term Maintenance Moderate4543.12 0 00.087 -0.087Acute Care Long-term Maintenance Moderate7719.71 967.57 967.57-0.644Acute Care Long-term Maintenance Severe Long-term Maintenance Severe7725.75 4403.61 5475.59-0.071 -0.071Hemorrhagic Stroke Mild Acute Care Long-term Maintenance Moderate-0.071 2696.51 2696.51 2696.51 2696.51 2696.51 2696.51 2696.51 2183.41 Severe Acute Care Long-term Maintenance Severe-0.071 2696.51 2696.51 2696.51 2696.51 2696.51 2183.41 Severe Acute Care Long-term Maintenance Severe Acute Care Long-term Maintenance Severe Acute Care Long-term Maintenance Sistemic Embolism Acute Care Long-term Maintenance Sistemic Embolism Acute Care Long-term Maintenance Sistemic Embolism Acute Care Long-term Maintenance-0.084 2125.68 3537.48-0.084 3537.48Fill GI Bleeds Non-ICH and Non-GI Related Major Bleeds2058.10-0.0582Mil Acute Care Long-term Maintenance2058.10-0.0582Mil Acute Care Long-term Maintenance2058.10-0.0582Mil Acute Care Long-term Maintenance2058.10-0.0582Mil Acute Care Long-term Maintenance2058.10-0.0582Mil Acute Care Long-term Maintenance2058.10<	INR-monitoring, warfarin only (Nurse visit	64.85 <sup>35,36</sup>	
Laboratory tests <sup>c</sup> 13.20 <sup>36</sup> 13.20 <sup>36</sup> Health states/Events <sup>18</sup> I0AF00.743Ischemic stroke Mild Acute Care Long-term Maintenance Moderate Acute Care Long-term Maintenance Severe Acute Care Long-term Maintenance Moderate Acute Care Long-term Maintenance Severe Acute Care Long-term Maintenance Mild Acute Care Long-term Maintenance Severe Acute Care Long-term Maintenance Fatal-0.087 453.12 0-0.087 -0.087 -0.198 -0.198 -0.198 -0.198Hemorrhagic Stroke Mild Acute Care Long-term Maintenance Severe Acute Care Long-term Maintenance Severe Acute Care Long-term Maintenance Severe Acute Care Long-term Maintenance Severe Acute Care Long-term Maintenance Severe Acute Care Long-term Maintenance Severe Acute Care Long-term Maintenance-0.071 2696.51 2696.51 2183.4	and INR-test)		
Health states/Events18IIAF0.743Ischemic stroke Mild Acute Care Long-term Maintenance Moderate4543.12 0-0.087 -0.198Acute Care Long-term Maintenance Moderate7719.71 967.57-0.198 -0.198Acute Care Long-term Maintenance Fatal7725.75 4403.61 5475.59-0.644Hemorrhagic Stroke Mild Acute Care Long-term Maintenance Fatal7725.75 2696.51 26	Laboratory tests <sup>c</sup>	13.20 <sup>36</sup>	
AFInterpretationInterpretationIschemic stroke Mild Acute Care Long-term Maintenance Moderate4543.12 0-0.087 4543.12 0Acute Care Long-term Maintenance Severe Acute Care Long-term Maintenance Severe Acute Care Long-term Maintenance Fatal7719.71 967.57 967.57 1000000000000000000000000000000000000	Health states/Events <sup>18</sup>		
Ar0.743Ischemic stroke-0.087Mild-0.087Acute Care4543.12Long-term Maintenance0Moderate7719.71Long-term Maintenance967.57Severe-0.644Acute Care7725.75Long-term Maintenance4403.61Fatal5475.59Hemorrhagic Stroke-0.071Mild-0.071Acute Care2696.51Long-term Maintenance2696.51Fatal-0.071Acute Care2696.51Long-term Maintenance2696.51Moderate2696.51Long-term Maintenance2696.51Long-term Maintenance2696.51Long-term Maintenance2183.41Severe-0.578Acute Care2183.41Long-term Maintenance3818.62Fatal5708.06Systemic Embolism-0.084Acute Care106.76Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)-0.168Other Major Bleeds (excluding ICH)-0.168GI Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48Mil-0.058.10Acute Care2058.10Long-term Maintenance5453.00Other Major Bleeds5453.00CRNM Bleeds5453.00Acute Care5453.00Long-term Maintenance5453.00Care5453.00Acute Care5453.00Acute Care <td< td=""><td>AE</td><td></td><td>0.742</td></td<>	AE		0.742
Ischemic strokeImage: strokeImag			0.743
Mild   -0.087     Acute Care   4543.12     Long-term Maintenance   0     Moderate   7719.71     Acute Care   7719.71     Long-term Maintenance   967.57     Severe   7725.75     Long-term Maintenance   4403.61     Fatal   5475.59     Hemorrhagic Stroke   -0.071     Mild   -0.071     Acute Care   2696.51     Long-term Maintenance   2696.51     Mild   -0.352     Acute Care   2696.51     Long-term Maintenance   2183.41     Moderate   -0.578     Acute Care   2183.41     Severe   3818.62     Long-term Maintenance   3818.62     Evere Maintenance   3818.62     Fatal   5708.06     Systemic Embolism   -0.168     Acute Care   106.76     Long-term Maintenance   106.76     Other Major Bleeds (excluding ICH)   -0.168     Gi Bleeds   3537.48     Non-ICH and Non-GI Related Major Bleeds   3537.48     Mil   <	Ischemic stroke		
Acute Care4543.12Long-term Maintenance0Moderate7719.71Moderate967.57Acute Care967.57Severe7725.75Long-term Maintenance4403.61Fatal5475.59Hemorrhagic Stroke-0.071Mild2696.51Long-term Maintenance2696.51Long-term Maintenance2696.51Long-term Maintenance2696.51Moderate2696.51Long-term Maintenance2696.51Long-term Maintenance2183.41Severe9455.08Long-term Maintenance2183.41Severe9641.27Acute Care2183.41Long-term Maintenance3818.62Fatal5708.06Systemic Embolism-0.084Acute Care106.76Other Major Bleeds (excluding ICH)753.748Gl Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds353.748Mil-0.0582Acute Care2058.10Long-term Maintenance5453.00Gl Bleeds353.748Non-ICH and Non-GI Related Major Bleeds353.748Non-ICH and Non-GI Related Major Bleeds353.748Acute Care5453.00Long-term Maintenance5453.00Acute Care5453.00Acute Care5453.00Acute Care5453.00Acute Care5453.00Acute Care5453.00Acut	Mild		-0.087
Long-term Maintenance     0     -0.198       Moderate     7719.71     -0.198       Acute Care     967.57     -0.644       Long-term Maintenance     967.57     -0.644       Acute Care     7725.75     -0.644       Acute Care     4403.61     -0.071       Experim Maintenance     4403.61     -0.071       Fatal     5475.59     -0.071       Acute Care     2696.51     -0.071       Long-term Maintenance     2696.51     -0.352       Acute Care     2696.51     -0.352       Acute Care     2183.41     -0.578       Long-term Maintenance     2183.41     -0.578       Acute Care     9641.27     -0.578       Acute Care     3818.62     -0.578       Acute Care     106.76     -0.168       Long-term Maintenance     106.76     -0.168       Systemic Embolism     -0.168     -0.168       Acute Care     106.76     -0.168       Long-term Maintenance     3537.48     -0.168       Gl Bleeds     3537.48	Acute Care	4543.12	
Moderate-0.198Acute Care7719.71Long-term Maintenance967.57Severe7725.75Acute Care7725.75Long-term Maintenance4403.61Fatal5475.59Hemorrhagic Stroke-0.071Mild-0.071Acute Care2696.51Long-term Maintenance2696.51Moderate2696.51Long-term Maintenance9455.08Long-term Maintenance2183.41Severe9455.08Long-term Maintenance3818.62Long-term Maintenance3818.62Long-term Maintenance3818.62Systemic Embolism-0.084Acute Care2125.68Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)4366.50Other Major Bleeds (excluding ICH)-0.168Gl Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds353.748Acute Care2058.10Long-term Maintenance363.50CRNM Bleeds2058.10Acute Care5453.00Long-term Maintenance538.50	Long-term Maintenance	0	
Acute Care7719.71Long-term Maintenance967.57Severe7725.75Acute Care7725.75Long-term Maintenance4403.61Fatal5475.59Hemorrhagic Stroke-0.071Mild-0.071Acute Care2696.51Long-term Maintenance2696.51Moderate-0.352Acute Care2183.41Severe9455.08Long-term Maintenance2183.41Severe9641.27Acute Care9641.27Long-term Maintenance3818.62Fatal5708.06Systemic Embolism-0.084Acute Care2125.68Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)4366.50Other Major Bleeds (excluding ICH)3537.48Gi Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48Acute Care2058.10Long-term Maintenance5453.00Gi Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48Acute Care5453.00Long-term Maintenance538.50	Moderate		-0.198
Long-term Maintenance967.57Severe-0.644Acute Care7725.75Long-term Maintenance4403.61Fatal5475.59Hemorrhagic Stroke-0.071Mild-0.071Acute Care2696.51Long-term Maintenance2696.51Moderate-0.352Acute Care2183.41Severe9455.08Long-term Maintenance2183.41Severe9641.27Acute Care2183.41Long-term Maintenance3818.62Fatal5708.06Systemic Embolism-0.084Acute Care2125.68Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)3537.48Other Major Bleeds (excluding ICH)5337.48Gi Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48Acute Care2058.10Long-term Maintenance5453.00Gi Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48Acute Care5453.00Acute Care5453.00Acute Care5453.00Acute Care538.50Acute Care5453.00Acute Care538.50	Acute Care	7719.71	
Severe-0.644Acute Care7725.75Long-term Maintenance4403.61Fatal5475.59Hemorrhagic Stroke-0.071Mild-0.071Acute Care2696.51Long-term Maintenance2696.51Moderate2696.51Acute Care2183.41Severe-0.578Acute Care9641.27Long-term Maintenance3818.62Fatal5708.06Systemic Embolism-0.084Acute Care106.76Long-term Maintenance106.76Gther ICH (excluding hemorrhagic stroke)-0.168Other ICH and Non-Gl Related Major Bleeds3537.48Non-ICH and Non-Gl Related Major Bleeds2058.10Acute Care2058.10Long-term Maintenance2058.10Other ICH maintenance3537.48Non-ICH and Non-Gl Related Major Bleeds3537.48Acute Care2058.10Long-term Maintenance2058.10Other ICH maintenance2058.10Other ICH maintenance3537.48Non-ICH and Non-Gl Related Major Bleeds3537.48Acute Care2453.00Long-term Maintenance2453.00Subsection5453.00Acute Care5453.00Acute Care5453.00Acute Care5453.00Acute Care5453.00Acute Care5453.00Acute Care5453.00Acute Care5453.00Acute Care5453.00Acute Care5453.00	Long-term Maintenance	967.57	
Acute Care7725.75Long-term Maintenance4403.61Fatal5475.59Hemorrhagic Stroke-0.071Mild-0.071Acute Care2696.51Long-term Maintenance2696.51Moderate-0.352Acute Care9455.08Long-term Maintenance2183.41Severe-0.578Acute Care9641.27Long-term Maintenance3818.62Fatal5708.06Systemic Embolism-0.084Acute Care106.76Long-term Maintenance106.76Systemic Embolism-0.168Acute Care106.76Joher ICH (excluding hemorrhagic stroke)-0.168Other ICH (excluding hemorrhagic stroke)3537.48Non-ICH and Non-GI Related Major Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds2058.10Acute Care2058.10Long-term Maintenance5453.00Gi Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48Acute Care2058.10Long-term Maintenance2453.00State Care5453.00Acute Care <td>Severe</td> <td></td> <td>-0.644</td>	Severe		-0.644
Long-term Maintenance4403.61Fatal5475.59Hemorrhagic Stroke-0.071Mild-0.071Acute Care2696.51Long-term Maintenance2696.51Moderate9455.08Long-term Maintenance2183.41Severe9455.08Long-term Maintenance2183.41Severe9641.27Long-term Maintenance3818.62Fatal5708.06Systemic Embolism-0.084Acute Care106.76Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)4366.50Other Major Bleeds (excluding ICH)-0.168GI Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds2058.10MI-0.005Acute Care2058.10Long-term Maintenance5453.00CRNM Bleeds538.50MI-0.058Acute Care5453.00Long-term Maintenance538.50	Acute Care	//25./5	
Fatal54/5.59Hemorrhagic Stroke0.071Mild0.071Acute Care2696.51Long-term Maintenance2696.51Moderate0.352Acute Care9455.08Long-term Maintenance2183.41Severe0.578Acute Care9641.27Long-term Maintenance3818.62Fatal5708.06Systemic Embolism0.071Acute Care106.76Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)4366.50Other Major Bleeds (excluding ICH)-0.168Gl Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48MI-0.058Acute Care2058.10Long-term Maintenance5453.00CRNM Bleeds5453.00Acute Care5453.00Long-term Maintenance538.50	Long-term Maintenance	4403.61	
Hemorrhagic StrokeIIMildI-0.071Acute Care2696.51-0.352Long-term Maintenance2696.51-0.352Acute Care9455.08-0.352Long-term Maintenance2183.41-0.578Severe9451.07-0.578Acute Care9641.27-0.578Long-term Maintenance3818.62-0.578Fatal5708.06-0.084Acute Care2125.68-0.168Long-term Maintenance106.76-0.168Other ICH (excluding hemorrhagic stroke)3537.48-0.168Other Major Bleeds (excluding ICH)-0.168-0.168Gl Bleeds3537.48537.48-0.0582MIAcute Care2058.10-0.0582Long-term Maintenance5453.00-0.005Acute Care5453.00-0.005Acute Care5453.00<	Fatal	54/5.59	
Mild   -0.071     Acute Care   2696.51     Long-term Maintenance   2696.51     Moderate   2696.51     Moderate   2696.51     Acute Care   2696.51     Long-term Maintenance   2183.41     Severe   2183.41     Acute Care   9641.27     Long-term Maintenance   3818.62     Fatal   5708.06     Systemic Embolism   -0.084     Acute Care   2125.68     Long-term Maintenance   106.76     Other ICH (excluding hemorrhagic stroke)   4366.50   -0.168     Other ICH (excluding hemorrhagic stroke)   3537.48   -0.0582     GI Bleeds   3537.48   5357.48   -0.0582     MI   2058.10   -0.0582     MI   -0.005   -0.005     Acute Care   5453.00   -0.005     Long-term Maintenance   538.50   -0.005	Hemorrhagic Stroke		
Acute Care2696.51Long-term Maintenance2696.51Moderate2696.51Acute Care9455.08Long-term Maintenance2183.41Severe2183.41Acute Care9641.27Long-term Maintenance3818.62Fatal5708.06Systemic Embolism-0.084Acute Care106.76Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)4366.50Other Major Bleeds (excluding ICH)-0.168Gl Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48CRNM Bleeds2058.10Acute Care2058.10Long-term Maintenance5453.00CRNM Bleeds5453.00Acute Care5453.00Acute Care538.50CRNM Gleade Care538.50Acute Care5453.00Acute Care5453.00Acu	Mild		-0.071
Long-term Maintenance2696.51Moderate-0.352Acute Care9455.08Long-term Maintenance2183.41Severe2183.41Acute Care9641.27Long-term Maintenance3818.62Fatal5708.06Systemic Embolism-0.084Acute Care2125.68Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)4366.50Other ICH (excluding hemorrhagic stroke)3537.48Other Major Bleeds (excluding ICH)3537.48Gl Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48CRNM Bleeds2058.10MI-0.0582MI-0.005Acute Care5453.00Long-term Maintenance538.50	Acute Care	2696.51	
Moderate-0.352Acute Care9455.08Long-term Maintenance2183.41Severe-0.578Acute Care9641.27Long-term Maintenance3818.62Fatal5708.06Systemic Embolism-0.084Acute Care2125.68Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)4366.50Other Major Bleeds (excluding ICH)-0.168GI Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48CRNM Bleeds2058.10Acute Care5453.00Long-term Maintenance5453.00CRNM Bleeds538.50	Long-term Maintenance	2696.51	
Acute Care9455.08Long-term Maintenance2183.41Severe2183.41Acute Care9641.27Long-term Maintenance3818.62Fatal5708.06Systemic Embolism-0.084Acute Care2125.68Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)4366.50Other Major Bleeds (excluding ICH)-0.168GI Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48CRNM Bleeds2058.10Acute Care2058.10Long-term Maintenance5453.00CRNM Bleeds538.50	Moderate		-0.352
Long-term Maintenance2183.41Severe-0.578Acute Care9641.27Long-term Maintenance3818.62Fatal5708.06Systemic Embolism-0.084Acute Care2125.68Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)4366.50Other Major Bleeds (excluding ICH)-0.168GI Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48CRNM Bleeds2058.10Acute Care-0.0582MI-0.005Acute Care5453.00Long-term Maintenance538.50	Acute Care	9455.08	
Severe-0.578Acute Care9641.27Long-term Maintenance3818.62Fatal5708.06Systemic Embolism-0.084Acute Care106.76Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)4366.50Other Major Bleeds (excluding 1CH)-0.168Gl Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48CRNM Bleeds2058.10Acute Care5453.00Long-term Maintenance5453.00Other GM Unovication of the strike538.50	Long-term Maintenance	2183.41	0.570
Acture Care7641.27Long-term Maintenance3818.62Fatal5708.06Systemic Embolism-0.084Acute Care2125.68Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)4366.50Other Major Bleeds (excluding ICH)-0.168Gl Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48CRNM Bleeds2058.10Acute Care5453.00Long-term Maintenance538.50	Severe	0(4) 27	-0.578
Long-term Praintenance3618.82Fatal5708.06Systemic Embolism-0.084Acute Care2125.68Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)4366.50Other Major Bleeds (excluding ICH)-0.168Gl Bleeds3537.48Non-ICH and Non-Gl Related Major Bleeds3537.48CRNM Bleeds2058.10Acute Care5453.00Long-term Maintenance538.50	Acute Care	2010/2	
Systemic Embolism-0.084Acute Care2125.68Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)4366.50Other Major Bleeds (excluding ICH)-0.168GI Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48CRNM Bleeds2058.10Acute Care5453.00Long-term Maintenance538.50		5010.02	
Systemic Embolism0.084Acute Care2125.68Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)4366.50Other Major Bleeds (excluding ICH)-0.168Gl Bleeds3537.48Non-ICH and Non-Gl Related Major Bleeds3537.48CRNM Bleeds2058.10Acute Care5453.00Long-term Maintenance538.50		3708.06	
Acute Care2125.68Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)4366.50Other Major Bleeds (excluding ICH)-0.168GI Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48CRNM Bleeds2058.10Acute Care5453.00Long-term Maintenance538.50	Systemic Embolism		-0.084
Long-term Maintenance106.76Other ICH (excluding hemorrhagic stroke)4366.50Other Major Bleeds (excluding ICH)-0.168GI Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48CRNM Bleeds2058.10CRNM Bleeds-0.0582MI-0.005Acute Care5453.00Long-term Maintenance538.50	Acute Care	2125.68	
Other ICH (excluding hemorrhagic stroke)4366.50-0.168Other Major Bleeds (excluding ICH)3537.48-0.168GI Bleeds3537.483537.48Non-ICH and Non-GI Related Major Bleeds3537.48-0.0582CRNM Bleeds2058.10-0.0582MI-0.005-0.005Acute Care5453.00-0.005Long-term Maintenance538.50-0.005	Long-term Maintenance	106.76	
Other Major Bleeds (excluding ICH)-0.168GI Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48CRNM Bleeds2058.10CRNM Bleeds-0.0582MI-0.005Acute Care5453.00Long-term Maintenance538.50	Other ICH (excluding hemorrhagic stroke)	4366.50	-0.168
GI Bleeds3537.48Non-ICH and Non-GI Related Major Bleeds3537.48CRNM Bleeds2058.10-0.0582MI-0.005Acute Care5453.00Long-term Maintenance538.50	Other Major Bleeds (excluding ICH)		-0.168
Non-ICH and Non-GI Related Major Bleeds3537.48CRNM Bleeds2058.10-0.0582MI Acute Care Long-term Maintenance-0.005States SM Lines (clinication)5453.00Other SM Lines (clinication)2771.02	GI Bleeds	3537.48	
CRNM Bleeds     2058.10     -0.0582       MI     -0.005     -0.005       Acute Care     5453.00     -0.005       Long-term Maintenance     538.50     -0.005	Non-ICH and Non-GI Related Major Bleeds	3537.48	
MI–0.005Acute Care5453.00Long-term Maintenance538.50		2058.10	-0.0582
Acute Care 5453.00   Long-term Maintenance 538.50	м		-0.005
Long-term Maintenance 538.50	Acute Care	5453.00	
	Long-term Maintenance	538.50	
Other CV Hospitalization 37/4.93	Other CV Hospitalization	3774.93	

(Continued)

#### Table 2 (Continued).

	Cost <sup>b</sup>	Utility
Adverse events	125 49	
<b>D</b> /spepsia	123.17	

**Notes:** <sup>a</sup>Drug costs were assessed based on the cost of the reimbursed price of the pack with approximately one-month supply for each product in 2/2021.<sup>37</sup> <sup>b</sup>Conversion from previous years (to year 2019 real values) were made using the communal health care price index by statistics Finland.<sup>38</sup> <sup>c</sup>Basic blood count, thrombocytes, creatinine, estimated glomerular filtration rate, alanine aminotransferase.<sup>39</sup> <sup>d</sup>Assumption: GP visit, pantoprazole 20 mg 100 tablets. The proportion of patients experiencing dyspepsia was obtained from the respective clinical trials and were 1.7% for apixaban and rivaroxaban, 3.5% for dabigatran, and 1.8% for warfarin.

are determined from the wholesale prices using a stepwise scheme that provides the dispensing pharmacies with diminishing retail margins as the wholesale price increases. Thus, the Finnish pricing scheme tends to artificially increase the cost of products with smaller pack sizes relative to those with larger pack sizes.<sup>42</sup> Rivaroxaban is currently available in packs containing tablets for 28-day and 98-day treatment whereas dabigatran and apixaban packs provide tablets for a 30-day treatment.

A probabilistic sensitivity analysis (PSA) was performed to explore the impact of parameter uncertainty on the outcomes of the assessment. The values of the key input parameters were varied based on their probability distributions over 2000 simulations and a cost–effectiveness plane was drawn of the findings to illustrate the observed differences in costs and effects between apixaban and its comparators. Probabilities of cost-effectiveness were separately reported for strong dominance (<0  $\epsilon$ /QALY) and cost-effectiveness (<35,000  $\epsilon$ /QALY).

## Results

## Base Case Analysis

Apixaban increased life-expectancy and quality-adjusted life-expectancy compared to warfarin, dabigatran, and rivaroxaban (Table 3). The discounted gain in life-years and QALYs with apixaban was 0.05 and 0.03 years compared to rivaroxaban, and 0.13 and 0.11 when compared to dabigatran, respectively. Highest total treatment costs over patients' lifetime were seen in the warfarin group (22,033 euros) followed by dabigatran (18,746 euros), rivaroxaban (18,385 euros), and apixaban (18,342 euros). Apixaban dominated both warfarin and other DOACs. With WTP of 35,000 euros per QALY gained, NMB of apixaban was 8723, 4168, and 1129 euros per patient when

	LY	QALY	Costs				Difference, Apixaban vs Comparator			ICER (€/QALY)
			Total	Monitoring	Event	Drug	Cost	LY	QALY	
Apixaban	8.37	6.09	18,342	1068	13,356	3918				
Rivaroxaban	8.32	6.05	18,385	1057	13,703	3625	-43	0.05	0.03	Apixaban dominates
Dabigatran	8.23	5.98	18,746	1046	14,588	3112	-404	0.13	0.11	Apixaban dominates
Warfarin	8.22	5.94	22,033	5877	16,008	149	-3691	0.15	0.14	Apixaban dominates

Table 3 Results of the Cost-Effectiveness Analysis

compared to warfarin, dabigatran, and rivaroxaban, respectively.

## Sensitivity Analyses

The cost-effectiveness plane illustrating differences between apixaban and its comparators over 2000 model simulations are shown in Figure 2. Based on these simulations, the average total treatment costs were 18,384 euros for apixaban, 18,552 euros for rivaroxaban, 19,074 euros for dabigatran and 22,237 euros for warfarin during patient's lifetime. The respective average QALYs were 6.09 for apixaban, 6.06 for rivaroxaban, 5.98 for dabigatran and 5.95 for warfarin. Apixaban dominated warfarin in 99.8% of the simulations and was cost-effective in all. As WTP increases from 35,000 to 100,000 euros/QALY, the probability of apixaban being cost-effective increases from 64% to 79% and from 92% to 99% versus rivaroxaban and dabigatran (Figure 3), respectively.

Apixaban remained cost-effective versus warfarin and other DOACs in all conducted deterministic sensitivity analyses (Table 4). Cost-savings associated with apixaban ranged from 824 to 4257 euros when compared to warfarin and from 270 to 507 euros when compared to dabigatran. When compared to rivaroxaban the cost difference ranged from -109 to 279 euros. Apixaban was associated with additional costs only in the scenario where drug costs were based on the largest pack size and retail prices where the distorting impact of the Finnish Pharmaceutical Pricing Scheme used for determining retail prices is visible.

## Discussion

Apixaban improved the outcomes of Finnish AF patients, when compared to warfarin, dabigatran, and rivaroxaban in our real-world based analysis. At the current Finnish drug prices, apixaban was also a less costly treatment alternative than warfarin despite the higher drug acquisition costs. Apixaban dominated warfarin, dabigatran, and rivaroxaban as it was both less costly and more effective. Apixaban's incremental NMB was positive in all comparisons indicating that the cost to derive the benefits associated with apixaban treatment is less than what the society is willing to pay (here 35,000 euros/QALY gained). Our analysis confirmed the importance of even slight differences in relative efficacy between the different DOACs and suggests that, overall, apixaban has slightly better overall efficacy than rivaroxaban and dabigatran. While it also appears to have slightly higher overall drug costs during patients' lifetime due to improved survival, the analysis demonstrated that the investment "pays itself back" through reduced event costs.

The results of our analysis were insensitive to changes in most modelling assumptions and the probability of apixaban being cost-effective was high in all comparisons. However, due to the Finnish pharmaceutical pricing scheme that provides the dispensing pharmacies with a lower relative retail margin for larger pack sizes, apixaban was slightly more expensive than rivaroxaban in a scenario using the retail price of the largest available rivaroxaban pack as the acquisition cost for rivaroxaban. In this scenario, apixaban remained nevertheless a costeffective treatment alternative when compared to rivaroxaban.

In our analysis, we applied the relative risk estimates for the primary efficacy and safety outcomes from the NAXOS study (Evaluation of Apixaban in Stroke and Systemic Embolism Prevention in Patients With Nonvalvular Atrial Fibrillation).<sup>7</sup> NAXOS was a French real-world study including altogether 321,501 patients initiating VKA (35.0%), apixaban (27.2%), rivaroxaban (31.1%) or dabigatran (6.6%) with the aim to compare the safety, effectiveness, and mortality of the alternative anticoagulants in oral anticoagulant-naive patients with nonvalvular atrial fibrillation. Even though the NAXOS study was conducted in France we chose it for the current analyses, since it had a highly representative sample of AF patients in France and it is one of the largest real-world



Incremental effectiveness for apixaban vs. rivaroxaban, QALYs

Figure 2 The cost-effectiveness plane for apixaban versus (A) warfarin, (B) dabigatran and (C) rivaroxaban. Green line depicts ICER threshold of 35,000 euros per QALY gained. The points lying below the line represent the simulations where apixaban was either cost-effective or dominating (South-West quadrant) versus its comparator.



Figure 3 The cost-effectiveness acceptability curve for apixaban versus warfarin, dabigatran and rivaroxaban.

studies ever conducted in Europe. Even though large realworld studies have been conducted in the United States,<sup>8,9</sup> there are remarkable differences in the organization of health care between the United States and European countries. The modelled life-expectancy of AF patients in our analyses is well in line with what would be expected for the population in Finland. The remaining life-expectancy for Finns aged 76 years is currently 12 years,<sup>43</sup> and the mortality risk in AF is approximately two-fold compared to the general population.<sup>44</sup>

In cost-effectiveness assessments the compared treatments are assumed to be perfect substitutes which means that they would be interchangeable. However, in real-world setting the treatments may nevertheless be prescribed to slightly different kind of patients. In Finland<sup>19</sup> and elsewhere<sup>45</sup> DOAC-treated patients have been younger and with less comorbidities than patients treated with warfarin. In Finland, patients using apixaban and warfarin appear to be at increased risk of thromboembolic complications (based on CHA2DS2-VASc) when compared to users of dabigatran and rivaroxaban.<sup>19</sup> In addition, apixaban is more frequently prescribed to women and patients with prior history of myocardial infarction.<sup>19</sup> Even factors not related to patient's disease characteristics may influence treatment choices in real life. Current evidence suggests that there may be inequalities in access to DOACs as DOAC-treated patients tend to have higher education and income levels.<sup>45,46</sup> In our NAXOS-study based analysis, the relative treatment effects were obtained from analyses where apixaban was separately compared to each of the comparators using propensity score matching.<sup>7</sup> Because of this, the current analysis should not be interpreted to provide evidence of the relative cost-effectiveness between the other three anticoagulants. Furthermore, it should be kept in mind that the results may not be generalizable to other countries since country-specific differences (eg drug prices, costs of other health care resources) influence the results.

Our assessment was conducted from a health care payer perspective. Therefore, the analysis excluded productivity losses as well as time and travelling costs. In the elderly patient population, the inclusion of productivity losses would have been unlikely to have a significant impact on the results of the analysis. However, the total costs associated with warfarin treatment may have been underestimated due to the exclusion of time and travelling costs which have been estimated to form 26.6% of total therapy costs for warfarin in Finland.<sup>47</sup>

## Conclusion

Apixaban use increased the life-expectancy and qualityadjusted life-expectancy of Finnish AF-patients when

Scenario	Treatment	QALY	Costs	Net Cost	Net QALY	ICER (€/QALY), Apixaban vs Comparator
No discounting	Apixaban	7.34	22,511			
	Rivaroxaban	7.30	22,553	-42	0.04	Apixaban dominates
	Dabigatran	7.20	22,951	-440	0.14	Apixaban dominates
	Warfarin	7.15	26,769	-4257	0.19	Apixaban dominates
5-year timeframe	Apixaban	3.06	8028			
	Rivaroxaban	3.05	8095	-68	0.01	Apixaban dominates
	Dabigatran	3.04	8298	-270	0.02	Apixaban dominates
	Warfarin	3.03	10,263	-2235	0.04	Apixaban dominates
Retail prices, largest available pack	Apixaban	6.09	18,342			
	Rivaroxaban	6.05	18,063	279	0.03	8978
Wholesale prices, largest available	Apixaban	6.09	17,329			
pack	Rivaroxaban	6.05	17,438	-109	0.03	Apixaban dominates
	Dabigatran	5.98	17,837	-507	0.11	Apixaban dominates
	Warfarin	5.94	21,987	-4658	0.14	Apixaban dominates
Warfarin monitoring cost reduced <sup>a</sup>	Apixaban	6.09	18,342			
	Warfarin	5.94	19,166	-824	0.14	Apixaban dominates

#### Table 4 Results of the Deterministic Sensitivity Analyses

**Notes**: <sup>a</sup>Phone call, 13.08 euros, instead of a nurse visit. Dominates = more effective, less costly.

Abbreviations: AF, atrial fibrillation; DOAC, direct oral anticoagulant; GI, gastrointestinal; HS, hemorrhagic stroke; ICH, intracranial hemorrhage; ICER, incremental costeffectiveness ratio; IS, ischemic stroke; MI, myocardial infarction; NMB, net monetary benefit; SE, systemic embolism; PSA, probabilistic sensitivity analysis; QALY, qualityadjusted life-year; WTP, willingness-to-pay.

compared to warfarin, dabigatran, and rivaroxaban in a real-world setting. Apixaban dominated warfarin, dabigatran and rivaroxaban as these gains were obtained with lower total health care costs.

## Disclosure

TH and ES are partners and employees, and CA is an employee of ESiOR Oy, which was commissioned by Pfizer Oy to perform this study. ESiOR has carried out commissioned studies and health-economic analyses for several other pharmaceutical companies, food industry companies, device companies, research groups, health care organizations, and hospitals. ML is an employee of Aalto University. PE, SS and MK are employees of Pfizer Oy. This study was sponsored by Pfizer and Bristol Myers Squibb. Medical Writing support was provided by Taru Hallinen at ESiOR Oy and was funded by Pfizer and Bristol Myers Squibb. The authors report no other conflicts of interest in this work.

# References

- 1. Hindricks G, Potpara T, Dagres N, et al; ESC Scientific Document Group. 2020 ESC guidelines for the diagnosis and management of atrial fibrillation developed in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS). *Eur Heart J*. 2021;42(5):373–498.
- Working group set up by the Finnish Medical Society Duodecim and the Finnish Cardiac Society. Atrial fibrillation. Current care guidelines. Helsinki: The Finnish Medical Society Duodecim; 2017. (referred February 5, 2021). Available from: www.kaypahoito.fi. Accessed March 16, 2021.
- Granger C, Alexander J, McMurray J, et al. Apixaban versus warfarin in patients with atrial fibrillation. N Engl J Med. 2011;365 (11):981–992. doi:10.1056/NEJMoa1107039
- Patel M, Mahaffey K, Garg J, et al. Rivaroxaban versus warfarin in nonvalvular atrial fibrillation. N Engl J Med. 2011;365(10):883–891. doi:10.1056/NEJMoa1009638
- 5. Connolly SJ, Ezekowitz MD, Yusuf S, et al. Dabigatran versus warfarin in patients with atrial fibrillation. *N Engl J Med.* 2009;361 (12):1139–1151. doi:10.1056/NEJMoa0905561
- Giugliano RP, Ruff CT, Braunwald E, et al; ENGAGE AF-TIMI 48 Investigators. Edoxaban versus warfarin in patients with atrial fibrillation. N Engl J Med. 2013;369(22):2093–2104. doi:10.1056/ NEJMoa1310907

- Van Ganse E, Danchin N, Mahé I, et al. Comparative safety and effectiveness of oral anticoagulants in nonvalvular atrial fibrillation: the NAXOS Study. *Stroke*. 2020;51(7):2066–2075. doi:10.1161/ STROKEAHA.120.028825
- Lip GYH, Keshishian A, Li X, et al. Effectiveness and safety of oral anticoagulants among nonvalvular atrial fibrillation patients. *Stroke*. 2018;49(12):2933–2944. doi:10.1161/STROKEAHA.118.020232
- Graham DJ, Baro E, Zhang R, et al. Comparative stroke, bleeding, and mortality risks in older medicare patients treated with oral anticoagulants for nonvalvular atrial fibrillation. *Am J Med.* 2019;132 (5):596–604.e11. doi:10.1016/j.amjmed.2018.12.023
- Dorian P, Kongnakorn T, Phatak H, et al. Cost-effectiveness of apixaban vs. current standard of care for stroke prevention in patients with atrial fibrillation. *Eur Heart J.* 2014;35(28):1897–1906. doi:10.1093/eurheartj/ehu006
- Lip GY, Kongnakorn T, Phatak H, et al. Cost-effectiveness of apixaban versus other new oral anticoagulants for stroke prevention in atrial fibrillation. *Clin Ther.* 2014;36(2):192–210.e20. doi:10.1016/j. clinthera.2013.12.011
- Wouters H, Thijs V, Annemans L. Cost-effectiveness of dabigatran etexilate in the prevention of stroke and systemic embolism in patients with atrial fibrillation in Belgium. *J Med Econ.* 2013;16 (3):407–414. doi:10.3111/13696998.2013.766200
- Kansal AR, Sorensen SV, Gani R, et al. Cost-effectiveness of dabigatran etexilate for the prevention of stroke and systemic embolism in UK patients with atrial fibrillation. *Heart.* 2012;98(7):573–578. doi:10.1136/heartjnl-2011-300646
- Mensch A, Stock S, Stollenwerk B, Müller D. Cost effectiveness of rivaroxaban for stroke prevention in German patients with atrial fibrillation. *Pharmacoeconomics*. 2015;33(3):271–283. doi:10.1007/ s40273-014-0236-9
- Kleintjens J, Li X, Simoens S, et al. Cost-effectiveness of rivaroxaban versus warfarin for stroke prevention in atrial fibrillation in the Belgian healthcare setting. *Pharmacoeconomics*. 2013;31 (10):909–918. doi:10.1007/s40273-013-0087-9
- Rognoni C, Marchetti M, Quaglini S, Liberato NL. Edoxaban versus warfarin for stroke prevention in non-valvular atrial fibrillation: a cost-effectiveness analysis. J Thromb Thrombolysis. 2015;39 (2):149–154. doi:10.1007/s11239-014-1104-3
- Hallinen T, Soini E, Tuovinen M. Dabigatraanin kustannusvaikuttavuus eteisvärinäpotilaiden aivohalvausten ehkäisyssä [Costeffectiveness of dabigatran compared to warfarin in the prevention of stroke and systemic embolism in Finnish AF patients]. *DOSIS*. 2012;28(2):130–144. Finnish.
- Hallinen T, Soini EJ, Linna M, Saarni SI. Cost-effectiveness of apixaban and warfarin in the prevention of thromboembolic complications among atrial fibrillation patients. *Springerplus*. 2016;5 (1):1354. doi:10.1186/s40064-016-3024-5
- Hellman T, Salo H, Kulathinal S, Eloranta P, Airaksinen J, Laine J. Oraalisten antikoagulanttien käyttö eteisvärinäpotilailla vuosina 2015–2017-rekisteritutkimus [Use of oral anticoagulants among patients with atrial fibrillation in 2015–2017 – a register study]. *Suom Laakaril*. 2020;75(47):2531–2535. Finnish.
- Guelker JE, Ilousis D, Kröger K, Santosa F, Kowall B, Stang A. Increasing use of anticoagulants in Germany and its impact on hospitalization for gastrointestinal bleeding. *Thromb Res.* 2019;181:135–140. doi:10.1016/j.thromres.2019.07.009
- 21. Kelasto statistical database [homepage on the Internet]. Reimbursements of medicine expenses: number of recipients and prescription data. The Social Insurance Institution of Finland, 2021. Available from: https://www.kela.fi/kelasto. Accessed May 28, 2021.
- 22. López-López JA, Sterne JAC, Thom HHZ, et al. Oral anticoagulants for prevention of stroke in atrial fibrillation: systematic review, network meta-analysis, and cost effectiveness analysis. *BMJ*. 2017;359: j5058. doi:10.1136/bmj.j5058

- Harrington AR, Armstrong EP, Nolan PE Jr, Malone DC. Costeffectiveness of apixaban, dabigatran, rivaroxaban, and warfarin for stroke prevention in atrial fibrillation. *Stroke*. 2013;44(6):1676–1681. doi:10.1161/STROKEAHA.111.000402
- 24. Thom HHZ, Hollingworth W, Sofat R, et al. Directly acting oral anticoagulants for the prevention of stroke in atrial fibrillation in England and Wales: cost-effectiveness model and value of information analysis. *MDM Policy Pract.* 2019;4(2):2381468319866828.
- Wisløff T, Hagen G, Klemp M. Economic evaluation of warfarin, dabigatran, rivaroxaban, and apixaban for stroke prevention in atrial fibrillation. *Pharmacoeconomics*. 2014;32(6):601–612. doi:10.1007/ s40273-014-0152-z
- 26. de Jong LA, Groeneveld J, Stevanovic J, et al. Cost-effectiveness of apixaban compared to other anticoagulants in patients with atrial fibrillation in the real-world and trial settings. *PLoS One*. 2019;14 (9):e0222658. doi:10.1371/journal.pone.0222658
- 27. Rannanheimo P, Hyvärinen A, Kiviniemi V. Atetsolitsumabi yhdessä nab-paklitakselin kanssa kolmoisnegatiivisen rintasyövän hoidossa [Atezolizumab in combination with nab-paclitaxel in the treatment of triple-negative breast cancer]. Finnish Medicines Agency Fimea. Fimea kehittää, arvioi ja informoi -julkaisusarja 9/2019. ISBN 978-952-7299-07-4. Finnish.
- 28. Wikman E, Rannanheimo P Isatuksimabi yhdessä pomalidomidin ja deksametasonin kanssa uusiutuneen multippelin myelooman hoidossa [Isatuximab together with pomalidomide and dexamethasone in the treatment of relapsed multiple myeloma]. Finnish Medicines Agency Fimea. Fimea kehittää, arvioi ja informoi -julkaisusarja 3/2020. ISBN 978-952-7299-10-4. Finnish.
- Joensuu JT, Huoponen S, Aaltonen KJ, Konttinen YT, Nordström D, Blom M. The cost-effectiveness of biologics for the treatment of rheumatoid arthritis: a systematic review. *PLoS One.* 2015;10(3): e0119683. doi:10.1371/journal.pone.0119683
- 30. National Institute for Health and Care Excellence. Guide to the methods of technology appraisal 2013. Process and methods [PMG9]; 2013. Available from: https://www.nice.org.uk/article/ PMG9/chapter/Foreword. Accessed May 20, 2021.
- 31. Linna M, Keto J, Piuhola J, Vesalainen R, Hällberg V, Laine J. Eteisvärinäpotilaan sosiaali- ja terveydenhuoltopalvelujen käyttö komplikaation jälkeen [Use of social and health care services by patients with complications of atrial fibrillation]. *Suom Laakaril.* 2017;35:1856–1862. Finnish.
- 32. Penttilä T, Lehto M, Niiranen J, et al. Differences in the risk of stroke, bleeding events, and mortality between female and male patients with atrial fibrillation during warfarin therapy. *Eur Heart J Cardiovasc Pharmacother*. 2019;5(1):29–36. doi:10.1093/ehjcvp/pvy026
- 33. Raatikainen MJP, Penttilä T, Korhonen P, Mehtälä J, Lassila R, Lehto M. The quality of warfarin therapy and CHA2DS2-VASc score associate with the incidence of myocardial infarction and cardiovascular outcome in patients with atrial fibrillation: data from the nationwide FinWAF registry. *Eur Heart J Cardiovasc Pharmacother*. 2018;4(4):211–219. doi:10.1093/ehjcvp/pvy009
- 34. Lip GY, Mitchell SA, Liu X, et al. Relative efficacy and safety of non-vitamin K oral anticoagulants for non-valvular atrial fibrillation: network meta-analysis comparing apixaban, dabigatran, rivaroxaban and edoxaban in three patient subgroups. *Int J Cardiol.* 2016;204:88–94. doi:10.1016/j.ijcard.2015.11.084
- 35. Kapiainen S, Väisänen A, Haula T. Terveyden- ja sosiaalihuollon yksikkökustannuksen Suomessa vuonna 2011 [Health and social care unit costs in Finland in 2011]. THL Raportti 3/2014. Juvenes Print, Suomen Yliopistopaino Oy. Tampere. Finnish.
- 36. T12 laboratoriotoimialue hinnasto [Tariffs for the laboratory division]. Turku: Hospital District of Southwest Finland; 2021. Finnish.
- 37. Official public tariff of reimbursable authorized medicinal products in Finland [homepage on the Internet]. Helsinki: Pharmaceuticals Pricing Board; 2021. Available from: https://www.hila.fi/luettelot/korvattavatmyyntiluvalliset-laakevalmisteet/. Accessed February 10, 2021.

- Official Statistics of Finland (OSF). Price index of public expenditure [e-publication]. Helsinki: statistics Finland. Available from: http:// www.stat.fi/til/jmhi/index\_en.html. Accessed November 9, 2020.
- 39. Lassila R. Suorat oraaliset antikoagulantit [Direct oral anticoagulants]. Lääkärin käsikirja; [updated January 8, 2021]. Available from: https://www.terveysportti.fi/apps/ltk/article/ ykt01922/search/suorat%20oraaliset%20antikoagulantit. Accessed April 8, 2021. Finnish.
- 40. Lääkkeiden hintalautakunta. Preparing a health economic evaluation to be attached to the application for reimbursement status and wholesale price for a medicinal product. Application instructions, health economic evaluation; December 17, 2019. Available from: https:// www.hila.fi/content/uploads/2020/01/Instructions\_TTS\_2019.pdf. Accessed March 16, 2021.
- 41. Lehto M, Niiranen J, Korhonen P, et al. Quality of warfarin therapy and risk of stroke, bleeding, and mortality among patients with atrial fibrillation: results from the nationwide FinWAF registry. *Pharmacoepidemiol Drug Saf.* 2017;26(6):657–665. doi:10.1002/ pds.4194
- 42. Hallinen T, Soini E. The impact of the pharmaceutical pricing system on cost-effectiveness results: finnish analysis. Open Pharmacoeconomics Health Econ J. 2011;3:6–10. doi:10.2174/ 1876824501103010006

- Official Statistics of Finland (OSF): Deaths [e-publication]. ISSN=1798-2545. Helsinki: statistics Finland; referred 26 May 2021. Available from: http://www.stat.fi/til/kuol/meta\_en. html. Accessed May 20, 2021.
- 44. Benjamin EJ, Wolf PA, D'Agostino RB, Silbershatz H, Kannel WB, Levy D. Impact of atrial fibrillation on the risk of death: the Framingham Heart Study. *Circulation*. 1998;98(10):946–952. doi:10.1161/01.CIR.98.10.946
- 45. Zhu J, Alexander GC, Nazarian S, Segal JB, Wu AW. Trends and variation in oral anticoagulant choice in patients with atrial fibrillation, 2010–2017. *Pharmacotherapy*. 2018;38(9):907–920. doi:10.1002/phar.2158
- 46. Gurusamy VK, Brobert G, Vora P, Friberg L. Sociodemographic factors and choice of oral anticoagulant in patients with non-valvular atrial fibrillation in Sweden: a population-based cross-sectional study using data from national registers. *BMC Cardiovasc Disord*. 2019;19(1):43. doi:10.1186/s12872-019-1029-z
- 47. Leminen A, Pyykönen M, Tynkkynen J, Tykkyläinen M, Laatikainen T. Modeling patients' time, travel, and monitoring costs in anticoagulation management: societal savings achievable with the shift from warfarin to direct oral anticoagulants. *BMC Health Serv Res.* 2019;19(1):901. doi:10.1186/s12913-019-4711-z

ClinicoEconomics and Outcomes Research

#### **Dove**press

**Dove**Press

Publish your work in this journal

ClinicoEconomics and Outcomes Research is an international, peerreviewed open-access journal focusing on Health Technology Assessment, Pharmacoeconomics and Outcomes Research in the areas of diagnosis, medical devices, and clinical, surgical and pharmacological intervention. The economic impact of health policy and health systems

organization also constitute important areas of coverage. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/clinicoeconomics-and-outcomes-research-journal