Diagnostic Usefulness of MLPA Techniques for Recurrent Copy Number Variants Detection in Global Developmental Delay/Intellectual Disability

Diana Miclea^{1,2}
 Adriana Szucs^{1,2}
 Andreea Mirea^{1,2}
 Delia-Maria Stefan^{1,2}
 Florina Nazarie^{1,2}
 Simona Bucerzan^{2,3}
 Cecilia Lazea^{2,3}
 Alina Grama^{2,3}
 Tudor Lucian Pop^{2,3}
 Marius Farcas⁴
 Gabriela Zaharie^{3,4}
 Melinda Matyas^{3,4}
 Monica Mager Snr^{2,3}
 Mihaela Vintan^{2,3}
 Radu Popp¹
 Camelia Alkhzouz^{2,3}

¹Department of Molecular Sciences, "Iuliu Hatieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania;

²Emergency Clinical Hospital for Children, Cluj-Napoca, Romania;

³Department of Mother and Child, "Iuliu Hatieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania;

⁴County Emergency Clinical Hospital, Cluj-Napoca, Romania

Correspondence: Diana Miclea
 Department of Molecular Sciences, "Iuliu Hatieganu" University of Medicine and Pharmacy, Pasteur Street, No. 6, Cluj-Napoca, 400349, Romania
 Email diana.miclea@umfcluj.ro

Background: Genetic testing has become a standardized practice in the diagnosis of patients with global developmental delay/intellectual disability (GDD/ID). The aim of this study is to observe the frequency of recurrent copy number variations (CNVs) in patients diagnosed with GDD/ID, using MLPA technique.

Methods: A total of 501 paediatric patients with GDD/ID were analysed using SALSA MLPA probemix P245 Microdeletion Syndromes-1A, and the technical steps were performed according to the MRC Holland MLPA general protocol.

Results: Twenty-five of 501 patients (5%) were diagnosed with a microdeletion/microduplication syndrome. Amongst them, 7 of 25 (30%) with clinical suggestion have a confirmed diagnosis, for the other cases the clinical features were not evocative for a specific syndrome.

Conclusion: This study showed that in cases with a specific clinical diagnosis the MLPA technique could be a useful alternative, less expensive and more efficient to indicate as first intention of a targeted diagnostic test, as it is the case of Williams syndrome, Prader-Willi syndrome or DiGeorge syndrome.

Keywords: MLPA, global developmental delay, intellectual disability, diagnostic

Background

Genetic testing has become a standardized practice in the diagnosis of patients with global developmental delay/intellectual disability (GDD/ID). Based on its definition from the Diagnostic and Statistical Manual of Mental Disorders 5 (DSM5) ID is characterized by significant intellectual disability and deficits in conceptual, social and practical adaptative functions, occurring in the growth period.¹ GDD is confirmed if the values from the standardized tests are lower than two standard deviations below average, in a minimum of two areas of the following: global or fine motility, language and speech, cognition, personal or social field, daily activities.² Children under the age of five are diagnosed with GDD, and after this age threshold with ID. GDD/ID is regarded as a pathology that belongs to the neurodevelopmental disorders.³

Genetic causes are responsible for over 50% of GDD/ID cases.⁴⁻⁶ Amongst them, the most frequent are numerical chromosomal anomalies, such as Down Syndrome, which is responsible for up to 10% of cases^{6,7} and structural chromosomal anomalies, frequently microdeletion/microduplication syndromes, known as recurrent copy number variations (CNVs). Anomalies given by microdeletion/microduplication syndromes are small chromosomal alterations, lower than 5 Mb,

which cannot be detected using standard karyotyping due to its low-resolution capacity. The frequency of microdeletions/microduplications was found to be around 20% when using high-resolution genomic techniques, such as chromosomal microarray analysis (CMA).⁵

Karyotyping has been the golden standard in the diagnosis of chromosomal anomalies for a long time. However, in recent years, CMA has become a very useful technique and is used as the first intention diagnostic test when either numerical or structural anomalies are suspected.⁵ At a genomic level, certain CNVs are more frequent than others and these can be identified as recurrent CNVs. These recurrent CNVs can be observed in around 10% of GDD/ID cases⁸ and usually occur through non-allelic homologue recombination (NAHR) between region-specific low copy repeats (LCRs) in meiosis.⁹ These anomalies can be identified using multiplex ligation-dependent probe analysis (MLPA) or quantitative PCR techniques, which imply relatively low costs. It is known that MLPA evaluates fewer target regions than CMA, but these regions are very well chosen, generally according to their frequent involvement in human pathology. Thus, the less frequent regions associated with disorders could be omitted in MLPA but seen in CMA. However, the advantages of using MLPA instead of other high-resolution genomic techniques consist of the omission regarding incidental modifications such as consanguinity, malignancy predisposition, or other changes not related to the disorder.

The aim of this study is to observe the frequency of the recurrent CNVs for patients diagnosed with GDD/ID, using MLPA technique.

Materials and Methods

A total of 501 paediatric patients with GDD/ID were analysed. They were diagnosed at the Emergency Clinical Hospital for Children from Cluj-Napoca, Romania, from 1st October 2017 to 1 April 2019. Each patient was investigated by anamnesis, clinical exam, basic biochemical investigation in GDD/ID (blood count, serum iron and ferritin, alanine transaminase – ALT, aspartate aminotransferase – AST, serum creatinine, blood urea nitrogen – BUN, creatine phosphokinase, uric acid, thyroid stimulating hormone – TSH, free thyroxine – free T4, free triiodothyronine – free T3, blood glucose level), ammonia and lactic acid. Depending on the clinical context, neurological, ophthalmological, ENT consult or others were indicated. Also, for the investigation of any internal malformations, an ultrasound examination was performed when needed.

Ethical Issues

For each patient, an informed consent regarding their participation in the study, was obtained from their parents, as a signed consent form. The study was approved by the Ethics Committee of Emergency Clinical Hospital for Children, Cluj-Napoca. The study was conducted in accordance with the Declaration of Helsinki.

MLPA Genetic Testing

Three millilitres of blood in a vacutainer containing EDTA was collected from each of the patients. DNA extraction was performed using a DNA extraction kit (Wizard Genomic DNA Purification Kit, Promega, Madison, WI, USA). SALSA MLPA probemix P245 Microdeletion Syndromes-1A was used, and the technical steps were performed according to the MRC Holland MLPA general protocol.¹⁰ The probes detect sequences involved in a distinct subset of the most common microdeletion and microduplication disorders.

The specific chromosomal regions evaluated by the MLPA kit are described in Table 1. Statistical analysis was done using IBM SPSS Statistics software (IBM Corp., Armonk, NY, USA).

Results

The average age for diagnosis was 10.6 years (SD = 6.05 years); 160/501 (32%) of the patients were aged 2 years or younger, 144/501 (29%) of the patients were older than 2 years but younger or equal to 10 years, and 197/501 (39%) were older than 10 years. Concerning sex distribution, 223/501 patients (44.5%) were females and 278/501 (55.5%) were males. 315/501 (63%) had associated varying signs of craniofacial dysmorphism. 128/501 (25%) of the patients presented associated malformation of internal organs.

25/501 patients (5%) were diagnosed with a microdeletion/microduplication syndrome. Amongst them, 7/25 (30%) had been clinically diagnosed, for the other cases the clinical features were not evocative for a specific syndrome. The aetiology observed in these cases is described in Table 2.

Considering a clinical approach, 7 out of 50 patients (14%) who had a different clinical diagnosis, possible to be evaluated in our panel, were also genetically confirmed. The other patients, who did not have a specific clinical diagnosis, were referred only with isolated or syndromic GDD/ID. Two of 11 patients (18%) who presented a clinical picture of DiGeorge syndrome were confirmed by genetic testing. For Prader-Willi syndrome, the clinical suggestion was confirmed in 2/15 patients (13%). For Williams syndrome, the clinical picture

Table I Chromosomal Region Assessed by SALSA MLPA P245-BI Microdeletion Syndromes-IA

Genetic Syndrome	Chromosomal Region
1p36 deletion syndrome	1p36
22p16.1-p15 microdeletion syndrome	2p16.1-p15
2q23.1 microdeletion/microduplication syndrome	2q23.1
Glass syndrome	2q32-q33
3q29 microdeletion/microduplication syndrome	3q29
Wolf-Hirschhorn syndrome	4p16.3
Cri-du-Chat syndrome	5p15
Sotos syndrome	5q35.3
Williams-Beuren/duplication syndrome	7q11.23
Langer-Giedion syndrome	8q24.11-q24.13
9q22.3 microdeletion syndrome	9q22.3
DiGeorge syndrome	10p13-p14
Prader-Willi/Angelman syndrome	15q11.21
Witteveen-Kolk/15q24 microdeletion syndrome	15q24
Rubinstein-Taybi syndrome	16p13.3
Miller-Dieker syndrome	17p13.3
Lissencephaly-I	17p13.3
Smith-Magenis syndrome	17p11.2
Potocki-Lupski syndrome	17p11.2
NF1 microdeletion syndrome	17p11.2
Koolen-de Vries syndrome	17q21.31
17q21.31 microduplication syndrome	17q21.31
DiGeorge syndrome/22q11.2 duplication syndrome	22q11.21
Distal 22q11.2 deletion syndrome	22q11.2
Phelan-McDermid syndrome	22q13
Rett MECP2 duplication syndrome	Xq28

Notes: Table derived from product description version BI-08.¹¹

was confirmed in 2/5 patients (40%). *GATA3* deletion was confirmed for the only patient who was clinically suspected, and whose clinical phenotype included hypoparathyroidism,

deafness and renal dysplasia, a specific picture for this abnormality.

Discussions

The main result of this research is the 5% diagnostic rate in the investigation of microdeletion/microduplication syndromes using the MLPA technique, in patients with isolated/syndromic GDD/ID. This percentage was observed performing a “genotype first” approach. 14% of patients with suggestive clinical diagnosis, “phenotype first approach”, were confirmed after genetic testing. Patients with quite specific clinical diagnosis, such as DiGeorge syndrome (3 patients) or Prader-Willi syndrome (1 patient) were detected by MLPA without clinical suggestion, recommending MLPA as a possible first low cost investigation to rule out recurrent CNVs. However, although the MLPA test is useful for recurrent CNVs detection, rare CNVs, non-recurrent could not be seen by this technique, requiring further investigations (CMA or exome/genome sequencing with CNVs analysis), in the case of an undiagnosed patients with GDD/ID after this first test.

Regarding the different types of genetic testing used to establish an aetiological diagnosis in GDD/ID, an extensive review of the literature regarding the evaluation possibilities of the children affected by GDD determined the overall karyotype detection rate to be 3.7%, and the most frequent encountered anomalies were Down syndrome, sex chromosome aneuploidies and unbalanced translocations/deletion syndromes.²

A meta-analysis aiming to highlight the importance of the CMA in learning disability and congenital anomalies determined an overall diagnostic rate of 10% from 19 studies and nearly 14,000 cases.⁷ A study investigating GDD/ID with CMA and conventional karyotyping, reported a 32.2% diagnostic rate for CMA and 18.1% diagnostic rate for karyotyping.¹² This comparison between the two tests regarding developmental delay diagnosis (among others), had already been stated in a prior study, in which the chromosomal microarray detection rate was rather modest – around 9%, but this rate was still twice as good as the karyotype diagnostic rate.¹³ This CMA and karyotype connection in this pathology was also approached by Siggberg et al.¹⁴ in a study that described CMA diagnostic rates of 10% with low-resolution and 15.8% with high-resolution in cases in which the karyotype had been negative.¹⁴

Trying to assess the diagnostic efficiency using karyotyping and MLPA, a study evaluating recurrent microdeletions/

Table 2 CNVs Observed in Studied Patients with GDD/ID

Observed CNVs	Syndrome	Patients with Positive Genetic Testing (n = 501 Patients Tested)	Patients with Clinical Diagnosis Confirmed by Genetic Testing
Del 22q11.21 – del <i>CLD5</i> , <i>SNAP29</i> , <i>GPIBB</i>	DiGeorge syndrome	5	2
Del 7q11.23 – del <i>ELN</i>	Williams syndrome	2	2
Dup 7q11.23 – dup <i>ELN</i>	7q11.23 microduplication syndrome	1	–
Dup 22q13.33 – dup <i>SHANK3</i> , <i>RABL2B</i>	Phelan–McDermid syndrome	1	–
Dup 16p13.3 – dup <i>CREBBP</i>	16p13.3 microduplication syndrome	1	–
Del 15q11.2 – del <i>SNRPN</i> , <i>UBE3A</i>	Prader–Willi syndrome	3	2
Del 10p14 – del <i>GATA3</i>	DiGeorge syndrome 2	1	1
Del 17q11.2 – del <i>NFI</i>	NFI microdeletion syndrome	2	–
Dup 17p11.2 – dup <i>RAI1</i> , <i>DRC3</i> , <i>LLGL1</i>	Potocki–Lupski syndrome	6	–
Dup 9q22.32 – dup <i>PTCH1</i> , <i>FANCC</i>	9q22.3 microduplication	1	–
Del 5q35.3 – del <i>NSD1</i>	Sotos syndrome	1	–
Del 15q24 – del <i>CYP11A1</i> , <i>SEMA7A</i>	Witteveen–Kolk syndrome	1	–

Abbreviations: del, deletion; dup, duplication.

microduplications and karyotype indicated a diagnostic rate of 19%.¹⁵ Another study evaluating by MLPA the microdeletions and subtelomeric regions in children with GDD/ID with normal karyotype, reported a 9% detection rate.¹⁶ Another investigation on intellectual disability assessing only subtelomeric regions revealed a 4.2% prevalence of subtelomeric rearrangements.¹⁷ A cohort of 150 patients tested with MLPA subtelomere kits and MLPA kit for microdeletions, revealed 14% diagnostic rate, 7.3% subtelomeric rearrangements and 6.6% microdeletions, the most frequent, as in our study, being DiGeorge, Prader–Willi, Angelman, Langer–Giedion syndromes and 17q21.31, 15q24 microdeletions.¹⁸ Other MLPA assessment was applied in intellectual disability using the MLPA telomere kit. The diagnostic rate was 6.7%, but it was nearly doubled – 12.4% when a clinical selection was performed pre-test.¹⁹

A 10-year retrospective analysis on 36,325 cases with GDD/ID targeted the diagnostic outcomes from CMA, karyotyping and FISH. While the diagnostic yield of array-based tests was estimated to be a minimum of 19%, the karyotype detection rate was 4.5%, and 3.5% for FISH.²⁰ The most frequent CNVs detected by CMA platforms were associated with microdeletion/microduplication syndromes: 15q11.2-q13.1 – Prader–Willi/Angelman; *GRIA3* gene on X chromosome; 22q13.3 deletion; 17p11.2 deletion.²¹ Similarly, a CMA and MLPA investigation indicated a 15.6% diagnostic rate for CMA and a 2.1% for MLPA.²²

The syndromes which were clinically suggested from the start, were those presenting strongly defined clinical

features, such as Prader–Willi syndrome, Williams syndrome and velo-cardio-facial syndrome. Regarding the other cases presenting non-specific clinical features, a certain aetiological diagnosis was not established.

This study regarding the frequency of the main microdeletion/microduplication syndromes in a group of 501 patients with isolated or syndromic GDD/ID is important and indicates that an attentive clinical assessment could elevate the diagnostic rate. Genomic evaluation of CNVs (using chromosomal microarray or next-generation sequencing) has become more and more useful for a better diagnosis, with a diagnostic efficiency of more than 50%. However, in cases with a specific clinical diagnosis, it could be less expensive and more efficient to indicate as first intention a low cost diagnostic test, as in the case of Williams syndrome, Prader–Willi syndrome or DiGeorge syndrome.

Ethical Issues

For each patient, an informed consent regarding their participation in the study, was obtained from their parents, as a signed consent form. The study was approved by the Ethics Committee of Emergency Clinical Hospital for Children, Cluj-Napoca.

Author Contributions

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed to submit to the current

journal; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Funding

There is no funding to report.

Disclosure

The authors report no conflicts of interest.

References

1. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. American Psychiatric Association; 2013:271–279. Available from: <https://psychiatryonline.org/doi/book/10.1176/appi.books.9780890425596>. Accessed March 1, 2020.
2. Shevell M, Ashwal S, Donley D, et al. Practice parameter: evaluation of the child with global developmental delay: report of the quality standards subcommittee of the American Academy of Neurology and The Practice Committee of the Child Neurology Society. *Neurology*. 2003;60(3):367–380. doi:10.1212/01.wnl.0000031431.81555.16
3. Moeschler JB, Shevell M, Saul RA, et al. Comprehensive evaluation of the child with intellectual disability or global developmental delays. *Pediatrics*. 2014;134(3):e903–18. doi:10.1542/peds.2014-1839
4. Lavillaureix A, Nava C, Mignot C, et al. New generation sequencing for the diagnosis of intellectual disabilities: exome sequencing or large panel? *Eur J Paediatr Neurol*. 2017;21:e155. doi:10.1016/j.ejpn.2017.04.1148
5. Miller DT, Adam MP, Aradhya S, et al. Consensus statement: chromosomal Microarray is a first-tier clinical diagnostic test for individuals with developmental disabilities or congenital anomalies. *Am J Hum Genet*. 2010;86(5):749–764. doi:10.1016/j.ajhg.2010.04.006
6. Karam SM, Riegel M, Segal SL, et al. Genetic causes of intellectual disability in a birth cohort: a population-based study. *Am J Med Genet A*. 2015;167(6):1204–1214. doi:10.1002/ajmg.a.37011
7. Sagoo GS, Butterworth AS, Sanderson S, Shaw-Smith C, Higgins JP, Burton H. Array CGH in patients with learning disability (mental retardation) and congenital anomalies: updated systematic review and meta-analysis of 19 studies and 13,926 subjects. *Genet Med*. 2009;11(3):139–146. doi:10.1097/GIM.0b013e318194ee8f
8. Wolfe K, Strydom A, Morrough D, et al. Chromosomal microarray testing in adults with intellectual disability presenting with comorbid psychiatric disorders. *Eur J Hum Genet*. 2017;25(1):66–72. doi:10.1038/ejhg.2016.107
9. Lee JA, Lupski JR. Genomic rearrangements and gene copy-number alterations as a cause of nervous system disorders. *Neuron*. 2006;52(1):103–121. doi:10.1016/j.neuron.2006.09.027
10. MLPA[®]. General Protocol Instructions For Use MLPA (Multiplex Ligation-dependent Probe Amplification) General Protocol for the detection and quantification of DNA sequences; 2019. Available from: www.mrcholland.com, www.mlpa.com. Accessed March 1, 2020
11. MLPA[®]. MRC-Holland Product Description SALSA[®] MLPA[®] Probemix P245-B1 Microdeletion Syndromes-1A; 2020. Available from: <https://www.mrcholland.com/products/30024/Product%20Description%20P245-B1%20Microdeletion%20Syndromes-1A-v08.pdf>. Accessed August 02, 2021
12. Lee CL, Lee CH, Chuang CK, et al. Array-CGH increased the diagnostic rate of developmental delay or intellectual disability in Taiwan. *Pediatr Neonatol*. 2019;60(4):453–460. doi:10.1016/j.pedneo.2018.11.006
13. Bejjani BA, Shaffer LG. Application of array-based comparative genomic hybridization to clinical diagnostics. *J Mol Diagn*. 2006;8(5):528–533. doi:10.2353/jmoldx.2006.060029
14. Siggberg L, Ala-Mello S, Jaakkola E, et al. Array CGH in molecular diagnosis of mental retardation - a study of 150 finnish patients. *Am J Med Genet A*. 2010;152A(6):1398–1410. doi:10.1002/ajmg.a.33402
15. David M, Dieterich K, Billette de Villemeur A, et al. Prevalence and characteristics of children with mild intellectual disability in a French county. *J Intellect Disabil Res*. 2014;58(7):591–602. doi:10.1111/jir.12057
16. John N, Rajasekhar M, Girisha K, Sharma PSN, Gopinath P. Multiplex ligation-dependant probe amplification study of children with idiopathic mental retardation in South India. *Indian J Hum Genet*. 2013;19(2):165–170. doi:10.4103/0971-6866.116115
17. Medina A, Piñeros L, Arteaga C, et al. Multiplex ligation-dependent probe amplification to subtelomeric rearrangements in idiopathic intellectual disability in Colombia. *Pediatr Neurol*. 2014;50(3):250–254. doi:10.1016/j.pediatrneurol.2013.10.017
18. Pohovski LM, Dumić KK, Odak L, Barisic I. Multiplex ligation-dependent probe amplification workflow for the detection of submicroscopic chromosomal abnormalities in patients with developmental delay/intellectual disability. *Mol Cytogenet*. 2013;6(1):7. doi:10.1186/1755-8166-6-7
19. Koolen DA, Nillesen WM, Versteeg MH, et al. Screening for subtelomeric rearrangements in 210 patients with unexplained mental retardation using multiplex ligation dependent probe amplification (MLPA). *J Med Genet*. 2004;41(12):892–899. doi:10.1136/jmg.2004.023671
20. Hochstenbach R, van Binsbergen E, Engelen J, et al. Array analysis and karyotyping: workflow consequences based on a retrospective study of 36,325 patients with idiopathic developmental delay in the Netherlands. *Eur J Med Genet*. 2009;52(4):161–169. doi:10.1016/j.ejmg.2009.03.015
21. Rodríguez-Revenga L, Vallespín E, Madrigal I, et al. A parallel study of different array-CGH platforms in a set of Spanish patients with developmental delay and intellectual disability. *Gene*. 2013;521(1):82–86. doi:10.1016/j.gene.2013.02.043
22. Shin S, Yu N, Choi JR, Jeong S, Lee KA. Routine chromosomal microarray analysis is necessary in Korean patients with unexplained developmental delay/mental retardation/autism spectrum disorder. *Ann Lab Med*. 2015;35(5):510–518. doi:10.3343/alm.2015.35.5.510

International Journal of General Medicine

Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-general-medicine-journal>

across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress