Hearing Status of Low-Income, Middle-Aged and Elderly Women in Northern China: A Population-Based, Cross-Sectional Study

Yi Xu,^{1,*} Jie Liu,^{2-4,*}
Chao Dong,¹ Sitong Guo,¹
Hui Cao,¹ Jing Zhang,¹
Mingxin Wang,¹
Jinghua Wang,²⁻⁴
Xin Li,⁵ Dong Yang¹

¹Department of Otorhinolaryngology, Tianjin Medical University General Hospital, Tianjin, 300052, People's Republic of China;

²Department of Neurology, Tianjin Medical University General Hospital, Tianjin, 300052, People's Republic of China;

³Laboratory of Epidemiology, Tianjin Neurological Institute, Tianjin, 300052, People's Republic of China; ⁴Tianjin Neurological Institute, Key Laboratory of Post-Neuroinjury Neuro-Repair and Regeneration in Central Nervous System, Ministry of Education and Tianjin City, Tianjin, 300052, People's Republic of China;

⁵Department of Otorhinolaryngology, Beijing Tsinghua Changgung Hospital, School of Clinical Medicine, Tsinghua University, Beijing, 102218, People's Republic of China

*These authors contributed equally to this work

Correspondence: Dong Yang
Department of Otorhinolaryngology,
Tianjin Medical University General
Hospital, 154 Anshan Road, Heping
District, Tianjin, 300052, People's
Republic of China
Tel +86-22-60814958
Fax +86-22-60362400
Email zyyentyd@163.com

Xin Li
Department of Otorhinolaryngology,
Beijing Tsinghua Changgung Hospital,
School of Clinical Medicine, Tsinghua
University, No. 168, Litang Road,
Changping District, Beijing, 102218,
People's Republic of China
Tel +86 13263358503
Fax +86-10-56118969
Email entlixin@sina.com

Objective: Hearing impairment (HI) has become one of the most common causes of disability worldwide. To date, few studies have examined the hearing of women in these frequently rural regions. Thus, we explored the HI prevalence and risk factors among low-income, middle-aged, and elderly women in Tianjin, China.

Methods: Between October and November 2013, female residents aged ≥ 45 years of rural Tianjin, China were recruited into the study. The participants completed questionnaire surveys, physical examinations, and hearing tests. The hearing at frequencies of 500 Hz, 1000 Hz, 2000 Hz, and 4000 Hz was used to analyze the hearing characteristics of specific frequency bands, and HI was defined as the better ear pure tone averages (PTA) > 25 dB HL.

Results: Among the 1416 participants, the prevalence of HI was 46.0%. Among those aged 45–54-years, most (65.3%) demonstrated normal hearing; in other age groups, slight HI accounted for the largest proportions of individuals. Compared with women who did not drink, the odds ratio (OR) of HI among women who consumed alcohol was 4.2 (95% confidence interval [CI]: 1.844–9.574; $P = 0.001$). Compared with pre-menopausal women, the OR of HI among postmenopausal women was 1.8 (95% CI: 1.261–2.667; $P = 0.001$). Further, each 1-year increase in age in women resulted in a 7.1% increase in HI risk ($P < 0.001$).

Conclusion: The burden of HI among women is heavy in rural northern China, especially among those who experienced menopause. Additionally, the results suggest that to further reduce the risk of developing HI, women in rural areas should stop consuming alcohol. The problem of HI among women in rural areas should be taken seriously; moreover, the measures implemented to prevent HI in high-risk women should be strengthened.

Keywords: hearing impairment, women, risk factors, epidemiology, population-based study

Introduction

Globally, hearing impairment (HI) has become the third most common cause of disability in the elderly.^{1,2} As an age-related disease, HI accounts for almost 5% of the total years lived with disability³ and imposes a great burden on healthcare systems due to its long-term social, functional, and psychological complications.⁴ In 2018, there were approximately 466 million individuals with HI worldwide,⁵ with approximately 90% of those living in low- and middle-income countries.⁶ There are differences in hearing between men and women.⁷ In low-income countries or regions, women frequently have a lower social status than men. Therefore, more attention should be paid to hearing function in women.

Previous studies indicated that the risk of HI is higher among postmenopausal women.^{8,9} Estrogen can be produced by auditory neurons,¹⁰ which can directly affect neurotransmission and indirectly affect cochlear blood flow.^{11,12} Regardless of the mechanism, few studies have focused on HI in rural women. Further, relevant population studies from China have not reported on the hearing abilities of rural women nor on the impact of menopause on the hearing of this population.

Thus, this study explored the hearing status of middle-aged and elderly women in rural Tianjin, China and investigated factors affecting HI prevalence among these women.

Methods

Study Population

This population-based study was performed between September and November 2013 in rural Tianjin, China. The participants were recruited from the Tianjin Brain Study, which is a chronic disease surveillance study of low-income individuals, monitoring the chronic disease prevalence of long-term residents in the area since 1991. In brief, the participants were recruited from 18 administrative villages in rural Tianjin, China. Most (95%) residents were low-income farmers with an annual per capita income of <\$100 US dollars (USD) in 1991 and <\$2500 USD in 2018.¹³ All women aged over 45 years who live in Yangjinzhuang Town, Jizhou District, Tianjin were qualified in this study. Those women who worked out of Tianjin, or were disability due to diseases, were excluded in this study.

Information Collection and Risk Factor Definition

The data for this study were obtained and recorded by trained researchers using face-to-face interviews. Demographic information, including participant sex, age, and educational level, were captured using pre-designed questionnaires. The participants were grouped into four age categories: 45–54 years, 55–64 years, 65–74 years, and ≥75 years. Participant educational levels were classified into three groups, based on their years of formal education: illiterate (0 years), 1–6 years, and >6 years.

Individual and family medical histories, including information regarding hypertension, diabetes mellitus (DM), stroke, and coronary heart disease (CHD), were obtained from participant self-reports or existing medical records. Cigarette smoking was defined as smoking >1

cigarette/day for ≥1 year; participants were categorized as non-smokers, current smokers, or previous smokers. Alcohol consumption was defined as drinking one alcoholic beverage/week for ≥1 year; participants were categorized as non-drinkers, past drinkers, or current drinkers. Data regarding eating habits and menopause details were obtained through participant self-reports.

Physical Examination

Systolic blood pressure (SBP), diastolic blood pressure (DBP), height, weight, waist circumference, and hip circumference measurements were performed by local general practitioners. Further, fasting levels of blood glucose (FBG), total cholesterol (TC), triglycerides (TG), high-density lipoprotein cholesterol (HDL-C), and low-density lipoprotein cholesterol (LDL-C) were determined within 2 hours after the blood sample was taken at the Jizhou People's Hospital (Tianjin, China), which has an international standard blood laboratory with reliable test results. Body mass index (BMI) was calculated as the individual's weight (kg) divided by the square of their height (m²); weight classifications were based on BMI (low-weight, <18.5 kg/m²; normal, 18.5–23.9 kg/m²; overweight, 24.0–27.9 kg/m²; and obese, ≥28.0 kg/m²).¹⁴

Hearing Test

Professional otolaryngology medical technicians performed bilateral otoscopy and audiometry evaluations on each participant in a quiet, soundproof room; participants with suspected HI were referred to audiologists for final diagnoses. Air conduction hearing thresholds at 500 Hz, 1000 Hz, 2000 Hz, and 4000 Hz frequencies were obtained using a Denmark Xeta Audiometer (EN60645-1, type 3). When measuring the hearing of one ear, masking was used on the other ear. Hearing at frequencies of 500 Hz, 1000 Hz, 2000 Hz, and 4000 Hz was used to analyze the hearing characteristics of specific frequency bands.

Pure tone averages (PTAs), representing speech frequencies and reported as decibels of hearing level (dB HL), were calculated using the average audiometric thresholds of 500 Hz, 1000 Hz, 2000 Hz, and 4000 Hz for the bilateral ears. HI was defined as the better PTA >25 dB HL; and HI was stratified using the bilateral PTA results as slight (>25 and ≤40 dB HL), moderate (>40 and ≤60 dB HL), severe (>60 and ≤80 dB HL), and severe to profound (>80 dB HL).¹⁵ HI was defined based on the better hearing ear (using PTA).

Statistical Analyses

Continuous variables (age, BMI, SBP, DBP, WHR, FBG, TC, TG, HDL-C, LDL-C, and PTA) are presented as means and standard deviations; between-group comparisons of these values were performed using Student's *t*-tests. Categorical variables of age group; education group; BMI group; smoking history; alcohol consumption history; and previous histories of hypertension, DM, stroke, and CHD are presented as number of patients and prevalence. The between-group comparisons for categorical variables were performed using the chi-squared test. Binary logistic regression was chosen to evaluate the association between HI and its potential risk factors. In the multivariate analysis, we selected variables that had *P* values <0.05 in two-tailed tests in univariate analysis. Finally, the dependent variable was binary (HI or normal hearing), and the independent variables were education group, alcohol consumption history, menopause history, age, and SBP. Moreover, multivariate analysis results are presented as odds ratios (ORs) with 95% confidence intervals (CIs). All analyses were conducted using SPSS for Windows (version 25.0; SPSS, Chicago, IL, USA); *P* < 0.05 was considered statistically significant.

Results

Total of 1881 participants were recruited among 2452 permanent women after excluded those worked outside residents (*n*=200), disability due to diseases (*n*=59). Of these, 1416 women finished the hearing test. As menopausal history was included in the multivariate analysis, 81 (5.7%) women with missing menstrual history data were excluded from the model. Finally, 1335 participants were assessed the prevalence and risk factors of HI.

Patient Demographics

Of the 1416 included women, 652 (46.0%) demonstrated HI, 617 (44.4%) were illiterate, 588 (41.5%) had hypertension, 143 (10.1%) had DM, and 220 (15.6%) had CHD (Table 1).

Factors Associated with HI in the Univariate Analysis

The prevalence of HI increased significantly with increasing age, but decreased with increasing levels of education (both, *P* < 0.001). The HI prevalence was higher among women who consumed alcohol than among those who did not and among menopausal women than among premenopausal women (both, *P* < 0.001). The mean menopausal

age and DBP were significantly lower in participants with HI than in those without HI (both, *P* < 0.05); however, mean SBP was significantly higher in participants with HI (*P* = 0.001) (Table 1).

Hearing Test Results, by Age

With increasing age, the percentage of participants with normal hearing declined; the prevalence of moderate, severe, and extremely severe HI was significantly higher in the older age groups than in the 45–54-years-old group (all, *P* < 0.001). In addition to right ear HI at 500 Hz, the percentage of patients with bilateral HI at 1000 Hz, 2000 Hz, 4000 Hz, and PTA increased with age (all, *P* < 0.001) (Table 2).

Factors Associated with HI in the Multivariate Analysis

After adjusting for education and SBP, older age, alcohol consumption, and menopausal history were independent risk factors for HI (all, *P* < 0.05). The prevalence of HI was 4.2-fold higher among women who consumed alcohol than among women who never did so (95% CI: 1.84–9.556; *P* = 0.001). Compared with premenopausal women, the OR of HI among postmenopausal women was 1.8 (95% CI: 1.261–2.677; *P* = 0.001); each 1-year increase in age resulted in 7.1% decrease in the risk of HI (*P* < 0.001) (Table 3).

Discussion

This large-scale population study explored the hearing status of women aged ≥45 years who resided in a rural area of northern China. The HI prevalence among the study population was 46.0% and increased with increasing age. In addition, the OR for developing HI was 4.2-fold higher for women who consumed alcohol than for those who did not. Postmenopausal women were 1.8 times more likely to develop HI than were premenopausal women.

A previous study reported that in the United States, the prevalence of HI was 28.5% among women ≥55 years old.¹⁶ Similarly, a study conducted in the Netherlands showed that the prevalence of HI was 29% among female participants ≥65 years old.¹⁷ In Finland, a population-based HI study involving 55–64-year-old adults reported that the prevalence of HI was 18.4%.¹⁸ However, the prevalence of HI among ≥65 years old women in Brazil was 39.4%,¹⁹ and the total prevalence of HI was 38.8% in rural India,²⁰ which was much higher than that in

Table I Associated Factors of Hearing Impairment in the Univariate Analysis

Category	Total	Normal Hearing	Hearing Impairment	P
Total:	1416	764 (54.0)	652 (46.0)	
Age group, n (%)				<0.001
45~54 years	123 (8.7)	25 (3.3)	98 (15.0)	
55~64 years	325 (23.0)	139 (18.2)	186 (28.5)	
65~74 years	591 (41.7)	354 (46.3)	237 (36.3)	
≥75 years	377 (26.6)	246 (32.2)	131 (20.1)	
Education*, n (%)				<0.001
0 years	617 (44.4)	293 (39.1)	324 (50.5)	
1~6 years	399 (28.7)	214 (28.6)	185 (28.8)	
> 6 years	275 (19.8)	242 (32.3)	133 (20.7)	
Menopause*, n (%)				<0.001
No	215 (16.1)	134 (62.3)	81 (37.7)	
Yes	1120 (83.9)	589 (52.3)	531 (47.4)	
Smoking status, n (%)				0.372
Never smoking	1356 (95.8)	735 (54.2)	621 (45.8)	
Now or ever smoking	60 (4.2)	29 (48.3)	31 (51.7)	
Alcohol consumption, n (%)				<0.001
Never drinking	1380 (97.4)	755 (54.7)	625 (45.3)	
Now or ever drinking	36 (2.6)	9 (25.0)	27 (75.0)	
Stroke, n (%)				0.999
No	1340 (94.6)	723 (54)	617 (46)	
Yes	76 (5.4)	41 (53.9)	35 (46.1)	
Hypertension, n (%)				0.152
No	828 (58.5)	460 (55.6)	368 (44.4)	
Yes	588 (41.5)	304 (51.7)	284 (48.3)	
High-carbohydrate diet, n (%)				0.081
No	1376 (97.2)	737 (53.6)	639 (46.4)	
Yes	40 (2.8)	27 (67.5)	13 (32.5)	
Diabetes, n (%)				0.205
No	1273 (89.9)	694 (90.8)	579 (88.8)	
Yes	143 (10.1)	70 (9.2)	73 (11.2)	
CHD, n (%)				0.085
No	1196 (84.4)	657 (54.9)	539 (45.1)	
Yes	220 (15.6)	107 (48.6)	113 (51.4)	
BMI, n (%)				0.173
Low-weight	265 (4.6)	142 (53.6)	123 (46.4)	
Normal	561 (39.6)	317 (56.5)	244 (43.5)	
Overweight	42 (3.0)	17 (40.5)	25 (59.5)	
Obesity, n (%)	548 (38.7)	288 (52.6)	260 (47.4)	
Age, means (SD), years	60.83 (9.04)	58.69 (7.60)	63.34 (9.92)	<0.001
BMI, means (SD), Kg/m ²	24.91 (3.67)	24.99 (3.50)	24.82 (3.85)	0.405
WHR, means (SD)	0.90 (0.26)	0.89 (0.06)	0.91 (0.38)	0.191
Menopause age, mmol/L	49.18 (4.11)	49.44 (3.82)	48.89 (4.40)	0.025

(Continued)

Table 1 (Continued).

Category	Total	Normal Hearing	Hearing Impairment	P
FBG, means (SD), mmol/L	5.54 (1.81)	5.5 (1.76)	5.58 (1.86)	0.373
TC, means (SD), mmol/L	4.78 (0.94)	4.75 (0.93)	4.82 (0.95)	0.143
TG, means (SD), mmol/L	1.58 (0.97)	1.59 (0.96)	1.57 (0.97)	0.756
HDL-C, means (SD), mmol/L	1.40 (0.46)	1.39 (0.43)	1.41 (0.49)	0.302
LDL-C, means (SD), mmol/L	2.71 (0.85)	2.69 (0.84)	2.75 (0.86)	0.180
SBP, means (SD), mmHg	151.57 (24.04)	149.57 (23.73)	153.9 (24.22)	0.001
DBP, means (SD), mmHg	90.55 (12.36)	91.21 (12.12)	89.77 (12.59)	0.030

Note: *Presented data with missing case for education (n=25) and menopause (n=81).

Abbreviations: SD, standard deviation; SBP, systolic blood pressure; DBP, diastolic blood pressure; BMI, body mass index; FBG, fasting blood glucose; TC, total cholesterol; TG, triglycerides; HDL-C, high density lipoprotein cholesterol; LDL-C, low density lipoprotein cholesterol.

Table 2 Hearing Test Results and Hearing Grading of Women by Age Groups

Category	45–54 Years	55–64 Years	65–74 Years	≥75 Years	P
Hearing classification, n (%):					<0.001
Normal	246 (65.3)	354 (59.9)	139 (42.8)	25 (20.3)	
Slight	126 (33.4)	210 (35.5)	156 (48.0)	61 (49.6)	
Moderate	5 (1.3)	26 (4.4)	27 (8.3)	35 (28.5)	
Severe	–	1 (0.2)	3 (0.9)	2 (1.6)	
Hearing test (dBHL), means(SD):					
L 500 Hz	28.16 (9.86)	28.67 (9.85)	31.48 (12.15)	35.73 (14.48)	<0.001
L 1000 Hz	27.37 (9.59)	28.64 (10.12)	32.08 (13.24)	36.79 (13.72)	<0.001
L 2000 Hz	26.29 (8.19)	27.64 (9.33)	31.92 (13.08)	37.52 (15.11)	<0.001
L 4000 Hz	26.59 (8.39)	29.85 (23.90)	34.23 (15.39)	44.76 (18.38)	<0.001
R 500 Hz	27.32 (8.64)	27.78 (10.40)	41.58 (19.56)	37.64 (31.92)	0.057
R 1000 Hz	27.27 (8.27)	28.47 (11.66)	31.58 (13.16)	37.44 (13.90)	<0.001
R 2000 Hz	25.74 (7.45)	27.32 (11.36)	31.72 (17.52)	38.66 (14.40)	<0.001
R 4000 Hz	26.21 (8.79)	28.69 (12.00)	33.86 (15.15)	44.02 (17.40)	<0.001
RPTA	26.63 (7.47)	28.06 (10.49)	34.69 (50.36)	39.44 (14.63)	<0.001
LPTA	27.10 (8.35)	28.70 (10.66)	32.43 (12.40)	38.70 (14.07)	<0.001

developed countries. Previous studies have indicated that the level of economic development affects the prevalence of HI.^{16,17} Socioeconomic status has been confirmed to be a main factor predicting HI prevalence.^{4,21–23} From previous studies, it can be found that the prevalence of HI in developed countries is much lower than in developing countries.^{8,17} Moreover, study populations from different regions have different living and eating habits that need to be considered more in depth to determine their effect on hearing. Thus, this may explain why the prevalence of HI among the rural women reported in this study was higher than that reported in more developed areas.

Between 1980 and 2010, as the global population has aged, the prevalence of HI in developed countries has doubled.²⁴ Previous studies have indicated that HI is an age-related disease.^{25–27} Similarly, in this study, the values of the pure-tone hearing tests increased with increasing age, and the risk of developing HI also increased with age. Prior studies have shown that mitochondrial DNA mutations²⁸ and oxidative damage²⁹ are mechanisms contributing to HI in aging people.

Controversy exists about the relationship between alcohol consumption and HI. One previous study reported that drinking increases the risk of HI,³⁰ while others either

Table 3 Adjusted OR (95% CI) of Risk Factors Association with Hearing Impairment

Risk Factors	OR (95% CI)	P
Education groups:		
0 years	1.241 (0.910, 1.692)	0.172
1–6 years	1.184 (0.863, 1.625)	0.295
> 6 years	1.00	–
Alcohol consumption		
No	1.00	–
Yes	4.202 (1.844, 9.574)	0.001
Menopause history		
No	1.00	–
Yes	1.834 (1.261, 2.667)	0.001
Age	1.071 (1.052, 1.090)	< 0.001
SBP	1.002 (0.997, 1.007)	0.504

indicated a U-shaped association between alcohol consumption and HI^{31,32} or did not find a significant association.^{33,34} The present study showed that the prevalence of HI was significantly higher among women who consumed alcohol than among those who never consumed alcohol. The mechanism for alcohol consumption's effect on hearing is unclear, and further research is needed.

The relationship between menopause and HI is also unclear. Although some studies have shown that ovarian hormones, including estrogen, may have a negative effect on the cochlear nerve,³⁵ most studies have reported that estrogen exerts a protective effect on hearing.^{36,37} Previous studies have also demonstrated the expression of estrogen in the inner ear.¹⁰ Moreover, estrogen can be produced by auditory neurons per se and directly affects neurotransmission.¹¹ In addition, estrogen has also been shown to indirectly affect cochlear blood flow, which may play a role in auditory function.¹² In this study, compared with premenopausal women, the prevalence of HI increased among postmenopausal women. The mechanism by which menopause may impact HI remains unclear, and more research is needed to strengthen the conclusions.

The impact of blood pressure on HI is also inconclusive. Some studies have indicated that hypertension is a risk factor for HI,^{38,39} while another study reported no significant relationship between SBP and HI.⁴⁰ Another study showed that the relationship between age-related hearing loss and blood pressure differs according to age. There is no correlation between blood pressure and hearing in women aged 70–75 years, but in women aged over

79 years, there is a high correlation between SBP and low-frequency hearing loss.⁴¹ In the present study, a significant relationship between SBP and HI was not observed. Further, the mechanism by which blood pressure may impact HI remains unclear. High blood pressure may harm hearing due to pressure-induced microvascular atherosclerosis, which affects the auditory blood flow and causes hearing loss.³⁶ Long-term follow-up studies are required to validate this relationship.

This study has some limitations. First, this was a single-center study and a multi-center study is needed to verify the present results. Second, due to the lack of the exact amount of alcohol consumed within a certain period, we used categorical variables to evaluate the impact of alcohol consumption on HI. The relationship between the amount of alcohol consumed and any associated HI needs further exploration. Third, this study did not employ otoscopy and tympanometry test on patients because the conditions of the study were limited. In the follow-up, we will implement more rigorous hearing-related testing. Finally, as this was a cross-sectional study, causal relationships between the risk factors and HI cannot be determined; follow-up studies involving similar populations are required.

Conclusion

This was the first large-scale population study to investigate the hearing status of rural women ≥ 45 years old in northern China. There is a substantial burden of HI among women in rural northern China. Alcohol consumption and older age appear to increase the prevalence of HI. Moreover, postmenopausal women have a significantly increased prevalence of HI. Therefore, targeted health guidance should be formulated regarding the factors influencing HI to reduce the prevalence of HI among women residing in rural areas of northern China; in particular, attention should be given to postmenopausal women.

Abbreviations

HI, hearing impairment; DM, diabetes mellitus; CHD, coronary heart disease; SBP, systolic blood pressure; DBP, diastolic blood pressure; FBG, fasting levels of blood glucose; TC, total cholesterol; TG, triglycerides; HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol; BMI, body mass index; PTA, pure tone average; SD, standard deviation; OR, odds ratio.

Data Sharing Statement

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethics Approval and Informed Consent

The study was approved by the ethics committee of Tianjin Medical University General Hospital; informed consent was obtained from each participant. This study was conducted in accordance with the Declaration of Helsinki.

Acknowledgments

We thank all participants of the Tianjin Brain Study, and local medical care professionals for their valuable contributions.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

There is no funding to report.

Disclosure

The authors declare that they have no competing interests.

References

- World report on disability. Available from: <https://www.who.int/health-topics/disability>. Accessed September 7, 2021.
- Kiely KM, Gopinath B, Mitchell P, Luszcz M, Anstey KJ. Cognitive, health, and sociodemographic predictors of longitudinal decline in hearing acuity among older adults. *J Gerontol A Biol Sci Med Sci*. 2012;67(9):997–1003. doi:10.1093/gerona/gls066
- World Health Organization. Addressing the rising prevalence of hearing loss. Available from: <http://www.who.int/pbd/deafness/estimates/en/>. Accessed September 7, 2021.
- Cruickshanks KJ, Nondahl DM, Tweed TS, et al. Education, occupation, noise exposure history and the 10-yr cumulative incidence of hearing impairment in older adults. *Hear Res*. 2010;264(1–2):3–9. doi:10.1016/j.heares.2009.10.008
- GBD 2017 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018;392(10159):1789–1858. doi:10.1016/S0140-6736(18)32279-7
- Davis AC, Hoffman HJ. Hearing loss: rising prevalence and impact. *Bull World Health Organ*. 2019;97(10):646. doi:10.2471/BLT.19.224683
- Pearson JD, Morrell CH, Gordon-Salant S, et al. Gender differences in a longitudinal study of age-associated hearing loss. *J Acoust Soc Am*. 1995;97(2):1196–1205. doi:10.1121/1.412231
- Caruso S, Maiolino L, Agnello C, Garozzo A, Di Mari L, Serra A. Effects of patch or gel estrogen therapies on auditory brainstem response in surgically postmenopausal women: a prospective, randomized study. *Fertil Steril*. 2003;79(3):556–561. doi:10.1016/S0015-0282(02)04763-5
- Khaliq F, Tandon OP, Goel N. Auditory evoked responses in postmenopausal women on hormone replacement therapy. *Indian J Physiol Pharmacol*. 2003;47(4):393–399.
- Stenberg AE, Wang H, Fish J, Schrott-Fischer A, Sahlin L, Hultcrantz M. Estrogen receptors in the normal adult and developing human inner ear and in Turner's syndrome. *Hear Res*. 2001;157(1–2):87–92. doi:10.1016/S0378-5955(01)00280-5
- Pinaud R, Tremere LA. Control of central auditory processing by a brain-generated oestrogen. *Nat Rev Neurosci*. 2012;13(8):521–527. doi:10.1038/nrn3291
- Laugel GR, Dengerink HA, Wright JW. Ovarian steroid and vasoconstrictor effects on cochlear blood flow. *Hear Res*. 1987;31(3):245–251. doi:10.1016/0378-5955(87)90194-8
- China statistical yearbook 2019; 2019. Available from: <http://www.stats.gov.cn/tjsj/ndsj/>. Accessed September 7, 2021.
- Zhou BF. Effect of body mass index on all-cause mortality and incidence of cardiovascular diseases—report for meta-analysis of prospective studies open optimal cut-off points of body mass index in Chinese adults. *Biomed Environ Sci*. 2002;15(3):245–252.
- World Health Organization. Grades of hearing impairment. Prevention of blindness and deafness; 1997. Available from: https://www.who.int/pbd/deafness/hearing_impairment_grades/en/. Accessed September 7, 2021.
- Fischer ME, Cruickshanks KJ, Pinto A, Klein BE, Klein R, Dalton DS. Hearing impairment and retirement. *J Am Acad Audiol*. 2014;25(2):164–170. doi:10.3766/jaaa.25.2.5
- Homans NC, Metselaar RM, Dingemans JG, et al. Prevalence of age-related hearing loss, including sex differences, in older adults in a large cohort study. *Laryngoscope*. 2017;127(3):725–730. doi:10.1002/lary.26150
- Hannula S, Mäki-Torkko E, Majamaa K, Sorri M. Hearing in a 54- to 66-year-old population in northern Finland. *Int J Audiol*. 2010;49(12):920–927. doi:10.3109/14992027.2010.510146
- Mattos LC, Veras RP. The prevalence of hearing loss in an elderly population in Rio de Janeiro: a cross-sectional study. *Braz J Otorhinolaryngol*. 2007;73(5):654–659. doi:10.1016/S1808-8694(15)30126-9
- Garg S, Kohli C, Mangla V, Chadha S, Singh MM, Dahiya N. An epidemiological study on burden of hearing loss and its associated factors in Delhi, India. *Ann Otol Rhinol Laryngol*. 2018;127(9):614–619. doi:10.1177/0003489418781968
- Luo Y, Gao J, Zheng X. Individual-level and area-level socioeconomic inequalities in hearing impairment among Chinese adults: a population-based study. *J Epidemiol Community Health*. 2020;74(6):541–547. doi:10.1136/jech-2019-213437
- Emmett SD, Francis HW. The socioeconomic impact of hearing loss in U.S. adults. *Otol Neurotol*. 2015;36(3):545–550. doi:10.1097/MAO.0000000000000562
- Ramsey T, Svider PF, Folbe AJ. Health burden and socioeconomic disparities from hearing loss: a global perspective. *Otol Neurotol*. 2018;39(1):12–16. doi:10.1097/MAO.0000000000001630
- Knipper M, Van Dijk P, Nunes I, Rüttiger L, Zimmermann U. Advances in the neurobiology of hearing disorders: recent developments regarding the basis of tinnitus and hyperacusis. *Prog Neurobiol*. 2013;111:17–33. doi:10.1016/j.pneurobio.2013.08.002

25. Stevens KN, Lang IA, Guralnik JM, Melzer D. Epidemiology of balance and dizziness in a national population: findings from the English Longitudinal Study of Ageing. *Age Ageing*. 2008;37(3):300–305. doi:10.1093/ageing/afn019
26. Yamasoba T, Lin FR, Someya S, Kashio A, Sakamoto T, Kondo K. Current concepts in age-related hearing loss: epidemiology and mechanistic pathways. *Hear Res*. 2013;303:30–38. doi:10.1016/j.heares.2013.01.021
27. Park HJ, Yoo MH, Woo SY, Kim SW, Cho YS. Prevalence of hearing loss and associated factors in subjects with normal otoscopy: a national cross-sectional study. *Int J Audiol*. 2017;56(12):951–957. doi:10.1080/14992027.2017.1373866
28. Kujoth GC, Hiona A, Pugh TD, et al. Mitochondrial DNA mutations, oxidative stress, and apoptosis in mammalian aging. *Science*. 2005;309(5733):481–484. doi:10.1126/science.1112125
29. Someya S, Xu J, Kondo K, et al. Age-related hearing loss in C57BL/6J mice is mediated by Bak-dependent mitochondrial apoptosis. *Proc Natl Acad Sci U S A*. 2009;106(46):19432–19437. doi:10.1073/pnas.0908786106
30. Rosenhall U, Sixt E, Sundh V, Svanborg A. Correlations between presbycusis and extrinsic noxious factors. *Audiology*. 1993;32(4):234–243. doi:10.3109/00206099309072939
31. Gopinath B, Flood VM, McMahon CM, Burlutsky G, Smith W, Mitchell P. The effects of smoking and alcohol consumption on age-related hearing loss: the Blue Mountains Hearing Study. *Ear Hear*. 2010;31(2):277–282. doi:10.1097/AUD.0b013e3181c8e902
32. Fransen E, Topsakal V, Hendrickx JJ, et al. Occupational noise, smoking, and a high body mass index are risk factors for age-related hearing impairment and moderate alcohol consumption is protective: a European population-based multicenter study. *J Assoc Res Otolaryngol*. 2008;9(3):264–276. doi:10.1007/s10162-008-0123-1
33. Brant LJ, Gordon-Salant S, Pearson JD, et al. Risk factors related to age-associated hearing loss in the speech frequencies. *J Am Acad Audiol*. 1996;7(3):152–160.
34. Curhan SG, Eavey R, Wang M, Stampfer MJ, Curhan GC. Prospective study of alcohol consumption and self-reported hearing loss in women. *Alcohol*. 2015;49(1):71–77. doi:10.1016/j.alcohol.2014.10.001
35. Willott JF. Effects of sex, gonadal hormones, and augmented acoustic environments on sensorineural hearing loss and the central auditory system: insights from research on C57BL/6J mice. *Hear Res*. 2009;252(1–2):89–99. doi:10.1016/j.heares.2008.12.002
36. Trott S, Cline T, Weihing J, Beshear D, Bush M, Shinn J. Hormones and hearing: central auditory processing in women. *J Am Acad Audiol*. 2019;30(6):493–501. doi:10.3766/jaaa.17123
37. Orendorz-Frączkowska K, Tempore H. Organ of hearing and balance in peri- and postmenopausal women. Effects of hormone replacement therapy on hearing and balance in peri- and post-menopausal women: the current state of knowledge. *Adv Clin Exp Med*. 2020;29(6):751–755. doi:10.17219/acem/121935
38. Gates GA, Cobb JL, D'Agostino RB, Wolf PA. The relation of hearing in the elderly to the presence of cardiovascular disease and cardiovascular risk factors. *Arch Otolaryngol Head Neck Surg*. 1993;119(2):156–161. doi:10.1001/archotol.1993.01880140038006
39. Reed NS, Huddle MG, Betz J, et al. Association of midlife hypertension with late-life hearing loss. *Otolaryngol Head Neck Surg*. 2019;161(6):996–1003. doi:10.1177/0194599819868145
40. Fischer ME, Schubert CR, Nondahl DM, et al. Subclinical atherosclerosis and increased risk of hearing impairment. *Atherosclerosis*. 2015;238(2):344–349. doi:10.1016/j.atherosclerosis.2014.12.031
41. Rosenhall U, Sundh V. Age-related hearing loss and blood pressure. *Noise Health*. 2006;8(31):88–94. doi:10.4103/1463-1741.33539

Journal of Multidisciplinary Healthcare

Dovepress

Publish your work in this journal

The Journal of Multidisciplinary Healthcare is an international, peer-reviewed open-access journal that aims to represent and publish research in healthcare areas delivered by practitioners of different disciplines. This includes studies and reviews conducted by multidisciplinary teams as well as research which evaluates the results or conduct of such teams or healthcare processes in general. The journal

covers a very wide range of areas and welcomes submissions from practitioners at all levels, from all over the world. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-inflammation-research-journal>