

Allied health assistants and what they do: A systematic review of the literature

Lucylynn Lizarondo¹
Saravana Kumar¹
Lisa Hyde²
Dawn Skidmore²

¹International Centre for Allied Health Evidence, School of Health Sciences, University of South Australia, Adelaide; ²South Australian Health Allied and Scientific Health Office, Adelaide, Australia

Objective: Allied health assistants (AHAs) are an emerging group in allied health practice with the potential to improve quality of care and safety of patients. This systematic review summarizes the evidence regarding the roles and responsibilities of AHAs and describes the benefits and barriers to utilizing AHAs in current health care settings.

Methods: A systematic process of literature searching was undertaken. A search strategy which included a range of electronic databases was searched using key terms. Studies which examined the roles and responsibilities of AHAs (across all allied health disciplines) were included in the review. Only publications written in the English language were considered, with no restriction on publication date. Two reviewers independently assessed eligibility of the articles. Data extraction was performed by the same reviewers. A narrative summary of findings was presented.

Results: Of the initial 415 papers, 10 studies were included in the review. The majority of papers reported roles performed by general health care assistants or rehabilitation assistants who work in multiple settings or are not specifically affiliated to a health discipline. All current AHAs duties have elements of direct patient care and indirect support via clerical and administrative or housekeeping tasks. Benefits from the introduction of the AHA role in health care include improved clinical outcomes, increased patient satisfaction, higher-level services, and more “free” time for allied health professionals to concentrate on patients with complex needs. Barriers to the use of AHAs are related to blurred role boundaries, which raises issues associated with professional status and security.

Conclusions: There is consensus in the literature that AHAs make a valuable contribution to allied health care. Whilst there are clear advantages associated with the use of AHAs to support allied health service delivery, ongoing barriers to their effective use persist.

Keywords: allied health assistants, health care assistants, rehabilitation assistants, allied health workforce

Introduction

In recent years, there have been numerous changes in health care service delivery, increasing the pressure and demand placed on primary and secondary health care services, both nationally and internationally. There has also been a widespread move towards the implementation of health care practices underpinned by research evidence and guided by principles of safety, effectiveness, patient-centeredness, timeliness, efficiency, and equity.¹ Along with these changes are emerging issues related to the demographic shift, science and technological advancements, increased patient expectations, and a shortage of health care professionals. The international trend of an ageing population means that health care consumers are increasingly presenting with chronic and complex diseases. Patients' expectations have changed as well, with

Correspondence: Lucylynn Lizarondo
International Centre for Allied Health Evidence, C7-62 City East Campus,
University of South Australia, Adelaide,
SA 5000, Australia
Tel +618 8302 2099
Fax +618 8302 2766
Email lucylynn.lizarondo@unisa.edu.au

patients becoming active participants rather than passive receivers of care. With these changes, health care services are increasingly faced with the need to ensure that there is an adequate number of health professionals and paraprofessionals who can provide the most appropriate and timely services to patients.

There is evidence that highly qualified health care providers are increasingly allocating tasks to other practitioners in order to allow management of patients with more complex conditions and needs.^{3,4} The boundaries between groups of health professionals are shifting, for example, between doctors and physiotherapists in orthopedic clinics. Specially trained physiotherapists work in an extended role by being involved with the assessment and management of referrals from orthopedic surgeons.⁵ This concept of extended scope practice combines “role enhancement” and “role substitution”. Role enhancement is defined as increasing the depth of the job by extending the roles or skills of a particular group of workers. Role substitution, on the other hand, involves expanding the breadth of a job in particular, by working across professional divides or exchanging one type of worker for another.⁶ Extended scope of practice has led to task redistribution resulting in other staff fulfilling roles traditionally performed by other health professionals.^{7,8}

Currently, the health system is challenged to accommodate these changes in the workforce and to provide an adequate number of appropriately skilled health care workers to meet the changing needs of health care consumers and organizations. The development of the allied health assistant (AHA) role is one method to address these challenging issues of health care delivery. In the United Kingdom, for instance, the introduction of “assistant practitioners” to complement the work of professionals is one of the initiatives which emerged out of policies that encouraged modernization of the professions and challenged traditional working practices.^{9–12} Traditional health care professionals are increasingly allocating tasks to allied health assistants or support workers, freeing the highly qualified practitioners to manage clients with more complex issues.^{13–15}

While there is an increasing recognition of the importance of the role of AHAs, to date there has not been a systematic analysis of this workforce. Therefore, the aim of this systematic review is to provide evidence on the roles and responsibilities of AHAs as reported in the literature, and where possible, briefly describe the benefits of and barriers to utilizing AHAs in current health care settings.

Methods

For the purpose of this review, an AHA is defined as a person who assists or provides any type of support to the work of a qualified allied health professional. Studies which reported AHAs or other terminologies such as generic assistants, community rehabilitation assistants, multidisciplinary assistants, therapy assistants/supports, aides, technicians and support workers aligned with defined allied health professional groups, and indigenous support workers, were included in this review. Because the aim of this systematic review was targeted towards allied health assistants particularly, publications which described people working in the paramedical ambulance context, medical assistants, physician assistants, nursing assistants, drug and alcohol support workers, and postnatal and midwifery support workers were excluded.

This systematic review considered only peer reviewed literature in which the primary objective or the main focus was related to the roles or responsibilities of an AHA. Articles which have training, education and supervision as their focus were excluded from this review. Only those publications which were written in the English language were included, and no publication date restrictions were set.

Search strategy

A three-step search strategy was utilized in this systematic review. An initial search using the key words “health care assistant, allied health assistant, and health technician” was done in Medline, followed by examination of the titles and abstracts of relevant hits to identify related terms and synonyms. A second extensive search using all identified search terms was then undertaken in all of the following databases: Cochrane, AMED, Medline, Ageline, Ovid, EMBASE, PEDro, PubMed, CINAHL, and Web of Science. An extensive list of key search terms, grouped into two key concepts, was utilized for literature searching. Concept one represented key words within the category “allied health”. Concept two represented key words within the category “assistant”. These two concepts were combined in the electronic search in order to capture the most number of relevant articles.

Concept One: Physiotherap*; Physical Therap*; Occupational Therap*; Speech Therap* or speech pathology*; Diet* OR nutrition*; Allied Health; Social work*; Podiatr*/chiroprody; Radiograph*/medical radiation/diagnostic imaging/ nuclear medicine; Indigenous worker OR Aboriginal support workers; Audiology; Prosthetic* OR orthotic*; Pharmacy; Psychology; Orthoptic*.

Concept Two: Support work*; Health work*; Assistant Aid*; Technician; Helper; Health care assistant; Auxiliary personnel.

Finally, the reference lists of retrieved papers were scrutinized for additional studies that may not have been indexed in any of the electronic databases.

Selection of studies

The titles and abstracts identified from the above search strategy were independently assessed by the first two authors (LL, SK). Full texts of potentially relevant papers were then retrieved for a more detailed examination. The decision to include or exclude studies based on the criteria set was made independently by the same authors. Differences in opinion regarding adherence to inclusion criteria were resolved by discussion.

The methodologic quality of the included papers was not examined for two reasons. First, this review was aimed at exploring the evidence regarding the roles and responsibilities of AHAs rather than a review of evidence of effectiveness. Second, the diversity of the evidence sources found for this review did not allow the reviewers to identify a critical appraisal tool which would be applicable to a range of study designs.

Data extraction

Data were extracted from each of the included studies, such as author, year of publication, type of study, country of origin, roles and responsibilities, barriers and benefits of introducing AHA in the health workforce. A narrative summary of the synthesized findings is presented.

Results

The electronic search yielded a total of 896 citations from the databases searched. After removal of duplicate records, there were 415 potentially eligible papers for inclusion. All 415 papers were evaluated for inclusion by the two reviewers based on title and abstract. A total of 402 were found to be irrelevant to this review. Reasons for exclusion included not satisfying the inclusion criteria ($n = 365$), focused on education, training and competencies ($n = 31$), and related to supervision and ethical standards ($n = 15$). Thirteen papers were retrieved for full examination, three of which discussed the roles of nursing assistants and were therefore excluded. The reference lists of the 10 included papers were carefully examined for additional literature. No further studies were found eligible, yielding a total of 10 papers for this systematic review. Figure 1 illustrates the process involved in the selection of articles for review.




Figure 1 Publication selection process.

Types of allied health assistants

There was a great deal of variability in the retrieved literature regarding the affiliation of AHAs with the different allied health professions. As indicated in Figure 2, the majority of papers were related to the roles performed by general health care assistants or rehabilitation assistants who worked in multiple settings or were not specifically affiliated to a health discipline.

The majority of reported roles concerned AHAs in outpatient and community care settings. Discipline-specific studies that have analyzed the workforce have identified that the largest percentage of AHAs are employed in the geriatric field, although a variety of clinical settings have also been utilized. These include acute care, pediatrics, acquired brain injury, and orthopedics. There is a range of generic or multidisciplinary support worker roles within the context of acute, intermediate, and community care.




Figure 2 Health discipline affiliation of assistants.

A brief summary of the main findings from individual studies considered in the review is provided in Table 1.

Findings for each type of allied health assistant

General health care assistant/rehabilitation assistant

Conway and Kearin surveyed 21 assistants, and their results have shown consistency in the activities they most frequently performed.¹⁶ These included bulk cleaning, general assistance or housekeeping duties and providing physical assistance to nurses, medical, and physiotherapy staff, primarily for manual handling of heavy or difficult patients. These tasks are devoid of direct therapy or clinical components. In another survey, 98 multidisciplinary therapy assistants practicing in rural and remote areas participated in the study.¹⁷ Therapy assistants provided assistance to allied health professionals, and spent time on different program delivery such as individual therapy, group therapy, administration duties, and health promotion. Similar findings were reported by Pullenayegum et al following a small scale audit to assess the role and effectiveness of rehabilitation assistants.¹⁸ The assistants operated only on weekends to maintain the rehabilitation process over a seven-day period. The role of a rehabilitation assistant was created after a negative change in a patient's condition (mobility, motivation, and attitude towards rehabilitation) was observed by nurses and therapy staff when therapy ceased over the weekend. Their responsibility was to facilitate the carry-over and maintenance of the rehabilitation program.

A qualitative, ethnographic approach was undertaken by Stanmore and Waterman to examine the role of rehabilitation assistants.¹⁹ A total of 55 semistructured interviews of patients, associated professionals, and rehabilitation assistants were conducted to examine this new role. Common elements of the role, irrespective of workplace or organization, included working with patients towards individual rehabilitation goals, supporting and supervising patients in activities of daily living, carrying on therapy as delegated by professionals, promoting independence, promoting patients' rights and identity, monitoring progress, providing feedback to professionals on patient progress and service provision, assisting clinicians in the safe use of equipment for patients/carers, and maintaining records of work undertaken with patients. Variations in the role appeared to be a consequence of the demands of the organizational context and the relationships between rehabilitation assistants and the associated professionals and support staff.

A descriptive evaluation using a case study approach of 13 rehabilitation assistants was carried out by Knight et al.²⁰ The assistants' roles in different rehabilitation teams (general medical, general surgical, hospital and community, orthopedics and rheumatology, outpatients, stroke, and vascular and general rehabilitation) mostly involved facilitating mobility, washing and dressing, and activities of daily living of patients. Whilst there were similarities in the tasks, the role of the rehabilitation assistant differed according to the team focus, structure, and process within which the team operated. There were some assistants who were required to spend time on administrative duties which, in turn, reduced the time spent on clinical duties. An interesting finding from this study was that all of the rehabilitation assistants displayed the ability to operate at a level beyond simply following instructions and seemed to have a good overview of the rehabilitation process.

Support worker roles in intermediate care were reported by Nancarrow et al.²¹ This study presented data from a survey of 33 intermediate care services which employed 794 support workers and 368 professionally qualified staff. The roles constituted working in multidisciplinary settings, meeting rehabilitation needs, providing personal care, and enablement. Some workers were also involved in providing administrative support.

Psychology assistant

Woodruff and Wang examined the roles and tasks of assistant psychologists through a functional job analysis which determined both functions within the organization (organizational analysis) and at the individual level (personal analysis).²² The roles included assisting in the provision of clinical psychology services, maintenance of equipment and resources, liaising with other staff, and audit activities. The authors have reported that while the roles of a psychology assistant were clearly articulated in the job description, in practice the roles varied a great deal due to the differing interpretations and perceptions of supervisors and managers. Another possible reason for the variation was the competency and experience of the psychology assistants themselves.

Occupational therapy assistant

Nancarrow and Mackey described the roles and responsibilities of the occupational therapy support worker.²³ Focus group interviews with four groups of stakeholders, namely assistant practitioners (n = 5), supervisors (n = 5), managers (n = 4), and service users (n = 3) were conducted. The role changed according to the setting in which they worked

Table 1 Summary of findings from individual studies

Author, Date	Type of Study	Country	Role	Barriers	Benefits
General health care assistant/rehabilitation assistant					
Conway and Kearin ¹⁶	Survey	Australia	<ul style="list-style-type: none"> • Cleaning, general assistance, and housekeeping • Assistance to nursing, physiotherapist, and medical staff with heavy and difficult to manage patients 	<ul style="list-style-type: none"> • Lack of clarity regarding role • Being expected to achieve cleaning tasks and provide patient support, assist with aggressive patients, work with other assistants 	
Lin et al ¹⁷	Survey	Australia	<ul style="list-style-type: none"> • Facilitates individual therapy • Facilitates group therapy • Performs administrative duties • Participates in health promotion 		
Pullenayegum et al ¹⁸	Audit	UK	<ul style="list-style-type: none"> • Assists in speech therapy • Communication • Lip seal and facial exercises • Swallowing practice • Specific fluid thickness • Use of communication board • Nonverbal communication methods Assists in physiotherapy		Rehabilitation assistants provide consistent and goal-directed rehabilitation to patients who received treatment from a multidisciplinary team
Stannore and Waterman ¹⁹	Qualitative ethnographic	UK	<ul style="list-style-type: none"> • Stroke positioning • Safe transfers with aids • Passive ROM and massage for UL • Posture correction and alignment • Orthotic use • UL and LL strengthening and ROM exercises • Transfer practice • Standing practice Assists in occupational therapy	<ul style="list-style-type: none"> • Bilateral UL exercise • Handgrip • Dressing • Wash and dress practice • Massage for edema • Equipment usage • Feeding aids <ul style="list-style-type: none"> • Works with patients towards individual rehabilitation goals, as prescribed by clinicians, covering nursing, occupational 	

(Continued)

Table 1 (Continued)

Author, Date	Type of Study	Country	Role	Barriers	Benefits
			therapy, and physiotherapy aspects of their treatment		
			<ul style="list-style-type: none"> • Supports and supervises patients in ADL • Carry on therapy as delegated by professionals • Promotes independence • Promotes patients' rights and identity • Monitors progress • Monitors the activities of other care providers • Feeds back to professionals on patient progress and service provision • Assists clinicians in identification, provision, fitting, and safe use of equipment for patients and carers • Maintains records of work undertaken with patients • Variations in the roles between types of setting (eg, more personal patient care in hospital, liaison with a range of professionals and other agencies, promote patient independence and social recovery, less personal care in the community); may vary depending on clinical speciality within the area of work 		
Knight et al ²⁰	Mixed	UK	<ul style="list-style-type: none"> • Performs a wide range of tasks, eg, mobility, washing, dressing, and ADL • Roles depend on team focus, structure, and process • Some have administrative duties 	<ul style="list-style-type: none"> • Confusion over role demarcation and time management issues • Difficulty with complex patient care packages 	<ul style="list-style-type: none"> • Role was organized to link all the disciplines within the multidisciplinary team together and to integrate community rehabilitation • Role offered patients a more continuous, holistic and patient focused service (pass information on from one therapist to another and helping provide an adaptive health care environment consisting of enhanced continuity and patient focused holistic care) • The role allowed therapists extra freedom to carry out more complex cases
Nancarrow et al ²¹	Survey	UK	<ul style="list-style-type: none"> • Working individually and as teams to carry out formalized, competency-based rehabilitation programs in the patient's 		

<p>own home; to provide a holistic approach to rehabilitation; promote maximum independence for the service user with regard to all aspects of care, lifestyle and independence; encourage service users to adhere to their rehabilitation program</p> <ul style="list-style-type: none"> • Assists with all aspects of patient care • Focus on enablement, ie, provide enablement to promote independence 			<ul style="list-style-type: none"> • Assists in the provision of clinical psychology service and assists in client and carer assessments by using psychometrics, assists in statistical analysis, assists in observations, and involvement in report writing. • Responsible for maintenance of equipment and resources, liaising with other staff and professionals, undertaking literature searches and audit activities, and psychologic treatment under supervision. • Assistants would be required to carry out "... any other duties, appropriate to grade, which may be required by ..." the supervisor. 		<ul style="list-style-type: none"> • OTs become free to undertake other tasks • OT assistant helps with referral of clients to appropriate services • OT assistant can identify with patients better because they use the same language as patient • Patients valued the increased time spent with them by the AHA
<p>Psychology assistant Woodruff and Wang²²</p>	<p>Mixed (survey, literature review and qualitative)</p>	<p>UK</p>			
<p>Occupational therapy assistant Nancarrow and Mackey²³</p>	<p>Qualitative approach</p>	<p>UK</p>	<ul style="list-style-type: none"> • Varies according to the setting in which they work (health or social care setting); their relationship with their supervising OT, and the supervisory arrangement within the service • Discharge home visits were the only tasks that participants identified as a role that OT could do that an assistant practitioner could not deliver. 	<ul style="list-style-type: none"> • Other service providers and users did not understand the role, and supervising OTs were concerned that assistants may be seen as a cheap way of delivery of occupational therapy services. 	
<p>Physical therapy assistant Conti et al²⁴</p>	<p>Narrative</p>	<p>US</p>	<p>Facilitates treatment plan by physiotherapist</p>	<p>Successful patient outcomes: skin breakdown rates decreased, average ventilator days per patient and pneumonia rate declined, complete continuity of therapy, early intervention of each</p>	

(Continued)

Table 1 (Continued)

Author, Date	Type of Study	Country	Role	Barriers	Benefits
Ellis et al ²⁵	Survey	UK	Tasks routinely performed by more than 75% <ul style="list-style-type: none"> • Writing in records • Clerical administration tasks • Domestic tasks • Supervising a patient: walking, practicing transfers, performing exercises • Other patient care tasks • Mobilizations • Obtaining a history • Application of superficial heat/cold modalities 		patient and family, a dramatic increase in mobilization and overall fewer complications

Abbreviations: ADL, activities of daily living; OT, occupational therapist; UL, upper limb; LL, lower limb; ROM, range of movement.

(health or social care setting), their relationship with their supervising occupational therapist and the supervisory arrangement within the service. Discharge home visits were the only task identified that an assistant could not do. This task needs to be performed by an occupational therapist.

Physical therapy assistant

Conti et al described the role of a physical therapy assistant (PTA) who was a new inclusion in their critical care team.²⁴ The PTA facilitated the treatment plan designed by the physical therapist and treated the patients daily. Similar findings were obtained by Ellis et al but in addition to direct patient care, PTAs also engaged in clerical/administration work, record keeping, and domestic tasks.²⁵

Synthesis of roles and responsibilities

Overall, a narrative synthesis of the literature indicates that the roles of AHAs can be divided into two key categories, ie, clinical and nonclinical duties. Table 2 summarizes the different duties assumed by AHAs in each category. Clinical duties encompassed tasks which required direct patient contact, such as administration of clinical services, preparation of patients, patient education, and supervision of patients. It is not surprising to note that many of these duties undertaken by AHAs correspond to duties undertaken by allied health professionals as well. This is to be expected because assistants' roles are

Table 2 Clinical and nonclinical duties and responsibilities of AHAs as reported in the literature

Clinical duties	Nonclinical duties
<ul style="list-style-type: none"> • Assist allied health professional • Physical and social support to patient • Administer clinical services and modalities • Transfer patients • Communication of patient progress/communication with other staff • Assist with mobility and gait • Provision of equipment • Patient education • Provision of health care to patients • Supervise/conduct exercise classes • Prepare patients for treatment • Individual or group therapy • Coordinate and assist in the operation of services • Assist and coordinate health service 	<ul style="list-style-type: none"> • Administration • Stock ordering/requisition • Prepare/maintain environment • Equipment maintenance • Health promotion • Monitor and update health care-specific database • Recording/statistics/database • Housekeeping • Cleaning

inherently linked to the health professionals' roles. Nonclinical duties, on the other hand, do not require direct patient contact and include administrative and clerical duties.

Key terms used in describing AHA duties were assisting, supporting, administrating, monitoring, and maintaining. This was in stark contrast with the key terms used by allied health professional staff such as evaluating, assessing, diagnosing, planning, and implementing. Key terms used to denote AHAs' duties in direct patient care are reflective of their scope of practice. Duties such as diagnosing and planning of treatments are beyond the scope of AHAs and are hence exclusive to allied health professionals.

Benefits of introducing AHAs within the allied health workforce

Positive changes to processes and outcomes of health care were reported as a result of introducing AHAs. Knight et al evaluated the role of rehabilitation assistants and found that the assistant's role served to link all the disciplines within the multidisciplinary team and integrate community rehabilitation.²⁰ The team leaders felt that inclusion of assistants in the team would improve the quality of service by being more patient-focused. Rehabilitation assistants, on the other hand, saw their role mainly as being organized to offer patients a more continuous, holistic and patient-focused service while at the same time allowing health professionals extra time to carry out more complex tasks. Similarly, the results of an evaluation conducted by Pullenayegum et al demonstrated the valuable contribution of rehabilitation assistants in providing consistent and goal-directed rehabilitation to patients who received treatment from a multidisciplinary team.¹⁸ Improved communication and interdisciplinary working between nurses, therapists, and rehabilitation assistants has created a well-coordinated and integrated approach to rehabilitation.

Improved outcomes were also described from patients and allied health professionals' perspectives. Conti et al described the improvements in patient clinical outcomes as a result of the introduction of a physical therapy assistant in the critical care team.²⁴ These included reduction in skin breakdown rates, ventilator days per patients, ventilator pneumonia rate, and overall fewer complications. In a qualitative study by Nancarrow and Mackey, patients expressed satisfaction with the amount of time spent with them by the staff member which was facilitated through the introduction of occupational therapy assistants.²³ The ability to identify better with patients, because of a similarity in background and less use of complicated language, was also identified as

contributing to improved patient satisfaction. From a health professional perspective, Nancarrow and Mackey reported reduced burden on occupational therapists because occupational therapy assistants could manage their own case load, which allowed them to undertake other tasks.²³

Barriers to introduction of AHAs

The literature also points to key barriers associated with AHA roles. In the study by Nancarrow and Mackey, concerns were raised by supervising occupational therapists that assistant practitioners may be seen by service users as a cheap way of delivering occupational therapy services.²³ Whilst the occupational therapists felt that they could benefit from delegating a wide range of tasks to the assistant practitioners, the roles were not clear enough to comfortably "let go" much of their work. Findings from a survey conducted by Conway and Kearin revealed the lack of clarity amongst staff in terms of the scope of the assistant's role in supporting direct patient care.¹⁶ This could lead to an unrealistic expectation for AHAs to provide care for which they have had no training and/or which is beyond the scope of their role. There were other challenges voiced by assistants, such as being expected to achieve both allocated cleaning tasks and provide patient support, being requested to assist with aggressive patients, and working with other assistants. The issue of role confusion was also reported by Knight et al.²⁰ These authors also highlighted other barriers, such as time management and feelings of inadequacy, due to the complex and often multidisciplinary nature of the work.

Discussion

The development of AHA roles has emerged as a response to meet the challenges associated with changes in health care demand and service delivery. The aim of this systematic review was to provide evidence on the roles and responsibilities of AHAs and the benefits and barriers to introducing AHAs in current health care settings. This review found consistent evidence that all current AHA duties have elements of direct patient care and indirect support via clerical and administrative or housekeeping tasks. This is similar to other assistant roles in health care such as nursing and medical assistants.^{26,27} Based on the evidence, the scope of practice for AHAs is limited to duties of assisting, supporting, monitoring, and maintaining rather than evaluating, assessing, diagnosing, and planning. These latter duties are typically expected of allied health professionals. The commonly reported duties that relate to direct patient care fit within this AHA's scope of practice. The balance between direct

patient care duties and indirect support varied considerably and was influenced by several variables, including work setting, the profession, the relationship between assistant and therapist, perceived competency of the AHA, and the organizational hierarchy. Recent initiatives in Australia to establish formal training and qualifications through registered training organizations for AHAs may minimize variability and formalize duties undertaken by AHAs.²⁸

Evidence from the literature highlights health care benefits from introducing AHAs, in terms of both process and service outcomes. These include increased patient satisfaction, increased intensity of clinical care, more free time for allied health professionals to concentrate on complex tasks, and improved clinical outcomes. Some of these benefits have also been reported in the medical literature. Medical assistants were described to optimize patient flow which enables physicians to check more patients and conduct more robust visits.²⁷ There are, however, some barriers to introducing AHAs in health care settings. These include ongoing uncertainty regarding the scope of AHA roles and responsibilities, protectionism of allied health professions, and feelings of inadequacy by AHAs themselves. Such concerns may have implications to the quality of care being delivered which may potentially affect patient safety. This is consistent with the barriers described by Bosley and Dale et al who reported that the boundaries between assistant and nursing roles are unclear and that the development of the health care assistant's role challenges the nurse's professional identity.² A clear demarcation of the roles and responsibilities should therefore address the issue of professional status and security, which can lead to adequate and appropriate utilization of AHA services, and ultimately safe and high quality health care for patients.

Implications for policy and practice

The existing literature highlights many advantages associated with the use of AHA roles to support allied health service delivery. The employment of AHAs provides a strategic approach to dealing with current and projected allied health workforce shortages. However, there continues to be ambiguity in terms of role definition, accountability, and clarity. As the capacity of the allied health workforce requires expansion to meet the future needs of the community, these issues concerning AHAs need to be addressed.

This review also highlights the importance of recognizing local settings and contexts because AHA roles

and responsibilities seem to be driven by local needs and organizational requirements. Therefore, a "one size fits all" approach may not be appropriate across all settings. Key determinants, such as staffing mix of AHAs and allied health professionals, need to consider local contexts.

Implications for research

Currently, there are significant knowledge gaps pertinent to AHAs and further research is required to address these gaps and inform policy and practice. Knowledge gaps include how AHAs are used to supplement, complement, or replace allied health professionals, the optimal mix of assistants to professional staff, impact on outcomes as a result of changing roles in patient care, and how best to ensure AHAs gain appropriate competencies.

Cost effectiveness of alternate workforce models which incorporate AHAs are rarely reported in the literature. It is important that future research consider cost effectiveness as an integral measure of outcome to demonstrate the impact (or lack thereof) of AHAs in the workforce.

Given the variety of roles assumed by AHAs, it is possible that the level of education and training received by AHAs is also variable. Future research should explore the competencies required for this practice and establish an educational program that utilizes a skills escalation framework that will provide career opportunities to AHAs.

It is apparent that much of the research underpinning AHAs is based upon small-scale, quality improvement (case study) approaches rather than large-scale, multicenter research initiatives. While large scale research initiatives are to be encouraged, small-scale case study approaches do provide key learnings for similar allied health settings and can contribute to the growing body of evidence for AHAs. Therefore, a mixture of small-scale case studies and large-scale multicenter studies are equally valuable.

Conclusion

There are different types of allied health assistants described in the literature and most of the evidence relates to the roles performed by rehabilitation assistants who are not exclusively affiliated with a health discipline. There is consistency in the literature regarding the roles assumed by AHAs, and these can be divided into clinical and nonclinical or administrative duties. There is emerging evidence that introduction of AHAs in the allied health workforce can improve the processes and outcomes of health care. However, barriers to their use persist and need to be addressed to maximize the benefits.

Acknowledgments

This review was commissioned by the Allied Health Workforce Reform Project – a collaborative project involving several government agencies and nongovernment organizations in the health and community services sectors in South Australia between February 2008 and November 2009. The project was managed by the South Australian Health and Community Services Skills Board, with the South Australian Health Allied and Scientific Health Office playing a leading role in project activities. The Allied Health Workforce Reform Project was jointly funded through the Department of Education, Employment and Workplace Relations, Targeting Skills Needs in Regions Programme, and the Department of Further Education, Employment, Science and Technology through the South Australia Works Initiative.

Disclosure

The authors report no conflicts of interest in this work.

References

- Institute of Medicine. *Crossing the Quality Chasm: A New Health System for the 21st Century*. Washington, DC: National Academy Press; 2001.
- Bosley S, Dale J. Healthcare assistants in general practice: Practical and conceptual issues of skill-mix change. *Br J Gen Pract*. 2008;58(547):118–124.
- Gardiner J, Wagstaff S. Extended scope physiotherapy. The way towards consultant physiotherapists? *Physiotherapy*. 2001;87(1):2–3.
- Kersten P, McPherson K, Lattimer V, George S, Breton A, Ellis B. Physiotherapy extended scope of practice: Who is doing what and why? *Physiotherapy*. 2007;93(4):235–242.
- Daker-White G, Carr AJ, Harvey I, et al. A randomised controlled trial: Shifting boundaries of doctors and physiotherapists in orthopaedic outpatient departments. *J Epidemiol Community Health*. 1999;53:643–650.
- Sibbald B, Shen J, McBride A. Changing the skill-mix of the healthcare workforce. *J Health Serv Res Policy*. 2004;9(1):28–38.
- Woodford AJ. An investigation of the impact/potential impact of a four-tier profession on the practice of radiography: A literature review. *Radiography*. 2006;12(4):318–326.
- McPherson K, Kersten P, George S, et al. A systematic review of evidence about extended roles for allied health professionals. *J Health Serv Res Policy*. 2006;11(4):240–247.
- Department of Health. Meeting the Challenge: A Strategy for the Allied Health Professions. London: Department of Health; 2000.
- Department of Health. The NHS Plan (No. CM 4818-I). London: Department of Health; 2000.
- Department of Health. HR in the NHS Plan – A Consultation Document. London: Department of Health; 2002.
- Changing Workforce Programme. Developing Support Worker Roles in Rehabilitation in Intermediate Care Services. London: NHS Modernisation Agency; 2003.
- Robinson AJ, DePalma MT, McCall M. Physical therapist assistants' perceptions of the documented roles of the physical therapist assistant. *Phys Ther*. 1995;75(12):1054–1066.
- Cooper RA. Health care workforce for the twenty-first century: The impact of non-physician clinicians. *Annu Rev Med*. 2001;52:51–61.
- Ford P. The role of support workers in the department of diagnostic imaging – service managers' perspectives. *Radiography*. 2004;10:259–267.
- Conway J, Kearin M. The contribution of the patient support assistant to direct patient care: An exploration of nursing and PSA role perceptions. *Contemp Nurse*. 2007;24:175–188.
- Lin I, Goodale B, Vilanueva K, Spitz S. Supporting an emerging workforce: Characteristics of rural and remote therapy assistants in Western Australia. *Aust J Rural Health*. 2007;15:334–339.
- Pullenayegum S, Fielding B, Du Plessis E, Peate I. The value of the role of the rehabilitation assistant. *Br J Nurs*. 2005;14(14):778–784.
- Stanmore E, Waterman H. Crossing professional and organizational boundaries: The implementation of generic rehabilitation assistants within three organizations in the northwest of England. *Disabil Rehabil*. 2007;29(9):751–759.
- Knight K, Lerner S, Waters K. Evaluation of the role of the rehabilitation assistant. *Int J Ther Rehabil*. 2004;11(7):311–317.
- Nancarrow SA, Shuttleworth P, Tongue A, Brown L. Support workers in intermediate care. *Health Soc Care Community*. 2005;13(4):338–344.
- Woodruff G, Wang M. Assistant psychologists and their supervisors: Role or semantic confusion? *Clin Psychol*. 2005;48:33–36.
- Nancarrow S, Mackey H. The introduction and evaluation of an occupational therapy assistant practitioner. *Aust Occup Ther J*. 2005;52:293–301.
- Conti S, LaMartina M, Petre C, Vitthuhn K. Introducing a vital new member to the critical care team. Our physical therapy assistant. *Crit Care Nurse*. 2007;27(4):67–68.
- Ellis B, Connell NAD, Ellis-Hill C. Role, training and job satisfaction of physiotherapy assistants. *Physiotherapy*. 1998;84(12):608–616.
- Thornley C. A question of competence? Re-evaluating the roles of the nursing auxiliary and healthcare assistant in the NHS. *J Clin Nurs*. 2000;9:451–458.
- Tache S, Chapman S. The expanding roles and occupational characteristics of medical assistants. *J Allied Health*. 2006;35:233–237.
- Chief Health Professions Office. Discussion Paper: Allied Health Assistants, Assistants in Allied Health and Health Science Workforce Project. Perth: Department of Health; 2008.

Journal of Multidisciplinary Healthcare

Publish your work in this journal

The Journal of Multidisciplinary Healthcare is an international, peer-reviewed open-access journal that aims to represent and publish research in healthcare areas delivered by practitioners of different disciplines. This includes studies and reviews conducted by multidisciplinary teams as well as research which evaluates the results or conduct of such teams or

Submit your manuscript here: <http://www.dovepress.com/journal-of-multidisciplinary-healthcare-journal>

healthcare processes in general. The journal covers a wide range of areas and welcomes submission from practitioners at all levels, from all over the world. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.