

Development of an UPLC-MS/MS Method for the Quantitative Analysis of Upadacitinib in Beagle Dog Plasma and Pharmacokinetics Study

Meng-Jie Wang¹
Yu-Hang Zhao²
Chen Fan³
Ying-Jie Wang³
Xin-Qi Wang³
Xiang-Jun Qiu³
Rui-Le Shen¹

¹Department of Neurology, The First Affiliated Hospital, and College of Clinical Medicine of Henan University of Science and Technology, Luoyang, Henan, 471003, People's Republic of China; ²School of Medical Imaging, Hangzhou Medical College, Hangzhou, 310051, Zhejiang, People's Republic of China; ³Department of Pharmacy, School of Basic Medical Sciences, Henan University of Science and Technology, Luoyang, 471023, Henan, People's Republic of China

Background: Upadacitinib, a novel selective Janus kinase 1 (JAK1) inhibitor, has been recently approved by the US FDA for the treatment of adult patients with moderately to severely active rheumatoid arthritis (RA). An ultra-performance liquid chromatography tandem mass spectrometry (UPLC-MS/MS) method for the quantitative analysis of upadacitinib in beagle dog plasma was developed and validated.

Methods: Upadacitinib and fedratinib (internal standard, IS) were extracted with ethyl acetate under alkaline condition and then separated and detected. The chromatographic column was Waters Acquity UPLC BEH C18 column (2.1 mm × 50 mm, 1.7 μm), the mobile phase was acetonitrile and 0.1% formic acid in water with gradient elution procedure, and the flow rate was 0.40 mL/min. Under the positive ion mode, upadacitinib and IS were monitored by multiple reaction monitoring (MRM) as the following mass transition pairs: m/z 447.00 → 361.94 for upadacitinib and m/z 529.82 → 141.01 for IS.

Results: In the concentration range of 1–500 ng/mL, upadacitinib had good linearity, and the lower limit of quantification (LLOQ) was 1 ng/mL. The RSD of the intra- and inter-day precision was less than 10.03%, and the RE of accuracy was –3.79% to 2.58%. The extraction recovery of upadacitinib was more than 80%, the matrix effect was around 100%, and upadacitinib was found to be stable.

Conclusion: The novel optimized UPLC-MS/MS assay was an effective tool for the determination of upadacitinib and had been successfully applied to the pharmacokinetic study of upadacitinib in beagle dogs, and this method would also be used to study DDIs.

Keywords: upadacitinib, UPLC-MS/MS, pharmacokinetic, beagle dog

Introduction

Rheumatoid arthritis (RA) is a chronic systemic inflammatory autoimmune disease, which can lead to severe pain, disability and progressive joint destruction. Without adequate treatment, irreversible joint destruction and loss of function can lead to severe disability and impaired quality of life.^{1,2} In recent 30 years, with the development of biological treatment methods such as anti TNF, anti IL-6, CTLA4 Ig and anti CD20, the treatment of RA has undergone revolutionary changes. The effectiveness of these treatments provides patients with a variety of treatment methods.³ Based on this clinical understanding, drugs targeting a variety of pro-inflammatory mechanisms, such as the Janus kinase (JAK) family of intracellular signaling enzymes, are being developed for the treatment of RA.³

Correspondence: Xiang-Jun Qiu
Department of Pharmacy, School of Basic Medical Sciences, Henan University of Science and Technology, Luoyang, 471023, Henan, People's Republic of China
Email lyxiangjun@126.com

Rui-Le Shen
Department of Neurology, The First Affiliated Hospital, and College of Clinical Medicine of Henan University of Science and Technology, Luoyang, 471003, Henan, People's Republic of China
Email yfshenrui@126.com

JAK enzymes are tyrosine kinases that can bind to the cytoplasmic region of cytokine receptors and are activated after the cytokines bind to their cellular receptors. The JAK family (JAK 1, 2, 3 and tyrosine kinase 2 [TYK 2]) are important mediators of a variety of cytokine signaling pathways involved in normal cellular processes, and also important mediators in the pathogenesis of RA and other immune-mediated inflammatory diseases.^{2,4}

Oral JAK inhibitors belong to the class of targeted synthetic disease-modifying antirheumatic drugs (DMARD). At present, they are recommended as a treatment option for RA patients worldwide. Although these patients have received traditional synthetic DMARD (csdmard) treatment, they still have moderate or high disease activity, or they fail to treat with biological drug DMARD (bdmard).^{5,6} It has been hypothesized that selectively inhibiting JAK1 block intercepts pathogenic cytokine signaling in RA while retaining other signaling pathways required for normal physiological functions.⁷

Upadacitinib (Figure 1A), a novel selective JAK1 inhibitor, was recently approved by the US FDA for the treatment of adult patients with moderately to severely active RA.⁸ Upadacitinib 15 mg demonstrated a favorable benefit-risk profile and filled an unmet medical need for patients with RA, providing significantly greater rates of disease outcomes (remission and LDA) compared with established therapies (methotrexate, adalimumab, and abatacept), and inhibits structural joint damage.⁹ Upadacitinib could be a favorable long-term treatment option in patients with PsA who are refractory to biologic therapy.¹⁰

Upadacitinib is mainly metabolized by CYP3A and a small part by CYP2D6 in vitro.¹¹ The AUC and C_{max} of Upadacitinib are significantly affected by strong CYP3A4

inhibitors and strong CYP3A4 inducers. Therefore, in patients receiving long-term treatment with strong CYP3A4 inhibitors, it is recommended to use upadacitinib carefully, and it is not recommended to use upadacitinib with strong CYP3A4 inducer.¹² Therefore, it is necessary to develop a quantitative analysis method of upadacitinib to study the pharmacokinetics and drug-drug interactions (DDI).

UPLC-MS/MS was a rapid, sensitive and accurate method for the determination of drug concentration in biological samples.^{13,14} There have been reports on the human pharmacokinetics of upadacitinib, but the description of the detection of plasma upadacitinib concentration is incomplete.^{11,15,16} At present, there is no literature on the complete methodological validation of the detection of plasma upadacitinib concentration. Therefore, in this experiment, we developed and validated a sensitive and quick UPLC-MS/MS assay for the determination of upadacitinib in beagle dog plasma, and fedratinib was used as the internal standard (IS, Figure 1B). And, the novel developed and validated UPLC-MS/MS method was successfully employed to study the pharmacokinetics of upadacitinib in beagle dogs.

Materials and Methods

Chemicals Materials

The purity of upadacitinib and fedratinib were >98% and were purchased from Beijing sunflower technology development CO., LTD (Beijing, China). Methanol and acetonitrile in this study were HPLC grade and were purchased from Tianjin Kermel Chemical Reagent Co., Ltd (Tianjin, China). Analytically pure formic acid was purchased from Tianjin Kermel Chemical Reagent Co., Ltd (Tianjin, China).




Figure 1 The chemical structure upadacitinib (A) and fedratinib (B).

Instruments

Waters ACQUITY UPLC I-Class instrument equipped with online degasser, automatic sampler, four element pump (Waters, USA). Waters XEVO TQD triple quadrupole mass spectrometer equipped with electrospray ionization (ESI) source (Waters, USA). Other instruments included electronic analytical balance and vortex mixer, ultra-pure water equipment, etc.

Solutions Preparation

Accurately weigh 10 mg of upadacitinib in a volumetric flask, dissolve it with methanol and fix the volume to obtain 1 mg/mL standard stock solution. 1 mg/mL IS stock solution was prepared using the same method. The stock solution of 1 mg/mL upadacitinib was diluted 10 times with methanol to obtain the standard solution of 100 µg/mL, 10 µg/mL and 1 µg/mL for calibration curve and quality control (QC) samples.

Blank beagle dog plasma was used to dilute the standard solution and prepare different concentrations of plasma standard solution. The calibration curves with eight different concentrations of upadacitinib were prepared, and the concentration of calibration curve in plasma was as follows: 1, 2.5, 5, 10, 25, 50, 100, 200 ng/mL. QC samples of low, medium and high concentrations (2.5, 50 and 150 ng/mL) were prepared in the same way. The IS working solution at a concentration of 100 ng/mL was obtained by dilution of its stock solution with methanol.

Plasma Sample Treatment

100 µL of beagle dog plasma to be tested was taken into a 1.5 mL EP tube, 20 µL of internal standard working solution was added, and mixed well. 200 µL of 1 mol/L sodium hydroxide (NaOH) solution was added and mixed well. Then, 1 mL ethyl acetate was added and vortexed for 1 min. Took the upper organic phase into another 1.5 mL EP tube and blew dry with nitrogen flow. The residue was dissolved with 100 µL mobile phase, take 50 µL into the sample of an automatic injector, and 5 µL supernatant was injected into UPLC-MS/MS for detection.

Analytical Conditions

The chromatographic column was Waters Acquity UPLC BEH C18 column (2.1 mm × 50 mm, 1.7 µm), and the mobile phase was 0.1% formic acid in water (A) and acetonitrile (B) with gradient elution procedure, the flow rate was 0.40 mL/min. The gradient elution procedure was

as follows: 0–0.5 min A 90→10%, 0.5–1.0 min A 10%, 1.0–1.1 min A 10→90%, and 1.1–2.0 min A 90%. The column was maintained at 40°C, the temperature of the autosampler tray was set at 4°C. The volume of each injection was 5.0 µL.

Under the positive ion mode, upadacitinib and IS were monitored by multiple reaction monitoring (MRM) as the following mass transition pairs: m/z 380.95 → 255.97 for fedratinib and m/z 525.12 → 98.00 for IS. The collision energy and cone voltage of upadacitinib and IS were 25 V and 30 V, respectively. The desolvation temperature was 1000°C, the capillary voltage was 2.0 kV. Masslynx 4.1 (waters, USA) was used for data acquisition and instrument control.

Method Validation

Methodology validation of UPLC-MS/MS included specificity, standard curve, LLOQ, precision, recovery, matrix effect and stability. According to the principles of Industry Bioanalytical Method Validation proposed by FDA, and the technical guidelines for nonclinical pharmacokinetics of chemical drugs (China Food and Drug Administration, CFDA), the UPLC-MS/MS method was validated.^{17,18}

By comparing the chromatograms of blank beagle dog plasma samples from different sources, beagle dog blank plasma spiked with upadacitinib and IS, and beagle dog plasma sample after administration of upadacitinib, the specificity of UPLC-MS/MS method was evaluated.

The plasma standard solution with the concentration of 1, 2.5, 5, 10, 25, 50, 100, 200 ng/mL of upadacitinib was prepared and detected according to the plasma sample preparation method. The peak areas of upadacitinib and IS were recorded, respectively. The ratio of the peak area of upadacitinib to the IS peak area was taken as the ordinate y , the concentration of corresponding points was taken as the abscissa x , and the standard curve of upadacitinib was drawn with the least square method with a weighted ($1/x^2$). The lowest concentration of the standard curve was the LLOQ.

Plasma quality control samples of low, medium and high concentrations levels (2.5, 50 and 150 ng/mL) were taken and injected according to the item of “plasma sample treatment”. Six samples of each concentration were measured in parallel and detected within one day to calculate the intra-day precision and accuracy. The same method was used for 3 consecutive days to calculate the inter-day precision and accuracy. The precision was

expressed by relative standard deviation (RSD, %) and the accuracy was expressed by relative error (RE, %).

At low, medium and high concentration levels (2.5, 50 and 150 ng/mL), the extraction recovery was determined by comparing the ratio of the peak area of upadacitinib before and after the extraction, respectively. Matrix effect (ME) was also analyzed in 6 replicates by comparing the response of upadacitinib in plasma matrix after extraction with that in neat solution.

At low, medium and high concentration levels (2.5, 50 and 150 ng/mL), the stability of plasma samples was investigated under four different storage conditions: room temperature for 4 h, processed samples at 4°C in auto-sampler tray for 6 h, three freeze-thaw cycles (−20°C to 25°C), −20°C for 4 weeks.

Animal Experiments

Six beagle dogs (weight 7.5 ~ 9.5 kg) were purchased from Hubei yizhicheng Biotechnology Co., Ltd (Shiyan, HUBEI), and the animal license number was SCXK (HUBEI) 2016–0020. The six beagle dogs were raised in the Laboratory Animal Center of Henan University of Science and Technology (Luoyang, China). All the experimental behaviors and operations were approved by the Institutional Ethics Committee of Henan University of Science and Technology (Luoyang, China), and the ethical approval number of the animal experiment was 202012003. The experimental operation was carried out according to the rules for the Care and Use of Laboratory Animals.

All the beagle dogs were free to access the water and had a 12 h fasting before the experiment. Upadacitinib was dissolved in 0.5% carboxymethyl cellulose sodium (CMC-Na) solution and was orally administered to beagle dogs at a dose of 1 mg/kg. At the different time points of 0, 0.33, 0.67, 1, 1.5, 2, 3, 4, 6, 9, 12, 24, and 48 h, approximate 1.0 mL blood samples were collected from the veins of the anterior and posterior limbs and taken into 1.5 mL heparinized EP tubes. The plasmas were separated by centrifugation for 10 minutes at 10,000 rpm, and the plasmas were collected and kept frozen at −20°C.

Plasma Sample Detection

Using batch processing method, the concentration of upadacitinib in beagle dog plasma was detected by the developed UPLC-MS/MS technique in this study. The analytical batch included standard curve and QC samples.

Statistical Analysis

DAS (Drug and Statistics, version 2.0) was used to calculate the important pharmacokinetic parameters of upadacitinib through statistical moment method. The main pharmacokinetic parameters of upadacitinib were as follows: T_{max} , C_{max} , $t_{1/2}$, MRT, CL, Vd, and AUC, where T_{max} and C_{max} were actual measured values. All data were expressed as mean ± standard deviation (SD).

Results and Discussion

Method Validation and Improvement

UPLC-MS/MS has the advantages of high sensitivity, strong specificity, short analysis time and good




Figure 2 The Mass spectra of upadacitinib (A) and fedratinib (IS, B).




Figure 3 Representative chromatograms of upadacitinib and IS in beagle dog plasma. **(A)** A blank beagle dog plasma sample. **(B)** A beagle blank dog plasma sample spiked with upadacitinib 1 ng/mL (LLOQ) and IS. **(C)** A sample obtained from a beagle dog at 2 h after oral administration of 1.0 mg/kg upadacitinib.

reproducibility. Therefore, it is often used in the detection of drug concentration in biological samples, the study of pharmacokinetics and DDI.^{19,20}

Fedratinib and upadacitinib were JAK inhibitors. Fedratinib was a selective JAK2 inhibitor for the treatment of adult patients with myelofibrosis (MF). Under the chromatographic and mass spectrometric conditions of this experiment, the peak shape of fedratinib and upadacitinib was good, the retention time was close and did not interfere with each other. At the same time, mass spectrometry detection had a good response. Therefore, fedratinib was selected as the internal standard of this experiment.

In this study, both ESI positive and negative were explored to discover the most sensitive ionization mode for upadacitinib and IS. It is demonstrated that the positive ion mode exhibits higher mass response than the negative ion mode. In positive full mass scan (as indicated in Figure 2), upadacitinib showed the most abundant protonated parent ion and daughter ion at m/z 380.95 and 255.97, respectively, and IS showed the most abundant parent ion and daughter ion at m/z 525.12 and 98.00, respectively. Therefore, the parent ion to the daughter ion of quantifier transitions were m/z 380.95 \rightarrow 255.97 for upadacitinib, and m/z 525.12 \rightarrow 98.00 for IS, respectively.

Table 1 The Precision and Accuracy of Upadacitinib in Beagle Dog Plasma (n = 6)

Added (ng/mL)	Intra-Day			Inter-Day		
	Found (ng/mL)	RSD%	RE%	Found (ng/mL)	RSD%	RE%
1	0.97 ± 0.10	10.03	-2.83	1.01 ± 0.10	9.68	0.95
2.5	2.55 ± 0.18	6.96	1.84	2.56 ± 0.16	6.26	2.58
50	48.10 ± 2.87	5.96	-3.79	49.34 ± 3.27	6.62	-1.37
150	149.71 ± 5.88	3.93	-0.19	153.26 ± 7.77	5.07	2.17

Table 2 Recovery and Matrix Effect of Upadacitinib in Beagle Dog Plasma (n=6)

Added (ng/mL)	Recovery (%)		Matrix Effect (%)	
	Mean ± SD	RSD (%)	Mean ± SD	RSD (%)
2.5	81.78 ± 3.43	4.20	101.09 ± 3.26	3.23
50	82.69 ± 3.21	3.88	103.42 ± 3.32	3.21
150	84.42 ± 4.05	4.80	98.48 ± 4.77	4.85

Through the mass spectrometry standard sample injection, adjusting the mass spectrometry conditions, the characteristic parent ion and daughter ion to qualitative and quantitative were found out, and the specificity and accuracy were higher.

The mobile phase should meet the requirements of HPLC and LC/MS. Volatile salts should be added to the mobile phase as far as possible, surfactants and other non-volatile buffers should not be used as far as possible, such as phosphoric acid buffer. Phosphate and other non-volatile buffer salts will precipitate in the ion source and plug the capillary. In this experiment, formic acid, water and ammonium formate were first selected to investigate the influence of mobile phase on peak pattern and peak intensity. Finally, 0.1% formic acid and acetonitrile were used as mobile phase, which could increase the ionization degree of the sample, increase the signal intensity and improve the peak pattern.

In the field of pharmaceutical analysis, sample pretreatment and purification was an important step. Improving the pretreatment method could not only protect the

detection instrument from pollution and prolong its service life but also reduce the detection matrix and improve the sensitivity and selectivity. In this study, through the exploration of a series of pretreatment methods, the ethyl acetate liquid-liquid extraction method was selected to pretreat the plasma samples, and treatment with NaOH during extraction process could alkalize plasma and significantly improve the extraction recovery rate of upadacitinib and IS. The influence of matrix interference on the determination was reduced.

Selectivity

Under the above experimental conditions, upadacitinib and IS were well separated completely, and the endogenous substances did not interfere with the detection. The representative chromatograms of a blank beagle dog plasma sample (A), a beagle blank dog plasma sample spiked with upadacitinib and IS (B), and a sample obtained from a beagle dog at 1.5 h after oral administration of 1.0 mg/kg upadacitinib (C) are shown in Figure 3. The retention times of upadacitinib and IS were 0.68 and 0.65 min, respectively. The total running time for each sample was 2.0 min.

Calibration Curve and LLOQ

At the concentration range of 1–200 ng/mL for upadacitinib, the typical regression equations of upadacitinib were $y = 0.0754 * x + 0.1276$ ($r^2 = 0.9994$), which exhibited an excellent linearity. The sensitivity of the method was detected by LLOQ and established as 1.0 ng/mL, and the

Table 3 Stability Results of Upadacitinib in Plasma Under Different Conditions (n=6)

Added (ng/mL)	Room Temperature, 12 h			Autosampler 4 °C, 6 h			Three Freeze-Thaw			-20°C, 4 Weeks		
	Mean ± SD	RSD (%)	RE (%)	Mean ± SD	RSD (%)	RE (%)	Mean ± SD	RSD (%)	RE (%)	Mean ± SD	RSD (%)	RE (%)
2.5	2.46 ± 0.20	8.21	-1.73	2.48 ± 0.22	8.93	-0.73	2.52 ± 0.15	6.15	0.87	2.42 ± 0.14	5.79	-3.40
50	51.14 ± 2.63	5.14	2.28	49.66 ± 4.33	8.73	-0.69	50.11 ± 4.02	8.02	0.21	48.56 ± 1.79	3.68	-2.87
150	148.54 ± 9.50	6.39	-0.98	151.80 ± 6.19	4.08	1.20	153.82 ± 6.43	4.18	2.55	149.63 ± 7.42	4.96	-0.25




Figure 4 Mean plasma concentration–time curves of upadacitinib in beagle dogs after oral administration of upadacitinib at a single dose of 1.0 mg/kg (n=6).

precision was below 10.03%, whereas the accuracy ranged from −2.83% to 0.95% (Table 1).

Precision and Accuracy

The results obtained for the intra- and inter-day precision and accuracy of upadacitinib are shown in Table 1. The precision (% RSD) did not exceed 6.96%. Accuracy (% RE) was in the

range from −3.79% to 2.58% at low, medium and high concentrations and met the requirements of validation.

Recovery and Matrix Effect

At LQC, MQC and HQC three different concentration levels, the mean recovery of upadacitinib was within the range of 81.78% ~ 84.42% (Table 2), and the matrix effect values for upadacitinib were 98.48% ~ 103.42% (Table 2). Matrix effect did not affect the detection of samples.

Stability

All results of the stability are summarized in Table 3, and it is found that upadacitinib in beagle dog plasma is stable under the conditions described above.

Pharmacokinetic Application

The novel established UPLC-MS/MS method was successfully applied to a pharmacokinetic study after a single 1.0 mg/kg upadacitinib oral administration to six beagle dogs. The average plasma concentration–time curves of upadacitinib in beagle dogs are displayed in Figure 4, and the main pharmacokinetic parameters of upadacitinib are summarized in Table 4.

Table 4 The Main Pharmacokinetic Parameters of Upadacitinib in Beagle Dog Plasma After Oral Administration of Upadacitinib at a Single Dose of 1.0 mg/kg (n=6, Mean ± SD)

Parameters	Upadacitinib
T_{max} (h)	1.75 ± 0.27
C_{max} (ng/mL)	89.38 ± 21.31
$t_{1/2}$ (h)	11.55 ± 1.36
$MRT_{0 \rightarrow t}$ (h)	9.27 ± 1.08
$MRT_{0 \rightarrow \infty}$ (h)	11.55 ± 1.34
CLz/F (L/h/kg)	1.76 ± 0.27
Vz/F (L/kg)	29.05 ± 4.32
$AUC_{0 \rightarrow t}$ (ng/mL·h)	558.09 ± 94.32
$AUC_{0 \rightarrow \infty}$ (ng/mL·h)	581.85 ± 98.67

Abbreviations: T_{max} , time of peak concentration; C_{max} , peak concentration; $t_{1/2}$, half-life; $MRT_{(0 \rightarrow t)}$, mean residence time of 0-t time; $MRT_{(0 \rightarrow \infty)}$, mean residence time of 0-infinity time; CLz/F , clearance rate; Vz/F , apparent volume of distribution; $AUC_{(0 \rightarrow t)}$, area under curve of 0-t time; $AUC_{(0 \rightarrow \infty)}$, area under curve of 0-infinity time.

Conclusions

A robust, quick and reliable UPLC-MS/MS assay was fully optimized and firstly developed to detect the plasma concentration of upadacitinib in beagle dogs. The optimized method had the advantage of short analysis time (2.0 min), high recovery, good precision and strong specificity. After validation, the method had been successfully applied to the pharmacokinetic study of upadacitinib in beagle dogs. In future studies, this UPLC-MS/MS method would be applied to the study of DDIs, including the effects of Chinese herbal medicine or western medicine on the pharmacokinetics of upadacitinib.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Disclosure

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- Smolen JS, Aletaha D, Barton A, et al. Rheumatoid arthritis. *Nat Rev Dis Primers*. 2018;4:18001. doi:10.1038/nrdp.2018.1
- McInnes IB, Schett G. Pathogenetic insights from the treatment of rheumatoid arthritis. *Lancet* 2017;389:2328–2337. doi:10.1016/S0140-6736(17)31472-1
- Parmentier JM, Voss J, Graff C, et al. In vitro and in vivo characterization of the JAK1 selectivity of upadacitinib (ABT-494). *BMC Rheumatol*. 2018;2(1):23. doi:10.1186/s41927-018-0031-x
- Alunno A, Padjen I, Fanouriakis A, Boumpas DT. Pathogenic and therapeutic relevance of JAK/STAT signaling in systemic lupus erythematosus: integration of distinct inflammatory pathways and the prospect of their inhibition with an oral agent. *Cells*. 2019;8(8):898. doi:10.3390/cells8080898
- Kameda H, Fujii T, Nakajima A, et al. Japan college of rheumatology guideline for the use of methotrexate in patients with rheumatoid arthritis. *Mod Rheumatol*. 2019;29(1):31–40. doi:10.1080/14397595.2018.1472358
- Smolen JS, Landewe RBM, Bijlsma JWW, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2019 update. *Ann Rheum Dis*. 2020;79(6):685–699. doi:10.1136/annrheumdis-2019-216655
- Norman P. Selective JAK inhibitors in development for rheumatoid arthritis. *Expert Opin Investig Drugs*. 2014;23(8):1067–1077. doi:10.1517/13543784.2014.918604
- AbbVie. AbbVie receives FDA approval of RINVOQ™ (upadacitinib), an oral JAK inhibitor for the treatment of moderate to severe rheumatoid arthritis; 2019. Available from: <https://news.abbvie.com/news/press-releases/abbvie-receives-fda-approval-rinvoq-upadacitinib-an-oral-jak-inhibitor-for-treatment-moderate-to-severe-rheumatoid-arthritis.htm>. Accessed November 28, 2019.
- Conaghan PG, Mysler E, Tanaka Y, et al. Upadacitinib in rheumatoid arthritis: a benefit–risk assessment across a Phase III program. *Drug Saf*. 2021;44(5):515–530. doi:10.1007/s40264-020-01036-w
- Mease PJ, Lertratanakul A, Papp KA, et al. Upadacitinib in patients with psoriatic arthritis and inadequate response to biologics: 56-week data from the randomized controlled Phase 3 SELECT-PsA 2 study. *Rheumatol Ther*. 2021;8(2):903–919. doi:10.1007/s40744-021-00305-z
- Mohamed MF, Camp HS, Jiang P, Padley RJ, Asatryan A, Othman AA. Pharmacokinetics, safety and tolerability of ABT-494, a novel selective JAK 1 inhibitor, in healthy volunteers and subjects with rheumatoid arthritis. *Clin Pharmacokinet*. 2016;55(12):1547–1558. doi:10.1007/s40262-016-0419-y
- Mohamed-Eslam F, Klünder MB, Othman AA. Clinical pharmacokinetics of Upadacitinib: review of data relevant to the rheumatoid arthritis indication. *Clin Pharmacokinet*. 2020;59(5):531–544. doi:10.1007/s40262-019-00855-0
- Xuegu X, Luo S, Yang Q, et al. Development and validation of the quantitative determination of avapritinib in rat plasma by a bioanalytical method of UPLC-MS/MS. *Arab J Chem*. 2021;14(6):103152. doi:10.1016/j.arabjc.2021.103152
- Zhang Y, Zhu M, Xie S, Ye X, Xuegu X. Simultaneous determination of amiodarone, dronedarone, and their principal metabolites in SD rat plasma by UPLC-MS/MS and its application in pharmacokinetics. *Arab J Chem*. 2021;14(8):103300. doi:10.1016/j.arabjc.2021.103300
- Mohamed MF, Zeng J, Marroum PJ, Song IH, Othman AA. Pharmacokinetics of Upadacitinib with the clinical regimens of the extended-release formulation utilized in rheumatoid arthritis Phase 3 trials. *Clin Pharmacol Drug Dev*. 2019;8(2):208–216. doi:10.1002/cpdd.462
- Mohamed MF, Trueman S, Feng T, Anderson J, Marbury TC, Othman AA. Characterization of the effect of renal impairment on Upadacitinib pharmacokinetics. *J Clin Pharmacol*. 2019;59(6):856–862. doi:10.1002/jcph.1375
- US Food and Drug Administration. Guidance for industry: bioanalytical method validation. Rockville, MD, USA: US Department of Health and Human Services, US FDA, Center for Drug Evaluation and Research; 2018. Available from: <https://www.fda.gov/regulatory-information/search-fda-guidance-documents/bioanalyticalmethodvalidation-guidance-industry>. Accessed August 10, 2018.
- The technical guidelines for non clinical pharmacokinetics of drugs. Available from: <http://www.nmpa.gov.cn/gsz05106/15.pdf>. Accessed September 22, 2021
- Hu J, Su X-J, Si H-L, et al. Simultaneous determination of Celecoxib, Dezocine and Dexmedetomidine in Beagle Plasma using UPLC-MS/MS method and the application in pharmacokinetics. *Drug Des Devel Ther*. 2021;15:2529–2541. doi:10.2147/DDDT.S314562
- Zhou C-J, Wang H-J, Zhou C-Y, et al. Establishment and verification of UPLC-MS/MS technique for pharmacokinetic drug–drug interactions of Selinexor with Posaconazole in rats. *Drug Des Devel Ther*. 2021;15:1561–1568. doi:10.2147/DDDT.S303928

Drug Design, Development and Therapy

Dovepress

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which has also

been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/drug-design-development-and-therapy-journal>