



Implementation of PROMIS[®] in an Optometry Clinic

Randall S Porter ¹

Kathleen Holt ²

Rajeev S Ramchandran ³

¹Pediatric Ophthalmology and Ocular Genetics Service, University of Rochester Flaum Eye Institute, Rochester, NY, USA;

²University of Rochester Clinical & Translational Science Institute, Rochester, NY, USA; ³Retina and Vitreous Service, University of Rochester Flaum Eye Institute, Rochester, NY, USA

Purpose: The full utility of general health Patient-Reported Outcomes Measurement Information System[®] (PROMIS) surveys in the eye care setting has not been previously demonstrated. This report demonstrates the feasibility of implementing PROMIS in an eye care clinic.

Patients and Methods: Over 2 months, general health and functioning PROMIS surveys were offered to all patients in an optometric clinic in Rochester, NY. Demographic and clinical variables were recorded along with percent completion and time to completion of the survey.

Results: Across 651 patients, 258 chose to attempt PROMIS. Patients with low visual acuity were less likely to attempt the survey ($p=0.049$), and younger patients were more likely to complete the survey ($p=0.025$); no other patient characteristics were found to differ between those who did and did not participate in, nor complete, PROMIS. A total of 193 patients completed the survey (74.8%) in a mean time of 6.36 minutes (range = [1.43, 51.92] minutes; standard deviation = 5.62 minutes). Time to completion did not vary significantly across any groups.

Conclusion: Our relatively high completion rate among those who attempted PROMIS indicates that PROMIS surveys are feasible to implement in an optometry clinic. While most patients completed the survey in little time, the large range of time to completion may indicate that some patients had difficulty completing the survey. Furthermore, the significant difference in visual acuity between those who participated in the survey and those who did not highlights the need to address the way PROMIS is delivered in order to foster greater inclusion.

Keywords: patient-reported outcome, PRO, ophthalmology, optometry, clinic

Introduction

The National Institutes of Health-funded Patient-Reported Outcomes Measurement Information System[®] (PROMIS) consists of a set of computerized adaptive testing (CAT) questionnaires used to create a standardized set of patient-reported outcomes (PROs) which are freely available for use in clinical practice and research.¹ Based on recent research in psychometrics, item response theory, and qualitative health surveys, PROMIS has become a standardized set of CAT-administered PROs. It is utilized widely across clinical practice and research in the US and internationally as a validated measure of general health and wellbeing in both adult and pediatric patients, reported in more than 400 peer-reviewed publications to date.^{2,3} This body of research validates the importance of such PROs as crucial to understanding a patient's health-related quality of life, which has been demonstrated to require patient, rather than physician, estimation.⁴ PROMIS is available to researchers and clinicians free of charge, and provides quick and reliable evaluations of these quality of health measures.⁵⁻⁸

Correspondence: Randall S Porter
Flaum Eye Institute, Box 659, 601
Elmwood Ave, Rochester, NY, 14642,
USA
Tel +1 585 275-7039
Fax +1 585 275-2352
Email Randall_Porter@urmc.rochester.edu

In many institutions, PROMIS scores are linked to the electronic health record.⁹ This linked interface allows providers to view a variety of PROMIS scores and compare them with reference standards to support decision-making while remaining informed of the patient's perspective. It also allows tracking of patient responses within functional domains over time and across visits to their various health care providers.⁹ Ideally, providers would have access to this resource, and know that these data are applicable to their specialty. Research into the feasibility and usefulness of PROMIS in many specialties has consistently found it to be of benefit when utilized appropriately.^{10,11}

The utility of PROMIS has not been documented in the eye care setting. Prior studies have described several vision-oriented questionnaires, some of which have been shown to reliably correlate vision-specific medical diagnoses with visual function and detriments to vision-related quality of life;^{4,12–27} however, few studies have used non-vision-specific PROs in the eye care setting,^{28–31} and the use of general PROs to identify individuals at greater risk for comorbid medical or psychosocial life disturbances due to visual impairment, or of visual impairment due to other medical or psychosocial factors, remains less certain.

This report details the implementation of computer adapted PROMIS surveys in an optometry clinic, analyzes patient factors which may relate to the feasibility of completing such surveys, and discusses the implications of our results on the equitable administration of PROMIS for patients with vision problems. At the University of Rochester Medical Center (URMC), more than 80% of the patient population has participated in PROMIS due to a University-wide effort to promote patient-centered care through the implementation of a standardized PRO system. Each department was asked to implement PROMIS in its own clinical setting and required to administer at least three domains of PROMIS: physical function, pain interference, and depression. An additional two domains could be chosen at the discretion of the clinical department. Based on the potential benefits to providers' approach to treatment and patient supports, the URMC Department of Ophthalmology chose the additional domains of social isolation and cognition, as these are aspects commonly impacted by increased age and poorer visual acuity, both of which are prevalent in the general optometry clinic. We piloted these five PROMIS domains in the department's optometry clinics as part of the medical center's PRO system implementation.

Methods

Between October 28 and December 30, 2016, PROMIS surveys were distributed in an outpatient optometric clinic of the URMC Flaum Eye Institute in Rochester, NY. These surveys were offered to all patients by office staff members at check-in and were self-administered on electronic tablets following brief instruction by check-in staff. The tablet automatically recorded the time between opening the survey and finishing the survey, which we used directly as time to completion. Staffing and equipment constraints prevented the use of assistive technologies or other means of improving access to the questionnaire for patients with poorer visual acuity in this initial implementation period. In sum, 258 patients (of 651 patients to visit the clinic in total) agreed to complete the PROMIS survey. Some patients also participated in PROMIS in subsequent visits; in these cases ($n = 14$), we selected only the 1st encounter in which the survey was completed (or began, if the survey was never completed) to include in our data. The URMC Office of Human Protection granted Institutional Review Board approval for the administration of and collection of data from the surveys and the electronic medical record for said patients after obtaining informed consent for the use of this data from participating subjects. This report and the PROMIS implementation itself are adherent to the Health Insurance Portability and Accountability Act, the World Medical Association Declaration of Helsinki, and all federal and New York State laws.

Additional data extracted from the patients' electronic medical record specifically include race (white or non-white), ethnicity (Hispanic or non-Hispanic), age (continuous), diagnosis of diabetes mellitus (presence or absence), and visual acuity. Visual acuity was abstracted from physician notes and was categorized into a 3-tier impairment score. If the distance visual acuity in either eye were 20/20 or better, then visual acuity was categorized as high; acuities worse than 20/20 in both eyes but at least 20/30 in either eye were categorized as moderate acuity, and those with vision poorer than 20/30 in both eyes were categorized as low acuity.

Data regarding those given and those who completed the survey in comparison with categorical demographic data and diabetes diagnosis were analyzed using Cochran-Mantel-Haenszel tests. Survey completion time in comparison with categorical demographic variables and diabetes diagnosis were analyzed using pooled *t*-tests. Data regarding those given and those who completed the survey in comparison

with age were also analyzed using a pooled *t*-test. Age in comparison with time to completion was analyzed by linear regression. All statistical tests used a significance level of $\alpha=0.05$. Data were analyzed using SAS[®] version 9.4.³²

Results

Of the 258 patients who attempted the PROMIS survey, 78.8% were White. Due to the small percentage of patients in all other race categories, we chose to combine those categories into a single category, Non-White (21.2%) for the purposes of analysis. Hispanic patients made up only 3.1% of the sample; the Hispanic/Non-Hispanic ethnicity variable was not included in subsequent analyses due to the small sample size of Hispanic participants. Mean age was 58.9 years with a range of [20, 97] years. Most patients in our sample did not have a diagnosis of diabetes mellitus (80.6%).

Patients with low visual acuity were less likely to participate ($\chi^2=3.86$, $p=0.049$) in PROMIS, and the association of poorer visual acuity with not completing the survey once started trended towards significance ($p=0.077$). There was no significant association between age and participation in PROMIS ($t=-0.62$, $DF=645$, $p=0.534$), but older patients were less likely to complete the survey after starting ($t=2.25$, $DF=254$, $p=0.025$). There

were no statistically significant differences in race, ethnicity, sex, nor diagnosis of diabetes between those who did and did not participate in PROMIS, nor between those who did and did not completed PROMIS once started. A summary of these data may be found in Table 1.

Among the 258 patients who chose to participate in PROMIS, 193 (74.8%) completed the entire survey. Mean time to completion was 6.36 minutes, with a range of [1.43, 51.92] minutes and a standard deviation of 5.62 minutes. Of those who completed the survey, time to completion was not significantly different across any groups (race, ethnicity, sex, diagnosis of diabetes mellitus, age, nor visual acuity).

Discussion

Here we report an implementation of general health and functioning PROMIS surveys in an optometry clinic, and examine patient factors which may have led to less equitable participation in PROMIS or difficulties completing the surveys. General health and functioning PROs yield self-reported information on multiple domains, which could be of great importance to eye care providers. While only 39.6% of clinic patients agreed to participate in the PROMIS survey, our completion rate of 74.8%

Table 1 Demographic Table: Characteristics of the Population of All Patients to Visit the Clinic, Patients Who Participated in PROMIS, and Patients Who Completed PROMIS, by Percentage of the Respective Population or Sample

Characteristic	Total Population (n=651)	Participated in PROMIS (n=258)	Completed PROMIS (n=193)
	%	%	%
Younger than 65	61	60	64
65 and older	39	40**	36**
Female	56	61	64
Male	44	39	36
White	83	79	84
Non-White	17	21	16
Hispanic	2	3	2
Non-Hispanic	98	97	98
High Visual Acuity	67	70	73
Moderate Visual Acuity	26	26	23
Low Visual Acuity	7**	4**	4
Dx of Diabetes Mellitus	18	19	20

Note: Statistically significant differences are noted by double asterisks (**).

Abbreviations: PROMIS, Patient-Reported Outcomes Measurement Information System[®]; CAT, computerized adaptive testing; PRO, patient-reported outcome; URM, University of Rochester Medical Center.

among those who agreed to participate indicates that PROMIS is feasible to implement in the optometry clinic.

Previous patient outcome studies in ophthalmology or optometry clinics report completion times between 15 and 30 minutes,^{16,19,21,23} compared to an average of just over 6 minutes for our PROMIS surveys. Our low time-to-completion and high completion rate are especially encouraging as PROMIS is not a paid, proprietary resource, but rather a freely available resource, open to researchers and clinicians alike,¹ with a significant history of research to date validating its use in other settings.^{2,3,5–8} However, our data does indicate a large variability in time-to-completion, suggesting that, for at least some of our subjects, and particularly older individuals, survey completion may have been difficult. The difference in completion based on age may suggest that CAT questionnaires, or perhaps the methodology of administering the surveys via electronic tablets, are of lesser utility in older patients. The use of different means of survey administration that accounts for challenges older adults face when completing surveys may allow for greater survey completion rates, and should be addressed in future research.

Patients with better visual acuity also appeared more likely to complete the PROMIS survey. While this trend was not found to be significant in this report ($p=0.077$), it is noteworthy that only 4% of those who completed the survey were categorized as poor acuity, as those with poor vision were less likely to participate in the survey. While staff selection biases are one possible explanation, this would only be likely in the case of patients with profound vision loss requiring assistive devices or help from others to perform visual tasks, as check-in staff have no way of becoming aware of a patient's visual acuity beyond that which is obvious to the general public. Further, staff were instructed to offer PROMIS to all patients at check-in, and it seems unlikely that if they forgot to do so, it would be more prevalent specifically in patients of poor visual acuity. This could be clarified by allowing office staff a means to document declined participation; however, patients who declined participation did not give consent for any information to be included in the present report, and thus it is not possible to report such information here. The more likely explanation of the correlation between visual acuity and choosing to participate in PROMIS is patient self-selection. Delivering PROMIS in a format that allows for those with poorer visual acuity to feel more comfortable attempting PROMIS, such as by utilizing assistive technologies or having a staff member offer to

read the questions and enter the patient's verbal responses, may have increased the number of those with lower visual acuity who participated. Such aids may also have increased the completion rate among older adults attempting the PROMIS survey. As such, we recommend future studies to include such aids so as to avoid any inherent bias in the distribution of PROMIS, or in the clinical benefits its use may provide, due to the means of its distribution. This is perhaps especially important in optometry and ophthalmology clinics, where the prevalence of poor vision in patients is expected to be higher than in most clinical settings.

As little information was available for those patients declining to participate, it is difficult to speculate on the reasons for non-participation. Many studies do not report their overall completion rate, making direct comparison difficult.^{16,19} It is also difficult to put this in context as this report is one of the few from our institution that reports the number of patients declining to participate. Our rates of attempting and completing PROMIS once attempted are well within the ranges reported at the University of Rochester and elsewhere.^{33–41} It is important to qualify this, however, as PROMIS is a customizable survey instrument, and we could find no other study reporting the exact PROMIS survey used here.

Finally, PROMIS was only available in English for our implementation, which likely limited non-English speakers' participation. Future research incorporating the use of PROMIS in other languages would help to diversify the population participating in PROMIS and further generalize results to more diverse populations.

Conclusion

Our high completion rate and short time-to-completion demonstrate the feasibility of implementing such general PROMIS surveys in the eye care setting; however, the lower participation rate in patients of poorer visual acuity indicates a need for the use of alternate means of survey administration such as verbal aids or assistive devices in order not to neglect this vulnerable population. As our institution has captured both clinical and PROMIS data from this initial implementation, our next study aims to compare PROMIS results and clinical data relevant to eye health to determine if general wellness and overall functional ability is associated with vision and eye health status – which, if present, may highlight the further utility of using PROMIS in eye care settings.

Data Sharing Statement

The datasets generated and/or analyzed during the period reported on in this manuscript are not publicly available in order to maintain patient privacy. However, de-identified data are available from the corresponding author upon reasonable request.

Ethics Approval

Approval was obtained from the University of Rochester Medical Center Ethics Committee.

Consent for Publication

Informed consent that medical, demographic, and survey data may be used in research studies was obtained from each individual who elected to participate in PROMIS surveys. No individual personal information is reported in this manuscript.

Acknowledgments

We would like to acknowledge our office staff, who took on the work of administering the PROMIS surveys in addition to their preexisting clinical duties. We would also like to thank the National Institutes of Health (NIH) Clinical and Translational Science Institute and Research to Prevent Blindness (RPB) for funding this project.

Author Contributions

All authors made significant a contribution to the work reported, whether that was in the conception, design, execution, acquisition of data, analysis and interpretation, or in all these areas; have drafted, wrote, substantially revised, or critically reviewed the article; have agreed on the journal to which the article will be submitted; reviewed and agreed on all versions of the article before submission, during revision, the final version accepted for publication, and any significant changes introduced at the proofing stage; and agree to take responsibility and be accountable for the contents of the article, in accordance with the International Committee of Medical Journal Editors authorship guidelines.

Funding

The project described in this publication was supported by the University of Rochester Clinical and Translational Science Awards (CTSA), award number UL1 TR002001, from the National Center for Advancing Translational Sciences of the National Institutes of Health; as well as

by a University of Rochester CTSA award from Research to Prevent Blindness (RPB). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH or RPB.

Disclosure

The authors report no financial or non-financial competing interests in this work.

References

1. HealthMeasures.net. PROMIS instrument development and validation scientific standards version 2.0; 2013:1–72.
2. NIH. Patient-Reported Outcomes Measurement Information System (PROMIS); 2019.
3. Cella D, Yount S, Rothrock N, et al. The Patient-Reported Outcomes Measurement Information System (PROMIS): progress of an NIH roadmap cooperative group during its first two years. *Med Care*. 2007;45(5 Suppl 1):S3–S11. doi:10.1097/01.mlr.0000258615.42478.55
4. Paz SH, Slotkin J, McKean-Cowdin R, et al. Development of a vision-targeted health-related quality of life item measure. *Qual Life Res*. 2013;22(9):2477–2487. doi:10.1007/s11136-013-0365-1
5. Cella D, Gershon R, Bass M, Rothrock N. Assessment center SM; 2013.
6. Northwestern U. HealthMeasures; 2020.
7. Ader D. Developing the Patient-Reported Outcomes Measurement Information System (PROMIS). *Med Care*. 2007;45(5):S1–S2. doi:10.1097/01.mlr.0000260537.45076.74
8. Gershon RC, Cella D, Fox NA, Havlik RJ, Hendrie HC, Wagster MV. Assessment of neurological and behavioural function: the NIH toolbox. *Lancet Neurol*. 2010;9(2):138–139. doi:10.1016/s1474-4422(09)70335-7
9. Baumhauer JF. Patient-reported outcomes - Are they living up to their potential? *N Engl J Med*. 2017;377(1):6–9. doi:10.1056/NEJMp1702978
10. Jacobson RP, Kang D, Houck J. Can Patient-Reported Outcomes Measurement Information System® (PROMIS) measures accurately enhance understanding of acceptable symptoms and functioning in primary care? *J Patient Rep Outcomes*. 2020;4(1):39. doi:10.1186/s41687-020-00206-9
11. van der Meij E, Anema JR, Huirne JAF, Terwee CB. Using PROMIS for measuring recovery after abdominal surgery: a pilot study. *BMC Health Serv Res*. 2018;18(1):128. doi:10.1186/s12913-018-2929-9
12. Varma R, McKean-Cowdin R, Vitale S, Slotkin J, Hays RD. Vision assessment using the NIH toolbox. *Neurology*. 2013;80(11 Suppl 3):S37–S40. doi:10.1212/WNL.0b013e3182876e0a
13. Weih LM, Hassell JB, Keeffe J. Assessment of the impact of vision impairment. *Invest Ophthalmol Vis Sci*. 2002;43(4):927–935.
14. Hays RSK. National eye institute refractive error quality of life instrument (NEI-RQL-42TM), Version 1.0: a manual for use and scoring. National Eye Institute; 2002.
15. Vitale S, Schein OD, Meinert CL, Steinberg EP. The refractive status and vision profile: a questionnaire to measure vision-related quality of life in persons with refractive error. *Ophthalmology*. 2000;107(8):1529–1539. doi:10.1016/s0161-6420(00)00171-8
16. Rajagopalan K, Abetz L, Mertzani P, et al. Comparing the discriminative validity of two generic and one disease-specific health-related quality of life measures in a sample of patients with dry eye. *Value Health*. 2005;8(2):168–174. doi:10.1111/j.1524-4733.2005.03074.x
17. Steinberg EP, Tielsch JM, Schein OD, et al. The VF-14. An index of functional impairment in patients with cataract. *Arch Ophthalmol*. 1994;112(5):630–638. doi:10.1001/archophth.1994.01090170074026

18. Wolffsohn JS, Cochrane AL. Design of the low vision quality-of-life questionnaire (LVQOL) and measuring the outcome of low-vision rehabilitation. *Am J Ophthalmol.* 2000;130(6):793–802. doi:10.1016/s0002-9394(00)00610-3
19. Mangione CM, Lee PP, Gutierrez PR, Spritzer K, Berry S, Hays RD. Development of the 25-item National Eye Institute Visual Function Questionnaire. *Arch Ophthalmol.* 2001;119(7):1050–1058. doi:10.1001/archophth.119.7.1050
20. Sloane M, Ball K, Owsley C, Bruni JR, Roenker DL. The visual activities questionnaire: developing an instrument for assessing problems in everyday visual tasks. *Tech Dig Noninvas Assess Vis Sys.* 1992;1:26–29.
21. Mangione CM, Lee PP, Pitts J, Gutierrez P, Berry S, Hays RD. Psychometric properties of the National Eye Institute Visual Function Questionnaire (NEI-VFQ). NEI-VFQ field test investigators. *Arch Ophthalmol.* 1998;116(11):1496–1504. doi:10.1001/archophth.116.11.1496
22. Mangione CM, Phillips RS, Seddon JM, et al. Development of the “Activities of Daily Vision Scale”. A measure of visual functional status. *Med Care.* 1992;30(12):1111–1126. doi:10.1097/00005650-199212000-00004
23. Terwee CB, Gerding MN, Dekker FW, Prummel MF, Wiersinga WM. Development of a disease specific quality of life questionnaire for patients with Graves’ ophthalmopathy: the GO-QOL. *Br J Ophthalmol.* 1998;82(7):773–779. doi:10.1136/bjo.82.7.773
24. Carta A, Braccio L, Belpoliti M, et al. Self-assessment of the quality of vision: association of questionnaire score with objective clinical tests. *Curr Eye Res.* 1998;17(5):506–511. doi:10.1076/ceyr.17.5.50.6.5191
25. Owsley C, McGwin G Jr., Scilley K, Kallies K. Development of a questionnaire to assess vision problems under low luminance in age-related maculopathy. *Invest Ophthalmol Vis Sci.* 2006;47(2):528–535. doi:10.1167/iovs.05-1222
26. Ehrlich JR, Frank C, Smiley J, et al. Development of a rapid point-of-care patient reported outcome measure for cataract surgery in India. *Health Qual Life Outcomes.* 2018;16(1):25. doi:10.1186/s12955-018-0855-5
27. Ehrlich JR, Spaeth GL, Carozzi NE, Lee PP. Patient-centered outcome measures to assess functioning in randomized controlled trials of low-vision rehabilitation: a review. *Patient.* 2017;10(1):39–49. doi:10.1007/s40271-016-0189-5
28. Ehrlich JR, Hassan SE, Stagg BC. Prevalence of falls and fall-related outcomes in older adults with self-reported vision impairment. *J Am Geriatr Soc.* 2019;67(2):239–245. doi:10.1111/jgs.15628
29. Xiang X, Freedman VA, Shah K, Hu RX, Stagg BC, Ehrlich JR. Self-reported vision impairment and subjective well-being in older adults: a longitudinal mediation analysis. *J Gerontol A Biol Sci Med Sci.* 2020;75(3):589–595. doi:10.1093/gerona/glz148
30. Frank CR, Xiang X, Stagg BC, Ehrlich JR. Longitudinal associations of self-reported vision impairment with symptoms of anxiety and depression among older adults in the United States. *JAMA Ophthalmol.* 2019;137(7):793–800. doi:10.1001/jamaophth.137.7.793
31. Saydah S, Gerzoff RB, Taylor CA, Ehrlich JR, Saaddine J. Vision impairment and subjective cognitive decline-related functional limitations - United States, 2015–2017. *MMWR Morb Mortal Wkly Rep.* 2019;68(20):453–457. doi:10.15585/mmwr.mm6820a2
32. Ho B, Houck JR, Flemister AS, et al. Preoperative PROMIS scores predict postoperative success in foot and ankle patients. *Foot Ankle Int.* 2016;37(9):911–918. doi:10.1177/1071100716665113
33. SAS. SAS 9.4; 2020.
34. Anderson MR, Baumhauer JF, Digiovanni BF, et al. Determining success or failure after foot and ankle surgery using Patient Acceptable Symptom State (PASS) and Patient Reported Outcome Information System (PROMIS). *Foot Ankle Int.* 2018;39(8):894–902. doi:10.1177/1071100718769666
35. Chen RE, Papuga MO, Voloshin I, et al. Preoperative PROMIS scores predict postoperative outcomes after primary ACL reconstruction. *Orthop J Sports Med.* 2018;6(5):232596711877128. doi:10.1177/2325967118771286
36. Quinzi DA, Childs S, Kuhns B, Balkissoon R, Drinkwater C, Ginnetti J. The impact of total hip arthroplasty surgical approach on patient-reported outcomes measurement information system computer adaptive tests of physical function and pain interference. *J Arthroplasty.* 2020;35(10):2899–2903. doi:10.1016/j.arth.20.05.006
37. Vella MA, Warshauer A, Tortorello G, et al. Long-term functional, psychological, emotional, and social outcomes in survivors of firearm injuries. *JAMA Surg.* 2020;155(1):51. doi:10.1001/jamasurg.2019.4533
38. Waldman OV, Hao SP, Houck JR, Lee NJ, Baumhauer JF, Oh I. Operative intervention does not change pain perception in patients with diabetic foot ulcers. *Clin Diabetes.* 2020;38(2):132–140. doi:10.2337/cd19-0031
39. Esaa F, Prezzano J, Pentland A, Ryan Wolf J. The utility of PROMIS domain measures in dermatologic care. *Arch Dermatol Res.* 2021;313(1):17–24. doi:10.1007/s00403-020-02074-1
40. Flynn KE, Lin L, Weinfurt KP. Sexual function and satisfaction among heterosexual and sexual minority U.S. adults: a cross-sectional survey. *PLoS One.* 2017;12(4):e0174981. doi:10.1371/journal.pone.0174981
41. Waheed A, Djurdjev O, Dong J, Gill J, Barbour S. Validation of self-reported race in a Canadian provincial renal administrative database. *Can J Kidney Health Dis.* 2019;6:2054358119859528. doi:10.1177/2054358119859528

Patient Related Outcome Measures

Publish your work in this journal

Patient Related Outcome Measures is an international, peer-reviewed, open access journal focusing on treatment outcomes specifically relevant to patients. All aspects of patient care are addressed within the journal and practitioners from all disciplines are invited to submit their work as well as healthcare researchers and patient support groups.

Submit your manuscript here: <http://www.dovepress.com/patient-related-outcome-measures-journal>

Dovepress

The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.