Maackiain Protects the Kidneys of Type 2 Diabetic Rats via Modulating the Nrf2/HO-1 and TLR4/NF- κ B/Caspase-3 Pathways

Jiahong Guo¹
Junying Li²
Hua Wei¹
Zhaozhi Liang¹

¹Department of Nephrology, Xinxiang Central Hospital, The Fourth Clinical College of Xinxiang Medical University, Xinxiang, 453000, People's Republic of China; ²Department of Nephrology, The Affiliated Hospital of Qingdao University Pingdu district, Pingdu City, Qingdao, Shandong, 266000, People's Republic of China

Background: Type 2 diabetes (T2D) is a global health burden that accounts for about 90% of all cases of diabetes. Injury to the kidneys is a serious complication of type 2 diabetes. Maackiain, apterocarpin extracted from roots of *Sophora flavescens*, has been traditionally used for various disease conditions. However, nothing is known about its possible potential effect on HFD/STZ-T2D-induced nephrotoxicity.

Methods: In this study, T2D rat model is created by high-fat diet (HFD) for 2 weeks with injection of a single dose of streptozotocin (35mg/kg body weight). T2D rats were orally administered with maackiain (10 and 20mg/kg body weight) for 7 weeks.

Results: Maackiain suppressed T2D-induced alterations in metabolic parameters, lipid profile and kidney functionality markers. By administering 10 and 20mg/kg maackiain to T2D rats, it was able to reduce lipid peroxidation while improving antioxidant levels (SOD, CAT, and GSH). Furthermore, the present study demonstrated the molecular mechanisms through which maackiain attenuated T2D-induced oxidative stress (mRNA: *Nrf2*, *Nqo-1*, *Ho-1*, *Gclc* and *Gpx-1*; protein: NRF2, NQO-1, HO-1 and NOX-4), inflammation (mRNA: *Tlr*, *Myd88*, *Ikb α* , *Mcp-1*, *Tgf- β* , *col4*, *Icam1*, *Vcam1* and *E-selectin*; Protein: TLR4, MYD88, NF- κ B, Ikb α , MCP-1; levels: TNF- α and MCP-1) and apoptosis (mRNA: *Bcl-2*, *Bax*, *Bad*, *Apaf-1*, *Caspase-9* and *Caspase-3*; protein: Bcl-2, Bax, Caspase-3 and Caspase-9) mediated renal injury. Additionally, significant improvement in kidney architecture was observed after treatment of diabetic rats with 10 or 20mg/kg maackiain.

Conclusion: Maackiain protects the kidney by decreasing oxidative stress, inflammation, and apoptosis to preserve normal renal function in type 2 diabetes.

Keywords: high-fat diet, streptozotocin, oxidative stress, inflammation, apoptosis, kidney

Introduction

Diabetes is a serious public health threat that respects no socioeconomic status globally in developed and developing countries.¹ Diabetes is a metabolic disorder that occurs when the pancreas is unable to produce insulin or the body is unable to use insulin efficiently. According to the most recent assessments of the Global Diabetes Organization, 463million adults now have diabetes, with this number expected to rise to 578million by 2030 and 700million by 2045.² Type 2 diabetes is a more common condition in adults where the body cannot effectively use insulin to make glucose into energy. It accounts for around 90% of all diabetes cases and is linked to multiple comorbidities and complications. Hyperglycemia affects the heart, blood vessels, eyes, nerves and teeth, including kidneys. Microvascular alterations in the kidneys caused by

Correspondence: Junying Li
Department of Nephrology, The Affiliated Hospital of Qingdao University Pingdu District, No.369 Shanghai Road, Qingdao, Pingdu City, Shandong, 266000, People's Republic of China
Email lijunying2020@sina.cn

Graphical Abstract

various mechanisms often result in diabetic kidney disease, also known as diabetic nephropathy.^{3,4}

Among various pathophysiological processes involved in the pathogenesis of diabetic nephropathy, oxidative stress, inflammation, and apoptosis have been considered significant contributors.⁵ Earlier evidence pointed out the contributory roles of elevated levels of reactive oxygen species and suppressed antioxidant defenses in the development of diabetic kidney disease.^{6,7} Recent reports also highlighted the role of Nrf2/Keap1/ARE pathway at the transcriptional level to regulate cellular oxidative and anti-oxidative status.⁸ Furthermore, nuclear factor erythroid 2-related factor 2 (Nrf2) knockout in streptozotocin-induced mice showed high levels of reactive oxygen species (ROS) with greater oxidative DNA damage and renal injury.⁹ Studies also reported the involvement of diabetes-mediated oxidative stress in instigating the inflammatory pathways for the progression of kidney damage.^{5,10} Inflammation has been recognized as an underlying mechanism that plays a critical role in the progression and pathogenesis of diabetic kidney disease.¹¹ It is also noteworthy to mention the involvement of TLR4/MyD88/NF- κ B signaling pathway and its effector mediators such as proinflammatory cytokines, chemokines, pro-fibrotic factors and adhesion molecules inactivation of inflammation.^{12,13} However, oxidative stress and inflammation work in coordination to activate mitochondria-dependent apoptotic pathway, which further damages the diabetic kidney and ultimately results in renal failure.⁵ So agents which counteract oxidative stress, inflammation and apoptosis might prevent diabetes-mediated renal damage.

Although there are many treatment modalities, including western medications, available to treat and manage type 2 diabetes-mediated complications and kidney damage, management with lesser side effects at affordable cost is still a big challenge for researchers. Nowadays, much emphasis has been given to herbal medication due to its relatively safe nature, less side effects and availability at low cost.^{14,15} *Sophora flavescens* is one such well-known Chinese herbal medicine. It has also history in traditional medicines of Japan, Korea, India, and some countries of Europe that are prescribed for various ailments such as skin burns, asthma, and jaundice dysentery, ulcers, fever and inflammatory disorders.¹⁶ Maackiain, apterocarpan isolated from the roots of *S. flavescens*, has a broad spectrum of biological activities including anti-cancer,¹⁷ anti-allergic,¹⁸ and inhibitory activity on monoamine oxidase B¹⁹ and anti-inflammatory properties.²⁰

Taking into consideration the diabetes-induced kidney problems as well as the safety properties of maackiain, the current study was designed to investigate the possible protective effect of maackiain on type 2 diabetes-induced renal dysfunctions and to characterize the anti-oxidative, anti-inflammatory, and anti-apoptotic mechanisms through which it protects renal injury using arat as an exploratory model.

Materials and Methods

Chemicals

Maackiain (MN) (<98%) was procured from Ruicong Ltd (Shanghai, China) and streptozotocin (STZ; <98%) was procured from Sigma-Aldrich, Inc., (St. Louis, MO, USA).

All of the other chemicals and reagents used in this research were obtained locally.

Animals

Adult male healthy Swiss Albino mice and Sprague-Dawley (SD) rats with body weights ranging from 25 ± 03 and 200 ± 20 g, respectively, were obtained from the Institute of Experimental Animals, Qingdao No.9 People's Hospital, China. Initially, after procurement, animals were shifted to aquarantine room to check the health status of animals by the veterinarian. After one week of acclimation to experimental room conditions, healthy animals were subjected to experiments. Animals were maintained in standard experimental room conditions ($22 \pm 3^\circ\text{C}$ temperature, $50 \pm 10\%$ humidity, 12 hours lighting) and had to access to *ad libitum* irradiated standard rodent diet and autoclaved water before dietary manipulation. All animal experiments were approved and carried out in compliance with Institutional Animal Care Guidelines (approval no.202009673590; Experimental Animal Ethics Committee, Qingdao No.9 People's Hospital, Shandong, China).

Induction of Type 2 Diabetes in Experimental Animals

Type 2 diabetes was induced in rats as described previously by Srinivasan et al²¹ and Mahmoud et al.²² Induction of type 2 diabetes in rats using a high fat diet [29.5% tallow of beef; 22% casein; 23% starch; 17.9% cellulose; 4% l-cysteine; 0.3% choline chloride; 1.8% vitamin mixture; 1.5% salt mixture (AIN-93 Vx)] for 2 weeks and a single low dose of STZ, 35mg/kg body weight dissolved in freshly prepared 0.1 M cold citrate buffer (pH 4.5) through intraperitoneal route administered intraperitoneally. After a week of STZ injection, rats with fasting blood glucose levels more than 12.5 mmol/L were classified as type 2 diabetic and used in

the study. However, control rats received a normal diet (56% starch; 18.5% protein; 8% fat; 12% fibre) and a freshly prepared 0.1 M cold citrate buffer (pH 4.5) through intraperitoneal route.

Acute Oral Toxicity Study

The acute toxicity of maackiain was investigated using OECD [1998] Guideline 425 in Swiss Albino male mice after oral administration up to a maximum dosage of 2000mg/kg. The animals were constantly monitored for two hours to assess their behavioral, neurological, and autonomic features.²³

Experimental Design

The rats were divided randomly into four groups of ten individuals each:

Group 1: Normal control rats received 1% sodium carboxymethyl cellulose (Na-CMC) vehicle

Group 2: Type 2 diabetic rats received 1% Na-CMC vehicle

Group 3: Type 2 diabetic rats received 10mg/kg/bw of maackiain

Group 4: Type 2 diabetic rats received 20mg/kg/bw of maackiain

Doses of maackiain were selected based on earlier study.²⁴ Meanwhile, groups 2 to 4 proceed with HFD for 7 weeks of treatment. Maackiain was suspended in 1% Na-CMC and administered orally with an oral gavage tube everyday for 7 weeks. The details are presented in Figure 1.

Assessment of Bodyweights, Feed and Water Intake and FBG Levels

Daily feed and water consumption and weekly body weight and fasting blood glucose levels were observed

Figure 1 Schematic representation of experimental design of the study.

Abbreviations: HFD, high-fat diet; STZ, streptozotocin; HFD, high-fat diet; FBG, fasting blood glucose.

throughout the study period. On the last day of the experiment, urine was collected in metabolic cages and animals were anesthetized (xylazine-12.5mg/kg/bw and ketamine-87.5mg/kg/bw, i.p), blood was collected via cardiac puncture and centrifuged at 2000g for 15 minutes to separate serum for biochemical analysis, and animals were sacrificed via cervical dislocation.

According to the manufacturer's instructions, insulin serum concentrations were determined using an enzyme-linked immunoassay (ELISA) test kit (Elabscience Biotechnology Co. Ltd., Wuhan, China). Glycated hemoglobin (HbA1c) and the serum concentrations of total cholesterol (TC), triglycerides (TG), high-density lipoprotein (HDL), low-density lipoprotein (LDL) (Nanjing Jiancheng Bioengineering Institute, Nanjing, China) were measured by using the automatic Beckman Coulter analyzer. Homeostatic Model Assessment of Insulin Resistance (HOMA-IR) and HOMA β -cell function was calculated using the formula.²⁵

Kidneys were taken immediately after necropsy and fixed in 10% neutral formalin buffer for histological examinations and biochemical and molecular studies stored at -80°C till further use.

Determination of Renal Functional Biomarkers

Renal functional assessment biomarkers such as serum creatinine ($\mu\text{mol/L}$), albumin (g/dL), urea (mg/dL) and uric acid ($\mu\text{mol/L}$) were evaluated using commercially available colorimetric assay kits (Elabscience Biotechnology Co. Ltd., Wuhan, China). BUN is calculated by using the following formulae:

$$\text{Urea}[\text{mmol/L}] = \text{BUN}[\text{mg/dL}] \times 10[\text{dL/L}] / 14 \times 2 \\ [\text{mg N/mmol urea}]$$

Further, Urinary protein concentrations (mg/day) were evaluated by bicinchoninic acid (BCA) assay method using BSA as standard (Thermo Fisher Scientific Inc., Waltham, MA, USA).

Determination of Oxidative and Antioxidative Status in Kidneys

The levels of lipid peroxidation products (MDA) and antioxidant defenses such as superoxide dismutase (SOD), catalase (CAT), and glutathione (GSH) were measured in 10% kidney tissue homogenate supernatant samples using commercially available kits (Nanjing Jiancheng

Bioengineering Institute, Nanjing, China). The Bradford protein assay kit assessed total protein concentrations (Sigma-Aldrich, St. Louis, MO, USA).

Reverse Transcription Quantitative Real-Time PCR (RT-qPCR) Analysis

Following the animals' euthanasia, the extracted kidneys were stored in RNA lysis solution to preserve the RNA's integrity. Total RNA was extracted using the TRIzol[®] Reagent, quantified with an nanodrop, and Complementary DNA (cDNA) was synthesized using a High-Capacity RNA-to-cDNA kit (Applied Biosystems, CA, USA). PCR was carried out using SYBR[™] Green PCR Master Mix (Applied Biosystems, Foster City, CA, USA) and was performed in StepOnePlus Real-time PCR system (Applied Biosystems, Foster City, CA, USA). Table 1 contains the primer sets used in RT-qPCR. After normalization to the housekeeping gene, ie, β -actin, the relative expression of each gene was measured. Three duplicates of each reaction were maintained to ensure the reproducibility of the findings.²⁶

Determination of Levels of TNF- α and MCP-1 Levels and NF- κ B p65 DNA Binding Activity

Using ELISA kits, cytokines (tumor necrosis factor; TNF- α and monocyte chemoattractant protein-1; MCP-1) levels were quantified in kidney homogenates according to the manufacturer's instructions (Elabscience Biotechnology Co. Ltd., Wuhan, China). According to the manufacturer's instructions, the DNA-binding activity of NF- κ B was determined using the TransAM[®] NF- κ B p65 transcription factor assay kit (Cat. No.40596, Active Motif, Tokyo, Japan). The optical density of protein-bound NF- κ B was measured at 450 nm. The TransAM format is perfect for assaying transcription factor binding to a consensus-binding site.

Histopathological Analysis

Left kidneys from both control and treatment groups were immediately fixed in 10% neutral formalin buffer until further investigation. On the day of processing, tissues were appropriately trimmed, subjected to a series of alcohol washes, embedded in paraffin, microtomed into 5 μm thick sections, and then stained with hematoxylin and eosin (H&E), Picro-Sirius Red (PSR) (ab246832) stains, and atrichrome stain kit (ab150686) (Abcam, Cambridge,

Table I PCR Primers Used in This Paper

<i>Nrf2</i>	F: 5'CCTATGCGTGAATCCCAAT3' R: 5'TGTGAGATGAGCCTCTAAGCG3'
<i>Keap1</i>	F: 5'ATGTTGACACGGAGGATTGG3' R: 5'TCATCCGCCACTCATTCT3'
<i>Nqo-1</i>	F: 5'GCGAGAAGAGCCCTGATTGT3' R: 5CTTCAGCTCACCTGTGATGTCAT3'
<i>Ho-1</i>	F: 5'CAAGCCGAGAATGCTGAGTT3' R: 5'CAGGGCCGTGTAGATATGGTA3'
<i>Gcl</i>	F: 5' TCGCCTCCGATTGAAGATG3' R: 5' TACTATTGGGTTTACCTGTGCC3'
<i>Gpx1</i>	F: 5' CAGTTCGGACATCAGGAGAAT3' R: 5' AGAGCGGGTGAGCCTTCT3'
<i>Tlr4</i>	F: 5' GTGGAAGTTGAACGAATGGA3' R: 5' TGGATGATGTTGGCAGCA3'
<i>Myd88</i>	F: 5' TCGCGCATCGGACAAACG3' R: 5' GCAATGGACCAGACACAGGT3'
<i>Col4</i>	F: 5' TACTGGCAGAGCCCTTGAGCC3' R: 5' CATTCTTTCTGGATTAGTGAAGC3'
<i>Tgf-β</i>	F: 5' GACCTCAATTGCGAGCTTTCT3' R: 5' AGTCCTCCTCCGCTTTAG3'
<i>Mcp-1</i>	F: 5'TCTCTTCTCCACCACTATGCA3' R: 5'GGCTGAGACAGCACGTGGAT3'
<i>Vcam-1</i>	F: 5'TGACAAGTCCCCATCGTTGA-3' R: 5'ACCTCGCGACGGCATAATT3'
<i>Icam-1</i> :	F: 5'CCTGTTTCCTGCCTCTGAA3' R: 5'GTCTGCTGAGACCCCTCTTG3'
<i>E-selectin</i> :	F: 5'TGAAGTGAAGGGATCAAGAAGACT3' R: 5'GCCGAGGGACATCATCACAT3'
<i>Bcl2</i>	F: 5' CCTGAGAGCAACCGAACG3' R: 5' CCTGAGAGCAACCGAACG3'
<i>Bax</i>	F: 5' CACCAGCTCTGAACAGATC3' R: 5' CTTCTTCCAGATGGTGAGC3'
<i>Bad</i>	F: 5' CCCCCCAATCTCTGGGCAGCG3' R: 5' TCACTGGGAGGGGTGGAGCC3'
<i>Apaf-1</i>	F: 5' GTAGACGGCTTTCTCCGCTC3' R: 5' CGGATCCAGGACACAAAAGC3'
<i>Caspase-9</i>	F: 5' ATGCAGGTCCCTGTCATG3' R: 5' GCTTGAGGTGGTTGTGGA3'
<i>Caspase-3</i>	F: 5' GAGCTTGGAACGCGAAGAAA3' R: 5' CCATTGCGAGCTGACATTCC3'
<i>β-actin</i>	F: 5'GTGCTATGTTGCTCTAGACTTCG3' R: 5'ATGCCACAGGATCCATACC3'

UK). An Olympus phase contrast microscope was used to examine and capture the pictures.

Immunohistochemistry

Immunohistochemistry was performed using the method described by Giribabu et al.²⁷ Kidney sections were deparaffinized, rehydrated, and antigen-retrieved using a 10 mM sodium citrate buffer solution (pH 6.0) for immunohistochemical investigations. Endogenous peroxidase activity was suppressed with 0.3% Hydrogen peroxide (H₂O₂) prior to sections being incubated overnight at 4°C with primary antibodies such as Anti-Nrf2 (ab62352), anti-NOX-4 (ab109225), anti-Keap-1 (ab218815), anti-NF-κB (ab16502), anti-MCP-1 (ab25124), anti-BCL-2 (ab194583) and anti-BAX (ab32503), anti-Caspase-9 (ab184786) (1:1000; Abcam, Cambridge, UK) diluted in 5% bovine serum albumin (BSA). After incubation, slides were rinsed three times in PBS, incubated for 1 hour with the appropriate secondary antibody for 60 min, stained with 3'-Diaminobenzidine (DAB), and counterstained with hematoxylin. The images were examined and captured using an Olympus phase-contrast microscope.

Immunofluorescence

Immunofluorescence was performed in accordance with the method described by Khalil et al.²⁸ On the kidney sections, immunofluorescence staining was performed after deparaffinization, rehydration, antigen retrieval with 10 mM sodium citrate buffer (pH 6.0), and blocking with 5% BSA. Following blocking, sections were incubated (anti-NQO-1: ab80588; anti-HO-1: ab189491; anti-TLR4: ab217274; anti-MYD88: ab219413; anti-IKB alpha: ab32518; and anti-Caspase-3: ab184787) overnight at 4°C with 1:1000 dilutions of primary antibodies in the serum. Following PBS washes, kidney sections were incubated with a secondary antibody conjugated with DyLight 488/594 (1:1000; Abcam, Cambridge, UK) at room temperature in the dark. The slices were mounted using UltraCruz (Santa Cruz, CA, USA) media containing 4',6-diamidino-2-phenylindole (DAPI) for nuclear staining and sealed with a coverslip. Finally, images were viewed and captured using fluorescence microscopy (Leica DM IRB, Germany).

Data Analysis

The data were analyzed using SPSS Statistics 22 software (IBM, Armonk, NY, USA). The data were expressed as mean

standard deviation (S.D). One-way analysis of variance (ANOVA) and Tukey's post hoc test was used to determine the significance of differences between groups. A probability value of less than 0.05 was considered significant.

Results

Effects of Maackiain on Acute Oral Toxicity Test

Maackiain was found to be safe following oral administration up to 2000mg/kg. No mice mortality was observed within 24 hours of the observational study.

Effects of Maackiain on Body Weights, Feed and Water Intake of HFD/STZ-Induced Type 2 Diabetic Rats

The changes in weekly body weights and feed and water consumption of control and experimental rats are shown in Figure 2A–C. When HFD-fed rats were compared to normal diet-fed rats, their body weights increased significantly ($p < 0.001$), while body weights dropped dramatically after STZ injection. As diabetic rats were given 10 ($p < 0.05$) or 20 ($p < 0.05$) mg/kg bw maackiain, their body weights increased substantially when compared to non-diabetic rats. Diabetic rats consumed significantly more feed ($p < 0.001$) and water ($p < 0.001$) than control rats, while treatment with maackiain 10 ($p < 0.05$) or 20 ($p < 0.001$) mg/kg bw dramatically decreased diabetic rats' feed and water intake.

Effects of Maackiain on Metabolic Parameters in Type 2 Diabetic Rats

Weekly FBG levels in diabetic rats did not decrease significantly when compared to control rats until the end of the experiment, whereas 7 weeks of treatment with 10 or 20 mg/kg maackiain resulted in a significant ($p < 0.001$) decrease in FBG levels to near normal levels when compared to diabetic rats at the same levels (Figure 2D). Serum insulin levels in diabetic rats were significantly ($p < 0.001$) lower than in control rats, however, treatment of 10 ($p < 0.05$) or 20 ($p < 0.05$) mg/kg maackiain resulted in a significant rise in insulin levels when compared to diabetic animals (Figure 2E). Diabetic rats also showed a dramatic increase in HOMA-IR ($p < 0.001$) (Figure 2F) and drastic decline in HOMA- β cell function ($p < 0.001$) (Figure 2G) with respect to controls, however, after treating diabetic rats with either 10 ($p < 0.001$) or 20 ($p < 0.001$) mg/kg maackiain showed a significant drop in HOMA-IR and rise in HOMA- β cell function.

The percentage of glycated hemoglobin (HbA1c) (Figure 2H) is within normal limits in controls, abnormally high in diabetic rats ($p < 0.001$), while treatment with either 10 ($p < 0.05$) or 20 ($p < 0.05$) mg/kg maackiain resulted in significant improvement.

Effects of Maackiain on Lipid Markers in Type 2 Diabetic Rats

When diabetic rats were compared to normal control rats, there were significant increases in total cholesterol ($p < 0.001$) (Figure 2I), triglycerides ($p < 0.001$) (Figure 2J), and LDL ($p < 0.05$) (Figure 2L), as well as a significant decrease in HDL ($p < 0.05$) (Figure 2K). Treatment with 10 ($p < 0.05$) or 20 ($p < 0.001$ for TC and TG; $p < 0.05$ for LDL and HDL) mg/kg maackiain for seven weeks significantly reduced the lipid metabolic alterations observed in diabetic rats.

Effects of Maackiain on Kidney Function Biomarkers in Type 2 Diabetic Rats

Diabetic rats exhibited significantly higher levels of renal functioning markers such as BUN ($p < 0.001$) (Figure 3A), serum creatinine ($p < 0.001$) (Figure 3B), 24 hrs urinary protein ($p < 0.001$) (Figure 3C), serum albumin ($p < 0.001$) (Figure 3D), serum urea ($p < 0.001$) (Figure 3E) and urinary uric acid ($p < 0.001$) (Figure 3F) than control rats. Treatment of diabetic rats with 10 ($p < 0.05$) or 20 ($p < 0.001$) mg/kg maackiain significantly improved renal function by significantly decreasing the above-mentioned parameters.

Effect of Maackiain on Histopathology of Kidney

Histopathological analysis of kidney tissue by H&E (Figure 4A), PSR (Figure 4B), MT (Figure 4C) staining depicted in Figure 4. Histological sections of the normal kidney displayed normal architecture with glomeruli and tubules in the kidney cortex. Diabetic rat kidney sections presented with thickened glomerular basement membrane, increased glomerular space, degenerated glomeruli, sclerotic glomeruli, tubular degeneration and interstitial fibrosis. However, kidney sections from 10 mg/kg maackiain treated diabetic rats revealed moderate improvement through reduced thickening of the glomerular basement membrane, reduction in sclerotic glomeruli and interstitial fibrosis. While, kidney sections from 20 mg/kg maackiain treated diabetic rats showed significant improvement, showing glomerular and tubular structures near normal limits.

Figure 2 The effects of maackiain metabolic parameters in HFD & low STZ induced diabetic rats. (A) Weekly body weights; (B) Food intake; (C) Water intake; (D) Fasting blood glucose levels (FBG); (E) serum insulin; (F) HOMA-IR; (G) HOMA-β cell function; (H) Glycated hemoglobin (HbA1c); (I) Total cholesterol (TC); (J) Triglycerides (TG); (K) High density lipoproteins (HDL); (L) Low density lipoproteins (LDL). The data were expressed as the mean ± S.D, n=6, **p < 0.001 and *p < 0.05 versus NC; ^p < 0.001 and ^p < 0.05 versus DC.

Abbreviations: NC, normal control; DC, diabetic control; DC+10 maackiain, 10 mg/kg/bw maackiain treated diabetic rats; DC+20: 20 maackiain, 20 mg/kg/bw maackiain treated diabetic rats.

PSR and MT observed collagen deposition in the tissues; the blue and red staining indicated collagen involvement, respectively. Collagen fibers were not seen in the renal interstitial control group, but higher collagen fibers were observed in diabetic groups. However, 10 and 20mg/kg maackiain treated diabetic rats were shown to have reduced collagen deposition.

Effect of Maackiain on Oxidative and Antioxidative Status in Type 2 Diabetic Rat's Kidney

The findings of this research showed dysregulation of oxidative and anti-oxidative state in diabetic rats, as shown by asignificant (p<0.05) rise in MDA (Figure 5A)

Figure 3 The effects of maackiain serum and urinary renal markers in HFD & low STZ induced diabetic rats. (A) Blood urea nitrogen (BUN); (B) Serum creatinine; (C) 24-hour urine protein; (D) Serum albumin; (E) Serum urea; (F) Urinary uric acid. The data were expressed as the mean $\pm$ S.D, n=6, **p < 0.001 versus NC; ^p < 0.001 and ^p < 0.05 versus DC.

Abbreviations: NC, normal control; DC, diabetic control; DC+10 maackiain, 10 mg/kg/bw maackiain treated diabetic rats; DC+20: 20 maackiain, 20 mg/kg/bw maackiain treated diabetic rats.

and as significant reduction in SOD (Figure 5C) and CAT (Figure 5D) and GSH (Figure 5B) levels. On the other hand, diabetic rats treated with either 10 or 20 mg/kg maackiain demonstrated significant ($p < 0.05$) improvement in the form of decreased lipid peroxidation products (MDA) and increased antioxidant defense activities such as SOD, CAT and GSH levels.

Figure 5E depicts the gene expression of the Nrf2-Keap-1 signaling pathway from control and experimental rats. The mRNA levels of *Nrf2*, *Nqo1*, *Ho-1*, *Gclc* and

Gpx1 were significantly ($p < 0.05$) reduced in diabetic rats, while *Keap1* ($p < 0.001$) was significantly elevated. Besides, diabetic rats treated with 10 or 20 mg/kg maackiain showed significantly ($p < 0.05$) increased mRNA expression levels of *Nrf2*, *Ho-1*, *Gclc* and *Gpx* with decreased *Keap1* as compared to diabetic rats.

Immunohistochemistry (Nrf2, NOX-4, and Keap1) and immunofluorescence (NQO-1 (red color) and HO-1 (green color)) studies confirmed the gene expression data, revealing decreased Nrf2 (Figure 6A), NQO-1 (Figure 6B), HO-1

Figure 4 The effects of maackiain histopathology in HFD & low STZ induced diabetic rats. **(A)** Hematoxylin and Eosin (H&E) staining; **(B)** Picro Sirius Red (PSR) staining; **(C)** Masson's Trichrome (MT) staining; H& E (Black color arrow shows necrosis, Green color arrow shows thickened glomerular basement membrane, blue color arrow shows increased glomerular space). PSR and MT (Blue color triangle shows collagen deposition and yellow color arrow also shows collagen deposition). Magnification = $\times 40$; Scale bar = $100\mu\text{m}$.

Abbreviations: NC, normal control; DC, diabetic control; DC+10 maackiain, 10 mg/kg/bw maackiain treated diabetic rats; DC+20: 20 maackiain, 20 mg/kg/bw maackiain treated diabetic rats.

(Figure 6B), NOX-4 (Figure 6C), and with increased Keap-1 (Figure 6D) protein distributions in the cortical and glomerular regions. In contrast, diabetic rats receiving either 10 or 20 mg/kg maackiain, Nrf2, NQO-1, HO-1, and NOX-4 proteins were increased, while the distribution of Keap-1 protein was decreased.

Effect of Maackiain on Inflammatory Status in Type 2 Diabetic Rat's Kidney

The inflammatory markers such as relative mRNA expression levels assessed by RT-qPCR, levels of cytokines and chemokines and activity of NF- κ B p65 in kidney were depicted in Figures 7 and 8. The mRNA expression levels of *Tlr4* (Figure 7A), *Myd88* (Figure 7B), *ikba* (Figure 7C), *Col-4* (Figure 8A), *Tgf- β* (Figure 8B), *Vcam* (Figure 8C), *Mcp-1* (Figure 8D), *E-Selectin* (Figure 8E) and *Icam* (Figure 8F) and levels of TNF- α (Figure 7E) and MCP-1 (Figure 7F) and activity of NF- κ B p65 (Figure 7D) were significantly elevated in

diabetic rats in comparison with same parameters in the control group. Contrary to that, those inflammatory markers' expression levels and activities were significantly ($p < 0.05$) reduced after treating diabetic animals with either 10 or 20 mg/kg maackiain than diabetic rats.

Additionally, immunofluorescence and immunohistochemistry experiments were used to determine the protein distribution pattern shown in Figure 9. The protein expression levels of TLR4 (red color) (Figure 9A), MYD88 (green color) (Figure 9A), NF- κ B p65 (Figure 9B), I κ B α (red color) (Figure 9C) and MCP-1 (Figure 9D) were elevated in diabetic rats, but those protein distributions were decreased in diabetic rats treated with either 10 or 20 mg/kg maackiain.

Effect of Maackiain on Apoptotic Status in Type 2 Diabetic Rat's Kidney

In diabetic rats' kidneys, *Bax* (Figure 10B), *Bad* (Figure 10C), *Apaf-1* (Figure 10D), *Caspase-3* (Figure 10E) and *Caspase-9*

Figure 5 The effects of maackiain on renal oxidative stress and antioxidant status in HFD & low STZ induced diabetic rats. **(A)** Lipid peroxidation product malondialdehyde (MDA) levels; **(B)** Reduced glutathione (GSH) content; **(C)** Superoxide dismutase (SOD) activity levels; **(D)** Catalase (CAT) activity levels in experimental rats kidney; **(E)** mRNA levels of *Nrf2*, *Keap1*, *Nqo1*, *Ho1*, *Gclc*, *Gpx1* in experimental rats kidney. The data were expressed as the mean $\pm$ S.D, n=6, **p < 0.001 and *p < 0.05 versus NC; ^^p < 0.001 and ^p < 0.05 versus DC.

Abbreviations: NC, normal control; DC, diabetic control; DC+10 maackiain, 10 mg/kg/bw maackiain treated diabetic rats; DC+20: 20 maackiain, 20 mg/kg/bw maackiain treated diabetic rats.

(Figure10F) gene expression levels were upregulated, while *Bcl-2* (Figure10A) gene expression levels were reduced. When compared to untreated diabetic rats, 10 or 20mg/kg maackiain treatment resulted in a reduction in Bax, Bad, Apaf-1, Caspase-3, and Caspase-9 mRNA expression but an increase in *Bcl-2* mRNA expression. In comparison to untreated diabetic rats, maackiain administration at 10 or 20mg/kg resulted in a decrease in Bax, Bad, Apaf-1, Caspase-3 and Caspase-9 but an increase in *Bcl-2* mRNA expression levels.

The findings of gene expression investigations in diabetic rats' kidneys were reproduced in immunohistochemistry [decreased Bcl-2 protein distribution (Figure11A) and

elevated Bax (Figure11B) and Caspase-9 (Figure11D)] and immunofluorescence [increased caspase-3 protein distribution] (Figure11C) in diabetic rat's kidney. However, treatment with either 10 or 20mg/kg maackiain increased Bcl-2 protein distribution levels in the kidneys of only diabetic rats while decreasing Bax, Caspase-3, and Caspase-9 protein distribution levels.

Discussion

The present study attracts special attention due to the unavailability of information on the possible effects of maackiain against HFD/STZ-induced type 2 diabetes-

Figure 6 The effects of maackiain on Nrf2/Keap-1 pathway protein destruction in HFD & low STZ induced diabetic rats. **(A)** Immunohistochemistry results of Nrf2; **(B)** Immunofluorescence double staining results of NQO-1 (red color) and HO-1 (green color); **(C)** Immunohistochemistry results of NOX-4; and **(D)** Keap-1. Magnification = $\times 40$; Scale bar = 100 μ m.

Abbreviations: NC, normal control; DC, diabetic control; DC+10, 10 mg/kg/bw maackiain treated diabetic rats; DC+20, 20 mg/kg/bw maackiain treated diabetic rats.

induced renal complications. The study provided clear evidence of significant protection against diabetes-mediated alterations in metabolic profile, lipid profile, and kidney dysfunction. Furthermore, maackiain treatment resulted in significant attenuation in oxidative stress, inflammation and apoptosis in kidneys of diabetic rats. Additionally, this study demonstrated the molecular mechanisms by which maackiain protects against HFD/

STZ type 2 diabetes-induced oxidative stress (Nrf2/Keap1/ARE), inflammation (TLR4/MYD88/NF- κ B) and apoptosis (intrinsic pathway) mediated renal damage.

Type-2 diabetes is characterized by chronic hyperglycemia and hypoinsulinemia and enhanced insulin resistance and β -cell dysfunction.^{29,30} Elevated glucose and glycated haemoglobin levels throughout the experimental period and reduced serum insulin levels, arise in HOMA-

Figure 7 The effects of maackiain on renal inflammatory marks in HFD & low STZ induced diabetic rats. mRNA levels of (A); *Tlr4*; (B) *Myd88* (C) *Ikba*; (D) renal NF-κB p65 (ELISA results); (E) renal TNF-α (ELISA) levels; (F) renal MCP-1 (ELISA) levels. The data were expressed as the mean ± S.D, n=6, **p < 0.001 versus NC; ^p<0.001 and ^^p<0.05 versus DC.

Abbreviations: NC, normal control; DC, diabetic control; DC+10 maackiain, 10 mg/kg/bw maackiain treated diabetic rats; DC+20: 20 maackiain, 20 mg/kg/bw maackiain treated diabetic rats.

IR, and areduction in HOMA-cell function were all indicators of hyperglycemia hypoinsulinemia, high insulin resistance, and low β-cell function, respectively. Similar findings were observed in a study conducted by Mahmoud et al²² in HFD/STZ-induced diabetic rats. However, a significant dose-dependent reversion in all these parameters was observed in HFD/STZ-induced diabetic rats after treating animals with 10 or 20mg/kg, indicating the anti-diabetic effect of Maackiain. The possibility of the anti-diabetic effect of flavonoids from *S.flavescens* is revealed from in silico molecular modeling and docking studies by interrupting Na⁺-glucose cotransporter activity in type 2 diabetes.³¹ The results of this study are well supported by

findings of Shao et al,³² which revealed an inhibitory effect of *S.flavescens* extract against a high-fat diet and low-dose streptozotocin-induced metabolic alterations (increase in fasted blood glucose levels and worsened lipid profile, glycosylated serum protein, glycosylated hemoglobin index with pancreas damage). Additionally, the anti-diabetic effect of maackiain was evidenced via AMP-activated protein kinase activity.³³

Dyslipidemia is another complication that is commonly associated with type 2 diabetic condition.^{34,35} Dyslipidemia in the diabetic milieu is responsible for developing diabetic nephropathy³⁰ and is considered a potential risk factor for cardiovascular diseases.³⁶ Usually, in

Figure 8 The effects of maackiain on renal inflammatory marks in HFD & low STZ induced diabetic rats. mRNA levels of (A) *Col4* (B) *Tgf-β*; (C) *Vcam*; (D) *Mcp-1*; (E) *E-Selectin*; (F) *Icam* gene in renal tissue of experimental rats. The data were expressed as the mean $\pm$ S.D, n=6, **p < 0.001 versus NC; ^p<0.001 and ^p<0.05 versus DC. **Abbreviations:** NC, normal control; DC, diabetic control; DC+10 maackiain, 10 mg/kg/bw maackiain treated diabetic rats; DC+20: 20 maackiain, 20 mg/kg/bw maackiain treated diabetic rats.

biological conditions, insulin acts on lipoprotein lipase to convert triglycerides to fatty acids and glycerol. Fatty acids generated from triglycerides undergo either re-esterification or oxidation for storage in adipose tissue or for fuel in muscle, respectively. It is reported that insulin resistance suppresses lipoprotein lipase activity and leads to a condition of triglyceridemia.³⁷ Furthermore, LDL transports cholesterol from the liver to other bodily tissues³⁸ and vice versa HDL transports cholesterol from bodily tissues to liver³⁹ to prevent cholesterol deposition. In the present study, HFD/STZ-induced diabetic rats presented with significantly increased total cholesterol, triglycerides and LDL levels with a significant decrease in HDL levels. The results are in consonance with the findings of

earlier reports.^{40,41} Furthermore, Hirano⁴² reported abnormal lipoprotein metabolism in diabetic nephropathy patients and managing dyslipidemia is particularly important otherwise, it leads to cardiovascular disease. Treatment of diabetic rats with either 10 or 20mg/kg maackiain resulted in significant improvement in dyslipidemia through a significant decrease in total cholesterol, triglycerides, and LDL levels with significant increase in HDL levels due to its insulinotropic effect or insulin secretagogue activity. These findings provided evidence that maackiain protects against diabetic kidney disease by improving lipid profile. The present findings support the earlier observations of Kim et al,⁴³ who reported hypolipidemic effects (significant reduction in elevated TC, TG,

Figure 9 The effects of maackiain TLR4/MYD88/NF-κB pathway protein destruction in HFD & low STZ induced diabetic rats. **(A)** Immunofluorescence double staining results of TLR4 (red color) and MYD88 (green color); **(B)** Immunohistochemistry staining results of NF-κB p65; **(C)** Immunofluorescence single staining results of IκBα (red color); **(D)** Immunohistochemistry staining results of MCP-1. Magnification = x40; Scale bar = 100μm.

Abbreviations: NC, normal control; DC, diabetic control; DC+10 maackiain, 10 mg/kg/bw maackiain treated diabetic rats; DC+20 maackiain, 20 mg/kg/bw maackiain treated diabetic rats.

and LDL-C levels and significant elevation in reduced HDL-C levels) of *S.flavescens* in poloxamer 407-induced hyperlipidemic and cholesterol-fed rats.

Evaluation of functional kidney biomarkers gives essential information related to the functioning of kidneys.^{44,45} In the present study, dysfunctioning of kidneys in diabetic rats was evident from the findings of kidney functionality assessment markers such as increased serum levels of BUN, urea, creatinine, albumin, 24 hrs urinary protein and urinary uric acid. The findings are in

line with earlier observations of Wen et al,⁴⁶ who reported that type 2 diabetic rats experienced kidney dysfunction through impairment in functional kidney markers. However, 10 or 20mg/kg maackiain treatment to diabetic rats dose-dependently decreased the serum levels of BUN, urea, creatinine, albumin, 24 hrs urinary protein, and urinary uric acid depicting significant improvement in kidney functional alterations induced by type 2 diabetic condition.

Numerous scientific studies have shown that oxidative stress is responsible for the pathophysiology of the

Figure 10 The effects of maackiain on renal apoptosis marks in HFD & low STZ induced diabetic rats. mRNA levels of (A) *Bcl-2* (B) *Bax*; (C) *Bad*; (D) *Apaf-1*; (E) *Caspase-3*; (F) *Caspase-9* gene in renal tissue of experimental rats. The data were expressed as the mean $\pm$ S.D, n=6, **p < 0.001 versus NC; ^^p < 0.001 and ^p < 0.05 versus DC. **Abbreviations:** NC, normal control; DC, diabetic control; DC+10 maackiain, 10 mg/kg/bw maackiain treated diabetic rats; DC+20: 20 maackiain, 20 mg/kg/bw maackiain treated diabetic rats.

diabetic kidneys.^{47,48} Oxidative stress is a condition that occurs when free radicals overwhelm antioxidant defenses.⁴⁹ Malondialdehyde (MDA), a lipid peroxidation product, induces kidney damage at cellular and tissue level and antioxidants such as SOD, CAT, and GSH protect against oxidative stress-mediated injuries.^{50,51} In this study, significantly increased levels of MDA and significantly decreased activities of SOD, catalase and levels of GSH were observed in kidneys of diabetic rats. Similar observations were reported in kidneys of experimentally induced type 2 diabetic rat model.⁵² In contrast, treatment of diabetic rats with either 10 or 20 mg/kg maackiain offered significant protection to oxidative stress in kidneys through the reduction in MDA

levels and elevation in SOD and catalase enzyme activities and GSH levels. Earlier it was reported that flavonoids from roots of *S. flavescentis* showed significant scavenging activities on 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical and ONOO-.⁵³ Furthermore, the antioxidant activity of maackiain was revealed by the significant reduction in 6-OHDA-induced elevation in levels of ROS.⁵⁴

Nowadays, the mechanism by which oxidative stress is mitigated has remained a more significant challenge. Recent studies claimed that the kidney cells are equipped with Nrf2, a transcriptional activator of antioxidant genes, to protect against oxidative stress.⁵⁵ At normal physiological state, Nrf2 is coupled with its negative regulator

Figure 1 The effects of maackiain on renal apoptosis marks in HFD & low STZ induced diabetic rats. Immunohistochemistry protein distribution of (A) Bcl-2; (B) Bax; (C) Immunofluorescence staining results of caspase-3 (green color); (D) Immunohistochemistry protein distribution of caspase-9 protein in renal tissue of experimental rats. Magnification = $\times 40$; Scale bar = $100\mu\text{m}$.

Abbreviations: NC, normal control; DC, diabetic control; DC+10 maackiain, 10 mg/kg/bw maackiain treated diabetic rats; DC+20: 20 maackiain, 20 mg/kg/bw maackiain treated diabetic rats.

Keap1 in the cytoplasm and at induced state, Nrf2 dissociates from Keap1 and enters into the nucleus to activate a battery of its regulated antioxidant genes such as *Nqo1*, *Ho-1*, *Gclc* and *Gpx1*.⁵⁶ To further know whether maackiain activates Nrf2 signaling pathway to act against diabetes-induced oxidative stress, we examined the mRNA and protein expression levels of Nrf2 and its downstream

genes and proteins. This study revealed that diabetic rats showed downregulation in mRNA (*Nrf2*, *Nqo-1*, *Ho-1*, *Gclc* and *Gpx-1*) and protein (Nrf2, NQO-1, NOX-4 and HO-1) expressions of Nrf2 pathway in association with an upregulation in mRNA and protein expressions of Keap1 in kidneys. Similar results of downregulation in mRNA and protein expressions of Nrf2 and its regulated genes

and proteins and upregulation in mRNA and protein expression level of Keap1 were observed in experimentally induced type 2 diabetic rat model.⁴⁶ Additionally, the importance of Nrf2 is evidenced from aNrf2 knockout study in which streptozotocin-induced Nrf2 knockout mice experienced hyperglycemia-induced oxidative stress and kidney damage.^{9,57} Interestingly, treating diabetic rats with maackiain at 10 or 20mg/kg body weight illustrated upregulation in Nrf2 and its regulated genes and proteins and downregulation in Keap1 gene and protein expression, demonstrating the activation of Nrf2 signaling pathway to counteract the diabetes-induced oxidative stress in kidneys.

Inflammation, in addition to oxidative stress, is a common feature of chronic kidney disease and a key mediator in its progression.⁵⁸ In recent years, the involvement of TLR4/MYD88/NF- κ B pathway in aggravating inflammation in kidneys has surfaced.⁵⁹ Studies have also been reported the implication of TLR4/NF- κ B pathway in acute and chronic kidney injuries, including diabetic nephropathy.^{59,60} At normal physiological state, NF- κ B is located in the cytoplasm with its negative regulator, I κ B α , but at induced state, I κ B α is degraded by IKK β and allows NF- κ B to enter into the nucleus to induce the expression of various effector molecules.¹² TLR4 through MYD88 (adaptor protein) activates NF- κ B, which in turn activates a range of effector molecules of inflammation such as pro-inflammatory cytokines, pro-fibrotic factors, chemokines and adhesion molecules.¹² In this study, HFD/STZ-induced type 2 diabetic condition caused stimulation in TLR4/MYD88/NF- κ B pathway through elevation in mRNA (TLR4, MYD88 and I κ B α) and protein (TLR4, MYD88, I κ B α and NF- κ B) expression levels and NF- κ B DNA binding activity. The activation of TLR4/MYD88/NF- κ B pathway elevates the levels of pro-inflammatory cytokine (TNF- α), chemokines (MCP-1) and mRNA expression levels of pro-fibrotic factors (MCP-1, TGF- β and Col-4), intercellular adhesion molecule (ICAM1) vesicular adhesion molecule (VCAM1) and selectin (E-selectin). Earlier it was reported that induction of levels of pro-inflammatory cytokines, chemokines and pro-fibrotic factors as a result of activation of NF- κ B leads to macrophage infiltration, extracellular matrix deposition and fibrosis and ended up with huge damage to the kidney.²⁷ In contrast, maackiain (10 or 20mg/kg) treatment showed anti-inflammatory effects in the form of reduced mRNA (*Tlr4*, *Myd88* and *Ikb α*) and protein (TLR4, MYD88, I κ B α and NF- κ B) expressions along with

decreased NF- κ B DNA binding activity as a consequence it limits the expression of various above mentioned inflammatory effector molecules and protects the kidney from diabetes-mediated inflammatory damage.

Previously, Ma et al²⁰ demonstrated the anti-inflammatory effects of extracts of *S.flavescens* residues both in-vivo (inhibition of ear and paw swelling and reduction in paw swelling production levels of PGE₂ in inflammatory tissues) and in-vitro (dose dependent inhibition in release of pro-inflammatory cytokines such as TNF- α , IL-6, NO and MCP-1 in LPS-induced RAW264.7 cells) study findings. Furthermore, the anti-inflammatory effects of maackiain is evidenced from its inhibitory action on LPS-induced NO production.⁶¹

A plethora of reports emphasized that oxidative stress and inflammation work together to activate apoptotic signals to further damage injury and lead to organ failure.^{5,62,63} The increased generation of ROS has been shown to upregulate NF- κ B and vice versa NF- κ B and/or its mediators have been shown to promote ROS production^{60,64} further. Inflammatory mediators or ROS generation in mitochondria due to hyperglycemia can release cytochrome c into cytosol which then forms apoptosome to activate executioner caspase ie, caspase-3.⁶⁵ The activated caspase-3 then cleaves DNA repair enzymes and cell stabilizing proteins, leading to apoptotic cell death.⁶⁵ Thus, suppressing elevated oxidative stress and inflammation can be ideal for preventing apoptosis in the diabetic kidney. In the present study, mRNA expression levels of *Bcl-2*, *Bax*, *Bad*, *Apaf-1*, *Caspase-9* and *Caspase-3* and protein expression levels of Bcl-2, Bax, Caspase-9 and Caspase-3 were upregulated in kidneys of diabetes-induced rats indicating cellular apoptosis in the kidney. These findings are consistent with the earlier findings of elevated expression of apoptotic mediators and subsequent apoptotic cell death in diabetic kidneys.^{5,66} Whereas, treatment of diabetic rats with either 10 or 20mg/kg maackiain significantly reduced the expression of pro-apoptotic factors (Bax and Bad), Apaf-1, initiator caspase (Caspase-9), executioner caspase (Caspase-3) and significantly improved anti-apoptotic factors (Bcl-2) signifying the anti-apoptotic potential of maackiain. The observed attenuation after maackiain treatment in renal apoptosis might be due to observed suppression in oxidative stress and inflammation. The inhibitory action of maackiain on apoptosis is revealed by its anti-apoptotic effect on 6-OHDA-induced apoptosis.⁵⁴ Supporting these observations, Kim et al⁶⁷ demonstrated protective effect of *S.flavescens* against

1-methyl-4-phenylpyridinium ion-induced intrinsic apoptosis by upregulating suppressed Bcl-2 expression and downregulating elevated Bax, cytochrome c and caspase-3 expression. Furthermore, supporting the observations of biochemical and molecular findings, histopathological findings revealed significant improvement in kidney architecture after treatment of diabetic rats with 10 or 20mg/kg maackiain.

Conclusions

In conclusion, HFD/STZ-induced type 2 diabetic rats showed substantial changes in metabolic parameters, lipid profile, renal functional markers, oxidative stress, inflammatory and apoptosis markers. However, treatment of HFD/STZ-induced type 2 diabetic rats with 20mg/kg body weight of maackiain showed more effect than 10mg/kg body weight of maackiain treatment in the reduction of metabolic, lipid profile and fictional kidney alterations, oxidative stress, inflammatory and apoptosis markers. The current research further showed maackiain nephroprotective benefits via modulating Nrf2/Keap1/ARE, TLR4/MYD88/NF- κ B, and Bcl2/Bax/Caspase-3/Caspase-9 pathways in HFD/STZ-induced type 2 diabetes. Based on these results, maackiain has excellent potential for use as an agent renal change in diabetic patients.

Data Sharing Statement

The data used to support the findings of this study are available from the corresponding author upon request.

Disclosure

The authors declare that there are no conflicts of interest.

References

- International Diabetes Federation. *IDF Diabetes Atlas*. 8th ed. Brussels, Belgium: International Diabetes Federation; 2017.
- SaeediP, SalpeaP, KarurangaS, et al. Mortality attributable to diabetes in 20–79 years old adults, 2019 estimates: results from the International Diabetes Federation Diabetes Atlas. *Diabetes Res Clin Pract*. 2020;162:108086. doi:10.1016/j.diabres.2020.108086
- AndersH-J, HuberTB, IsermannB, SchifferM. CKD in diabetes: diabetic kidney disease versus nondiabetic kidney disease. *Nat Rev Nephrol*. 2018;14(6):361–377.
- Anguiano GómezL, LeiY, Kumar DevarapuS, AndersH. The diabetes pandemic suggests unmet needs for ‘CKD with diabetes’ in addition to ‘diabetic nephropathy’—implications for pre-clinical research and drug testing. *Nephrol Dial Transplant*. 2018;33(8):1292–1304.
- Al HroobAM, AbukhalilMH, AlghonmeenRD, MahmoudAM. Ginger alleviates hyperglycemia-induced oxidative stress, inflammation and apoptosis and protects rats against diabetic nephropathy. *Biomed Pharmacother*. 2018;106:381–389.
- FernandesSM, CordeiroPM, WatanabeM, FonsecaCD, VattimoMF. The role of oxidative stress in streptozotocin-induced diabetic nephropathy in rats. *Arch Endocrinol Metabol*. 2016;60(5):443–449. doi:10.1590/2359-3997000000188
- MahmoodniaL, AghadavodE, BeigrezaeiS, Rafieian-KopaeiM. An update on diabetic kidney disease, oxidative stress and antioxidant agents. *JRenal Injury Prev*. 2017;6(2):153.
- VomundS, SchäferA, ParnhamMJ, BrüneB, Von KnethenA. Nrf2, the master regulator of anti-oxidative responses. *IntJMol Sci*. 2017;18(12):2772. doi:10.3390/ijms18122772
- JiangT, HuangZ, LinY, ZhangZ, FangD, ZhangDD. The protective role of Nrf2 in streptozotocin-induced diabetic nephropathy. *Diabetes*. 2010;59(4):850–860.
- BaoL, LiJ, ZhaD, et al. Chlorogenic acid prevents diabetic nephropathy by inhibiting oxidative stress and inflammation through modulation of the Nrf2/HO-1 and NF- κ B pathways. *IntImmunopharmacol*. 2018;54:245–253.
- Donate-CorreaJ, Luis-RodríguezD, Martín-NúñezE, et al. Inflammatory targets in diabetic nephropathy. *JClin Med*. 2020;9(2):458.
- SuryavanshiSV, KulkarniYA. NF- κ B: apotential target in the management of vascular complications of diabetes. *Front Pharmacol*. 2017;8:798.
- SuQ, LvX, SunY, et al. Role of TLR4/MyD88/NF- κ B signaling pathway in coronary microembolization-induced myocardial injury prevented and treated with nicorandil. *Biomed Pharmacother*. 2018;106:776–784.
- ModakM, DixitP, LondheJ, GhaskadbiS, DevasagayamTP. Recent advances in Indian herbal drug research guest editor: Thomas Paul Asir Devasagayam Indian herbs and herbal drugs used for the treatment of diabetes. *Nutrition*. 2007;40(3):163–173.
- WangJ, FengB, XiongX. Chinese herbal medicine for the treatment of obesity-related hypertension. *Evid Based Complement Altern Med*. 2013;2013. doi:10.1155/2013/757540
- HeX, FangJ, HuangL, WangJ, HuangXJ. Traditional usage, phytochemistry and pharmacology of an important traditional Chinese medicine. *Sophora Flavescens Ait*. 2015;172:10–29.
- CaoX, HeQ. Anti-tumor activities of bioactive phytochemicals in *sophora flavescens* for breast cancer. *Research*. 2020;12:1457.
- NariaiY, MizuguchiH, OgasawaraT, et al. Disruption of heat shock protein 90 (Hsp90)-protein kinase C δ (PKC δ) interaction by (–)-maackiain suppresses histamine H1 receptor gene transcription in HeLa cells. *JBiol Chem*. 2015;290(45):27393–27402.
- LeeHW, RyuHW, KangM-G, et al. Potent selective monoamine oxidase inhibition by maackiain, apterocarpan from the roots of *Sophora flavescens*. *Bioorgan Med Chem Lett*. 2016;26(19):4714–4719.
- MaH, HuangQ, QuW, et al. In vivo and invitro anti-inflammatory effects of *Sophora flavescens* residues. *JEthnopharmacol*. 2018;224:497–503.
- SrinivasanK, ViswanadB, AsratL, KaulC, RamaraoP. Combination of high-fat diet-fed and low-dose streptozotocin-treated rat: a model for type 2 diabetes and pharmacological screening. *Pharmacol Res*. 2005;52(4):313–320.
- MahmoudAM, AshourMB, Abdel-MoneimA, AhmedOM. Hesperidin and naringin attenuate hyperglycemia-mediated oxidative stress and proinflammatory cytokine production in high fat fed/streptozotocin-induced type 2 diabetic rats. *JDiabetes Complicat*. 2012;26(6):483–490.
- Organisation for Economic Co-operation and Development. *Test No.425: Acute Oral Toxicity: Up-And-Down Procedure*. OECD Publishing; 2008.
- MizuguchiH, NariaiY, KatoS, et al. Maackiain is a novel antiallergic compound that suppresses transcriptional upregulation of the histamine H1 receptor and interleukin-4 genes. *Pharmacol Res Perspect*. 2015;3(5):e00166.

25. RuntuweneJ, ChengK-C, AsakawaA, et al. Rosmarinic acid ameliorates hyperglycemia and insulin sensitivity in diabetic rats, potentially by modulating the expression of PEPCK and GLUT4. *Drug Design Dev Ther.* 2016;10:2193.
26. YelumalaiS, GiribabuN, Kamarulzaman KarimSZO, SallehNB, SallehNB. In vivo administration of quercetin ameliorates sperm oxidative stress, inflammation, preserves sperm morphology and functions in streptozotocin-nicotinamide induced adult male diabetic rats. *Archiv Med Sci.* 2019;15(1):240. doi:10.5114/aoms.2018.81038
27. GiribabuN, KarimK, KilariEK, SallehN. Phyllanthus niruri leaves aqueous extract improves kidney functions, ameliorates kidney oxidative stress, inflammation, fibrosis and apoptosis and enhances kidney cell proliferation in adult male rats with diabetes mellitus. *JEthnopharmacol.* 2017;205:123–137. doi:10.1016/j.jep.2017.05.002
28. KhalilASM, GiribabuN, YelumalaiS, ShahzadH, KilariEK, SallehN. Myristic acid defends against testicular oxidative stress, inflammation, apoptosis: restoration of spermatogenesis, steroidogenesis in diabetic rats. *Life Sci.* 2021;278:119605. doi:10.1016/j.lfs.2021.119605
29. HudishLI, ReuschJE, SusseIL. β cell dysfunction during progression of metabolic syndrome to type 2 diabetes. *JClin Invest.* 2019;129(9):4001–4008.
30. QiY, MaJ, LiS, WL. Applicability of adipose-derived mesenchymal stem cells in treatment of patients with type 2 diabetes. *Stem Cell Res Ther.* 2019;10(1):274.
31. GiriJaR, ArunaS, SangeethaR. In silico molecular modelling and docking studies of sophora flavescens derived flavonoids against SGLT2 for type 2 diabetes mellitus. *IntJBioinform Biol Sci.* 2018;6(2):71–76.
32. ShaoJ, LiuY, WangH, LuoY, ChenLJ. An integrated fecal microbiome and metabolomics in T2DM rats reveal antidiabetes effects from host-microbial metabolic axis of EtOAc extract from Sophora flavescens. *Oxid Med Cell Longev.* 2020;2020:1–25. doi:10.1155/2020/1805418
33. HuangY, HaoJ, TianD, et al. Antidiabetic activity of a flavonoid-rich extract from Sophora davidii (Franch.) Skeels in KK-Ay mice via activation of AMP-activated protein kinase. *Front Pharmacol.* 2018;9:760.
34. MatsuzakaT, ShimanoHJ. New perspective on type 2 diabetes, dyslipidemia and non-alcoholic fatty liver disease. *JDiabetes Investig.* 2020;11(3):532–534.
35. KawanamiD, MatobaK, UtsunomiyaK. Dyslipidemia in diabetic nephropathy. *Renal Replacement Ther.* 2016;2(1):16. doi:10.1186/s41100-016-0028-0
36. BurkhardtRJ. Hyperlipidemia and cardiovascular disease: new insights on lipoprotein (a). *Curr Opin Lipidol.* 2019;30(3):260–261.
37. QuispeR, MartinSS, JonesSR. Triglycerides to high-density lipoprotein-cholesterol ratio, glycemic control and cardiovascular risk in obese patients with type 2 diabetes. *Diabetes Obes.* 2016;23(2):150–156.
38. FerenceBA, KasteleinJJ, RayKK, et al. Association of triglyceride-lowering LPL variants and LDL-C-lowering LDLR variants with risk of coronary heart disease. *JAMA.* 2019;321(4):364–373.
39. XuY, HeZ, KingGL. Introduction of hyperglycemia and dyslipidemia in the pathogenesis of diabetic vascular complications. *Curr Diabetes Rep.* 2005;5(2):91–97.
40. ErejuwaOO, NwobodoNN, AkpanJL, et al. Nigerian honey ameliorates hyperglycemia and dyslipidemia in alloxan-induced diabetic rats. *Nutrients.* 2016;8(3):95. doi:10.3390/nu8030095
41. IroniEA, OboghG, AkindahunsiAA. Antidiabetic effects of Mangifera indica Kernel Flour-supplemented diet in streptozotocin-induced type 2 diabetes in rats. *Food Sci Nutr.* 2016;4(6):828–839. doi:10.1002/fsn3.348
42. HiranoT. Abnormal lipoprotein metabolism in diabetic nephropathy. *Clin Exp Nephrol.* 2014;18(2):206–209. doi:10.1007/s10157-013-0880-y
43. KimHY, JeongDM, JungHJ, JungYJ, YokozawaT, ChoiJS. Hypolipidemic effects of Sophora flavescens and its constituents in poloxamer 407-induced hyperlipidemic and cholesterol-fed rats. *Biol Pharm Bull.* 2008;31(1):73–78. doi:10.1248/bpb.31.73
44. Al-SawalhaNA, AlSariRR, KhabourOF, AlzoubiKH. Influence of prenatal waterpipe tobacco smoke exposure on renal biomarkers in adult offspring rats. *Inhal Toxicol.* 2019;31(5):171–179. doi:10.1080/08958378.2019.1624897
45. SeverinMJ, CampagnoRV, BrandoniA, TorresAM. Time evolution of methotrexate-induced kidney injury: a comparative study between different biomarkers of renal damage in rats. *Clin Exp Pharmacol Physiol.* 2019;46(9):828–836. doi:10.1111/1440-1681.13122
46. WenW, LinY, TiZ. Antidiabetic, antihyperlipidemic, antioxidant, anti-inflammatory activities of ethanolic seed extract of Annona reticulata L.in streptozotocin induced diabetic rats. *Front Endocrinol (Lausanne).* 2019;10:716. doi:10.3389/fendo.2019.00716
47. DuniA, LiakopoulosV, RoumeliotisS, PeschosD, DounousiE. Oxidative stress in the pathogenesis and evolution of chronic kidney disease: untangling Ariadne's thread. *IntJ Mol Sci.* 2019;20(15):3711. doi:10.3390/ijms20153711
48. LeeSJ, KangJS, KimHM, et al. CCR2 knockout ameliorates obesity-induced kidney injury through inhibiting oxidative stress and ER stress. *PLoS One.* 2019;14(9):e0222352. doi:10.1371/journal.pone.0222352
49. AlotaibiMF, Bin-JumahM, ElgebalyH, MahmoudAM, MahmoudAM. Olea europaea leaf extract up-regulates Nrf2/ARE/HO-1 signaling and attenuates cyclophosphamide-induced oxidative stress, inflammation and apoptosis in rat kidney. *Biomed Pharmacother.* 2019;111:676–685. doi:10.1016/j.biopha.2018.12.112
50. AbdelhalimMAK, QaidHA, Al-MohyYH, GhannamMM. The protective roles of vitamin eand α -lipoic acid against nephrotoxicity, lipid peroxidation, and inflammatory damage induced by gold nanoparticles. *IntJ Nanomedicine.* 2020;15:729–734. doi:10.2147/IJN.S192740
51. ParkWS, ParkMS, KangSW, et al. Hesperidin shows protective effects on renal function in ischemia-induced acute kidney injury (Sprague-Dawley Rats). *Transplant Proc.* 2019;51(8):2838–2841. doi:10.1016/j.transproceed.2019.02.055
52. MaheshwariR, BalaramanR, SenAK, ShuklaD, SethA. Effect of concomitant administration of coenzyme Q10 with sitagliptin on experimentally induced diabetic nephropathy in rats. *Ren Fail.* 2017;39(1):130–139. doi:10.1080/0886022X.2016.1254659
53. JungHJ, KangSS, WooJJ, ChoiJS. Anew lavandulylated flavonoid with free radical and ONOO- scavenging activities from Sophora flavescens. *Arch Pharm Res.* 2005;28(12):1333–1336. doi:10.1007/BF02977897
54. TsaiRT, TsaiCW, LiuSP, et al. Maackiain ameliorates 6-hydroxydopamine and SNCA pathologies by modulating the PINK1/parkin pathway in models of Parkinson's disease in Caenorhabditis elegans and the SH-SY5Y cell line. *IntJ Mol Sci.* 2020;21(12):4455. doi:10.3390/ijms21124455
55. LiX, ZouY, XingJ, et al. Pretreatment with roxadustat (FG-4592) attenuates folic acid-induced kidney injury through antiapoptosis via Akt/GSK-3 β /Nrf2 pathway. *Oxid Med Cell Longev.* 2020;2020:6286984.
56. AshrafizadehM, AhmadiZ, SamarghandianS, et al. MicroRNA-mediated regulation of Nrf2 signaling pathway: implications in disease therapy and protection against oxidative stress. *Life Sci.* 2020;244:117329. doi:10.1016/j.lfs.2020.117329
57. ChoiBH, KangKS, KwakMK. Effect of redox modulating NRF2 activators on chronic kidney disease. *Molecules (Basel, Switzerland).* 2014;19(8):12727–12759. doi:10.3390/molecules190812727

58. PedruzziLM, CardozoLF, DalepraneJB, et al. Systemic inflammation and oxidative stress in hemodialysis patients are associated with down-regulation of Nrf2. *JNephrol.* 2015;28(4):495–501. doi:10.1007/s40620-014-0162-0
59. TanJ, WanL, ChenX, et al. Conjugated linoleic acid ameliorates high fructose-induced hyperuricemia and renal inflammation in rats via NLRP3 inflammasome and TLR4 signaling pathway. *Mol Nutr Food Res.* 2019;63(12):e1801402. doi:10.1002/mnfr.201801402
60. ZhuS, TangS, SuF. Dioscin inhibits ischemic stroke-induced inflammation through inhibition of the TLR4/MyD88/NF- κ B signaling pathway in arat model. *Mol Med Rep.* 2018;17(1):660–666.
61. LeeW, KuS-K, BaeJ. Anti-inflammatory effects of Baicalin, Baicalein, and Wogonin invitro and in vivo. *Inflammation.* 2015;38(1):110–125.
62. KandemirFM, YildirimS, KucuklerS, CaglayanC, MahamaduA, DortbudakMB. Therapeutic efficacy of zingerone against vancomycin-induced oxidative stress, inflammation, apoptosis and aquaporin 1permeability in rat kidney. *Biomed Pharmacother.* 2018;105:981–991. doi:10.1016/j.biopha.2018.06.048
63. ChenX, WeiW, LiY, HuangJ, CiX. Hesperetin relieves cisplatin-induced acute kidney injury by mitigating oxidative stress, inflammation and apoptosis. *Chem Biol Interact.* 2019;308:269–278. doi:10.1016/j.cbi.2019.05.040
64. LingappanK. NF- κ B in oxidative stress. *Curr Opin Toxicol.* 2018;7:81–86. doi:10.1016/j.cotox.2017.11.002
65. FuchsY, StellerH. Live to die another way: modes of programmed-cell death and the signals emanating from dying cells. *Nat Rev Mol Cell Biol.* 2015;16(6):329–344. doi:10.1038/nrm3999
66. MarquesC, MegaC, GonçalvesA, et al. Sitagliptin prevents inflammation and apoptotic cell death in the kidney of type 2 diabetic animals. *Mediators Inflamm.* 2014;2014:538737. doi:10.1155/2014/538737
67. KimHY, JeonH, KimH, KooS, KimS. Sophora flavescens aiton decreases MPP(+)-induced mitochondrial dysfunction in SH-SY5Y cells. *Front Aging Neurosci.* 2018;10:119. doi:10.3389/fnagi.2018.00119

Drug Design, Development and Therapy

Dovepress

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which has also

been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/drug-design-development-and-therapy-journal>