**RESPONSE TO LETTER** 

# In vivo Simulation-Based Learning for Undergraduate Medical Students: Teaching and Assessment [Response to Letter]

Michail Sideris<sup>1,\*</sup> John Gerrard Hanrahan<sup>2,\*</sup> Marios Nicolaides<sup>3</sup> Jade Jagiello<sup>4</sup> Kathrine S Rallis<sup>3</sup> Elif Emin<sup>1</sup> Efthymia Theodorou<sup>3</sup> Rebecca Mallick<sup>5</sup> Funlayo Odejinmi<sup>6</sup> Nikolaos Lymperopoulos<sup>7</sup> Apostolos Papalois<sup>8</sup> George Tsoulfas <sup>(p)</sup>

<sup>1</sup>Women's Health Research Unit, Queen Mary University of London, London, UK; <sup>2</sup>Addenbrooke's Hospital, Cambridge University Hospitals, Cambridge, UK; <sup>3</sup>Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK; <sup>4</sup>University College, London Hospital, London, UK; <sup>5</sup>Princess Royal Hospital, Brighton and Sussex University Hospitals NHS Trust, Haywards Heath, UK; <sup>6</sup>Whipps Cross University Hospital, Barts Health NHS Trust, London, UK; <sup>7</sup>Guy's and St Thomas Hospitals NHS Foundation Trust, London, UK; <sup>8</sup>Special Unit for Biomedical Research and Education, Aristotle University of Thessaloniki, Thessaloniki, Greece; <sup>9</sup>Department of Surgery, Aristotle University of Thessaloniki, Thessaloniki, Greece

\*These authors contributed equally to this work

Correspondence: Michail Sideris Women's Health Research Unit, Queen Mary University of London, Yvonne Carter Building, 58 Turner Street, London, EI 2AB, UK Email m.sideris@qmul.ac.uk

# **Dear editor**

We read with interest the response to our article from Nicholls-Mindlin et al concluding that animal use should be eradicated from medical school curricula. The arguments around this conclusion are articulated around their own experience that several students find in vivo SBL demoralising and upsetting. An example given to support this statement is the investigation of the physiological effects of clinically relevant drugs like adrenaline in vivo during their pre-clinical years. Nicholls-Mindlin et al claim that although Oxford University follows 3R principles, still the additional educational value of animal use is negligible. In their letter, there is also mention of selection bias of the ESMSC course, as we selected the most motivated students and hence our experience would be obscured. They finally underline that each medical student has the right to follow the ethical pillars of autonomy and justice and therefore withdrawing participation from activities that include animals should not result in any assessment or progression failure in their studies.

The initial point raised describes a personal, single-centred experience which is perhaps as vulnerable to selection bias as the authors propose the ESMSC course is. We would suggest formalising these views, perhaps through a qualitative study, to capture and analyse the experiences of their colleagues – this could be meaningful, particularly as they have described an exercise in which we agree does not seem to mandate in vivo experimentation. Our study systematically reviews the literature and is informed by 5+ years of experience in delivering what we believe to be meaningful educational experiences, an aspect of which is in vivo SBL. Future reviews like ours would certainly be informed by formal characterisation of negative experiences, such as the one described, to allow appraisal of methods and quality of conclusions drawn.

The second point of surgically minded students preferring limited animal use, with what the authors quite rightly raised as a selectively biased cohort, is certainly not striking. It is well recognised that surgical educators are aiming to find valid alternatives that provide educational value. One only needs to observe the explosion of productivity in the simulation literature to see this. The ESMSC course itself evolved, with later iterations depending on fewer in vivo modules and more hyperrealistic simulations, mirroring this trend. However, the main thesis of the article is to highlight the place in vivo modules still have in surgical education. This position

© 2021 Sideris et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress.com/terms work you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs A2 and 5 of our Terms (https://www.dovepress.com/terms.php). is indeed waning; however, the review describes educational experiences outlining remaining value under specific circumstances.

We agree with the aspiration towards the admonishment of in vivo SBL from under and indeed post-graduate education. However, with documented benefits of their use for training, it is important that we have viable and cost-effective alternatives which maintain pre-clinical skill development but also motivate students to enter the operating theatre.

We would also like to clarify a couple of things regarding the ESMSC course, considering the issues raised in this letter:

ESMSC is a mixed fidelity course; the animal use is extremely limited and accounts for less than 10% of the educational time, which has reduced and is reducing in keeping with the ethical pillars of 3Rs. Further to this, students who sign up for the course are fully informed about the use of animals and must attend the induction talk, which provides a comprehensive overview of the course and use of animals. Our students are fully consented and are able to withdraw at any time.

In conclusion, Nicholls-Mindlin et al raise important points in their letter. Our position is that animal tissue use should be limited to necessary applications and should be guided by field experts. The role in medical school curricula should account for freedom of choice in terms of participation, and we should continue towards reducing and ultimately eliminating in vivo SBL once costeffective and efficacious alternatives are established.

## Disclosure

MS/AP are the leads of the ESMSC course; GT has extensive contribution in the eMERG project. The views expressed on this manuscript reflect the interpretation of the authors towards the topic and are based on previous work of the eMERG collaboration. The authors report no other conflicts of interest in this communication.

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Advances in Medical Education and Practice 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Advances in Medical Education and Practice editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

#### **Advances in Medical Education and Practice**

### **Dove**press

#### Publish your work in this journal

Advances in Medical Education and Practice is an international, peerreviewed, open access journal that aims to present and publish research on Medical Education covering medical, dental, nursing and allied health care professional education. The journal covers undergraduate education, postgraduate training and continuing medical education

including emerging trends and innovative models linking education, research, and health care services. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: http://www.dovepress.com/advances-in-medical-education-and-practice-journal

https://doi.org/10.2147/AMEP.S342865