Pulmonary Dysfunction Augmenting Bacterial Aerosols in Leather Tanneries of Punjab, Pakistan

Fiza Sarwar¹
Khan Alam²
Chung Wai Chow³
Muhammad Saeed¹
Riffat Naseem Malik¹

¹Environmental Biology and Ecotoxicology Laboratory, Department of Environmental Sciences, Quaid-i-Azam University, Islamabad, Pakistan;

²Department of Physics, University of Peshawar, Peshawar, Pakistan; ³Dalla Lana School of Public Health, Faculty of Medicine, University of Toronto, Toronto, Canada

Background: Particulate matter–associated microbes in the workplace are a burning issue in occupational toxicology. Studies have reported on respiratory infections among tannery cohorts. This study uniquely presents measurements of airborne bacterial concentrations associated with varied particulate-matter sizes, their exposure, and consequent severity in occupational respiratory problems, all for different microenvironments within leather tanneries.

Methods: Analyses included molecular identification of isolates, computation of mass median aerodynamic diameter of aerosols, tannery process–exposure dose (TPED) to bacterial aerosols, and spirometry and symptom assessment of impaired pulmonary function.

Results: The highest bacterial concentrations were for rawhide treatment and finishing units, showing 3.6×10^3 and 3.7×10^3 CFU/m³, respectively. Identified bacterial species included *Ochrobactrum pseudogrignonense*, *Neisseria bacilliformis*, *Enterobacter cloacae*, *Alcaligenes faecalis*, *Klebsiella pneumoniae*, and *Corynebacterium* spp. Maximum and minimum values of mass median aerodynamic diameter were 8.3 μ m and 0.65 μ m for buffing and snuffing and production units, respectively. The highest TPED was 1,516.9 CFU/kg for finishing units. Respiratory symptoms in order of incidence were dyspnea > phlegm > cough > wheezing and tachypnea (equivalent). Bronchodilator measurements of FEV₁, FVC, and PEF represent decline in lung function. Of 26 patients identified with COPD, most were working in rawhide treatment.

Conclusion: We conclude that exposure–infection synergy is also a cause of pulmonary ailments and COPD development, rather than the better-known exposure–smoking synergy.

Keywords: respiratory symptoms, bacteria, lung function, COPD, tannery process, MMAD, infection, airborne

Plain Language Summary

Industrial exposure to particulate matter (PM) is a notorious source of occupationally induced respiratory ailments. In the interests of finding the synergistic effect of this chemical multimix PM with work-related microorganisms, this study presents interesting data on leather tanneries, showing a strong relationship of respiratory pathogenic bacterial in inhaled PM with enhancement of occupational respiratory disorders (eg, COPD and decline in FEV₁ and FVC) and symptoms (dyspnea, phlegm, cough, wheeze and tachypnea) that exposed cohorts generally face even without bacterial infection. Further, instead of only airborne concentrations, mass median aerodynamic diameter of aerosols and tannery process–exposure doses are presented for different process sites within tanneries to identify the most risk-prone. A majority of identified bacteria in PM were opportunistic pulmonary pathogens, the origin of which may be attributed to hides and skins under treatment and indoor environmental conditions in the workplace.

Correspondence: Fiza Sarwar; Riffat Naseem Malik
Environmental Biology and Ecotoxicology Laboratory, Department of Environmental Sciences, Quaid-i-Azam University, Islamabad, Pakistan
Email fiza.sarwar.baloch@gmail.com; r_n_malik@qau.edu.pk

Graphical Abstract

Introduction

Organic dust originates from plants, animals, and microbes.¹ In workplaces, dispersed dust aerosols containing bacteria, viruses, and fungus spores located in the breathing zone of workers pose a particular danger to health.² The distinct impacts of inhaled bioaerosols and particulates are known to cause respiratory ailments. However, synergistic impacts include exacerbation of pulmonary diseases,³ irritation of mucous membrane, organic dust toxic syndrome, asthma, allergic rhinitis and alveolitis, and COPD.^{4,5} Bacteria can be observed in many occupational environments. Bacterial pathogens are classified as fine biological particles within indoor dust. Ultrafine particles do not include bacteria, because a single bacterium is more than a micrometer in diameter. Nevertheless, bacteria-derived substances like lipopolysaccharide are common in airborne ultrafine particles,⁶ comprising almost

5%–34% of indoor air pollutants.⁷ Gram-negative bacteria have lipopolysaccharide in cell walls that play a direct role in development of disease.⁸ Exposure to lipopolysaccharide and extracellular vesicles causes the creation of proinflammatory and immunomodulating factors.⁶ An invasion of neutrophils into the lower respiratory tract is a representation of this response, accompanied by production of inflammatory intermediaries. However, the inflammation declines after recurring exposure.⁹

A majority of occupational respiratory disorders are caused by dusts and gases, but many pulmonary infections have also been reported lately, eg, acquired by inhaling physical vectors like airborne particulates or droplets.¹⁰ A measure of bacterial concentration in an environment is not the only determinant of a pathogen's potential to induce problems among exposed personnel. In this regard, particle-size distribution and the bacterial concentration

that is bioavailable to reach target tissue in human body is important. The quantity of pollutants actually invading through tissue by crossing biological barriers is called the dose. Occupational exposure limits (OELs) for bioaerosol doses have not been established yet,¹¹ for there exists great variation in workplace characteristics (ventilation type, open or closed workstations), individual vulnerability, and personal exposure levels.¹²

In leather tanneries, bioaerosols come primarily from skin dehairing and defleshing processes. Organic particulates originating from animals (skin, hair, fur) are irritants of the upper respiratory tract, eyes, and skin. Small fractions are capable of reaching the lower respiratory tract and may cause diseases like bronchitis, asthma, conjunctivitis, and dermatitis,¹³ which have been reported among tannery employees. Animal hides facilitate growth of bacteria due to rich nutrients and high moisture content. Indoor air during tanning processes is associated with spread of microbes that lead to occupationally induced infections in employees. In buffing dust, fungal species have also been reported,¹⁴ which is also a major problem in hide-preservation units. In tanneries, major sources include fragments and fiber remnants from animal fur, skin, leather, brooms, protein-bound buffing dust, and organic tanning emissions. Animal skin provides an ideal medium for the propagation of microorganisms in both the leather-tannery and footwear industries.¹⁵ It has been found that the prevalence of symptoms suggestive of asthma, tuberculosis, and chronic bronchitis among tannery workers is higher than non-tannery workers.¹⁶ These symptoms include wheezing, obstructive lung disease, tightness of chest, difficulty in breathing, cough, and breathlessness after a walk.¹⁷ Increasing evidence suggests that the lung microbiome plays an important role in COPD severity. COPD is a pulmonary ailment that primarily develops with age, but environmental exposure and genetics are also considered significant disease-inducing agents.¹⁸

Acute exacerbations of COPD are sudden worsening of symptoms in which bacterial colonization is one major etiological factor.¹⁹ Bacteria have been isolated from the lower airways of stable COPD patients, and airway inflammation has been related to bacterial load and type. Bacterial colonization in COPD is an important factor in disease progression.²⁰ COPD is defined by the spirometric definition of postbronchodilator $FEV_1/FVC < 0.7$.²¹ When calculating FEV_1/FVC , the highest value of each after reversibility testing is used, as per ERS taskforce guidelines for epidemiological studies on COPD²² and supported by GOLD guidelines.²³ The second definition ($FEV_1/FVC < LLN$)

is not used, because compared to the first ($FEV_1/FVC < 0.7$), no substantial connection has been found with this definition when respiratory assessments for exposure to biological dust, mineral dust, gases, and fumes were made.²⁴

It is well proven that bacteria play an important role in pulmonary obstructions and infections. There exists a gap in the available research regarding process-wise identification of respiratory pathogenic bacterial strains and consequent disorders at leather tanneries. Further, the available literature focuses on airborne bacterial concentrations in occupational environments, but little has been reported on bacterial mass or particle-size association.⁴ We hypothesized that tannery workers engaged in areas of high-level bioaerosol emissions would experience exacerbated particulate matter (PM)-associated pulmonary dysfunctions, as respiratory pathogens might play a synergistic role. Therefore, we aimed to gauge the presence of such bacteria in a variety of tannery environments by measuring viable culturable counts of bacteria in ambient air, the mass median aerodynamic diameter (MMAD) of bacterial aerosols, and the tannery process-exposure dose (TPED) of bacterial aerosols, and assess pulmonary function among tannery workers, along with pulmonary symptoms and enhancement during bacterial infections, and identify COPD patients among employees for a number of tannery microenvironments with different processes in progress.

Methods

Leather tanneries in Pakistan are not complete industrial units. Therefore, different processes are found in different factories. These processes are classified in groups on the basis of their being performed under one roof and thus environmental samples not being segregated. The process clusters identified for this study are shown in Table 1, representing a variety of microenvironments with different occupational exposure. Measurements were made in summer (July–September). Though the season was not at its peak, all routine processes were in progress. Representative samples were taken from leather tanneries of Lahore, Sialkot, and Kasoor, all renowned tannery hubs in Punjab.

Bacterial Analysis

Air Sampling of Particulate Matter–Associated Bacteria

Sampling was carried out using a portable 11.B aerosol spectrometer (Grimm Aerosol Technik Ainring). This

Table 1 Tannery-site characteristics

	Processes	Type
Rawhide treatment	Soaking, liming, dehairing, defleshing, deliming, bating, and pickling	Sheltered
Wet machine operations	Chrome tanning, wet-blue storage, wet finishing (wet back, neutralization, retanning, washing, fat liquoring, dyeing, and washing)	Indoor
Buffing and snuffing	Buffing and snuffing	Indoor
Splitting and shaving	Splitting and shaving	Sheltered
Production unit	—	Indoor
Finishing unit	Hydraulic press, roller cutting, spray plant	Indoor
Packing and sales	Weighing, finished leather trim, sorting, and packing	Indoor

instrument has 31 size channels ranging from 0.25 to 32 μm . Its working principle is light scattering of a single particle with a semiconductor laser as a light source of 638 nm. Instead of filters, a greased glass slide was used as an impactor that collected aerosols. Samples were collected at a flow rate of 1.2 L/min. The sampler operated for 8 hours/day for 3 days at each site to collect triplicate samples. These glass-slide samples for each process under investigation were taken to the laboratory using sterile isolated containers.²⁵ The collected samples were incubated at 30°C, and colony-forming units (CFUs) were counted after 24 hours. Tryptic soy agar (Sigma-Aldrich) with 0.2% nystatin was used for collection of bacteria. Based on colony morphology, mixed-culture plates were then subcultured to obtain pure colonies.

Molecular Identification of Isolates

Confirmation of the isolated bacterial strains was done by molecular characterization through 16S rRNA gene sequencing. Genomic DNA of all bacterial strains was obtained using a DNA-extraction kit (Qiagen). Concentrations of DNA were qualified and quantified by spectrometry using a NanoDrop 2000 (Thermo Fisher Scientific). PCR amplification of extracted DNA was done using universal bacterial primers 27F' (5'-AGAGTTTGGATCTGGCTCAG-3') and 1492R' (5'-CTACGGCTACCTTGTACGA-3'). The PCR mixture

consisted of GoTaq green master mix (25 μL), primer 27F' (1 μL), primer 1492R' (1 μL), DNA extract (2 μL), and nuclease-free water (50 μL). An MJ Mini personal thermal cycler was used for the PCR reaction (Bio-Rad). Cycling conditions consisted of initial denaturation at 95°C for 3 minutes followed by 30 cycles of denaturation at 95°C for 1 minute, annealing at 55°C for 1 minute, and extension at 72°C for 1 minute. At 72°C, the final extension was performed for 7 minutes. Electrophoresis was done for PCR-product analysis with 1% agarose gel. Sequencing of the amplified PCR product was done by Macrogen Service Center (Seoul, South Korea).

MMAD and TPED

The mass held in each particle size defines a particle population. The MMAD basically splits the distribution of the mass in half. During respiration, there is maximum chance of particles $\leq 3 \mu\text{m}$ reaching the lower respiratory tract of an adult.²⁶

MMAD was calculated using:

$$MMAD = \sum m_i / M \quad 1$$

where m_i is the mass of all particles in one group i (where d is the midpoint diameter).

The TPED was measured for indoor and semi-indoor environments (CFU/kg).¹²

$$TPED = C \cdot IR \cdot IEF / BW \quad 2$$

where C is the bacterial aerosol concentration (CFU/ m^3), IEF the indoor exposure fraction (hours spent per day at work considering assorted activity patterns), IR the inhalation-rate coefficient characteristic of the selected activity levels (m^3/day),²⁷ and BW mean body weight (kg).

Pulmonary Assessment of Workforce Inclusion and Exclusion Criteria

A total of 250 male workers aged 18–60 years engaged at tannery works for at least 5 years in the same microenvironment were recruited as subjects. Those diagnosed with respiratory or other ailments like anemia, abnormal vertebral column or thoracic cage, diabetes, respiratory tuberculosis, asthma, and bronchitis and those severely ill were excluded from the study. Further the drug buffs, tobacco consumers were also omitted from the study. Smokers and nonsmokers were included and compared, as cigarette smoking is considered an important factor in the development of pulmonary ailments and occupational COPD.²⁸

Spirometry

Spirometry was performed in leather tanneries in Lahore, Kasur and Sialkot.

A handheld digital spirometer (SP10W, Contec) was used to record FEV₁, FVC, and FEV₁% (FEV₁/FVC), both pre- and postbronchodilation. For validation and accuracy of testing, equipment quality control was carried out with regular calibration checks before testing was performed, according to American Thoracic Society recommendations.²⁹ Bronchodilation was done with a combination of 0.4 mg salbutamol and 80 µg ipratropium bromide.²³ FEV₁ and FVC >80% is considered normal, FEV₁ and FVC <80% but FEV₁/FVC >0.7 represents restriction, and FEV₁ <80% with reduced FVC shows obstruction.³⁰ Subjects with COPD were classified into grades 1–4 according to GOLD.³¹ The GOLD classification of airflow-limitation severity in COPD (based on postbronchodilator FEV₁) is 1 (mild) = FEV₁ ≥ 80% predicted 2 (moderate) = 50% ≤ FEV₁ < 80% predicted, 3 (severe) 30% ≤ FEV₁ < 50% predicted, and 4 (very severe) FEV₁ < 30% predicted.

Pulmonary Symptom Assessment

A standard pulmonary symptom-identification questionnaire was filled in by all subjects. Symptoms included cough, phlegm, chest troubles, tachypnea, dyspnea, and wheezing.³² It was also modified to include if these symptoms were associated with routine exposure and/or enhanced with chest bacterial infections. Division of 8 working hours was also determined based on activity level via a walk-through survey for each process.

Statistical Analysis

Two sample *t*-tests were used to find statistical contrasts in pulmonary symptoms when conditions shifted from only dust-exposure days to the days when working cohorts experience increased respiratory infection with occupational dust exposure. Multiple linear (forward) regression models were used to identify predictor variables for cough, wheezing, tachypnea, dyspnea, and phlegm. For the chosen covariates, *P* < 0.1 was utilized to locate variables for the best-fit model to explain the highest variations in dependent variables. Differences were considered statistically significant at *P* < 0.05. SPSS 21.0 was employed for statistical analysis.

Results and Discussion

Bacteria in Tannery Dust

Tanneries in Pakistan perform different operations, ie, a complete chain of processes is not found in a single factory. That is why processes at all locations were clustered and considered as one workplace to collect data. Table 2 presents the bacteria identified along with reported characteristics from the literature and pulmonary pathogenicity.

Ten strains were observed, of which one was uncultured. *Noviherbaspirillum autotrophicum* is found in air of three units: wet operations, rawhide treatment, and splitting and shaving. This facultatively autotrophic bacterium was reportedly isolated from rice-paddy soil.³³

Alcaligenes spp. were found where splitting and shaving of hides and skins was in progress. These drug-resistant respiratory pathogens^{34,35} may worsen pulmonary infections. Compromised respiratory tracts become victims of this opportunistic pathogen. Three strains of this family were identified, of which two were *Alcaligene faecalis*. *Ochrobactrum pseudogrignonense* was isolated from aerosols collected from rawhide treatment, spraying and finishing, and packing and production units. This is a known human respiratory pathogen that has frequently been found in patients with cystic fibrosis.³⁶ This emergent pathogen is equally treacherously infectious for both immunocompromised and healthy humans.³⁷ *Neisseria bacilliformis* is also an opportunistic pathogen that causes infection in the upper respiratory tract.³⁸ In this study, it was found in skin-splitting and -shaving and leather-finishing units. *Enterobacter cloacae* is a Gram-negative anaerobic facultative bacteria that causes human infections of the respiratory tract, urinary tract, skin, and abdomen.³⁹ The basic source for its emergence in the tannery environment is the skin and hides of animals. It was found in the air of a finishing unit. *Klebsiella pneumoniae* is specifically a respiratory and urinary tract infectant; however, it may cause infection in any part of the body. It colonizes and multiplies in the gastrointestinal tract, and a majority of infections arise when bacteria move from the here.⁴⁰ *Corynebacterium* spp. are multidrug-resistant bacteria, of which the first epidemic was reported in of COPD patients in 2007. These organisms cause infection that worsens existing respiratory disorders.⁴¹ *Providencia rettgeri*, isolated from the air of a finishing unit is not a respiratory infectant, but causes urinary tract and skin infections in humans.⁴²

Table 2 Human pulmonary bacterial pathogens isolated from tannery air

	Family	Characteristics	Pulmonary symptoms from literature	References
<i>Noviherbaspirillum autotrophicum</i>	Oxalobacteraceae	Facultatively autotrophic bacterium isolated from rice paddy soil	NA	Ishii et al ³³
<i>Alcaligenes</i> spp.	Alcaligenaceae	Opportunistic respiratory pathogens	Adult respiratory distress syndrome Sepsis syndrome Cystic fibrosis	Aisenberg et al ³⁴
<i>Ochrobactrum pseudogrignonense</i>		Opportunistic respiratory pathogens No cross-contamination	Cystic fibrosis Respiratory failure Upper and lower respiratory tract infections	Lundy et al, Chmelař, Holý et al ^{36,64}
<i>Neisseria bacilliformis</i>	Neisseriaceae	Opportunistic pathogens	Upper respiratory tract infections	Han et al ³⁸
<i>Enterobacter cloacae</i>	Enterobacteriaceae		COPD Nosocomial pneumonia Lower respiratory tract pneumonia	Didilescu et al, Miyoshi-Akiyama et al ^{49,50}
<i>Alcaligenes faecalis</i> (two strains)	Alcaligenaceae	Pan-drug-resistant organism even to carbapenems, polymyxins, and tigecycline Respiratory dysfunction synergist	Respiratory depression (reduced SPO ₂) Bloodstream infections Acute respiratory distress syndrome	Kavuncuoglu et al, Hasan et al ^{35,63}
<i>Klebsiella pneumoniae</i>	Enterobacteriaceae	Opportunistic pathogen	Lung and urinary tract infection Gastrointestinal diseases	Kaur et al ⁶²
<i>Corynebacterium</i> spp.		Animal-skin flora Human pathogen	Respiratory infections	Díez-Aguilar et al ⁴¹
<i>Providencia rettgeri</i>	Enterobacteriaceae	—	Urinary tract infections, bacteremia, skin infections, diarrhea, and gastroenteritis	Yoh et al, Tada et al ^{42,61}

Four strains identified in this study have also been isolated from oropharyngeal mucosa of tannery workers in Mexico: *Alcaligenes faecalis*, *Neisseria bacilliformis*, *Enterobacter cloacae*, and *Klebsiella pneumoniae*,¹⁵ which confirms the association of these bacteria with the tannery environment. Table 3 gives a checklist of identified bacteria at different leather tanneries.

MMAD and TPED

MMAD values of the particle populations observed at defined sites are shown in Table 4. The maximum MMAD was measured for B&S — 8.3 µm. This means

50% of the total sample mass is present in particles with aerodynamic diameter <8.3 µm and the other 50% in particles >8.3 µm in diameter.⁴³ The minimum was 0.65 µm, for PU. Commonly, bioaerosol size is 0.02–100 µm,⁴⁴ but their association with particulates changes the aerodynamic diameter of PM,³ affecting their inhalation characteristics. The atmosphere shows a chemistry of two contrasting mechanisms in summer and spring. Bacteria grow quickly, but the particle-size range (fine) that is associated with bacterial mass tends to exterminate. Therefore, there are fewer fine particles (<3.3 µm) in total particulates.⁴⁵ The TPED results are also shown in

Table 3 Site-wise bacteria identified

	RHT	WMO	PU	FU-S	FU-HP	B&S	S&S	P&S
<i>Noviherbaspirillum autotrophicum</i>	x	x	-	-	-	-	x	-
<i>Alcaligenes</i> spp.	x	-	-	-	-	x	x	-
<i>Ochrobactrum pseudogrignonense</i>	x	-	x	x	-	x	-	x
<i>Neisseria bacilliformis</i>		-	-	x	x	x	x	x
<i>Enterobacter cloacae</i>	x	x	-	x	x	x	x	x
<i>Alcaligenes faecalis</i> (two strains)	x	-	-	x	x	-	x	x
<i>Klebsiella pneumoniae</i>	x	x	x	x	x	-	x	x
<i>Corynebacterium</i> spp.	x	-	-	x	x	x	x	x
<i>Providencia rettgeri</i>	x	-	-	x	x	-	x	x
Uncultured bacteria	x	-	-	-	-	-		-

Notes: x, presence; –, absence.

Table 4. Maximum TPED was 1,516.9 CFU/kg for FU-HP and the minimum was found for WMO — 92.78 CFU/kg. One reason for exposure limits not yet being established for bioaerosols of occupational origin is that individual exposure characteristics and vulnerabilities are exceedingly varied.¹²

Pulmonary Function

The process-wise distribution of the 250 tannery workers was PS 55, RHT 48, WMO 37, BS+SS 64, and FU 46. They were all male Muslims and residents of the areas where their workplaces were located. Two-thirds (66%) were married. Their awareness of occupational safety was medium (51%), low (45%), and high (4%). Demographic characteristics are shown in [Table 5](#). All were working for 8 hours a day.

Spirometry and COPD

Average process-based FEV₁, FVC, and PEF are shown in [Figure 1](#). There were clear differences in between pre- and postbronchodilation. However, the most compromised

lung function was found in BS+SS personnel, because of being exposed to buffing dust, toggling dust in the hang-dry process, and shaving and splitting producing fine and ultrafine dust. Their prebronchodilation FEV₁ was 2.10 ±0.667. However, considerable improvement was observed after bronchodilation — 2.62±0.492. Similarly, prebronchodilation FVC was 2.48±0.746 and became 2.96 ±0.951 after the reversibility test had been applied. This response to medication in the form of improvement in FEV₁, FVC, and PEF gives an indication of airway

Table 5 Demographic characteristics of participating workforce (n=250)

Parameter	Tannery workers N=250
Age (years)	35±10.55
BMI	23.73±4.77
Body weight (kg)	70±5.6
Smokers	28.4%
Nonsmokers	71.6%

Table 4 Bacterial concentrations, MMAD, and TPED

	RHT	WMO	PU	FU-S	FU-HP	B&S	S&S	P&S
Bacterial concentration (CFU/m ³)	3.6×10 ³	2.3×10 ²	2.6×10 ²	3.2×10 ³	3.7×10 ³	3.2×10 ³	2.8×10 ³	2.2×10 ³
Tolerable concentrations of airborne microbes in occupational environment: Mesophilic bacteria 100,000 and Gram-negative bacteria 20,000 Suggested standards for respirable aerosols are half those, ie, 50,000 CFU/m ³ for mesophilic bacteria and 10,000 CFU/m ³ for Gram-negative bacteria. ⁶⁵								
MMAD (μm)	7.740	0.741	0.659	7.199	7.050	8.351	7.960	7.265
TPED (CFU/kg)	1,464.4	92.78	104.9	1,295	1,516.9	1,299	1,130	923.9

Notes: Sheltered: (top covered corridor without walls); average work hours 8; average employee weight 70 kg.

Abbreviation: MMAD, median mass aerodynamic diameter; TPED, tannery process–exposure dose.

Figure 1 Process-based comparison of average pulmonary function in tannery workers. (A) PrBD and PoBD forced expiratory volume in one second (FEV₁); (B) PrBD and PoBD peak expiratory flow (PEF); (C) PrBD and PoBD forced vital capacity (FVC).

Abbreviations: PrBD, pre bronchodilation; PoBD, post bronchodilation.

obstruction.⁴⁶ These results are parallel to those reported in the literature.^{9,17,46–48} Moreover, these are the processes where respiratory pathogenic bacteria have also been recognized, and surveyed pulmonary symptoms back this finding as well.

Although highly severe cases of COPD were not identified, for those suffering from COPD, GOLD stages were

determined (Figure 2). Two COPD patients were identified in the B&S group, of which one (nonsmoker) was stage 1 and the other (smoker) stage 2. Similarly identified cases of COPD stages were shown for all sampled processes. Most were found in RHT (ten), including six smokers and two nonsmokers. A peculiar finding was one with stage 3 COPD being a nonsmoker. The overall trend was RHT (ten) > SS (5) > PU (4) > WMO, P&S, and BS (two each) > FU (one) > HD (none). The highest stage of COPD was identified as stage 3 of which four worked in RHT and two in S&S. These two environments presented the most COPD patients. It was also observed that the highest bacteria numbers, including pulmonary pathogens,^{49,50} were found in S&S and RHT units. Since cigarette smoking is a known COPD-exacerbating agent, distribution of COPD patients was determined using this factor. An important aspect to be noted is that the COPD patients identified were not necessarily smokers. It may be because cigarette smoking is not the only risk factor of pulmonary obstructions.^{51,52} This ailment may grow among nonsmokers due to their exposure to dust and infectious agents. Nonsmokers can also develop chronic airflow obstruction, and smokers show substantially inconsistent vulnerability when it comes to symptoms and signs of pulmonary obstruction,⁵³ which may be attributed to recurring exposure to a variety of tannery dusts along with pulmonary obstruction–aggravating bacteria. In a recent study on causative agents of COPD exacerbation, 55 noninfectious, 33 bacterial, and 12 bacterial agents were found. Of 33, nine patients had *Klebsiella pneumoniae*,⁵⁴ which was identified in PM sampled from all processes, except for B&S.

Bacterium-Induced Pulmonary Symptom Augmentation

All the tannery working groups showed respiratory symptoms with considerably high percentages, as shown in Table 6 (dyspnea > phlegm > cough > wheezing and tachypnea [equivalent]). These symptoms are induced occupationally by prolonged exposure to chemical-laden particulates emitted from different tannery processes.^{9,55–57} Respirational indicators like chest pain, cough, dyspnea, cough, and phlegm among tannery workers are symptoms of pulmonary dysfunction.⁵⁸ It was quite observable that among recorded respiratory symptoms, a majority were enhanced when exposure is synergized with chest infection. This supports that dust-induced occupational respiratory symptoms being boosted in coexistence with respiratory infection causing bacteria. Both chronic and developing pulmonary obstructions are intensified by

Figure 2 Stagewise (GOLD) distribution of COPD patients among tannery-worker cohorts: difference for smokers and nonsmokers. 0, no disease; 1, GOLD stage 1; 2, GOLD stage 2; 3, GOLD stage 3.

bronchial infections, and result in acute increases in symptoms like cough, dyspnea, and phlegm.⁵⁹ Further, upon bacterial respiratory infection, differences in symptoms between smokers and nonsmokers were recorded. The results showed increased magnitude of symptoms among nonsmokers (cough 24.8%, phlegm 18.0%, wheeze 15.2%, dyspnea 14.0%) compared to smokers (cough 1.6%, phlegm 6.4%, wheeze 5.2%, dyspnea 5.6%). Studies have reported symptoms among tannery workers of dry/productive cough, throat irritation, dyspnea, and chest troubles.^{9,60} So far, these symptoms have been reported without work/process specification or evaluation of exposure-induced symptoms vs symptoms synergized by bacterial infections. This study presents these assessments (Table 6A), where (except for tachypnea) differences in symptoms between two investigated conditions was statistically apparent. Further, it was identified that no independent variables, including process and smoking, had a significant impact on cough (only exposure) and wheeze (only exposure). The forward-regression model identified significant contributors to each pulmonary symptom, as presented in Table 6B. During exposure and infection, cough was found to be highly influenced by presence of bacteria and bacteria numbers ($R^2=0.023$, $P=0.01$). A similar trend was observed for the other pulmonary parameters. However, for tachypnea, two variables (department types and bacteria numbers) were found to have an influence on exposure to infection conditions ($R^2=0.033$, $P=0.002$). From these models, it can be inferred that the presence of bacteria was the highest contributor in enhancing respiratory symptoms like coughing, wheezing, and dyspnea.

This study is a pioneer in determining the synergy of occupational exposure of PM and bacterial pathogens in aggravating respiratory conditions in different microenvironments of leather tanneries. Implications of this study may be useful for studying synergy with chemical characteristics of PM. Further, the inclusion of both sexes as study subjects may widen the horizon of the subject matter.

Conclusion

From different microenvironments of leather tanneries, human respiratory pathogenic bacteria were associated with inhalable PM. According to MMAD computation, 50% bacterial mass was associated with average PM size $>7\ \mu\text{m}$, except for WMO and PU, where this association was also substantial with PM $<0.5\ \mu\text{m}$. Some areas present alarming airborne concentrations and TPED. Our data show that work-environment exposure along with chest infections enhance respiratory symptoms manyfold. This points to the synergy of opportunistic respiratory pathogens with exposure to incidental PM at work. Obstructive variations on pulmonary functions FEV₁, FVC, and PEF and identification of COPD patients with grade 3 COPD also prove the compromised lung function of tannery workers. Bacterial infections were a stronger synergist with occupationally induced pulmonary symptoms than smoking, signifying the importance of setting up protective measures focusing on respiratory health among exposed workers, especially in departments like rawhide treatment, shaving and splitting, buffing, finishing, and packing.

Data Sharing Statement

All generated or analyzed data in this study are included in this manuscript. Additional information/data will be furnished by the corresponding authors upon request.

Table 6 Statistical evidence of bacterial aerosol–tannery dust synergy enhancing pulmonary dysfunction

Respiratory symptoms among tannery workers (with and without bacterial infections)										
	Cough		Phlegm		Wheeze		Dyspnea		Tachypnea	
	DE	DE + BI	DE	DE + BI	DE	DE + BI	DE	DE + BI	DE	DE + BI
% Age	33.2	58.8	34	55.6	9.6	16.4	34.4	49.6	9.6	10.8
P	<0.0001*		<0.0001*		0.02*		<0.0001*		0.66	
Forward linear regression model: relationship of pulmonary symptoms with selected potential synergy										
			Unstandardized coefficients		Standardized coefficients					
Cough			β	SE	β			t	P	
Constant			0.438	0.65				0	0	
Bacteria numbers			0.30	0.011	0.163			2.604	0.01	
Smoking			−0.43	—	—			−1.094	0.99	
Department			−0.06	—	—			−0.056	0.852	
Model					F=6.781			P=0.010	^0.023	
Phlegm										
Constant			0.287	0.064				4.503	0	
Bacteria numbers			0.053	0.011	0.290			4.779	0	
Smoking			−0.15	—	—			−0.253	0.800	
Department			0.003	—	—			0.051	0.959	
Model					F=22.84			P=0	^0.08	
Wheeze										
Constant			0.067	0.049				1.364	0.174	
Bacteria numbers			0.019	0.009	0.141			2.235	0.026	
Smoking			−0.082	—	—			0.192	0.192	
Department			−0.039	—	—			0.573	0.852	
Model					F=4.966			P=0.026	^0.016	
Dyspnea										
Constant			0.260	0.065				4.009	0	
Bacteria numbers			0.047	0.011	0.253			4.121	0	
Smoking			−0.030	—	—			−4.84	0.629	
Department			−0.126	—	—			−0.120	0.852	
Model					F=16.986			P=0	^0.06	
Tachypnea										
Constant			−0.133	0.068				−0.176	0.860	
Departments			0.035	0.009	0.253			3.778	0	
Bacteria numbers			0.018	0.008	0.156			2.324	0.021	
Smoking			0.038	—	—			0.621	0.535	
Model					F=9.586			0.002	^0.033	

Notes: *P=0.05. ^aAdjusted R². Independent variables: bacteria numbers, smoking status, and department type. P-values in bold indicate maximum influence on pulmonary symptom.

Abbreviations: DE, dust exposure (days); BI, bacterial infection (days).

Ethics

This study was approved by the ethics review board of Quaid-i-Azam University, Islamabad. All participants were informed on the nature, purpose, and likely upshot of the study. Each participant signed a consent form prior to spirometry and questionnaire-based surveillance. The Declaration of Helsinki (revised 2013) was also complied with.

Consent for Publication

Informed consent was taken from all participants.

Author Contributions

All authors contributed to data analysis and drafting or revising the article, have agreed on submitting the article to this journal, gave final approval to the version to be published, and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this study.

References

- Peters S, Kromhout H, Olsson AC, et al. Occupational exposure to organic dust increases lung cancer risk in the general population. *Thorax*. 2012;67(2):111–116.
- Shkrabak V, Popov A, Enikeev V, Gavrikova E, Shkrabak R. Indoor air decontamination system and reduction of microorganism emissions into the atmosphere; Paper presented at: BIO Web of Conferences; 2020.
- Adhikari A, Reponen T, Grinshpun SA, Martuzevicius D, LeMasters G. Correlation of ambient inhalable bioaerosols with particulate matter and ozone: a two-year study. *Environ Pollut*. 2006;140(1):16–28.
- Bragoszewska E, Pawlak M. Health risks associated with occupational exposure to biological air pollutants occurring during the processing of biomass for energy purposes: a case study. *Energies*. 2021;14(8):2086.
- Gólofit-Szymczak, M., & Ławniczek-Walczyk, A. (2011). Biomass as a source of biological threats. *Occupational Safety: Science and Practice*, (12):17–19. Available from: <https://www.infona.pl/resource/bwmeta1.element.baztech-article-BPC1-0011-0009>
- Yang J, Kim EK, Park HJ, McDowell A, Kim Y-K. The impact of bacteria-derived ultrafine dust particles on pulmonary diseases. *Exp Mol Med*. 2020;52:1–10.
- Srikanth P, Sudharsanam S, Steinberg R. Bio-aerosols in indoor environment: composition, health effects and analysis. *Indian J Med Microbiol*. 2008;26(4):302.
- Kim Y-M, Kim Y-S, Jeon SG, Kim Y-K. Immunopathogenesis of allergic asthma: more than the th2 hypothesis. *Allergy Asthma Immunol Res*. 2013;5(4):189–196.
- Işsever H, Özdiilli K, Özyildirim BA, et al. Respiratory problems in tannery workers in Istanbul. *Indoor Built Environ*. 2007;16(2):177–183.
- Ho P, Becker M, Chan-Yeung M. Emerging occupational lung infections [State of the Art Series. Occupational lung disease in high-and low-income countries, edited by M. Chan-Yeung. Number 6 in the series]. *Int J Tuberc Lung Di*. 2007;11(7):710–721.
- Di Filippo P, Pomata D, Riccardi C, Buiarelli F, Perrino C. Fungal contribution to size-segregated aerosol measured through biomarkers. *Atmos Environ*. 2013;64:132–140.
- Bragoszewska E, Biedroń I, Kozielska B, Pastuszka JS. Microbiological indoor air quality in an office building in Gliwice, Poland: analysis of the case study. *Air Qual Atmos Health*. 2018;11(6):729–740.
- Buza C, Pacheco II, Robbie K. Nanomaterials and nanoparticles: sources and toxicity. *Biointerphases*. 2007;2(4):MR17–MR71.
- Emmanuel S, Adamu I, Ejila A, Ja'afaru M, Yabaya A, Habila B. Characterization of chrome buffing dust (CBD) generated from NILEST tannery associated with pathogenic fungi. *J Toxicol Environ Health Sci*. 2014;6(3):89–98.
- Castellanos-Arévalo DC, Castellanos-Arévalo AP, Camarena-Pozos DA, Colli-Mull JG, Maldonado-Vega M. Evaluation of microbial load in oropharyngeal mucosa from tannery workers. *Saf Health Work*. 2015;6(1):62–70.
- Kashyap GC, Sharma SK, Singh SK. Prevalence and predictors of asthma, tuberculosis and chronic bronchitis among male tannery workers: a study of Kanpur City, India. *Clin Epidemiol Glob Health*. 2020;9:71–77.
- Jamal A, Mehmood A, Putus T, Savolainen H, Liesivuori J. Prevalence of respiratory symptoms, bronchial asthma and obstructive lung disease among tannery workers. *Peertechz J Environ Sci Toxicol*. 2017;2(1):33–42.
- Martinez CH, Mannino DM, Divo MJ. Defining COPD-related comorbidities, 2004–2014. *J COPD Foundation*. 2014;1(1):51.
- Wang Z, Bafadhel M, Haldar K, et al. Lung microbiome dynamics in COPD exacerbations. *Eur Respir J*. 2016;47(4):1082–1092.
- Wilkinson TM, Patel IS, Wilks M, Donaldson GC, Wedzicha JA. Airway bacterial load and FEV1 decline in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2003;167(8):1090–1095.
- Song JH, Lee C-H, Kim DK, et al. Differences in prevalence of asthma–COPD overlap according to different criteria. *Medicine*. 2018;97(36):e12049.
- Bakke P, Rönmark E, Eagan T, et al. Recommendations for epidemiological studies on COPD. *Eur Respir Soc*. 2011;38:1261–1277.
- Hagstad S, Backman H, Bjerg A, et al. Prevalence and risk factors of COPD among never-smokers in two areas of Sweden—occupational exposure to gas, dust or fumes is an important risk factor. *Respir Med*. 2015;109(11):1439–1445.
- Alif SM, Dharmage SC, Bowatte G, et al. Occupational exposure and risk of chronic obstructive pulmonary disease: a systematic review and meta-analysis. *Expert Rev Respir Med*. 2016;10(8):861–872.
- Pathak B, Borah D, Khataniar A, Bhuyan P, Buragohain A. Characterization of bioaerosols in Northeast India in terms of culturable biological entities along with inhalable, thoracic and alveolar particles. *J Earth Syst Sci*. 2020;129(1):1–13.
- Dolovich MB, Dhand R. Aerosol drug delivery: developments in device design and clinical use. *Lancet*. 2011;377(9770):1032–1045.
- Health USEPAOo, Group EAEA. *Exposure Factors Handbook*. Vol. 90. Office of Health and Environmental Assessment, US Environmental Protection; 1989.
- Bepko J, Mansalis K. Common occupational disorders: asthma, COPD, dermatitis, and musculoskeletal disorders. *Am Fam Physician*. 2016;93(12):1000–1006.
- Miller M, Hankinson J, Brusasco V, et al. Standardization of spirometry, 1994 update. American thoracic society. *Am J Respir Crit Care Med*. 1995;152(3):1107–1136.
- Saleem I, Ali Z, Hussain H. Respiratory health implications among wet-blue leather tannery workers of Kasur. *Nat Sci*. 2021;20(3):e2021058.
- Marçôa R, Rodrigues DM, Dias M, et al. Classification of chronic obstructive pulmonary disease (COPD) according to the new Global Initiative for Chronic Obstructive Lung Disease (GOLD) 2017: comparison with GOLD 2011. *J Chron Obstruct Pulmon Dis*. 2018;15(1):21–26.

32. Ferris B. Epidemiology standardization project. II. Recommended respiratory disease questionnaires for use with adults and children in epidemiological research. *Am Rev Respir Dis.* 1978;118(6):7–53.
33. Ishii S, Ashida N, Ohno H, et al. *Noviherbaspirillum denitrificans* sp. nov., a denitrifying bacterium isolated from rice paddy soil and *Noviherbaspirillum autotrophicum* sp. nov., a denitrifying, facultatively autotrophic bacterium isolated from rice paddy soil and proposal to reclassify *Herbaspirillum massiliense* as *Noviherbaspirillum massiliense* comb. nov. *Int J Syst Evol Microbiol.* 2017;67(6):1841–1848.
34. Aisenberg G, Rolston KV, Safdar A. Bacteremia caused by *Achromobacter* and *Alcaligenes* species in 46 patients with cancer (1989–2003). *Cancer.* 2004;101(9):2134–2140.
35. Kavuncuoglu F, Unal A, Oguzhan N, Tokgoz B, Oymak O, Utas C. First reported case of *Alcaligenes faecalis* peritonitis. *Peritoneal Dial Int.* 2010;30(1):118–119.
36. Lundy F, Tunney M, Taggart C, Elborn J. Human Neutrophil Peptide-1 (HNP-1) and a truncated analogue display antimicrobial activity against respiratory pathogens. *J Cystic Fibros.* 2009;8:S47.
37. Cho HW, Byun J-H, Kim D, Lee H, Yong D, Lee KW. The first case of *Ochrobactrum pseudogrignonense* Bacteremia in Korea. *Ann Lab Med.* 2020;40(4):331–333.
38. Han XY, Hong T, Falsen E. *Neisseria bacilliformis* sp. nov. isolated from human infections. *J Clin Microbiol.* 2006;44(2):474–479.
39. Mezzatesta ML, Gona F, Stefani S. *Enterobacter cloacae* complex: clinical impact and emerging antibiotic resistance. *Future Microbiol.* 2012;7(7):887–902.
40. Struve C, Krogfelt KA. Pathogenic potential of environmental *Klebsiella pneumoniae* isolates. *Environ Microbiol.* 2004;6(6):584–590.
41. Díez-Aguilar M, Ruiz-Garbajosa P, Fernández-Olmos A, et al. Non-diphtheriae *Corynebacterium* species: an emerging respiratory pathogen. *Eur J Clin Microbiol Infect Dis.* 2013;32(6):769–772.
42. Tada T, Miyoshi-Akiyama T, Dahal RK, et al. NDM-1 Metallo- β -Lactamase and *ArmA* 16S rRNA methylase producing *Providencia rettgeri* clinical isolates in Nepal. *BMC Infect Dis.* 2014;14(1):56.
43. Jiménez AS, van Tongeren M, Aitken RJ. Guidance for collection of inhalable and respirable Ni dust. *IOM Rep.* 2012;538. Available from: https://www.nickelconsortia.eu/assets/files/library/Guidances/IOM%20Report%201_inhalation%20and%20respirably%20monitoring%20guidance%20for%20Ni_March%202012.pdf.
44. Raisi L, Lazaridis M, Katsivela E. Relationship between airborne microbial and particulate matter concentrations in the ambient air at a Mediterranean site. *Glob NEST J.* 2010;12(1):84–91.
45. Brągoszewska E, Mainka A, Pastuszka JS. Concentration and size distribution of culturable bacteria in ambient air during spring and winter in Gliwice: a typical urban area. *Atmosphere.* 2017;8(12):239.
46. Chandrasekaran V, Dilara K, Padmavathi R. Pulmonary functions in tannery workers—a cross sectional study. *Indian J Physiol Pharmacol.* 2014;58(3):206–210. PMID: 25906602.
47. Shahzad K, Akhtar S, Mahmud S. Prevalence and determinants of asthma in adult male leather tannery workers in Karachi, Pakistan: a cross sectional study. *BMC Public Health.* 2006;6(1):292.
48. Were FH, Moturi MC, Wafula GA. Chromium exposure and related health effects among tannery workers in Kenya. *J Health Pollut.* 2014;4(7):25–35.
49. Miyoshi-Akiyama T, Hayakawa K, Ohmagari N, Shimojima M, Kirikae T. Multilocus sequence typing (MLST) for characterization of *Enterobacter cloacae*. *PLoS One.* 2013;8(6):e66358.
50. Didilescu AC, Skaug N, Marica C, Didilescu C. Respiratory pathogens in dental plaque of hospitalized patients with chronic lung diseases. *Clin Oral Invest.* 2005;9(3):141–147.
51. Salvi SS, Barnes PJ. Chronic obstructive pulmonary disease in non-smokers. *Lancet.* 2009;374(9691):733–743.
52. Vogelmeier CF, Criner GJ, Martinez FJ, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive lung disease 2017 report. GOLD executive summary. *Am J Respir Crit Care Med.* 2017;195(5):557–582.
53. Hancox RJ, Gray AR, Poulton R, Sears MR. The effect of cigarette smoking on lung function in young adults with asthma. *Am J Respir Crit Care Med.* 2016;194(3):276–284.
54. Bardhan M, Sarkar M, Singh D, Negi R, Sharma S. Clinical and microbiological profile of patients with acute exacerbation of COPD. *NJR.* 2020;10(1):35–40.
55. Dalju I, Dessie A, Bogale L, Mekonnen TH. Occupational risk factors associated with respiratory symptoms among tannery workers in Mojo town, Southeast Ethiopia, 2018: a comparative cross-sectional study. *Multidiscip Respir Med.* 2019;14(1):1–10.
56. Jamal A, Mehmood A, Khatoon F, Putus T, Savolainen H, Liesivuori J. Prevalence of respiratory symptoms, bronchial asthma and obstructive lung disease among tannery workers. *Ann Environ Sci Toxicol.* 2017;2(1):033–042.
57. Stern FB. Mortality among chrome leather tannery workers: an update. *Am J Ind Med.* 2003;44(2):197–206.
58. Dalju I, Dessie A, Bogale L, Mekonnen TH. Occupational risk factors associated with respiratory symptoms among tannery workers in Mojo town, Southeast Ethiopia, 2018: a comparative cross-sectional study. *Multidiscip Respir Med.* 2019;14(1):27.
59. Miravittles M. Cough and sputum production as risk factors for poor outcomes in patients with COPD. *Respir Med.* 2011;105(8):1118–1128.
60. Rastogi SK, Pandey A, Tripathi S. Occupational health risks among the workers employed in leather tanneries at Kanpur. *Indian J Occup Environ Med.* 2008;12(3):132.
61. Yoh M, Matsuyama J, Ohnishi M, et al. Importance of *Providencia* species as a major cause of travellers' diarrhoea. *J Med Microbiol.* 2005;54(11):1077–1082.
62. Kaur CP, Vadivelu J, Chandramathi S. Impact of *Klebsiella pneumoniae* in lower gastrointestinal tract diseases. *J Dig Dis.* 2018;19(5):262–271.
63. Hasan MJ, Nizhu LN, Rabbani R. Bloodstream infection with pandrug-resistant *Alcaligenes faecalis* treated with double-dose of tigecycline. *IDCases.* 2019;18:e00600.
64. Chmelař D, Holý O, Kasáková I, et al. Antibiotic susceptibility and production of endotoxin by *Ochrobactrum anthropi* isolated from environment and from patients with cystic fibrosis. *Folia Microbiol (Praha).* 2019;64(6):861–865.
65. Grzyb J, Lenart-Boroń A. Bacterial bioaerosol concentration and size distribution in the selected animal premises in a zoological garden. *Aerobiologia.* 2019;35(2):253–268.

International Journal of Chronic Obstructive Pulmonary Disease

Dovepress

Publish your work in this journal

The International Journal of COPD is an international, peer-reviewed journal of therapeutics and pharmacology focusing on concise rapid reporting of clinical studies and reviews in COPD. Special focus is given to the pathophysiological processes underlying the disease, intervention programs, patient focused education, and self management

protocols. This journal is indexed on PubMed Central, MedLine and CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-chronic-obstructive-pulmonary-disease-journal>