Multiple Organ Dysfunction Syndrome Caused by Sepsis: Risk Factor Analysis

Guo-Dong Sun¹Yang Zhang²Shan-Shan Mo³Ming-Yan Zhao¹

¹Department of Critical Care Medicine, The First Affiliated Hospital of Harbin Medical University, Harbin, 150001, Heilongjiang Province, People's Republic of China; ²Department of Nursing, Central Hospital of Heilongjiang Province Prison, Harbin, 150001, Heilongjiang Province, People's Republic of China; ³Pharmacy Department of Heilongjiang Sailors General Hospital, Harbin, 150001, Heilongjiang Province, People's Republic of China

Purpose: To analyze the risk factors of multiple organ dysfunction syndromes (MODS) caused by sepsis.

Patients and Methods: A total of 180 patients with sepsis admitted to The First Affiliated Hospital of Harbin Medical University (No. 23, Post Street, Nangang District, Harbin 150001, Heilongjiang province, China) from July 2018 to June 2019 were selected and divided into a non-MODS group and a MODS group, with 90 cases in each group. Clinical data of the patients were retrospectively analyzed, and univariable and multivariable analyses were performed.

Results: The univariable analysis showed that there were no significant differences in terms of age, body temperature, heart rate, respiration, mean arterial pressure, RBC specific volume, blood sodium, serum potassium, and infection site ($P > 0.05$). Whereas significant differences were found between the groups in terms of gender, arterial blood pH, WBC count, Apache II score, blood glucose, creatinine, chronic medical history, surgery, and ventilator usage ($P < 0.05$). The growth of bacterial culture, the increase of creatinine level, chronic diseases and Apache II score were discovered to have significant effects on the occurrence of MODS through the multivariable logistic regression analysis.

Conclusion: Bacterial culture, serum creatinine level, history of chronic disease and Apache II score may be risk factors of MODS in sepsis patients.

Keywords: sepsis, multiple organ dysfunction syndrome, risk factors, systemic inflammatory response syndrome, sepsis patients, multivariable logistic regression analysis

Introduction

Sepsis is a systemic inflammatory response syndrome (SIRS) caused by either infectious or non-infectious factors.¹⁻³ It can easily develop into septic shock and multiple organ dysfunction syndromes (MODS) if not treated in a timely manner, and its fatality rate can be as high as 28–56%.⁴ Clinically, MODS can be categorized into Phase I immediate type and Phase II delayed type. To date, the preferred medical examination for MODS includes the examination of hemogram, blood, pathogenic bacteria and urine.^{5,6} Systemic inflammatory response syndrome may be viewed as the systemic expression of cytokine signals that normally function on an autocrine or paracrine level. Sepsis is defined as systemic inflammatory response syndrome caused by an infection. MODS may represent the end stage of severe systemic inflammatory response syndrome or sepsis. Many cells are involved, including endothelial cells and leukocytes and multiple proinflammatory and anti-inflammatory mediators (cytokines, oxygen free radicals, coagulation factors, and so forth). Various pathophysiologic mechanisms have been postulated. The most

Correspondence: Ming-Yan Zhao
Email shizhaomi0147@126.com

Received: 26 July 2021
Accepted: 28 September 2021
Published: 27 October 2021

International Journal of General Medicine 2021:14 7159–7164

7159

© 2021 Sun et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at <https://www.dovepress.com/terms.php> and incorporate the Creative Commons Attribution – Non Commercial (unported, v3.0) License (<http://creativecommons.org/licenses/by-nc/3.0/>). By accessing the work you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (<https://www.dovepress.com/terms.php>).

popular theory is that the inflammatory process loses its autoregulatory capacity.^{5,6} Existing studies recognize that immune dysfunction caused by uncontrolled systemic inflammatory response is the main pathophysiological mechanism of sepsis, while the underlying pathogenesis is poorly understood.⁵ Involving multiple organs and pathophysiological systems, these diseases are closely related to the systemic inflammatory network, bacterial endotoxin, inflammatory mediators, immune dysfunction, intestinal bacterial and endotoxin translocation, coagulation dysfunction, and gene polymorphism. The pathogenesis of sepsis needs further clarification.⁷ This study was undertaken with an attempt to further elucidate the risk factors related to MODS caused by sepsis, and to take targeted prevention and treatment measures.

Patients and Methods

General Material

A total of 180 patients presenting to The First Affiliated Hospital of Harbin Medical University (No. 23, Post Street, Nangang District, Harbin 150001, Heilongjiang province, China) due to sepsis from July 2018 to June 2019 were selected for this research. Among them, there were 113 males and 67 females, who aged between 18–79 years (mean, 41.23 ± 7.27) years old. There were 117 cases of acute abdominal disease (including suppurative cholangitis, acute severe pancreatitis, etc.), 6 cases of systemic infection, 19 cases of multiple injuries, 22 cases of abdominal trauma, and 16 other cases. Patients who were admitted to the intensive care unit (ICU) and then left or died in the ICU were included, while those who died within 24 hours after being sent to the ICU were excluded. The primary infection in patients with sepsis and diseases distribution were observed. The studies involving human participants were reviewed and approved by First Affiliated Hospital of Harbin Medical University. The patients provided their written informed consent to participate in this study. This study has been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans.

Diagnostic Criteria

Sepsis is defined in accordance with the diagnostic criteria in the Guidelines for Emergency Treatment of Sepsis/Septic Shock in China (2018) issued by the Emergency Physicians Branch of the Chinese Physicians Association.

When the sequential organ failure assessment (SOFA) score rises by over 2 points from baseline, it is considered as sepsis. See Table 1. The criteria for organ dysfunction lie with the MODS score developed by Marshall, a Canadian scholar. See Table 2.

Methods

Patients were equally divided into a non-MODS group and a MODS group means according to their conditions. MODS standard was interpreted by the kidney, lung, heart, blood, liver and brain. Each organ is associated with several indicators. For instance, creatinine is considered the indicator for kidney, respiration for lung, heart rate for heart, concentration for sodium ion, serum potassium ion and blood glucose the indicators for blood. Each clinical indicator of the patient was recorded, and the worst value of each indicator was input into the database.

Predictor Variables

- (1) General items: gender, age, history of chronic diseases.
- (2) Post-inoculation (HPI): body temperature, heart rate, respiration, mean arterial pressure, arterial blood pH value, infection site, red blood cell (RBC), white blood cell (WBC) count, APACHE II score, surgery, blood sodium, blood serum potassium, blood glucose, serum creatinine, bacteria growth, ventilator usage.

Statistical Methods

SPSS22.0 statistical software was used for univariable analysis. Measurement data were represented by ($\bar{x} \pm s$) and the differences between groups were analyzed by *t*-test. Enumeration data were expressed as n(%) and the differences between groups were analyzed by χ^2 test. Logistic regression was used for multivariable analysis. $P < 0.05$ was considered statistically significant.

Results

Comparison of Measurement Index Between the Groups

No statistically significant difference was found in terms of age, body temperature, heart rate, respiration, mean arterial pressure, RBC hematocrit, blood sodium, serum potassium between the two groups ($P > 0.05$). The arterial blood pH value, the WBC count, APACHE II score, blood glucose, and creatinine were statistically different between the groups ($P < 0.05$). See Table 3 for details.

Table 1 SOFA Scoring Criteria

System	Score/Points				
	0	1	2	3	4
Respiratory system PaO ₂ /FiO ₂ /mmHg(kPa)	≥400(53.3)	<400(53.3)	<300(40.0)	<200(26.7) + Mechanical ventilation	<100(13.3) + Mechanical ventilation
Coagulation system Blood platelet/ (10 ³ •μL ⁻¹)	≥150	<150	<100	<50	<20
Liver Bilirubin / [mg•dL ⁻¹ (μmol•L ⁻¹)]	<1.2(20)	1.2–1.9(20–32)	2.0–5.9(33–101)	<6.0–11.9(102–204)	≥12.0(204)
Cardiovascular system	MAP≥70mmHg	MAP<70mmHg	Dopamine <5 or dobutamine (any dose)	Dopamine 5.1–15.0 or epinephrine ≤0.1 or norepinephrine >0.1	Dopamine >15 or adrenaline >0.1 or norepinephrine >0.1
Central nerve system (GCS)	15	13–14	10–12	6–9	<6
Kidney Creatinine / [mg•dL ⁻¹ (μmol•L ⁻¹)]	<1.2(110)	1.2–1.9(110–170)	2.0–3.4(171–299)	3.5–4.9(300–400)	>4.9(440)
Urine Volume (mL•d ⁻¹)	–	–	–	<500	<200

Table 2 MODS Score (Marshall Standard)

Organ	Score				
	0	1	2	3	4
Respiratory system (PaO ₂ /FiO ₂)	>300	226–300	151–225	76–150	≤75
Kidney (Blood Cr μmol•L ⁻¹)	≤100	101–200	201–350	351–500	>500
Liver (hemobilirubin μmol•L ⁻¹)	≤20	21–60	61–120	121–240	>240
Cardiovascular system (PAR=HR×CVP/MAP)	≤10.0	10.1–15.0	15.1–20.0	20.1–30.0	>30.0
Blood (platelet count 10 ⁹ •L ⁻¹)	>120	81–120	51–80	21–50	≤20
Nerve system (Glasgow Coma score)	15	13–14	10–12	7–9	≤6

Comparison of Enumeration Indexes Between the Groups

There was no statistically significant difference between the two groups in the site of infection ($P > 0.05$), but significant differences were found in terms of frequencies of gender, history of chronic disease, surgery, bacterial culture and growth, and ventilator usage ($P < 0.05$). See [Table 4](#) for details.

Logistic Regression Analysis of Risk Factors for MODS in Sepsis Patients

The univariable analysis preliminarily screened out gender, arterial blood pH value, the WBC count, APACHE II

score, blood glucose, creatinine, chronic disease, bacterial culture growth, surgery, and ventilator usage, totaling 10 variables for binary logistic regression analysis. The result showed that the growth of bacterial culture, increased creatinine level, combined chronic diseases, a higher score of APACHE II are risk factors for MODS occurrence. See [Table 5](#) for details.

Site Composition of Primary Infection in Patients with Sepsis

Compared with the incidence of respiratory tract infection, the incidence of digestive tract, abdominal cavity,

Table 3 Comparison of Measurement Data Between the Two Groups ($\bar{x} \pm s$)

Observation Index	MODS Group (n=90)	Non-MODS Group (n=90)	P
Age (year old)	46.04 $\pm$ 17.42	43.83 $\pm$ 15.21	0.366
Body temperature ($^{\circ}$ C)	38.53 $\pm$ 0.91	38.58 $\pm$ 0.94	0.717
Heart rate (times/min)	113.80 $\pm$ 18.07	110.92 $\pm$ 16.81	0.269
Respiration (times/min)	29.44 $\pm$ 6.42	27.84 $\pm$ 5.05	0.065
MAP (mmHg)	88.15 $\pm$ 19.35	90.42 $\pm$ 18.01	0.416
Arterial blood Ph value	8.42 $\pm$ 0.23	7.45 $\pm$ 0.08	<0.001
RBC specific volume (%)	34.42 $\pm$ 9.95	36.76 $\pm$ 9.28	0.105
WBC count ($\times 10^9 \cdot L^{-1}$)	19.88 $\pm$ 5.34	16.86 $\pm$ 5.91	<0.001
APACHE II Score (points)	26.14 $\pm$ 11.36	16.13 $\pm$ 6.86	<0.001
Blood sodium (mmol $\cdot L^{-1}$)	138.88 $\pm$ 8.67	138.05 $\pm$ 7.11	0.483
Serum kalium (mmol $\cdot L^{-1}$)	5.38 $\pm$ 0.92	5.35 $\pm$ 0.64	0.799
Blood glucose (mmol $\cdot L^{-1}$)	21.33 $\pm$ 6.30	10.65 $\pm$ 3.92	<0.001
Creatinine (μ mol $\cdot L^{-1}$)	185.46 $\pm$ 235.16	84.03 $\pm$ 94.51	<0.001

Table 4 Comparison of Enumeration Data Between the Two Groups [n (%)]

Observation Index	MODS Group (n = 90)	Non-MODS Group (n = 90)	P
Gender (Male/Female)	71/19	58/32	0.032
History of chronic disease (Yes/No)	36/54	17/73	0.002
Infection site (Respiration system/Other than Respiration system)	73/17	81/9	0.090
Surgery (Surgery/Non-surgery)	39/51	54/36	0.025
Bacteria culture growth (Yes/No)	35/55	53/37	0.007
Ventilator usage (Applied/Not applied)	76/14	85/5	0.029

nervous system and other infections was significantly lower ($\chi^2=34.844$, $P < 0.05$). See Figure 1 for details.

Disease Distribution in the Two Groups

The diseases of the patients are mainly diffuse peritonitis, pancreatitis, hepatobiliary diseases, multiple injuries,

intestinal obstruction, chest diseases, limb diseases, etc. No significant difference was observed in the disease distribution between the two groups ($P > 0.05$). See Figure 2 for details.

Discussion

Sepsis serves as the organ dysfunction syndrome caused by the imbalance of the human body's response to infection.⁸ After infection, inflammatory response cells are activated by bacterial toxins and will release inflammatory mediators such as interleukin-6 (IL-6), interleukin-1 (IL-1) and tumor necrosis factor- α (TNF- α), triggering systemic or local inflammatory responses.⁹⁻¹¹ Its clinical manifestations include changes in mental state, shortness of breath, palpitation, fever, chills and other symptoms, and in severe cases, MODS will develop.¹² MODS refers to the clinical syndrome in which at least two systems or organs suffer from simultaneous or sequential dysfunction during major surgery, infection, shock, poisoning and severe trauma, thus undermining the stability of the internal environment.¹³⁻¹⁵ Early Goal-Directed Therapy (EGDT) regimen, early monitoring of risk factors and active fluid resuscitation can reduce the incidence and mortality of MODS, and plays a key role in early control of sepsis.¹⁶

Table 5 Logistic Regression Analysis of Risk Factors for MODS in Sepsis Patients

Factors	Estimated Parameter	P value	OR Value	95% CI
Bacterial culture growth	-1.776	0.032	0.275	0.045, 0.787
Increased creatinine level	0.061	<0.001	1.062	1.028, 1.075
Combined chronic diseases	-1.758	0.029	0.204	0.049, 0.768
Increased APACHE II score	0.339	<0.001	1.367	1.132, 1.394

Figure 1 Site composition of primary infection in patients with sepsis.

The univariable analysis of this research showed that the two groups of patients had significant differences in gender, arterial blood pH value, WBC count, APACHE II score, blood glucose, serum creatinine, chronic disease, bacterial culture growth, surgery, ventilation treatment. Additionally, it is documented that patients with sepsis caused by shock, infection, trauma and other factors will have pathophysiological changes in their circulatory, respiratory and nervous systems, including tissue and organ ischemia and hypoxia, renal function decline, electrolyte disturbance, shortness of breath, blood pressure drop, and acid-base imbalance.¹⁷ On account of these, MODS and even death may arise if early symptoms are not treated in time.

The multivariable logistic stepwise regression analysis showed that the growth of bacterial culture, increased creatinine level, combined chronic diseases, a higher score of APACHE II are risk factors for the occurrence of MODS. As a vital index for evaluating sepsis patients, APACHE II score reflects the association between the

body's physiological disorders and the severity of sepsis, which is positively correlated with MODS.¹⁸ Moreover, it has been pointed out by a prior trial that patients with sepsis combined with cardiopulmonary diseases, diabetes, hypertension and other chronic diseases have a higher risk of developing MODS than others.¹⁹ Serum creatinine is also an important indicator to observe the development of sepsis. Systemic inflammatory reactions owing to infection can easily cause changes in the circulatory system. It in turn dilates blood vessels, slows down blood flow and lowers blood pressure, and subsequently further decreases renal blood flow, impairs renal function and increases serum creatinine. Our study demonstrated that serum creatinine in the MODS group was significantly higher than that in the non-MODS group. Taken together, early protection of renal function and improvement of microcirculation are of paramount importance for the treatment of MODS.²⁰

Hence, we suggest that gender, arterial blood pH value, the WBC count, APACHE II score, blood glucose, serum creatinine, chronic disease, surgery, bacterial growth, ventilator usage and other factors were relevant to the development of MODS in patients with sepsis in the Emergency Department in The First Affiliated Hospital of Harbin Medical University (No. 23, Post Street, Nangang District, Harbin 150001, Heilongjiang province, China.). Among them, bacteria culture growth, increased creatinine level, combined chronic diseases, a higher score of APACHE II appear to be risk factors for the occurrence of MODS. Therefore, early monitoring of these factors and timely control measures can improve the

Figure 2 Disease distribution.

cure rate and reduce the fatality rate of sepsis patients. Moreover, the confounding and omitted variables of the results were critical and should be emphasized in future. Meanwhile, a multivariable model with a less selective set of explanatory variables should be performed in future studies.

Thus, we concluded that Bacterial culture, serum creatinine level, history of chronic disease and Apache II score might be risk factors of MODS in sepsis patients.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Srouji LS, Moore-Clingenpeel M, Hensley J, et al. shock severity modifies associations between RBC transfusion in the first 48 hours of sepsis onset and the duration of organ dysfunction in critically ill septic children. *Pediatr Crit Care Med*. 2020;21(8):e475–e484. doi:10.1097/PCC.0000000000002338
2. Zhang Q, Shan F. The role of p38 mitogen-activated protein kinase signaling pathway in the sepsis with multiple organ dysfunction syndrome. *Chin J Cell Biol*. 2018. Available from: <https://d.wanfangdata.com.cn/periodical/xbswxzz201802018>. Accessed October 19, 2021.
3. Bojana M, Roy H, Frédéric M, et al. Conservative management of pelvic sepsis with severe shock and multiple organ dysfunction syndrome after rubber-band ligation of internal haemorrhoids: surgery is not the only option. *J Surg Case Rep*. 2018;2018(8):8.
4. Ziesmann MT, Marshall JC. Multiple organ dysfunction: the defining syndrome of sepsis. *Surg Infect (Larchmt)*. 2018;19(2):184–190. doi:10.1089/sur.2017.298
5. Drobatz KJ, Hopper K, Rozanski E, et al. *Textbook of Small Animal Emergency Medicine || Systemic Inflammatory Response Syndrome, Sepsis, and Multiple Organ Dysfunction Syndrome*. John Wiley & Sons; 2018.
6. Zhang L, Zhang T, Bian K, Wu D. Clinical effect of CHVHF on severe sepsis complicated with multiple organ dysfunction syndrome and its effect on hemodynamics. *Clin Res Pract*. 2018. Available from: <https://wenku.baidu.com/view/bce38cc2e3bd960590c69ec3d5bbfd0a7956d5ae?fr=xueshu>. Accessed October 19, 2021.
7. Troia R, Giunti M, Goggs R. Plasma procalcitonin concentrations predict organ dysfunction and outcome in dogs with sepsis. *BMC Vet Res*. 2018;14(1):111. doi:10.1186/s12917-018-1427-y
8. Ramirez JA. A continuum of disease from community-acquired pneumonia to multiple organ dysfunction syndrome. *Univ Louisville J Respir Infect*. 2018;2:4–6. doi:10.18297/jri/vol2/iss2/1/
9. Alcamo A, Pang D, Bashir D, et al. Role of damage-associated molecular patterns and uncontrolled inflammation in pediatric sepsis-induced multiple organ dysfunction syndrome. *J Pediatr Intensive Care*. 2019;08(01):025–031. doi:10.1055/s-0038-1675639
10. Gatica S, Eltit F, Santibanez J, et al. Expression suppression and activity inhibition of TRPM7 regulate cytokine production and multiple organ dysfunction syndrome during endotoxemia: a new target for sepsis. *Curr Mol Med*. 2019;19(8):547–559. doi:10.2174/1566524019666190709181726
11. Kell DB, Pretorius E. To what extent are the terminal stages of sepsis, septic shock, systemic inflammatory response syndrome, and multiple organ dysfunction syndrome actually driven by a prion/amyloid form of fibrin? *Semin Thromb Hemost*. 2018;44(03):224–238. doi:10.1055/s-0037-1604108
12. Cheng Z, Abrams ST, Alhamdi Y, et al. Circulating histones are major mediators of multiple organ dysfunction syndrome in acute critical illnesses. *Crit Care Med*. 2019;47(8):e677–e684. doi:10.1097/CCM.0000000000003839
13. Zhang YJ, Zhuo XA, Zhou XM, Dai WJ. Prognostic value of lactate/albumin ratio in the elderly sepsis patients complicated with multiple organ dysfunction syndrome. *Chin J Multiple Organ Dis Elder*. 2018. Available from: <https://d.wanfangdata.com.cn/periodical/zhlndqgjbzz201804003>. Accessed October 19, 2021.
14. Man-Xue M, Zhi-Guang WU, Jin-Li F, et al. Overview of Chinese medicine for treatment of sepsis with multiple organ dysfunction. *J Guangzhou Univ Trad Chin Med*. 2019.
15. Vallejos A, Olivares P, Gatica S, et al. Endotoxemia-induced endothelial fibrosis inhibition improves hypotension, tachycardia, multiple organ dysfunction syndrome, cytokine response, oxidative stress, and survival. *Lab Invest*. 2019;99(8):1173–1192.
16. Huang W, Qin S, Sun Y, et al. [Establishment of multiple organ dysfunction syndrome early warning score in patients with severe trauma and its clinical significance: a multicenter study]. *Zhonghua W Zhong Bing Ji Jiu Yi Xue*. 2018;30(1):41–46. Chinese.
17. Xie J, Yang F, Chen J, Wang W, Yi C, Bai X. Application of vacuum sealing drainage in treatment of open fractures and multiple organ dysfunction syndrome. *J Trauma Surg*. 2018. Available from: http://qikan.cqvip.com/Qikan/Article/Detail?id=675978887&from=Qikan_Search_Index. Accessed October 19, 2021.
18. Uchida M, Abe T, Ono K, et al. Ulinastatin did not reduce mortality in elderly multiple organ failure patients: a retrospective observational study in a single center ICU. *Acute Med Surg*. 2018;5(1):90–97. doi:10.1002/ams2.304
19. Wu D, Shi S, Zhang R, et al. Expression and clinical significance of the novel inhibitory cytokine IL-3S in patients with sepsis and multiple organ dysfunction syndrome. *Chin J Emerg Med*. 2019;28(2):175–180.
20. Zhang WF, Chen DM, Wu LQ, et al. Clinical effect of continuous blood purification in treatment of multiple organ dysfunction syndrome in neonates. *Zhongguo Dang Dai Er Ke Za Zhi = Chinese Journal of Contemporary Pediatrics*. 2020;22(1):31–36.

International Journal of General Medicine

Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies

across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-general-medicine-journal>