#### ORIGINAL RESEARCH

Prevalence and Associated Factors of Kidney Dysfunction in Patients with Hypertension and/or **Diabetes Mellitus from a Primary Care Population** in Northwest China

Mengyue Lin Mulalibieke Heizhati<sup>2</sup> Lin Wang<sup>2</sup> Lin Gan <sup>[],2</sup> Mei Li<sup>2</sup> Wenbo Yang<sup>2</sup> Ling Yao<sup>2</sup> Zhongrong Wang<sup>2</sup> Zhikang Yang<sup>2</sup> Reyila Abudoyreyimu<sup>2</sup> Zihao Wu<sup>2</sup> Nanfang Li

<sup>1</sup>Xinjiang Medical University, Urumqi, People's Republic of China; <sup>2</sup>Hypertension Center of the People's Hospital of Xinjiang Uygur Autonomous Region; Xinjiang Hypertension Institute; National Health Committee Key Laboratory of Hypertension Clinical Research, Urumgi, Xinjiang, People's Republic of China

Hypertension Center of People's Hospital of Xinjiang Uygur Autonomous Region; Xinjiang Hypertension Institute; National Health Committee Key Laboratory of Hypertension Clinical Research, 91 Tianchi Road, Urumqi, Xinjiang, 830001, People's Republic of China Email Inanfang2016@sina.com

Purpose: The burden of kidney dysfunction (KD) is rapidly increasing in developing countries due to an ongoing epidemic of hypertension and diabetes. We aimed to evaluate the prevalence and associated factors of KD among population with hypertension and/or diabetes from primary care setting.

**Methods:** This study was part of a cross-sectional study conducted in Northwest China, which aimed to investigate the epidemiological status of hypertension and other chronic noninfectious diseases. Subjects aged  $\geq 18$  years old with hypertension and/or diabetes were included in this study. KD was defined as estimated glomerular filtration rate (eGFR) less than 60 mL/min/1.73m<sup>2</sup>. Least absolute shrinkage and selection operator and multivariable logistic regression were used to identify factors associated with KD. Sensitivity analysis was performed by comparing the results of different estimated equations for GFR.

Results: A total of 8899 participants with hypertension and/or diabetes were included. Prevalence of KD was 8.69% (n=144) among patients with both hypertension and DM, 3.36% (n=198) among patients with only hypertension, and 5.22% (n=70) among patients with only diabetes. Women showed higher prevalent KD than men. Modifiable factors independently associated with KD among patients with hypertension and/or diabetes included physical activity, duration of hypertension or DM, anemia, fasting plasma glucose and triglyceride. Results of different estimated equations showed similar trends of difference in the prevalence of KD among groups.

**Conclusion:** KD is common in patients with hypertension and/or DM in Northwest China. More attention should be paid to high-risk patients, especially to those with co-existence of hypertension and DM. Control of blood glucose and triglyceride may further improve KD management in this patient population.

**Keywords:** kidney dysfunction, hypertension, diabetes, epidemiology

#### Introduction

Chronic kidney disease (CKD) has been shown to be a risk factor for cardiovascular diseases and death.<sup>1</sup> Early-stage CKD is usually asymptomatic and is often ignored.<sup>2</sup> Most patients detected for CKD are already in an irreversible stage, which leads to occurrence of complications, and some of them have even progressed to end-stage renal disease (ESRD), requiring renal replacement therapy. Hence, the burden of CKD has become a huge public health challenge worldwide.

Correspondence: Nanfang Li

Prevalence of CKD varies widely among geographic regions, partly due to different lifestyles and economic conditions. Previous studies have shown a heavy burden of CKD in developing countries and low- and middleincome regions (LMIRs), which is responsible for more than two thirds of overall CKD,<sup>3</sup> with the prevalence of 17.95% in Iran, and 8.66% and 13.1% in Africa and South Asia, respectively.<sup>4</sup> Prevalence of CKD is about 10.8% in the general population of China,<sup>5</sup> and even higher in the elderly.<sup>6</sup> With changes in lifestyle and acceleration of aging, incidence of CKD is increasing as well, and risk factors are also changing, with an increasing trend of metabolic disease-related kidney disease.<sup>7</sup> Studies have demonstrated that hypertension and diabetes mellitus (DM) play strong roles in kidney disease. More than 40% of diabetic patients in America and Japan<sup>8,9</sup> and 25% of hypertensive patients in Turkey<sup>10</sup> are reported to be affected by kidney damage. Diabetic kidney disease and hypertensive nephropathy account for more than 40% of CKD in Chinese hospitalized patients.<sup>7</sup> Furthermore, it is noteworthy that hypertension and DM have become leading causes of CKD in Chinese elderly.<sup>11</sup> Rapid increase in incidence of hypertension and DM, even at younger age, over the past two decades<sup>12</sup> indicates that prevalence of CKD at population level may have been affected.

Kidney dysfunction (KD) is an important component of CKD. Although the prevalence of KD has previously been reported, few studies have focused on the population with hypertension and DM, especially in developing countries and LMIRs. Therefore, we aimed to investigate the prevalence and the related factors of KD in the population with hypertension and/or DM from the primary care setting of Xinjiang, a less-developed region in Northwest China, where hypertension and DM are highly prevalent and poorly managed (low awareness, treatment, and control rates).<sup>13,14</sup>

# Materials and Methods Study Design and Population

This study is part of a cross-sectional study conducted in Emin county between March and November 2019, and aimed to investigate epidemiological status of hypertension and other chronic noninfectious diseases. Emin county, in Northern Xinjiang, is home to a total population of over 160,000, living in urban, agricultural, and stockraising settings. Economic level of Emin county is close to the average of Xinjiang, making the study population characteristics well representative.

In the original investigation, we used multistage stratified random sampling method to obtain study population. At the first stage, six districts were randomly selected in Emin county. At the second stage, we selected 10 villages/ communities from each of district. At the third stage, individuals were randomly chosen from each village/community. Local inhabitants (residing at current address for  $\geq 6$  months) aged  $\geq 15$  years and agreed to participate and sign an informed consent form were investigated. The study was conducted in accordance with Declaration of Helsinki and approved by the Ethics Committee at the People's Hospital of Xinjiang Uygur Autonomous Region.

#### Eligibility Criteria in the Present Study

In this study, participants were included if they were aged  $\geq 18$  years with hypertension and/or diabetes. Hypertension was defined as systolic blood pressure (SBP)  $\geq 140$  mmHg, and/or diastolic blood pressure (DBP)  $\geq 90$  mmHg, and/or use of antihypertensive agents within two weeks, and/or self-reported history of hypertension. Diabetes mellitus was defined as self-reported history of diabetes and/or a FPG  $\geq 7$  mmol/L.

## Data Collection and Measurement

Data were collected by trained investigators using standardized questionnaires, including demographic characteristics, socioeconomical status, health-related behaviors, and medical history.

Before blood pressure (BP) measurement, each participant was asked to avoid alcohol, cigarettes, coffee or tea consumption. BP was measured thrice using with the OMRON HBP-1300 Professional Portable Blood Pressure Monitor (OMRON, Kyoto, Japan) on the right arm positioned at heart level after the participant was sitting at rest for five minutes, with 30 seconds between each measurement. The average of the three readings was used for analysis. Height and weight were measured to the nearest 0.1 cm and 0.1 kg, respectively, with participants in lightweight clothing and without shoes. Body mass index (BMI) was calculated by dividing weight by heightsquared  $(kg/m^2)$ . Waist circumference was measured at the midpoint between the lower rib and upper margin of the iliac crest to the nearest 0.1 cm at the end of a normal expiration.

All subjects were fasted for  $\geq 8$  hours, and a 5-mL fasting blood sample was collected. Serum creatinine was

measured by enzymatic method (sarcosine oxidase enzymatic analysis). Fasting plasma glucose (FPG) was tested using glucose oxidase method. Total cholesterol (TC) and triglycerides (TG) were measured with commercially available reagents.

# Definition of KD

KD was defined as eGFR <60 mL/min/per  $1.73m^2$ . eGFR was calculated using modified modification of diet in renal disease (MDRD) equation on the basis of data from Chinese adults.<sup>15</sup> As a sensitivity analysis, we also examined eGFR calculated using chronic kidney disease epidemiology collaboration (CKD-EPI) creatinine equation.<sup>16</sup>

#### Other Variables

Additional variables included age, gender, duration of hypertension and DM (years), smoker (yes/no), alcohol intake (yes/no), living regions (agricultural, stock-raising, or urban setting), educational level (primary and lower, junior high, or senior high and higher), occupational (manual or intelligent), ethnicity (Han, Kazakh, or others), marital status (single, married, or separated). Participants who have smoked at least 20 packets of cigarettes in their lifetime were defined as smoker. Alcohol intake was defined as consuming alcoholic beverage at least once per week in the past month. Central obesity was defined as waist circumference  $\geq$ 85 cm for women and  $\geq$ 90 cm for men. A positive cardiovascular disease (CVD) was considered as any self-reported myocardial infarction, coronary revascularization or stroke. Anemia was defined if hemoglobin <120 g/L for men or <110 g/L for women. Physical activity was evaluated by the Global Physical Activity Questionnaire and divided into three categories WHO's (2010)physical activity based on recommendations.<sup>17</sup> Sleep quality was assessed using Pittsburgh Sleep Quality Index (PSQI), and classified into very good, fair good, fair bad and very bad categories if PSQI score of 0–4, 5–9, 10–14, and  $\geq$ 15, respectively.<sup>18</sup>

## Statistical Analysis

Continuous variables were summarized as mean  $\pm$ standard deviation (SD) or median (interquartile range) and compared using Student's *t*-test or Mann–Whitney *U*-test, whereas categorical variables were expressed as absolute numbers with percentage and compared using chi-squared test. To evaluate associated factors of KD in patients with different disease status, data was divided into three analytic sets: set 1 including participants with both hypertension

and DM; set 2 including those with hypertension alone; set 3 including those with DM alone. First, we used least absolute shrinkage and selection operator (LASSO) regression to identify predictive features of KD. Then, multivariable logistic regression model including those variables was fitted to identify factors associated independently with KD, with results presented as odds ratio (OR) with 95% confidence interval (CI). CKD-EPI equation is also commonly used to estimate GFR and has been well confirmed for epidemiological investigation and clinical practice.<sup>16</sup> Therefore, we redefined KD based on CKD-EPI and repeated the above analysis process as sensitivity analysis for prevalence and associated factors of KD. Statistical analyses were conducted using SPSS 20.0 for Windows (IBM Corporation, Armonk, NY, USA) and R version 4.0.3. A two-sided P<0.05 was considered to be significant.

#### Results

## Prevalence of Kidney Dysfunction Based on MDRD Equation

A total of 41,089 people were recruited in the original investigation and 36,643 completed the survey with a response rate of 89.2%. After exclusion, we finally included 8899 participants with hypertension and/or DM in this study (Figure 1). Among the included participants, 5901 (66.3%) had hypertension alone, 1341 (15.1%) had DM alone, and 1657 (18.6%) had both hypertension and DM. KD was detected in 412 (4.63%) of study participants. As shown in Figure 2, prevalence of KD was as high as 8.69% (n=144) among patients with both hypertension and DM, 3.36% (n=198) among those with hypertension alone and 5.22% (n=70) among those with DM alone. Women had higher prevalent KD than men. Prevalence of KD in participants from both genders, different age groups, living region, ethnicity and educational status groups were shown in Table 1.

## Subject Characteristics

Among included participants, 7558 (84.9%) had hypertension and 2998 (33.7%) had DM. Mean age of all participants was  $56.61\pm12.11$  years, mean BMI was 27.22  $\pm4.33$  kg/m<sup>2</sup> and almost two thirds (65.1%) had central obesity and 49.3% participants were women. 881 (9.9%) participants reported a positive history of CVD. Compared with non-KD participants, patients with KD were older and had higher proportion of women (48.5% vs 66.3%,



Figure I Flow chart of screening and including subjects. Abbreviation: DM, diabetes mellitus.

P < 0.001). Proportion of coexistent hypertension and DM was higher in those with KD. Patients with KD were more likely to be diabetic and have higher FPG level. History of CVD and dyslipidemia were also higher in KD group. Details of characteristics for study population were shown in Table 2. Moreover, among participants with hypertension alone, treatment and control rates of hypertension were 46.1% and 18.5%, respectively; median duration of hypertension was 4.38 years. Among participants with DM alone, median duration of DM was 2.66 years.

## Associated Factors of KD

LASSO regression generated 15, 17, and six factors for analytic set 1 (for coexistent hypertension and DM), 2 (for hypertension alone) and 3 (for DM alone), respectively. In multivariable logistic regression models by including parameters produced by LASSO regression, among participants with both hypertension and DM, factors associated with KD included age, gender, duration of hypertension, anemia, FPG and TG (Table 3). Among participants with hypertension alone, age, gender, living region, physical activity, duration of hypertension and FPG were independently associated with KD (Table 4). Among participants with DM alone, associated factors for KD were duration of DM, living region and physical activity (Table 5).

## Sensitivity Analysis

Sensitivity analysis was performed by redefined KD based on CKD-EPI equation; results were shown as <u>supplemen</u> <u>tary Materials</u>. In CKD-EPI analysis, prevalence of KD was higher than MDRD analysis (7.09% vs 4.63%), and the lowest prevalence was found in patients with hypertension alone (5.37%), while the highest was in patients with both hypertension and DM (12.61%). Results of the two equations showed similar trends in difference in the prevalence of KD among groups (Figure S1 and Table 1). In



Figure 2 Prevalence of kidney dysfunction in patients with hypertension and/or diabetes based on MDRD equation. \*Indicates P<0.01. Abbreviations: HTN, hypertension; DM, diabetes mellitus.

multivariable analysis based on CKD-EPI, most of the associated factors identified by MDRD analysis remained consistent and significant (<u>Tables S1–S3</u>). Moreover, in CKD-EPI analysis, occupation, CVD and physical activity were significantly associated with KD among patients with both hypertension and DM; alcohol intake was associated with KD among patients with hypertension alone; age and female were associated with KD among patients with DM alone.

## Discussion

To our knowledge, this is the first study with large sample size to investigate the prevalence and associated factors of KD on population with hypertension and/or DM in a primary care setting in Northwest China. The present study showed that prevalence of KD was higher in the population with both hypertension and DM than in the population with hypertension or DM alone. There were significant differences between genders, with higher prevalence in women. Age was positively correlated with KD. In addition, there were also differences of the prevalence of KD among population from different living regions and ethnic groups. Age, gender, living region, physical activity, duration of hypertension or DM, anemia, FPG and TG were independently associated with KD in the population with hypertension and/or DM.

Previous studies investigating kidney disease in China focused mainly on general or elderly population. Zhang et al reported in a nationally representative investigation that prevalence of eGFR <60 mL/min/per  $1.73m^2$  in Chinese aged  $\geq 18$  years is 1.7%.<sup>5</sup> It is 1.7% and 2.1% in population from Beijing<sup>19</sup> and Tibet,<sup>20</sup> respectively. This is similar to other developing countries,<sup>21,22</sup> whereas lower than developed countries (6.11% in Sweden and 6.5% in America).<sup>23,24</sup> Hypertension and DM have been demonstrated as the main pathologies contributing to kidney disease.<sup>25</sup> In the past two decades, hypertension and DM has increased rapidly in China,<sup>12</sup> indicating that

Characteristics		Hypertens	ion and DM	Hypertension Only		DM	DM Only	
		Prevalence I	Prevalence 2	Prevalence I	Prevalence 2	Prevalence I	Prevalence 2	
Sex	Female	11.46 (9.36–13.84)	17.93 (15.36–20.73)	4.88 (4.13–5.73)	8.70 (7.70–9.79)	5.58 (3.96–7.61)	10.25 (8.05-12.82)	
	Male	5.73 (4.47–7.80)	7.41 (5.73–9.40)	1.87 (1.42–2.42)	2.14 (1.65–2.72)	4.87 (3.37–6.77)	5.46 (3.87–7.44)	
	P-value	<0.001	<0.001	<0.001	<0.001	0.557	0.001	
Age	< 30	0	0	1.87 (0.23–6.59)	0.93 (0.02-5.10)	3.51 (0.43–12.11)	3.51 (0.43–12.11)	
	30–59	5.14 (3.62–7.04)	5.85 (4.23-7.85)	2.15 (1.69–2.69)	3.02 (2.47-3.64)	4.15 (2.97-5.63)	5.11 (3.79–6.71)	
	≥ 60	11.36 (9.41–13.55)	17.67 (15.29–20.24)	5.20 (4.34-6.18)	9.04 (7.91–10.27)	8.43 (5.72–11.88)	15.99 (12.28–20.29)	
	P-value	<0.001	<0.001	<0.001	<0.001	0.008	<0.001	
Living region	Urban	11.18 (9.07–13.59)	14.76 (12.33–17.41)	5.38 (4.35–6.56)	7.27 (6.07–8.61)	8.63 (6.32–11.45)	12.85 (10.04–16.12)	
	Stock-raising	4.82 (1.33–11.88)	6.02 (1.98–13.5)	1.98 (1.14–3.19)	3.71 (2.52–5.26)	3.26 (0.68-9.23)	5.43 (1.79–12.23)	
	Agriculture	6.61 (4.97–8.57)	11.18 (9.07–13.59)	2.68 (2.16-3.28)	4.82 (4.13–5.60)	3.20 (2.06–4.72)	4.79 (3.38–6.57)	
	P-value	0.002	0.019	<0.001	<0.001	<0.001	<0.001	
Ethnicity	Han	9.61 (7.95–11.48)	14.32 (12.33–16.51)	3.76 (3.06–4.57)	5.97 (5.09-6.96)	5.65 (4.21–7.40)	8.88 (7.07-10.98)	
	Kazakh	7.17 (4.37–10.97)	9.06 (5.89-13.18)	2.84 (2.20-3.59)	4.70 (3.88–5.63)	4.42 (2.23–7.77)	5.62 (3.11-9.25)	
	Others	6.34 (3.74–9.96)	8.96 (5.82–13.03)	3.55 (2.47-4.92)	5.42 (4.06-7.05)	4.44 (2.15–8.02)	6.22 (3.44–10.22)	
	P-value	0.148	0.010	0.184	0.139	0.630	0.148	
Educational status	Primary and lower	9.14 (7.15–11.46)	14.73 (12.25–17.51)	3.38 (2.73-4.14)	6.35 (5.45–7.34)	5.66 (3.76–8.13)	9.22 (6.78–12.19)	
	Junior high	8.21 (6.08-10.80)	11.43 (8.91–14.36)	3.34 (2.59-4.23)	4.70 (3.81-5.73)	4.28 (2.73-6.35)	5.95 (4.10-8.29)	
	Senior high and higher	8.52 (5.86–11.87)	10.16 (7.26–13.74)	3.33 (2.41–4.48)	4.36 (3.30–5.64)	6.13 (3.79–9.32)	8.90 (6.04–12.53)	
	P-value	0.835	0.058	0.955	0.010	0.425	0.109	

Table I Prevalence of Kidney Dysfunction in the Study Population

Notes: Data was showed as percentage and 95% confidence interval. Prevalence 1: defined as eGFR <60 calculated by MDRD equation; prevalence 2: defined as eGFR <60 calculated by CKD-EPI equation. Abbreviation: DM, diabetes mellitus.

prevalence of kidney diseases at population level may have been affected. Cao et al reported that diabetes and hypertension have become leading causes of CKD in elderly Chinese patients.<sup>11</sup> It can be expected that prevalence of CKD will increase rapidly, with the acceleration of aging and the expansion of patients with hypertension and DM. Screening of CKD is cost-effective for patients with DM or hypertension in the US.<sup>26</sup> Therefore, it may be effective for reducing public health burden in the primary care-setting to take screening and preventative measures for kidney disease in this high-risk population in developing countries, such as China.

Age has been indicated to be independently associated with CKD in the general population; it is also an independent risk factor for KD in patients with hypertension and DM in our study. In normal individuals, there is progressive deterioration of renal function as decrease in renal mass and increased incidence of sclerotic glomeruli with age;<sup>27</sup> and this will be exacerbated when complicated with hypertension or DM. Moreover, extension of duration of hypertension and DM can also aggravate kidney damage.

Further studies to evaluate effects of management of hypertension and DM on occurrence and development of kidney disease in this population are needed.

Consistent with previous studies in other countries, which reported that women have a higher prevalence of KD, gender difference is also observed in our study. However, gender-different mechanism in KD remains unclear still. Difference in hormone metabolism and glomerular structure and hemodynamics between male and female may serve as part of the reason.<sup>28</sup> Moreover, imbalance in obesity between genders may also play a role. A meta-analysis showed that obesity in females is associated with a higher risk of kidney disease than in males.<sup>29</sup> In the present study, proportion of obesity is higher in women than in men as well (41.2% vs 35.3%).

National investigation in China showed that prevalence of CKD in rural regions is lower than that in urban regions.<sup>5</sup> Similar results are found in our study, which suggests that living regions may be associated with KD. Environmental factors may be part of the explanation for the higher prevalence in urban region. It has been demonstrated in cohort

 $\label{eq:table 2} \textbf{Table 2} \ \textbf{Characteristics of Participants with Hypertension and/or Diabetes}$ 

Characteristics	Total (n=8899)	eGFR ≥60 (n=8487)	eGFR <60 (n=412)	P-value
Age (years)	56.61±12.11	56.34±12.09	62.22±11.12	<0.001
BMI (kg/m <sup>2</sup> )	27.22±4.33	27.21±4.34	27.46±4.21	0.250
Female	4390 (49.3)	4117 (48.5)	273 (66.3)	<0.001
Systolic BP (mmHg)	143.7±21.5	143.8±21.4	141.2±23.4	0.024
Diastolic BP (mmHg)	87.5±13.1	87.7±13.0	83.9±13.9	<0.001
Waist circumference (cm)	92.0±11.4	92.0±11.4	92.4±11.2	0.464
Central obesity, n (%)	5795 (65.1)	5509 (64.9)	286 (69.4)	0.061
Hypertension, n (%)	7558 (84.9)	7216 (85.0)	342 (83.0)	0.264
Diabetes, n (%)	2998 (33.7)	2784 (32.8)	214 (51.9)	<0.001
Hypertension and DM, n (%)	1657 (18.6)	1513 (17.8)	144 (34.9)	<0.001
CVD, n (%)	881 (9.9)	810 (9.5)	71 (17.2)	<0.001
Region, n (%)				<0.001
Urban	2978 (33.5)	2756 (32.5)	222 (53.9)	
Stock-raising	983 (11.0)	960 (11.3)	23 (5.6)	
Agriculture	4938 (55.5)	4771 (56.2)	167 (40.5)	
Educational status, n (%)				0.734
Primary and lower	3872 (43 5)	3688 (43.5)	184 (447)	
	3076 (34.6)	2941 (347)	135 (32.8)	
Senior high and higher	1951 (21.9)	1858 (21.9)	93 (22.6)	
Occupation, n (%)				0.008
Manual	7650 (86.0)	7314 (86.2)	336 (81.6)	
Intelligent	1249 (14.0)	1173 (13.8)	76 (18.4)	
Ethnicity, n (%)				<0.001
Han	4570 (51.4)	4316 (50.9)	254 (61.7)	
Kazakh	2877 (32.3)	2780 (32.8)	97 (23.5)	
Others	1452 (16.3)	1391 (16.4)	61 (14.8)	
Marital status n (%)				0.004
Singlo	249 (2.8)	243 (2.9)	6 (15)	0.004
Magnie	74(1 (02.0)	ZTJ (Z.7)	221 (00 2)	
Farried	/401 (03.0)	7130 (84.0)	331 (00.3) 75 (19.3)	
Separated	1187 (13.4)	1114 (13.1)	75 (18.2)	
Smoke, n (%)	3060 (34.4)	2959 (34.9)	101 (24.5)	<0.001
Alcohol intake, n (%)	2888 (32.5)	2790 (32.9)	98 (23.8)	<0.001
Physical activity, n (%)				0.001
Low	1434 (16.1)	1347 (15.9)	87 (21.1)	
Median	4997 (56.2)	4759 (56.1)	238 (57.8)	
High	2468 (27.7)	2381 (28.1)	87 (21.1)	
Sleep quality, n (%)				<0.001
Very good	4515 (50.7)	4346 (51.2)	169 (41.0)	
Fair good	2992 (33.6)	2836 (33.4)	156 (37.9)	
Fair bad	1138 (12.8)	1063 (12 5)	75 (182)	
Very bad	254 (2.9)	242 (2.9)	12 (2 9)	
	2JT (2.7)	272 (2.7)	12 (2.7)	

(Continued)

#### Table 2 (Continued).

Characteristics	Total (n=8899)	eGFR ≥60 (n=8487)	eGFR <60 (n=412)	P-value
Hemoglobin (g/L)	139.5±20.6	139.7±20.6	136.4±20.3	0.001
Anemia, n (%)	570 (6.4)	535 (6.3)	35 (8.5)	0.076
Dyslipidemia, n (%)	3958 (44.5)	3729 (43.9)	229 (55.6)	<0.001
FPG (mmol/L)	6.44±2.80	6.38±2.75	7.54±3.59	<0.001
TC (mmol/L)	5.08±1.29	5.08±1.30	5.14±1.22	0.352
TG (mmol/L)	1.40 (0.99–2.00)	1.36 (0.93–1.94)	1.70 (1.18–2.40)	<0.001

Abbreviations: eGFR, estimated glomerular filtration rate; BP, blood pressure; DM, diabetes mellitus; CVD, cardiovascular disease; FPG, fasting plasma glucose; TC, total cholesterol; TG, triglyceride.

Table 3 Factors	Associated with	Kidney Dysfunctior	Among Patients with	Hypertension and Diabetes

Factors	Adjusted OR	95%CI	P-value
Age	1.037	1.016-1.059	0.001
Female	2.086	1.408–3.090	<0.001
BMI (vs <24.0)			
24.0–27.9	0.645	0.378–1.098	0.106
≥28.0	0.937	0.559–1.570	0.805
Region (vs urban)			
Stock-raising	0.608	0.206–1.791	0.366
Agriculture	0.731	0.483-1.105	0.137
Ethnicity (vs Han)			
Kazakh	0.985	0.551–1.664	0.879
Others	0.620	0.356-1.078	0.090
Marital status (vs married)			
Single	1.298	0.164–10.25	0.805
Separated	1.058	0.655–1.709	0.819
Occupation (vs manual)			
Intelligent	1.463	0.929–2.302	0.100
CVD	1.326	0.858–2.049	0.204
Physical activity (vs high)			
Median	1.268	0.757–2.123	0.366
Low	1.188	0.650-2.173	0.576
Control of hypertension	1.093	0.726-1.645	0.641
Duration of hypertension	1.022	1.001-1.042	0.036
Anemia	2.621	1.296–5.299	0.007
FPG	1.056	1.005–1.110	0.031
тс	0.908	0.782-1.055	0.208
TG	1.146	1.049–1.251	0.002

Abbreviations: BMI, body mass index; CVD, cardiovascular disease; FPG, fasting plasma glucose; TC, total cholesterol; TG, triglyceride. OR, odds ratio; CI, confidence interval.

 Table 4 Factors Associated with Kidney Dysfunction Among Patients with Only Hypertension

Factors	Adjusted OR	95%CI	P-value
Age	1.042	1.026–1.059	<0.001
Female	3.016	1.991–4.569	<0.001
BMI (vs <24.0)			
24.0–27.9	0.825	0.556-1.226	0.341
≥28.0	0.926	0.627–1.366	0.697
Systolic blood pressure	0.995	0.984–1.006	0.373
Diastolic blood pressure	0.999	0.984-1.015	0.932
Region (vs urban)			
Stock-raising	0.437	0.247–0.776	0.005
Agriculture	0.506	0.361–0.709	<0.001
Education (vs senior high and			
higher)			
Junior high	1.328	0.876–2.014	0.181
Primary and lower	1.020	0.666–1.563	0.928
Physical activity (vs high)			
Median	0.650	0.460–0.917	0.014
Low	1.056	0.693–1.609	0.800
Sleep quality (vs very good)			
Fair good	1.162	0.835-1.617	0.373
Fair bad	1.164	0.762–1.779	0.482
Very bad	0.567	0.221–1.455	0.238
Control of hypertension	1.253	0.797–1.968	0.328
Duration of hypertension	1.028	1.009–1.048	0.005
Alcohol	1.379	0.897–2.120	0.143
Anemia	I.854	1.147–2.996	0.012
Dyslipidemia	1.281	0.920-1.783	0.142
FPG	1.409	1.057–1.877	0.019
тс	0.987	0.868–1.122	0.842
TG	1.078	0.964–1.205	0.187

Abbreviations: BMI, body mass index; FPG, fasting plasma glucose; TC, total cholesterol; TG, triglyceride. OR, odds ratio; CI, confidence interval.

studies that increase in the  $PM_{2.5}$  is associated with higher risk of CKD and ESRD.<sup>30,31</sup> A study from Taipei also shows that there is positive association between traffic-related air pollution and kidney damage.<sup>32</sup> In addition, differences in lifestyle and dietary habits between living regions may also be involved in the process.

The relationship between physical activity and KD remains inconsistent, and it is varied among different patient groups in the present study. Exercise may protect renal function by lowering BMI and reducing the risk of hypertension and DM,<sup>33</sup> and this indirect effect may be attenuated in patients who have been suffering from hypertension and DM. Recently, a cohort study shows that highly active people have lower risk for developing CKD, compared with inactive people.<sup>34</sup> However, Tannor et al reported that there is no independent association between physical activity and CKD in Ghanaian patients with hypertension and DM.<sup>35</sup> Among the nondiabetic population, FPG and prediabetes are significantly associated with coronary artery disease and

Factors	Adjusted OR	95%CI	P-value
Age	1.023	0.999–1.048	0.064
Female	1.220	0.740-2.012	0.436
Duration of diabetes	1.040	1.001-1.081	0.047
Region (vs urban) Stock-raising Agriculture	0.427 0.423	0.127–1.438 0.250–0.715	0.169 0.001
Marital status (vs married) Single Separated	1.424 0.798	0.307–6.616 0.325–1.958	0.652 0.798
Physical activity (vs high) Median Low	2.409 2.000	1.115–5.207 0.805–4.966	0.025 0.135

Fable 5         Factors         Associated         with         Kidne	y Dysfunction Among	Patients with	Only Diabetes
---	---------------------	---------------	---------------

Abbreviations: OR, odds ratio; Cl, confidence interval.

stroke.<sup>36</sup> It is noteworthy that FPG is independently associated with CKD in hypertensive patients who are without DM in the present study. Glycemic control can help prevent onset of CKD in individuals with DM.<sup>37</sup> However, it needs to be further explored in hypertensive patients. Consistent with a previous study performed in elderly population,<sup>38</sup> the present study observed that anemia is significantly associated with KD. Anemia is a common comorbidity in patients with chronic kidney disease, mainly due to impaired production of erythropoietin by kidney damage; hepcidin and other factors are also involved.<sup>39</sup> Patients with anemia should be aware of presence of kidney disease. In addition, the association between anemia and the risk of incident CKD needs to be verified in a longitudinal study.

As expected, prevalence of KD in our study is significantly higher for CKD-EPI analysis than for MDRD analysis. In sensitivity analysis, occupation, history of CVD and alcohol intake are significantly associated with KD, which may be due to the larger sample size of KD cases based on CKD-EPI equation. These factors have been shown to be directly or indirectly associated with KD in patients with DM or kidney diseases.<sup>40,41</sup>

# Limitations

Although multistage random sampling in general population with a good sample size, several limitations warrant discussion for our study. First, causal inference between associated factors and KD needs to be further explored, because of the cross-sectional design of the present study. Second, only one county was selected, limiting the generalization of the results, although the county has a good representation. Third, single measurements of serum creatinine without repeated examination may have resulted in misclassification of individuals with KD. Fourth, because of the lack of urine test, we were only able to report prevalence of KD defined by eGFR, regardless of urinary protein. Fifth, although LASSO regression was used for screening potential factor, it is difficult to find all associated factors for KD; in addition, mediating effects between variables were also failed to resolve. Future study needs to take this into account. Sixth, although recognized methods applied for blood pressure and glucose measurements, diagnosis of hypertension and DM might be biased due to lack of repeated measurements from different days. In addition, oral glucose tolerance test should also be considered for DM diagnosis in future investigations.

# Conclusion

There is high prevalence of KD among patients with hypertension and DM in the primary care setting in Northwest China. Modifiable factors associated with KD among patients with hypertension and/or DM include living region, physical activity, anemia, FPG and TG, which may provide clues for prevention and intervention for kidney disease in this population in order to reduce the burden of public health care in less-developed regions.

# **Data Sharing Statement**

The datasets are available from the corresponding author upon reasonable request.

#### **Ethical Approval**

The present study was approved by the Ethics Committee at the People's Hospital of Xinjiang Uygur Autonomous Region (IRB approval number: 2019020327). Informed consent was obtained from all individual participants included in the study.

#### Acknowledgments

The authors are very much grateful to all study participants for their kind cooperation/participation in the study.

#### Funding

This work is partly funded by National Key R&D Program of China (grant number: 2018YFC1311503), and National Health Committee Key Laboratory of Hypertension Clinical Research (grant number: 2019[155]).

#### Disclosure

The authors report no conflicts of interest in this work.

#### References

- Herzog CA, Asinger RW, Berger AK, et al. Cardiovascular disease in chronic kidney disease. A clinical update from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney Int.* 2011;80 (6):572–586. doi:10.1038/ki.2011.223
- Jha V, Garcia-Garcia G, Iseki K, et al. Chronic kidney disease: global dimension and perspectives. *Lancet*. 2013;382(9888):260–272. doi:10.1016/S0140-6736(13)60687-X
- Mills KT, Xu Y, Zhang W, et al. A systematic analysis of worldwide population-based data on the global burden of chronic kidney disease in 2010. *Kidney Int.* 2015;88(5):950–957. doi:10.1038/ki.2015.230
- Hill NR, Fatoba ST, Oke JL, et al. Global prevalence of chronic kidney disease - A systematic review and meta-analysis. *PLoS One*. 2016;11(7):e0158765. doi:10.1371/journal.pone.0158765
- Zhang L, Wang F, Wang L, et al. Prevalence of chronic kidney disease in China: a cross-sectional survey. *Lancet*. 2012;379 (9818):815–822. doi:10.1016/S0140-6736(12)60033-6
- Wei H, Yan Y, Gong J, Dong J. Prevalence of kidney damage in Chinese elderly: a large-scale population-based study. *BMC Nephrol*. 2019;20(1):341. doi:10.1186/s12882-019-1525-5
- Huang YM, Xu D, Long J, et al. Spectrum of chronic kidney disease in China: a national study based on hospitalized patients from 2010 to 2015. *Nephrology*. 2019;24(7):725–736. doi:10.1111/nep.13489
- Laliberté F, Bookhart BK, Vekeman F, et al. Direct all-cause health care costs associated with chronic kidney disease in patients with diabetes and hypertension: a managed care perspective. J Manag Care Pharm. 2009;15(4):312–322. doi:10.18553/jmcp.2009.15.4.312
- Ohta M, Babazono T, Uchigata Y, Iwamoto Y. Comparison of the prevalence of chronic kidney disease in Japanese patients with Type 1 and Type 2 diabetes. *Diabet Med.* 2010;27(9):1017–1023. doi:10.1111/j.1464-5491.2010.03049.x
- Altun B, Süleymanlar G, Utaş C, et al. Prevalence, awareness, treatment and control of hypertension in adults with chronic kidney disease in Turkey: results from the CREDIT study. *Kidney Blood Press Res.* 2012;36(1):36–46. doi:10.1159/000339025

- Cao Y, Li W, Yang G, Liu Y, Li X. Diabetes and hypertension have become leading causes of CKD in Chinese elderly patients: a comparison between 1990–1991 and 2009–2010. *Int Urol Nephrol.* 2012;44(4):1269–1276. doi:10.1007/s11255-012-0194-0
- Zhang L, Wang H. Chronic kidney disease epidemic: cost and health care implications in China. *Semin Nephrol.* 2009;29(5):483–486. doi:10.1016/j.semnephrol.2009.06.012
- Zhou M, Astell-Burt T, Bi Y, et al. Geographical variation in diabetes prevalence and detection in China: multilevel spatial analysis of 98,058 adults. *Diabetes Care*. 2015;38(1):72–81. doi:10.2337/dc14-1100
- Wang Z, Chen Z, Zhang L, et al. Status of hypertension in China: results from the China hypertension survey, 2012–2015. *Circulation*. 2018;137 (22):2344–2356. doi:10.1161/CIRCULATIONAHA.117.032380
- Ma YC, Zuo L, Chen JH, et al. Modified glomerular filtration rate estimating equation for Chinese patients with chronic kidney disease. J Am Soc Nephrol. 2006;17(10):2937–2944. doi:10.1681/ ASN.2006040368
- Lv JC, Zhang LX. Prevalence and disease burden of chronic kidney disease. Adv Exp Med Biol. 2019;1165:3–15.
- World Health Organization. Global Recommendations on Physical Activity for Health. Geneva: World Health Organization Copyright © World Health Organization 2010; 2010.
- Buysse DJ, Reynolds CF 3rd, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res.* 1989;28(2):193–213. doi:10.1016/0165-1781(89)90047-4
- Zhang L, Zhang P, Wang F, et al. Prevalence and factors associated with CKD: a population study from Beijing. *Am J Kidney Dis.* 2008;51(3):373–384. doi:10.1053/j.ajkd.2007.11.009
- Chen W, Liu Q, Wang H, et al. Prevalence and risk factors of chronic kidney disease: a population study in the Tibetan population. *Nephrol Dial Transplant*. 2011;26(5):1592–1599. doi:10.1093/ndt/gfq608
- 21. Varma PP, Raman DK, Ramakrishnan TS, Singh P, Varma A. Prevalence of early stages of chronic kidney disease in apparently healthy central government employees in India. *Nephrol Dial Transplant*. 2010;25(9):3011–3017. doi:10.1093/ndt/gfq131
- 22. Alam A, Amanullah F, Baig-Ansari N, Lotia-Farrukh I, Khan FS. Prevalence and risk factors of kidney disease in urban Karachi: baseline findings from a community cohort study. *BMC Res Notes*. 2014;7(1):179. doi:10.1186/1756-0500-7-179
- Gasparini A, Evans M, Coresh J, et al. Prevalence and recognition of chronic kidney disease in Stockholm healthcare. *Nephrol Dial Transplant*. 2016;31(12):2086–2094. doi:10.1093/ndt/gfw354
- 24. Wang F, He K, Wang J, et al. Prevalence and risk factors for CKD: a comparison between the adult populations in China and the United States. *Kidney Int Rep.* 2018;3(5):1135–1143. doi:10.1016/j.ekir.201 8.05.011
- Nugent RA, Fathima SF, Feigl AB, Chyung D. The burden of chronic kidney disease on developing nations: a 21st century challenge in global health. *Nephron Clin Pract.* 2011;118(3):c269–c277. doi:10.1159/000321382
- Hoerger TJ, Wittenborn JS, Segel JE, et al. A health policy model of CKD: 2. The cost-effectiveness of microalbuminuria screening. *Am J Kidney Dis.* 2010;55(3):463–473. doi:10.1053/j.ajkd.2009.11.017
- Lindeman RD, Tobin J, Shock NW. Longitudinal studies on the rate of decline in renal function with age. J Am Geriatr Soc. 1985;33 (4):278–285. doi:10.1111/j.1532-5415.1985.tb07117.x
- Silbiger SR, Neugarten J. The role of gender in the progression of renal disease. Adv Ren Replace Ther. 2003;10(1):3–14. doi:10.1053/ jarr.2003.50001
- Wang Y, Chen X, Song Y, Caballero B, Cheskin LJ. Association between obesity and kidney disease: a systematic review and meta-analysis. *Kidney Int.* 2008;73(1):19–33. doi:10.1038/sj. ki.5002586

- 30. Chan TC, Zhang Z, Lin BC, et al. Long-term exposure to ambient fine particulate matter and chronic kidney disease: a cohort study. *Environ Health Perspect.* 2018;126(10):107002. doi:10.1289/ EHP3304
- Bowe B, Xie Y, Li T, Yan Y, Xian H, Al-Aly Z. Particulate matter air pollution and the risk of incident CKD and progression to ESRD. *J Am Soc Nephrol.* 2018;29(1):218–230. doi:10.1681/ASN.2017 030253
- Chen SY, Chu DC, Lee JH, Yang YR, Chan CC. Traffic-related air pollution associated with chronic kidney disease among elderly residents in Taipei City. *Environ Pollut*. 2018;234:838–845. doi:10.1016/ j.envpol.2017.11.084
- 33. Park S, Lee S, Kim Y, et al. Causal effects of physical activity or sedentary behaviors on kidney function: an integrated population-scale observational analysis and Mendelian randomization study. *Nephrol Dial Transplant*. 2021. doi:10.1093/ndt/gfab153
- 34. Parvathaneni K, Surapaneni A, Ballew SH, et al. Association between midlife physical activity and incident kidney disease: the Atherosclerosis Risk in Communities (ARIC) study. *Am J Kidney Dis.* 2021;77(1):74–81. doi:10.1053/j.ajkd.2020.07.020
- 35. Tannor EK, Sarfo FS, Mobula LM, Sarfo-Kantanka O, Adu-Gyamfi R, Plange-Rhule J. Prevalence and predictors of chronic kidney disease among Ghanaian patients with hypertension and diabetes mellitus: a multicenter cross-sectional study. J Clin Hypertens. 2019;21(10):1542–1550. doi:10.1111/jch.13672

- Mutie PM, Pomares-Millan H, Atabaki-Pasdar N, et al. An investigation of causal relationships between prediabetes and vascular complications. *Nat Commun.* 2020;11(1):4592. doi:10.1038/s41467-020-18386-9
- James MT, Hemmelgarn BR, Tonelli M. Early recognition and prevention of chronic kidney disease. *Lancet.* 2010;375 (9722):1296–1309. doi:10.1016/S0140-6736(09)62004-3
- 38. Ji A, Pan C, Wang H, et al. Prevalence and associated risk factors of chronic kidney disease in an elderly population from Eastern China. *Int J Environ Res Public Health.* 2019;16(22):22. doi:10.3390/ ijerph16224383
- Atkinson MA, Warady BA. Anemia in chronic kidney disease. *Pediatr Nephrol.* 2018;33(2):227–238. doi:10.1007/s00467-017-3663-y
- 40. Moazzeni SS, Arani RH, Hasheminia M, Tohidi M, Azizi F, Hadaegh F. High incidence of chronic kidney disease among Iranian diabetic adults: using CKD-EPI and MDRD equations for estimated Glomerular Filtration Rate. *Diabetes Metab J.* 2021;45 (5):684–697. doi:10.4093/dmj.2020.0109
- Kaze FF, Maimouna M, Beybey AF, et al. Prevalence and determinants of chronic kidney disease in urban adults' populations of northern Cameroon. *Clin Exp Nephrol.* 2021;25(7):718–726. doi:10.1007/ s10157-021-02036-5

#### **International Journal of General Medicine**

#### **Dove**press

Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/ testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/international-journal-of-general-medicine-journal