Refracture of the tibia 6 years after bilateral tibia nail removal in a patient later diagnosed with osteopetrosis

Joost H van Linge^{1,2}

JHJM Gert Bessems¹

P Koen Bos¹

¹Department of Orthopaedic Surgery, Erasmus MC, Rotterdam, the Netherlands; ²Reinier de Grant Gasthuis, Ga Delft, the Netherlands

Abstract: Osteopetrosis is a rare bone disease with a high fracture incidence and a risk for recurrent fractures. We describe a case report of an adult female with a refracture of her right tibia 6 years after routine bilateral tibia nail removal. The patient and 3 of her family members had a history of multiple (stress) fractures. Later we diagnosed autosomal dominant osteopetrosis in our patient and her family. In addition we give an overview of osteopetrosis and our considerations regarding routine removal of intramedullary implants in these patients.

Keywords: fracture, stress fracture, pathological fracture, intramedullary nailing

Introduction

Intramedullary nailing is widely accepted as standard treatment for closed, unstable tibial fracture, open tibial fractures, and tibial stress fractures.¹⁻³ There is no consensus, however, concerning absolute or relative indications for intramedullary nail removal after consolidation of the fracture. The decision to remove nails has largely been based on symptoms, routine treatment, or patient's preference.⁴

There is only one paper in literature describing serious peri- or intra-articular injuries following high velocity injuries in patients with a femoral nail in place.⁵ But does this justify routine removal? Anterior knee pain may form an indication for tibial nail removal.⁶ However, tibial nail removal for anterior knee pain may lead to disappointing results and can even provoke anterior knee pain in previous asymptomatic patients.^{7,8}

There is a potential harm in routine removal of metal implants such as wound infections, deep venous thrombosis, excessive blood loss, nerve damage to the infrapatellar branch of the saphenous nerve, and refracturing.⁸ Nail removal can be a difficult procedure due to bony overgrowth, broken screws, or unknown nails that were implanted elsewhere. To prevent further damage during extraction, nail retention has even been reported.^{7,9,10}

We present a case of refracture of the tibia 6 years after removal of an intramedullary nail, originally implanted for a fractured tibia. The patient had a history of stress fractures and a family history of recurrent fractures. Although the increased risk for refracture in metabolic or hereditary brittle bone disease is well known, in the intramedullary nail literature, there is little attention for evaluation of bone diseases during decision making for nail extraction.

Case report

A 27-year-old woman was treated with an intramedullary nail for a fractured left tibia in an area of a stress fracture (Figure 1). Her right tibia was treated with a nail one year later for a symptomatic stress fracture (Figure 2). Because of her young age and

Correspondence: J H van Linge
Erasmus MC, Department of
Orthopaedic Surgery,
Postbus 2040, 3000 CA Rotterdam,
the Netherlands
Tel: +31-10 10 – 703 3537
Fax: +31-10 10 – 703 2001
Email j.vanlinge@erasmusmc.nl

Figure 1 Fracture of left tibia and fibula at the site of a stress fracture 8 years prior to her presentation in 2005 **A)** before surgery, **B)** after intramedullary nailing, and **C)** 1 year after surgery showing tardy consolidation.

minor complaints, both implants were removed simultaneously after the fractures healed well.

Six years later, in 2005 at age 35 when she was 30 weeks pregnant, she was admitted following a minor fall in which she sustained a subtrochanteric femoral fracture on the left side and a refracture of her right tibia in the previous fractured area. The femoral fracture was stabilized with a long Gamma nail (Stryker, Mahwah, New Jersey, USA). This was a challenging procedure that took more than 3 hours because of complete obliteration of the femoral shaft. Because of her pregnancy and because we expected a demanding second procedure we decided to treat her tibia fracture conservatively with reposition and cast immobilization. Unfortunately, she developed a skin necrosis due to secondary angulation in the cast. 10 weeks later, after she gave birth to a son, the fracture of her right tibia was still not consolidated. We treated her fracture with a reamed intramedullary tibia nail (Figure 3). At the time of surgery we thought the infection healed but nevertheless it was complicated by a deep infection, prob-

ably due to the skin necrosis around the fracture site. This pre-tibial skin defect was treated by a fasciocutaneous flap according to Pontén. The fracture healed well but the nail had to be removed and the intramedullary infection was treated with intramedullary gentamicin beads. Because of the history of recurrent (stress) fractures and patient's preference to reduce the risk of future fractures, we decided to implant two reamed tibia nails. Recently, we treated her right femur with a long gamma nail because of a fracture after a minor fall (Figure 4).

Family history and counseling

With the fractures mentioned above and no significant trauma we decided to take a detailed history of the patient and her family. It appeared that she had suffered tibial stress fractures at age 22 and an insufficiency fracture of the foot at age 30. We also discovered that we treated her aunt some years ago because of multiple (stress) fractures. Despite the earlier efforts of the endocrinology and pathology department to

Figure 2 A) and B) Stress fracture of right tibia 1 year later after surgery. An intramedullary nail was placed and a biopsy was performed.

come to a diagnosis for her aunt, only an osteopetrosis-like disease was diagnosed mainly based on histology. The father and sister of our patient had multiple fractures as well. The patient's two brothers did not sustain any fractures.

Taking into account the patient's and family history with multiple fractures, we consulted the clinical genetics department. A detailed family history was taken, and additional X-rays were made to evaluate radiological signs. Medullary canal narrowing and subtle skull density was found, but no other typical radiological features were present. The Dutch Skeletal Dysplasia Group judged that there was enough evidence for diagnosing osteopetrosis with an autosomal dominant heritage pattern. Further evaluation did not reveal evidence for another cause of brittle bones (osteoporosis, endocrine pathology, or osteogenesis imperfecta).

Osteopetrosis

Osteopetrosis is a heterogeneous group of heritable conditions characterized by defective osteoclast resorption leading to

hard and brittle bone.^{11–14} Osteopetrosis is a rare bone disease (prevalence varies from 3.3 per million for the malignant form to 55 per million for the benign form); it is also known as Albers-Schönberg disease or marble bone disease.^{11,12} The malfunction of the osteoclast's capacity to resorb bone leads to osteosclerosis and increased brittleness of the bones. Before identification of genes that affect the function of osteoclast, three types of osteopetrosis were classified based on clinical aspects: the first type is the infantile or 'malignant' osteopetrosis, an early autosomal recessive form characterized by presentation at a young age and severe symptoms. The obliterated medullary cavity leads to an inadequate hematopoietic capacity. The skull bones show serious abnormalities with cranial nerve dysfunction. These infants die at very young age. The second type is the intermediate autosomal recessive form which gives a higher fracture risk and mild hematopoietic disturbance and less cranial nerve dysfunction than in the malignant form. The third type is the 'benign' or late form. This type is autosomal dominant and the most common. There

Figure 3 A) Right tibial refracture at presentation in 2005. B) Secondary dislocation in plaster cast 10 weeks later. C) After insertion of a reamed intramedullary nail.

are two subtypes: type I, with marked sclerosis of the vault but a normal fracture risk; and type II, with sclerosis of the base of the skull, rugger jersey vertebrae, and endobones in the pelvis. This subtype has a higher fracture risk.¹⁵ Autosomal dominant osteopetrosis has a genetic basis; however, some individuals do not develop clinical features because there is a 75% penetrance.¹² Genetic abnormalities in the chloride channel gene, proton pump and carbonic anhydrase II were identified. They have different effects on the osteoclast physiology, but all cause a decreased bone resorption.

On histology, osteopetrosis bone shows lack of remodeling during bone development. Lack of osteoclastic bone resorption leads to sclerotic cancellous bone. Calcified cartilage and thickened trabeculae in diaphyseal bone explain the mechanical inferior bone quality and increased fracture risk. Clinically, patients present with frequent fractures (commonly stress fractures; life-long prevalence of fractures varies from 40%–66.6%), bone deformity, osteoarthritis, osteomyelitis, spondylolysis, and cranial nerve palsies (present in 20%–25%). Fracture healing has been described as prolonged.^{12,16–18} Radiological findings include generalized sclerosis, bone-within-bone appearance (endobone),

medullary canal narrowing, skull base thickening skull density, and vertebral endplate thickening (rugger jersey spine).

Surgical fracture treatment is complicated by extremely hard and brittle bone. Drilling and reaming during internal fixation or arthroplasty can be very challenging. Therefore, nonoperative treatment should be considered when reasonably possible. Although intramedullary fracture fixation of long bones may be a challenge due to medullary canal obliteration, it may be the treatment of choice because of the long-term strength and stability.

Discussion

As we discussed above, the need for routine removal of (intramedullary) osteosynthesis devices remains controversial. We have shown that removal of an intramedullary tibia nail, in a patient later diagnosed with autosomal dominant osteopetrosis, resulted in a refracture after 6 years. The risk of these refractures leaves us a difficult clinical decision whether to remove the intramedullary nails or not. Our patient was seriously invalidated during the last weeks of her pregnancy, and the subsequent fracture treatment resulted in prolonged hospital stay and multiple

Figure 4 Right femoral fracture 3 years later: **A)** showing a narrow intramedullary canal and thickening of the cortex; and **B)** after surgery.

operations. In our department, the current attitude is that we don't routinely remove intramedullary nails. Relative indications for nail removal are pain related to hardware, young and active patients, and impending need for arthroplasty.

The increased risk for multiple fractures and refractures with brittle bone disease is well described. Until now, standard evaluation of possible presence of brittle bone disease (the risk for multiple fractures) was not emphasized in the intramedullary nail literature.

Conclusion

Based on this case history we advise to pay attention to the patient's and their family's fracture history before deciding to remove (intramedullary) implants. With a suspected metabolic or hereditary brittle bone disease we advise our patients to leave the device in situ to prevent recurrent fractures and subsequent morbidity.

Acknowledgment

The authors wish to thank Marleen E.H. Simon MD, Department of Clinical Genetics of the Erasmus MC.

Disclosure

The authors report no conflicts of interest in this work.

References

- Schmidt AH, Finkemeier CG, Tornetta P 3rd. Treatment of closed tibial fractures. *Instr Course Lect.* 2003;52:607–622.
- Court-Brown CM, McQueen MM, Quaba AA, Christie J. Locked intramedullary nailing of open tibial fractures. *J Bone Joint Surg Br.* 1991;73(6):959–964.
- Varner KE, Younas SA, Lintner DM, Marymont JV. Chronic anterior midtibial stress fractures in athletes treated with reamed intramedullary nailing. *Am J Sports Med.* 2005;33(7):1071–1076.
- Karladani AH, Ericsson PA, Granhed H, Karlsson L, Nyberg P. Tibial intramedullary nails – should they be removed? A retrospective study of 71 patients. *Acta Orthop.* 2007;78(5):668–671.

5. Sen RK, Gul A, Aggarwal S, Kumar SR, Nagi ON. Comminuted refracture of the distal femur and condyles in patients with an intramedullary nail: a report of 5 cases. *J Orthop Surg (Hong Kong)*. 2005;13(3):290–295.
6. Court-Brown CM, Gustilo T, Shaw AD. Knee pain after intramedullary tibial nailing: its incidence, etiology, and outcome. *J Orthop Trauma*. 1997;11(2):103–105.
7. Boerger TO, Patel G, Murphy JP. Is routine removal of intramedullary nails justified. *Injury*. 1999;30(2):79–81.
8. Karladani AH, Styf J. Percutaneous intramedullary nailing of tibial shaft fractures: a new approach for prevention of anterior knee pain. *Injury*. 2001;32(9):736–739.
9. Takakuwa M, Funakoshi M, Ishizaki K, Aono T, Hamaguchi H. Fracture on removal of the ACE tibial nail. *J Bone Joint Surg Br*. 1997;79(3):444–445.
10. Im GI, Lee KB. Difficulties in removing ACE tibial intramedullary nail. *Int Orthop*. 2003;27(6):355–358.
11. Tolar J, Teitelbaum SL, Orchard PJ. Osteopetrosis. *N Engl J Med*. 2004;351(27):2839–2849.
12. Landa J, Margolis N, Di Cesare P. Orthopaedic management of the patient with osteopetrosis. *J Am Acad Orthop Surg*. 2007;15(11):654–662.
13. Karol LA. Skeletal dysplasias, Chapter 30. In: Herring JA, editor. *Tachdjian's Pediatric Orthopaedics*. 4th ed. Elsevier. 2007.
14. Key LL Jr, Rodriguiz RM, Willi SM, et al. Long-term treatment of osteopetrosis with recombinant human interferon gamma. *N Engl J Med*. 1995;332(24):1594–1599.
15. Bhargava A, Blank R. Osteopetrosis. www.emedicine.com.
16. Ashby ME. Total hip arthroplasty in osteoporosis, a report of two cases. *Clin Orthop Relat Res*. 1992 Mar;(276):214–221.
17. Milgram JW, Jasty M. Osteopetrosis. A morphological study of twenty-one cases. *J Bone Joint Surg Am*. 1982;64(6):912–929.
18. Shapiro F. Osteopetrosis, current clinical considerations. *Clin Orthop Relat Res*. 1993 Sep;(294):34–44.

Open Access Surgery

Publish your work in this journal

Open Access Surgery is an international, peer-reviewed, open access journal that focuses on all aspects of surgical procedures and interventions. Patient care around the peri-operative period and patient outcomes post surgery are key topics. All grades of surgery from minor cosmetic interventions to major surgical procedures are covered. Novel techniques

Submit your manuscript here: <http://www.dovepress.com/open-access-surgery-journal>

Dovepress

and the utilization of new instruments and materials, including implants and prostheses that optimize outcomes constitute major areas of interest. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.