Medical Schools' Ophthalmology Course: An Appraisal by Ophthalmology Residents

Yahya Abdulrahman Al-Najmi¹

Ahmed Hussein Subki²

Nazih Suwalih Alzaidi³

Nadeem Shafique Butt⁴

Alaa Abdulhamid Alsammahi⁵

Firas Mohamed Madani⁵

Mohammed Saad Alsallum⁶

Rakan Salah Al-Harbi⁷

Nizar Mohammed Alhibshi⁵

¹Saggaf Eye Center, Jeddah, Saudi Arabia;

²Department of Internal Medicine, King Faisal Specialist Hospital and Research Center, Jeddah, Saudi Arabia;

³Ophthalmology Department, Prince Mansour Military Hospital, At Taif, Saudi Arabia; ⁴Department of Community Medicine, King Abdulaziz University, Jeddah, Saudi Arabia; ⁵Department of Ophthalmology, King Abdulaziz University Hospital, Jeddah, Saudi Arabia;

⁶Department of Neurology, King Abdulaziz Medical City, Jeddah, Saudi Arabia; ⁷Department of Family Medicine, King Fahd Armed Forces Hospital, Jeddah, Saudi Arabia

Objective: To investigate the perception and satisfaction of ophthalmology residents with the currently provided ophthalmology curricula to medical students.

Methods: A cross-sectional survey involving first to fourth year ophthalmology residents (N = 106) from all regions of Saudi Arabia was conducted between December 2018 and February 2019. An online questionnaire explored opinions about the ophthalmology course regarding three dimensions. Firstly, adequacy in covering essential parts of the specialty; secondly, improvements required; and thirdly, effectiveness. A score (0–21) was calculated, indicating the overall suitability of the ophthalmology course. In addition, factors of good overall suitability (score ≥ 10) were analyzed.

Results: Regarding adequacy, respondents opined that the ophthalmology course did not reasonably cover the basic part (35.8%), clinical part (61.3%), common disease (26.4%), and emergencies (39.6%). Concerning improvements required, more than 80% of the participants expressed that the course required to be improved for all its features, including duration (80.2%), objectives (85.8%), content (82.1%), organization (83.0%), and supervision (81.1%). As to effectiveness, half of them deemed the course unhelpful in familiarizing general practitioners with common ophthalmic diseases and emergencies. Overall, the ophthalmology course was generally deemed suitable (score ≥ 10) for only 27.4% of the participants, with no differences across gender, level, or region.

Conclusion: Ophthalmology residents perceived multiple deficits in the current Saudi ophthalmology teaching course. Significant improvements in ophthalmologic curricula are required, besides coping with unprecedented technological advancement in the ophthalmological field.

Keywords: ophthalmology, course, curriculum, academic, specialty, assessment, residents, opinions

Introduction

Patients with ophthalmologic complaints represent a considerable proportion of those presenting to general practice consultants (19%) and emergency departments (20%).^{1,2} Some of these patients potentially present with conditions that require critical evaluation by skilled general practitioners (GPs) or junior doctors to manage and refer patients before the development of unfavorable consequences. Additionally, the exponential growth of the aged population could have been associated with growing numbers of age-related eye diseases.³ Hence, the necessity to provide adequate ophthalmologic skills and knowledge to diagnose and manage ophthalmic pathologies. To achieve this, medical schools should improve the ophthalmologic competence of medical students.

Correspondence: Yahya Abdulrahman Al-Najmi
Saggaf Eye Center, Abdullah Salman St., Al Faiha'a Dist., P.O. Box: 31903, Jeddah, 21418, Saudi Arabia
Tel +966 564844281
Email dr.alnajmi@gmail.com

Nizar Mohammed Alhibshi
Department of Ophthalmology, King Abdulaziz University Hospital,
P.O. Box 50805, Jeddah, 21533, Saudi Arabia
Tel +966 505609950
Email alhibshi@doctor.com

However, the place of ophthalmology within the curricula of undergraduate medical students has been subject to controversial debates for approximately one century.⁴ The first descriptive report about ophthalmology teaching was published by the Council of British Ophthalmologists in 1919,⁴ when undergraduate ophthalmology curricula were investigated regarding their duration and knowledge assessment. Nonetheless, the scientific content of the curriculum, the methods, and the teaching environment were not explored. Since then, ophthalmologic education has received little attention, and the relevant undergraduate rotations have become less significant clinically.⁵ Rather, the clinical skills in general practice rotations have been emphasized at the expense of specialty attachments such as ophthalmology which may have resulted in a significant gap in the ability to perform a direct ophthalmologic examination by medical students. It is, in turn, reflected as a marked deficiency in eye examination performed by GPs.

In 2006, the seminal publication of a series of recommendations regarding ophthalmologic education was published to guide medical schools on their undergraduate curricula.⁶ The International Council of Ophthalmology (ICO)⁶ emphasized the need to develop evidence-based curricula in all medical schools, and their material should be able to support the basic skills and knowledge required for examination and referral to ophthalmology. In addition, the implementation of essential teaching methods, such as lectures, case studies, clinical practice, and integration with other medical subspecialties was also recommended. These subsequently shaped reliable standards for undergraduate education and eye care in multiple medical institutions globally.

In Saudi Arabia, medical practice has undergone drastic changes concomitant with the booming economic growth during the past few decades. New colleges of medicine and multiple modern hospitals have been established in various administrative regions in the kingdom, increasing the number of medical institutions from 3 to 21 colleges between 1985 and 2011.^{7,8} The practice of eye care could be further influenced by the continuous expansion of modern technologies in these institutions and the applied therapeutic advances.⁹ Besides, approximately one in 7 undergraduate medical students aspire for an ophthalmological career, and there was considerable interest in the early academic life regarding ophthalmology-related academic practices and community service activities.¹⁰ However, receiving inadequate education before

graduation may negatively influence future practice. The efficiency of ophthalmologic education in Saudi Arabia has been scarcely investigated. Rather than targeting medical students to assess their curricula, it is acceptable that ophthalmologic residents can better judge the scientific materials provided to undergraduate students. Thus, in line with the worldwide deficits in undergraduate education and the advanced skills required to handle the sophisticated equipment of eye examination effectively, we sought to investigate the perception and satisfaction of ophthalmology residents with the currently provided Saudi ophthalmology curricula to medical students.

Methods

Design and Participants

This cross-sectional study was conducted from December 2018 and February 2019. A nationwide electronic survey included first to fourth-year ophthalmology residents from all regions of Saudi Arabia. A convenience sampling was used to include all eligible participants during the study period. No stratification by region or level was applied. A minimum of 100 participants was acceptable as a sample size. The study was approved by the institutional review board of the bioethics committee of King Abdulaziz University, Jeddah, Saudi Arabia. We affirm that all participants provided informed consent. The study was conducted in accordance with the Declaration of Helsinki.

Study Questionnaire

An electronically administered, semi-structured questionnaire was designed for this study. It contained two parts: A) Participants' characteristics including level (R1, R2, R3, or R4), gender, nationality, university of graduation, and region; B) Assessment of ophthalmology course. Part B explored the participants' opinions about the ophthalmology course regarding three dimensions, including 1) adequacy of the course in covering essential parts of the specialty, namely basic science, clinical exposure, common diseases, and common emergencies (adequacy subscale, 4 items); 2) improvements required in content, objectives, duration, organization or supervision (improvements required subscale, 5 items); and 3) effectiveness of the course in familiarizing GPs with common diseases and common emergencies, prompting specialty choice, and providing helpful background among residents (effectiveness subscale, 4 items).

Scoring System

Questions from Part B were formulated as a Likert-type scale, with 2 (yes; no) or 3 (no; yes to some extent; yes) answering options. In the adequacy and effectiveness subscales, answering options were “no” (score = 0), “yes to some extent” (score = 1) and “yes” (score = 2). Adequacy (score range = 0, 8) and effectiveness (score range = 0, 8) subscale scores were calculated by adding up the respective item scores, with higher scores indicating better adequacy and greater effectiveness, respectively. In the improvements required subscale, a reverse scoring was used, where the answer “no” was scored as 1 and answer “yes” as 0; thus, a higher score (range = 0, 5) indicates lesser improvements required, namely more positive opinion. A total score (range = 0–21) was calculated as the sum of all 3 subscale scores, where a higher score indicates better overall suitability of the ophthalmology course.

Statistical methods

Statistical analyses were performed using the Statistical Package for Social Sciences version 21.0 for Windows (SPSS Inc., Chicago, IL, USA). Descriptive statistics were used to summarize the participants' characteristics and the patterns of answers to the questionnaire's Part B items. The reliability of the questionnaire scale (overall suitability) and subscales (adequacy, improvements required, and effectiveness) was analyzed by calculating Cronbach's alpha. Mean, standard deviation (SD), median, and 75th centile (P75) were calculated for all four variables. Kolmogorov–Smirnov and Shapiro Wilk's tests were used to analyze the distribution of the scores as mentioned above; none of these was normally distributed. Subsequently, the overall suitability score was compared by resident's level, gender, and region using nonparametric tests, including the four scores were analyzed as categorical variables using the following cutoffs: Adequacy (poor <4, good ≥4); improvements required (none [score = 0], one aspect [score = 1], 2+ aspects [score ≥2]); effectiveness (poor <4, good ≥4); and overall suitability (not suitable [score <10] and suitable [score ≥10]). Chi-square test was used to compare the percentage of overall suitability, adequacy level, number of improvements required, and effectiveness level between the resident's factors. A *p*-value of <0.05 was considered to reject the null hypothesis.

Our study was approved by the Institutional Review Board of the King Abdulaziz University Faculty of Medicine, Jeddah, Saudi Arabia, with reference number 205–17.

Results

Participants' Characteristics

We included 106 residents, and 53.8% were males. The majority were R1 (33.0%) or R2 (26.4%), and were from Central (45.3%) and Western (28.3%) regions (Table 1).

Reliability of the Questionnaire

The questionnaire showed high reliability with Cronbach's alpha = 0.877. Similarly, all three subscales showed good to high reliability (Table 2).

Assessment of Ophthalmology Course

Figure 1 presents the patterns of answers to different scale items, as classified by aspect. Regarding adequacy, respondents opined that the ophthalmology course did not reasonably cover the basic part (35.8%), clinical part (61.3%), common disease (26.4%), and emergencies (39.6%). Regarding

Table 1 Participants' characteristics

Parameter	Category	Frequency	Percentage
Level	R1	35	33.0
	R2	28	26.4
	R3	22	20.8
	R4	21	19.8
Gender	Male	57	53.8
	Female	49	46.2
Program region	Central	48	45.3
	Western	30	28.3
	Eastern	13	12.3
	Southern	15	14.2

Table 2 Reliability of the Questionnaire Scale and Subscales

Scale	No. Items (Score Range)	Cronbach's Alpha	Comment
Overall questionnaire	13 (0–20)	0.877	High reliability
Adequacy	4 (0–8)	0.841	High reliability
Improvements required	5 (0–5)	0.761	Good reliability
Effectiveness	4 (0–5)	0.805	High reliability

Figure 1 Assessment of ophthalmology course by residents: patterns of answers to the questionnaire items. Bars represent the percentage of participants who answered with the given option for the given aspect.

improvements needed, more than 80% of the participants stated that the course needed to be improved for all its features, which included duration (80.2%), objectives (85.8%), content (82.1%), organization (83.0%), and supervision (81.1%). Regarding effectiveness, 55.7% declared that the ophthalmology course contributed partially (42.5%) or exclusively (13.2%) in their choice of specialty, while half of them opined that it was not helpful for residents or in familiarizing GPs with common ophthalmic diseases and emergencies.

Metrical analysis (Table 3) of the overall score showed mean (SD)=6.84 (5.19) and median (P75) = 6.00 (10.00), which are less than half of the maximum possible score, indicating very low scores. Accordingly, the course is generally deemed suitable for only 27.4% of the participants. Similarly, the mean and median of the three sub-scales were less than half of the respective maximum possible scores.

Participants' Factors Associated with a Course Assessment

Females' overall scores were higher than males' (mean = 7.92 versus 5.91, median = 8.00 versus 5.00, respectively), and the difference was statistically significant using the Mann–Whitney *U*-test ($p = 0.022$). However, the difference in suitability rate between females (32.7%) and males (22.8%) was not statistically significant ($p = 0.257$). No statistically significant association of overall suitability score or suitability rate with the residents' level or region was observed (Table 4).

Analysis by aspect showed no statistically significant association of adequacy, improvements required, and effectiveness with the residents' gender, level, or region (Table 5).

Discussion

Currently, the provided undergraduate curricula in ophthalmology medical schools do not follow any national standards for teaching. By surveying ophthalmology residents, more than half of the respondents declared that ophthalmology undergraduate courses inadequately cover the required clinical aspects and do not adequately familiarize GPs with common ophthalmic diseases and emergencies. Also, the metrical analysis indicated low suitability scores since 72.6% of residents opined that the course was generally not suitable for undergraduate students. More specifically, the lowest scores were attained for the effectiveness of the course, and the majority of residents (94.3%) have agreed that the course required at least one improvement.

Our results were in agreement with those reported in the literature, although most of the other surveys employed medical students or non-ophthalmology specialists who had been receiving an ophthalmologic education. These studies revealed multiple deficient aspects in ophthalmologic education that would consequently affect graduate students. For example, a nationwide survey in Australia showed that the taught curricula were inefficient, and there was an inadequate emphasis on ophthalmology. The

Table 3 Assessment of Ophthalmology Course by the Residents

Scale	Level	Mean (SD)	Median (P75)	N	%
Adequacy	Score (0–8)	3.23 (2.45)	3.00 (5.00)		
	Poor (<4)			60	56.6
	Good (≥4)			46	43.4
Improvements required	Score (0–5)	0.88 (1.36)	0.00 (1.00)		
	None			6	5.7
	One aspect			19	17.9
	2+ aspects			81	76.4
Effectiveness	Score (0–8)	2.74 (2.35)	2.00 (5.00)		
	Poor (<4)			67	63.2
	Good (≥4)			39	36.8
Overall suitability	Score (0–20)	6.84 (5.19)	6.00 (10.00)		
	Not suitable (<10)			77	72.6
	Suitable (≥10)			29	27.4

Table 4 Overall Suitability of Ophthalmology Course by Residents' Gender, Level, and Region

Factor	Category	Overall Assessment				
		Mean (SD)	Median (P75)	p-value*	Suitable (N, %)	p-value [‡]
Level	R1	7.46 (5.14)	7.00 (11.00)	0.464	10 (28.6)	0.254
	R2	5.86 (4.96)	5.00 (8.75)		5 (17.9)	
	R3	6.73 (6.28)	5.50 (9.25)		5 (22.7)	
	R4	7.24 (4.44)	8.00 (11.00)		9 (42.9)	
Gender	Male	5.91 (5.23)	5.00 (9.00)	0.022 [†]	13 (22.8)	0.257
	Female	7.92 (4.98)	8.00 (11.50)		16 (32.7)	
Program Region	Central	7.13 (5.78)	6.50 (10.75)	0.989	16 (33.3)	0.562
	Western	6.60 (5.15)	5.00 (9.75)		7 (23.3)	
	Eastern	6.08 (3.01)	5.00 (8.00)		2 (15.4)	
	Southern	7.07 (5.12)	8.00 (10.00)		4 (26.7)	

Notes: The test used: *nonparametric tests (Mann-Whitney U-test or Kruskal-Wallis test, as applicable); [†]statistically significant result (p<0.05); [‡]Chi square test.

Table 5 Adequacy, Improvements Needed, and Effectiveness of Ophthalmology Course by Residents' Gender, Level, and Region

Factor	Category	Adequacy		Improvements Needed				Effectiveness	
		Good	p-value	None	1–3 Aspects	4–5 Aspects	p-value	Good	p-value
Level	R1	13 (37.1)	0.732	2 (5.7)	6 (17.1)	27 (77.1)	0.426	17 (48.6)	0.176
	R2	12 (42.9)		2 (7.1)	3 (10.7)	23 (82.1)		6 (21.4)	
	R3	10 (45.5)		2 (9.1)	2 (13.6)	17 (77.3)		8 (36.4)	
	R4	11 (52.4)		0 (0.0)	7 (33.3)	14 (66.7)		8 (38.1)	
Gender	Male	21 (36.8)	0.142	2 (3.5)	11 (19.3)	44 (77.2)	0.563	19 (33.3)	0.426
	Female	25 (51.0)		4 (8.2)	8 (16.3)	37 (75.5)		20 (40.8)	
Program Region	Central	24 (50.0)	0.316	5 (10.4)	4 (8.3)	39 (81.3)	0.192	19 (39.6)	0.341
	Western	10 (33.3)		1 (3.3)	8 (26.7)	21 (70.0)		13 (43.3)	
	Eastern	4 (30.8)		0 (0.0)	3 (23.1)	10 (76.9)		2 (15.4)	
	Southern	8 (53.3)		0 (0.0)	4 (26.7)	11 (73.3)		5 (33.3)	

Notes: Values are frequencies (percentages) of participants in the given factor category, whose opinions correspond to the given level of adequacy, improvement needed, and effectiveness. Chi-square test was used.

medical students and junior medical officers had low confidence with basic clinical eye examinations, such as slit lamp use and direct ophthalmoscopy.¹¹ Another cross-sectional survey of United Kingdom (UK) medical schools showed significant variations in the organization and teaching methods, and their taught curricula failed to meet the ICO recommendations.⁵ Furthermore, in Canada, it has been found that only 35.7% of schools have implemented a mandatory ophthalmology rotation; the duration of each mainly was less than two weeks.¹² Likewise, more than half of the students at New York University School of Medicine felt uncomfortable diagnosing eye emergencies, and they were less confident in testing visual acuity and using a direct ophthalmoscope.¹³

In the present investigation, no significant differences in suitability scores or rates were reported according to residency levels and the region of ophthalmology programs. The same applies to the subcategories of the assessment, including the adequacy and effectiveness of the program, as well as the required improvements. Indeed, this indicates nationwide consistency in the perception of ophthalmology residents, which was mainly criticizing the deficient aspects of the program. Moreover, these deficits have been consistent with the repeatedly reported negligence of ophthalmologic teaching in multiple settings, including ours. The deficiencies, as mentioned earlier in ophthalmologic teaching globally and locally, have indicated that medical students have not been adequately prepared to provide competent, high-quality, and confident eye care, regardless of the available resources.

Lack of adequate basic knowledge and clinical skills was significantly perceived by approximately two-thirds of our study residents. This might contribute to reducing the clinical competence of GPs and other specialties upon graduation. In the UK, the lack of adequate undergraduate education has led to marked deficiencies in basic ophthalmologic skills, particularly those who worked in emergency departments,¹⁴ and 63.9% of senior house officers had limited confidence in managing eye emergencies.¹⁵ Basic ophthalmologic examination, including direct ophthalmoscopy, should be effectively performed by non-specialist clinicians, especially when access to an ophthalmologist is not possible. In particular, the findings of direct ophthalmoscopy could indicate devastating systemic changes, such as the detection of papilledema (indicating accelerated hypertension or lesions occupying the cerebral space), as well as cytomegalovirus retinitis and endocarditis, which require further invasive investigations.^{16,17}

Identifying common preventable eye diseases and the subsequent referral of patients at early stages to eye care specialists is another crucial aspect of care, which can reduce the burden of community ophthalmological disorders.¹⁸ Therefore, there is an urgent need to make significant changes to undergraduate medical curricula to enhance patients' safety and increase the competency of future doctors.

Several proposed improvements could be considered to increase the competency of students. First, increasing the space of ophthalmology lectures to undergraduate students might support their competency and interests in such a specialty. In Canada, Li et al¹⁹ showed that instituting additional ophthalmology lectures to undergraduate year three has significantly increased their knowledge levels and has impacted their desire for teaching. Second, the contribution of all teaching staff to integrating basic scientific knowledge with clinical practice to small groups of students.²⁰ Third, training methods, such as didactic lectures, case presentations, and PowerPoint presentations, may have significant effects with varying efficacy outcomes.^{21,22} Fourth, the timing of educational training is another essential aspect. Previous studies have shown that distributing ophthalmologic training across four years of study would presumably be more effective than single- or two-year programs.^{13,23,24} Finally, providing feedback during teaching sessions would be invaluable for promoting examination skills. Therefore, such multi-optional approaches should be considered in future reformations of medical schools' undergraduate ophthalmology course, to promote a more effective process of teaching and improve students' ophthalmology curriculum and practice.

The technological advances in the ophthalmology sector should be borne in mind. Innovation and research have an integrative role in outlining the upcoming aspects of ophthalmology practice both at the national and international levels. Only technically prepared professionals will cope with the continuous progress achieved in the market, while low-skilled doctors will be excluded. Suitable strategies should be readily applied to meet the inevitable technological changes.²⁵

Our analysis selected residents because they have received adequate medical training and demonstrated high levels of clinical competency, as corroborated in a recent cross-sectional study.²⁶ The local residency training consists of a 4-year structured program, where training curricula are divided into monthly rotations in multiple specialties. Both clinical and surgical ophthalmological

skills are taught during the first two years and are extensively repeated in the following couple of years to support the learned skills. These programs were initially developed in 1982 as per the recommendations of a nationwide survey concerning the local etiological factors of blindness.²⁷ Currently, residency training programs are directly supervised by the Saudi Commission for Health Specialties, and their curricula have provided cognitive and technical skills comparable to those required by the ICO.^{6,26}

The present study has some limitations. Response bias is possible due to the cross-sectional nature of our analysis (namely, the accuracy of the obtained results and how they have truly reflected the population under study is unknown). The subjective nature of residents' judgments regarding the adequacy and effectiveness of the curricula may have been different than those really existing, by excluding student-related factors, such as the student's willingness to study ophthalmology, which would eventually change future competence and skills. Therefore, future nationwide studies which include both the residents and students are needed.

Conclusion

The present study showed multiple deficits in the teaching process of the ophthalmology course as perceived by residents. These were related to inadequate coverage of the basic and clinical skills required to manage patients by general practitioners and emergency physicians, and the lack of effective and informative materials that enable future students to be readily familiarized with common ophthalmic diseases. Lack of basic ophthalmological skills in future non-specialist physicians may pose significant risks to patients' eyes and systemic health. Significant improvements such as increasing the number of teaching hours, applying integrated teaching methods, and implementing an adequate feedback mechanism in ophthalmologic curricula are required.

Disclosure

The authors report no conflicts of interest for this work.

References

1. Succar T, Grigg J, Beaver HA, Lee AG. A systematic review of best practices in teaching ophthalmology to medical students. *Surv Ophthalmol*. 2016;61(1):83–94. doi:10.1016/j.survophthal.2015.09.001
2. Wenting SZ, Samin MM, Sanjay S, et al. A comparison of undergraduate clinical ophthalmology learning methods: smart phone television display versus slit-lamp teaching telescope. *Can J Ophthalmol*. 2017;52(4):385–391. doi:10.1016/j.jcjo.2016.11.032
3. Hashemi H, Khabazkhoob M, Nabovati P, et al. The prevalence of age-related eye disease in an elderly population. *Ophthalmic Epidemiol*. 2017;24(4):222–228. doi:10.1080/09286586.2016.1270335
4. Byers WG. The place of ophthalmology in the undergraduate medical curriculum. *Br Med J*. 1922;2(3209):4–6. doi:10.1136/bmj.2.3209.4
5. Hill S, Dennick R, Amoaku W. Present and future of the undergraduate ophthalmology curriculum: a survey of UK medical schools. *Int J Med Educ*. 2017;8:389–395. doi:10.5116/ijme.59ac.f69b
6. International Council of Ophthalmology. Principles and guidelines of a curriculum for education of the ophthalmic specialist. *Klin Monbl Augenheilkd*. 2006;223(Suppl 4):S3–S48.
7. Kilduff C, Lois C. Red eyes and red-flags: improving ophthalmic assessment and referral in primary care. *BMJ Open Qual*. 2016;5(1):u211608. w4680.
8. World Health Organization. Blindness and vision impairment. Geneva: WHO; 2019. Available from: <https://www.who.int/news-room/fact-sheets/detail/blindness-and-visual-impairment>. Accessed November 8, 2021.
9. Al-Essa RS, Al-Otaibi MD, Al-Qahtani BS, Masuadi EM, Omair A, Alkatan HM. Future ophthalmology practice pattern: a survey of Saudi Board of Ophthalmology residents. *Saudi J Ophthalmol*. 2019;33(1):1–6. doi:10.1016/j.sjopt.2019.01.005
10. Madani FM, Alasiri R, Howldar S, Zagzoog F, Alhibshi N. Perceptions and career expectation in ophthalmology among Saudi undergraduate medical students. *Middle East Afr J Ophthalmol*. 2018;25(3):142–149. doi:10.4103/meajo.MEAO_261_17
11. Zhang HH, Hepschke JL, Shulruf B, et al. Sharpening the focus on ophthalmology teaching: perceptions of medical students and junior medical officers. *Clin Exp Ophthalmol*. 2018;46(9):984–993. doi:10.1111/ceo.13342
12. Gostimir M, Sharma RA, Bhatti A. Status of Canadian undergraduate medical education in ophthalmology. *Can J Ophthalmol*. 2018;53(5):474–479. doi:10.1016/j.jcjo.2017.11.015
13. Cobbs L, Tsui E, Haberman ID, et al. Student perceptions of the ophthalmology curriculum in medical school. *J Acad Ophthalmol*. 2018;10(01):e79–e82. doi:10.1055/s-0038-1653974
14. Baylis O, Murray PI, Dayan M. Undergraduate ophthalmology education - a survey of UK medical schools. *Med Teach*. 2011;33(6):468–471. doi:10.3109/0142159X.2010.540594
15. Sim D, Hussain A, Tebbal A, Daly S, Pringle E, Ionides A. National survey of the management of eye emergencies in the accident and emergency departments by senior house officers: 10 years on—has anything changed? *Emerg Med J*. 2008;25(2):76–77. doi:10.1136/emj.2007.049999
16. Sergott RC. Headaches associated with papilledema. *Curr Pain Headache Rep*. 2012;16(4):354–358. doi:10.1007/s11916-012-0283-x
17. Yusuf IH, Salmon JF, Patel CK. Direct ophthalmoscopy should be taught to undergraduate medical students—yes. *Eye (Lond)*. 2015;29(8):987–989. doi:10.1038/eye.2015.90
18. Divya K, Suvetha K, Sen A, Sundar D. Needs assessment of ophthalmology education for undergraduate medical students - a study from a medical college in South India. *Educ Health (Abingdon)*. 2017;30(3):223–227. doi:10.4103/efh.Efh_52_16
19. Li B, Curtis D, Iordanous Y, Proulx A, Sharan S. Evaluation of Canadian undergraduate ophthalmology medical education at Western University. *Can J Ophthalmol*. 2016;51(5):373–377. doi:10.1016/j.jcjo.2016.04.024
20. Tsinopoulos IT, Symeonidis C, Tsaousis KT, et al. Contribution of integrated teaching in the improvement of an undergraduate ophthalmology curriculum. *Adv Med Educ Pract*. 2014;5:433–437.
21. Gupta RR, Lam WC. Medical students' self-confidence in performing direct ophthalmoscopy in clinical training. *Can J Ophthalmol*. 2006;41(2):169–174. doi:10.1139/I06-004

22. Gogate P, Deshpande M, Dharmadhikari S. Which is the best method to learn ophthalmology? Resident doctors' perspective of ophthalmology training. *Indian J Ophthalmol*. 2008;56(5):409–412. doi:10.4103/0301-4738.42419
23. Esparaz ES, Binder SB, Borges NJ. How prepared are medical students to diagnose and manage common ocular conditions. *J Educ Eval Health Prof*. 2014;11:29. doi:10.3352/jeehp.2014.11.29
24. Lippa LM, Boker J, Duke A, Amin A. A novel 3-year longitudinal pilot study of medical students' acquisition and retention of screening eye examination skills. *Ophthalmology*. 2006;113(1):133–139. doi:10.1016/j.ophtha.2005.09.003
25. Kara-Junior N. Technology, teaching, and the future of ophthalmology and the ophthalmologist. *Arq Bras Oftalmol*. 2018;81(3):V–vi. doi:10.5935/0004-2749.20180036
26. AlEnezi SH, Alfawaz AM, Al Owaifeer AM, Althiabi SM, Tabbara KF. Assessment of ophthalmology residency programs in Saudi Arabia: a trainee-based survey. *J Med Educ Curric Dev*. 2019;6:2382120519855060. doi:10.1177/2382120519855060
27. Tabbara KF, Ross-Degnan D. Blindness in Saudi Arabia. *JAMA*. 1986;255(24):3378–3384. doi:10.1001/jama.1986.03370240048035

International Journal of General Medicine

Dovepress

Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies

across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-general-medicine-journal>