Factors Affecting Final Surgical Outcome of Combined Trabeculotomy–Trabeculectomy in Primary Congenital Glaucoma

Marwa A Khairy

Said Kenawy

Khaled Mohamed Fawzi

Haitham Y Al-Nashar

Ophthalmology Department, Faculty of Medicine, Zagazig University, Zagazig, Egypt

Purpose: To evaluate different pre-operative variables on the success of combined trabeculotomy–trabeculectomy (CTT) surgery in patients with primary congenital glaucoma (PCG) to predict those at higher risk for surgical failure and for proper parent counseling.

Patients and Methods: Sixty-three eyes of patients with PCG were treated either with CTT without augmentation, or CTT augmented with mitomycin-C (0.2 mg/mL) in both subconjunctival space and under scleral flap for 3 minutes, or with CTT augmented with a collagen implant under both the scleral flap and the conjunctiva. Cases showed surgical failure was reported and evaluated in relation to different pre-operative variables.

Results: Complete success (IOP ≤ 21 mmHg) was achieved in 52 cases (82.5%). Cumulative success probability was calculated using Kaplan-Meier survival analysis, proving that higher pre-operative intraocular pressure (IOP) was associated with higher failure rates (28.6% for pre-operative IOP ≥ 30 mmHg versus 4.8% for IOP < 30 mmHg), with P value = 0.007.

Conclusion: CTT is an effective surgical intervention in PCG patients without sight threatening complications. Univariate survival analysis showed higher rates of surgical failure in patients with higher pre-operative IOP, while other pre-operative variables were irrelevant.

Keywords: buphthalmos, Kaplan–Meier survival analysis, surgical failure

Introduction

Congenital glaucoma is a major cause of childhood blindness worldwide, with variable birth prevalence in different populations worldwide,¹ and is responsible for 10.8% of all visual impairments in children.²

The disease is usually managed surgically, with medical treatment playing a temporary or an adjunctive role. Multiple surgical modalities have been described, however surgery in a buphthalmic eye with disturbed anatomy is challenging and is associated with higher failure rates and multiple complications than in adults. Early and proper surgical intervention can achieve a favorable outcome and preserve patients' vision.^{1,3}

A milder form of the disease associated with only trabeculodysgenesis can respond well to trabeculotomy surgery, however, the more advanced form with iridotrabeculodysgenesis, and corneotrabeculodysgenesis showed less favorable results to trabeculotomy.^{4,5} Combining trabeculotomy with trabeculectomy is one of the preferred surgical options, which showed favorable results in congenital glaucoma, and is advised mainly for treating moderate to severe forms of the disease, as the procedure

Correspondence: Marwa A Khairy
Ophthalmology Department, Faculty of Medicine, Zagazig University, Zagazig, Egypt
Tel +20-1111801956
Fax +20-552353738
Email ma_kh87@yahoo.com

provides dual pathways for aqueous drainage.⁶ The procedure is also recommended for severe familial form of the disease which occur mainly in Indian and Middle East populations which showed less favorable results to goniotomy compared to the Western population, another advantage is that in cases where Schlemm's canal cannot be identified and cannulated because of the disturbed limbal anatomy in larger eyes, the added trabeculectomy fistula can still function and lower the rates of failure.^{6–8}

Patients and Methods

The study was carried out in accordance with the principles of the Declaration of Helsinki, and approval from the Ethical Committee of Zagazig Faculty of Medicine and an Institutional Review Board (IRB) code was obtained (IRB No: ZU-IRB #5362/10-4-2019). Informed consent was obtained from the parents of all patients.

This clinical trial included 63 eyes of 36 patients with primary congenital glaucoma (PCG), eyes were randomized by random allocation computer program into 3 equal groups (21 eyes in each group); eyes in group I were treated with combined trabeculotomy–trabeculectomy (CTT), group II were treated with CTT augmented with mitomycin-C, while eyes in group III were treated with CTT augmented with collagen implant (Ologen).

Inclusion Criteria

Patients younger than 3 years diagnosed as having PCG were included.

Exclusion Criteria

Patients with previous ocular surgery, secondary glaucoma, or other ocular pathologies were excluded from the study. Also, those lost during the follow-up visits (lost for more than two visits during follow-up) were withdrawn from the study.

Pre-operative Assessment

Pre-operative patients' evaluation including demographic data (age, sex, consanguinity and family history). Also, examination under general anesthesia (EUGA) for intraocular pressure (IOP) calculation using Perkins applanation tonometer, horizontal corneal diameters using surgical caliper, anterior segment examination using surgical microscope for the degree of corneal edema and to exclude other pathologies, fundus examination to assess for cupping and exclude retinal pathology by indirect ophthalmoscope if corneal clarity allows for fundus visualization.

Surgical Techniques

Eyes in group I; were treated with CTT.

Eyes in group II; were treated with CTT augmented with mitomycin-C (0.2 mg/mL) in both subconjunctival space and under scleral flap for 3 minutes.

Eyes in group III; were treated with CTT augmented with a collagen implant (Ologen TM, Aeon Astron Europe B.V, Leiden, The Netherlands) divided unequally, with the smaller part implanted under the scleral flap and the larger part under the conjunctiva.

Post-Operative Treatment

All patients received topical antibiotic eye drops for two weeks and topical prednisolone acetate 1% eye drops five times per day for a week, with gradual withdrawal over another two weeks, also cyclopentolate drops twice daily for a week.

Post-Operative Follow Up

All patients were seen at the first day looking for shallow or lost anterior chamber, red reflex, IOP and any sign of ocular inflammation. Patients were seen again at one, and two weeks and at first, second, third, sixth, ninth and twelfth months post-operative.

Success Criteria

Complete success was defined as an IOP ≤ 21 mmHg without added antiglaucoma medications or needing another glaucoma surgery and without sight threatening complications. Qualified success was considered when the above criteria were achieved, but with added antiglaucoma medication. Failure was defined as IOP being >21 mmHg with maximum topical antiglaucoma medications, the requirement of another surgical intervention to lower IOP, or the appearance of sight threatening complications.

Statistical Analysis

All statistical analyses were performed using Statistical Package for the Social Sciences (SPSS, version 25.0). According to the type of data qualitative representation is as number and percentage, quantitative continuous group is represented by mean with standard deviation [SD]. The effect of individual clinical and demographic pre-operative variables on the final surgical outcome of treatment was analyzed using Kaplan–Meier survival analysis. P value <0.05 was considered significant.

Results

63 eyes of 36 patients with PCG were included in this study. 27 patients had bilateral disease, while 9 had a unilaterally affected eye. A random allocation computer program was used to assign each eye to one of the study groups. The included patients were 24 (66.7%) males, and 12 (33.3%) females. The mean age at time of surgery was 6.18 ± 2.8 months (Range 1.5–11 months). Positive family history of PCG was recorded in 9 patients (25%), while the remaining patients, 27 (75%), denied having any family member with PCG. Regarding consanguinity, 13 (36.1%) patients had positive consanguinity between both parents, while 23 (63.9%) eyes were negative. (Table 1).

Pre-operative examination data were as follows: mean pre-operative IOP was 27.65 ± 2.6 mmHg, with a range of (23–34 mmHg). Mean pre-operative corneal diameter was 13 ± 0.56 mm (range 12–14 mm). Pre-operative corneal edema was recorded in all eyes, with 22 (34.9%) eyes, with superficial epithelial edema, and 32 (50.8%) showed central edema involving both epithelium and parenchyma, while 9 (14.3%) had total diffuse edema obscuring fundus details (Table 1).

Table 1 Pre-operative Patient's Data

Variable	(Mean $\pm$ SD)/No (%)
Sex	
-Male patients	• 24 (66.7%)
-Female patients	• 12 (33.3%)
Patients' Age	6.18 $\pm$ 2.8 months Range (1.5–11 months)
Family history	
-Positive patients	• 9 (25%)
-Negative patients	• 27 (75%)
Consanguinity	
-Positive patients	• 13 (36.1%)
-Negative patients	• 23 (63.9%)
IOP	27.65 $\pm$ 2.6 mmHg Range (23–34 mmHg)
Corneal diameter	13 $\pm$ 0.56 mm Range (12–14 mm)
Corneal edema	
-Superficial epithelial edema	• 22 (34.9%)
-Central edema involving both epithelium and parenchyma	• 32 (50.8%)
-Total diffuse edema obscuring fundus details	• 9 (14.3%)

Total success (complete and qualified) was achieved in 55 cases (87.3%) and failure was reported in 8 cases (12.7%). Complete success was achieved in 52 cases (82.5%). Cumulative success probability was determined using Kaplan–Meier survival analysis, with univariate comparisons of final surgical outcome according to patient pre-operative characteristics (Table 2).

Comparing the type of surgical intervention to failure, 3 cases (14.3%) showed failure in group I, and 2 cases (9.5%) in group II, while group III had 3 failed cases (14.3%), with P value 0.882, which was statistically insignificant.

Comparing age at time of surgical intervention to failure, number of eyes below 6 months of age were 26 (41.3%), with 4 of them (15.4%) showing failure, while there were 37 eyes ≥ 6 months (58.7%), with 4 of them (10.8%) showing failure. P value = 0.553, which was statistically insignificant.

Regarding the patient's sex, 42 eyes (66.7%) were male, with 7 (16.7%) experiencing failure, while 21 eyes (33.3%) were female, with 1 (4.8%) failed eye. P value = 0.180, which was statistically insignificant.

Comparing family history to failure cases, 15 eyes (23.8%) had a positive family history of PCG. 2 (13.3%) of them experienced failure. While 48 eyes (76.2%) had a negative family history, 6 (12.5%) of them experienced failure. P value = 0.906 which was statistically insignificant.

Regarding consanguinity, 22 eyes (34.9%) had positive consanguinity, 4 (18.2%) of them showed failure. 41 eyes (65.1%) had negative consanguinity, with 4 (9.8%) of them failing.

Comparing pre-operative IOP to failure, 42 eyes (66.7%) had an IOP of less than 30 mmHg, of them 2 eyes (4.8%) showed failure. While 21 eyes (33.3%) had a pre-operative IOP ≥ 30 mmHg, with 6 eyes (28.6%) failing among them. P value = 0.007, which was statistically significant (Figure 1).

Regarding pre-operative corneal diameter, 25 eyes (39.7%) had a diameter of less than 13 mm, of them 3 eyes (12%) showed failure. While 38 eyes (60.3%) had a diameter ≥ 13 mm, with 5 (13.2%) of them showing failure. P value = 0.897, which was statistically insignificant.

Comparing pre-operative corneal clarity to failure showed 22 eyes (34.9%) with superficial epithelial edema, 4 (18.2%) of them showed failure. A central edema involving both epithelium and parenchyma was

Table 2 Probability Outcome of CTT Surgery in PCG Using Kaplan–Meier Survival Analysis, with Univariate Analysis Comparing Final Surgical Outcome According to Pre-Operative Variables

Total Number of Eyes According to Pre-Operative Variable (%)	Number of Failures	Probability of Failure	Probability of Success at 95% CI	Log Rank Test	Mean Survival Time (Days)	P value
Between groups -CTT = 21 (33.3%) -CTTM = 21 (33.3%) -CTTO = 21 (33.3%)	• 3 • 2 • 3	• 0.143 • 0.095 • 0.143	• 0.86 • 0.90 • 0.86	0.252	• 318.6 (269.9–367.3) • 333 (291.04–375.2) • 320 (272.8–367.2)	0.882
Age at surgery -< 6 months = 26 (41.3%) -≥ 6 months = 37 (58.7%)	• 4 • 4	• 0.154 • 0.108	• 0.85 • 0.89	0.351	• 313.46 (267–359.9) • 331.22 (299.9–362.5)	0.553
Sex -Male = 42 (66.7%) -Female = 21 (33.3%)	• 7 • 1	• 0.167 • 0.048	• 0.83 • 0.95	1.8	• 310.6 (273.8–347.4) • 350.5 (322.7–378.3)	0.180
Family history -Positive = 15 (23.8%) -Negative = 48 (76.2%)	• 2 • 6	• 0.133 • 0.125	• 0.87 • 0.88	0.014	• 320.3 (262.7–377.9) • 325 (295–354.9)	0.906
Consanguinity -Positive = 22 (34.9%) -Negative = 41 (65.1%)	• 4 • 4	• 0.182 • 0.098	• 0.82 • 0.90	0.95	• 305.5 (252.6–358.3) • 333.8 (304.7–362.9)	0.330
IOP -<30 mmHg = 42 (66.7%) -≥30 mmHg = 21 (33.3)	• 2 • 6	• 0.048 • 0.286	• 0.95 • 0.71	7.28	• 349.8 (329.1–370.1) • 272.1 (209.3–335)	0.007
Corneal diameter -<13 mm = 25 (39.7%) -≥13 mm = 38 (60.3%)	• 3 • 5	• 0.12 • 0.13	• 0.88 • 0.87	0.017	• 326 (284.6–367.4) • 322.5 (287.7–357.3)	0.897
Corneal clarity -SE = 22 (34.9%) -CE = 32 (50.8%) -TE = 9 (14.3%)	• 4 • 2 • 2	• 0.182 • 0.063 • 0.22	• 0.82 • 0.94 • 0.78	2.68	• 305.5 (252.6–358.3) • 345.9 (320.4–371.5) • 290.6 (199.6–381.5)	0.262

Abbreviations: CI, confidence interval; IOP, intraocular pressure; CTT, combined trabeculotomy–trabeculectomy; CTTM, combined trabeculotomy–trabeculectomy augmented with mitomycin-C; CTTO, combined trabeculotomy–trabeculectomy augmented with Ologen; SE, superficial epithelial edema; CE, central edema involving both epithelium and parenchyma; TE, total diffuse edema obscuring fundus details.

reported in 32 eyes (50.8%), with 2 (6.3%) of them experiencing failure. Total diffuse edema obscuring fundus details was reported in 9 eyes (14.3%), 2 (22.2%) of them experienced failure. P value = 0.262, which was statistically insignificant.

No sight threatening complications were recorded in any group, like endophthalmitis or retinal detachment. Only a few cases showed intraoperative hyphema that needed an anterior chamber wash. Also post-operative hypotony was recorded in the first and second group and was managed conservatively.

Cases with failed primary surgery needed another surgical intervention in the form of CTT augmented with

MMC. 5 eyes were controlled by the second procedure, while the other 3 had uncontrolled IOP and were scheduled for valve surgery.

Discussion

The definitive treatment of PCG is surgical, with wide varieties of surgical approaches, however, CTT showed promising results with higher success rates and better IOP control for a single procedure.⁸ One of the advantages of CTT is that it can be performed in eyes with significant ocular enlargement and large corneal diameter which was considered as a challenge to goniotomy surgery.^{7,9}

Figure 1 Kaplan-Meier curve showing the success probabilities in relation to pre-operative IOP.

The higher success rates of CTT were explained by the supplementary outflow pathway that the procedure creates, with less risk of surgical failure in cases where trabeculotomy was not done properly or Schlemm's canal cannot be identified.^{6,7,10} The addition of antimetabolites to augment CTT surgery is a matter of great debate among surgeons,^{7,11,12} it's believed by some surgeons to increase success rates as well as increase the rate of complications.^{6,12} In our study we reported a complete success in 52 cases (82.5%), with total success in 55 cases (87.3%) and failure in only 8 cases (12.7%). In univariate analysis to evaluate the role of different pre-operative risk factors on treatment outcome in the present study, we evaluated the type of surgery, either CTT alone or CTT augmented with the addition of MMC or with added Ologen implant on failure. P value was 0.882, which indicates that the three surgical modalities were equally effective.

Mandal et al reported a success probability of 85.2% at first year after CTT surgery (complete success at IOP <16 mmHg under general anesthesia or <21 mmHg in older patients examined by slit lamp, without cupping progression or increase in corneal diameter),⁸ Mullaney et al reported 78% success rates (IOP, <21 mmHg) in

congenital glaucoma patients without coexistent anterior segment anomalies after CTT surgery.¹¹ Also, Essuman et al reported a 79% overall surgical success rate of CTT.¹³ Our results are comparable to the previously mentioned studies, with equal efficacy in either augmented or nonaugmented procedure.

Other pre-operative risk factors including age at time of surgical intervention, patient's sex, family history of PCG, consanguinity, pre-operative corneal diameter or corneal clarity were all irrelevant compared to surgical failure. The only factor significantly associated with higher failure rates was pre-operative IOP, as cases with pre-operative IOP ≥ 30 mmHg were associated with higher failure rates (28.6%), compared to cases with IOP <30 mmHg (4.8%), with a P value equal to 0.007, which was statistically significant (Table 2). Our results were consistent with Mandal et al who found a strong correlation between pre-operative IOP and failure, with a higher risk of surgical failure among patients with higher pre-operative IOP values. Patients with pre-operative IOP >35 mmHg had 31.9% failure rates, while those with an IOP ≤ 35 mmHg showed only a 15.1% failure rate, with P value = 0.003.⁸

Mandal et al also, found other factors associated with failure in univariate analysis, which included patient's age at time of surgery, and a history of previous glaucoma surgery. He stated that patients older than 6 months of age at the time of surgery had higher failure rates (21.4%) compared to those ≤ 6 months (12.8%), with P value = 0.010. However, the author stated that the data regarding the age of disease onset was missing in 61.4% of cases, that's why he did not comment on that factor.⁸ In our study we reported higher failure rates in patients operated below 6 months of age (15.4%) compared to those operated at or older than 6 months (10.8%), however there was no statistically significant difference (P value = 0.553). This matches other studies which reported that the earlier onset of the disease is associated with a more severe form and worse surgical prognosis.^{14,15}

Another variable Mandal et al investigated was prior glaucoma surgery. The author found that prior surgical intervention was associated with higher failure rates of the second procedure (43.3%), compared to primary surgery (14.8%), with P value <0.0001.⁸ However, in our study we operated upon primary cases only, and excluded those with previous surgical intervention.

Limitations of our study are the small number of the patients included and the short follow up period.

Conclusion

In conclusion, the study showed that the only factor which was statistically significant regarding surgical failure in univariate survival analysis was pre-operative IOP, other factors did not show statistically significant differences.

Abbreviations

CTT, combined trabeculotomy–trabeculectomy surgery; PCG, primary congenital glaucoma; EUGA, examination under general anesthesia; IRB, institutional review board; IOP, intraocular pressure.

Acknowledgments

The authors would like to express their deep gratitude to the patients who participated in this study.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Badawi AH, Al-Muhaylib AA, Al Owaifeer AM, Al-Essa RS, Al-Shahwan SA. Primary congenital glaucoma: an updated review. *Saudi J Ophthalmol*. 2019;33:382–388. doi:10.1016/j.sjopt.2019.10.002
2. Haddad MA, Sei M, Sampaio MW, Kara-José N. Causes of visual impairment in children: a study of 3210 cases. *J Pediatr Ophthalmol Strabismus*. 2007;44:232–240. doi:10.3928/01913913-20070701-04
3. Papadopoulos M, Edmunds B, Fenerty C, Khaw PT. Childhood glaucoma surgery in the 21st century. *Eye (Lond)*. 2014;28:931–943. doi:10.1038/eye.2014.140
4. Hoskins HD Jr, Shaffer RN, Hetherington J. Anatomical classification of the developmental glaucomas. *Arch Ophthalmol*. 1984;102(9):1331–1336. doi:10.1001/archophth.1984.01040031081030
5. Luntz MH. The choice of surgical procedure in congenital, infantile, and juvenile glaucoma. *J Ophthalmic Nurs Technol*. 1990;9(3):100–101.
6. Al-Hazmi A, Awad A, Zwaan J, Al-Mesfer SA, Al-Jadaan I, Al-Mohammed A. Correlation between surgical success rate and severity of congenital glaucoma. *Br J Ophthalmol*. 2005;89:449–453. doi:10.1136/bjo.2004.047761
7. Mandal A, Matalia J, Nutheti R, et al. Combined trabeculotomy and trabeculectomy in advanced primary developmental glaucoma with corneal diameter of 14 mm or more. *Eye*. 2006;20:135–143. doi:10.1038/sj.eye.6701817
8. Mandal AK, Gothwal VK, Nutheti R. Surgical outcome of primary developmental glaucoma: a single surgeon's long-term experience from a tertiary eye care centre in India. *Eye (Lond)*. 2007;21:764–774. doi:10.1038/sj.eye.6702324
9. Helmy H. Ab interno goniotomy combined with Ab externo trabeculectomy in advanced primary congenital glaucoma patients: 2-year follow-up. *Clin Ophthalmol*. 2021;15:565–574. doi:10.2147/OPTH.S292168
10. Sood D, Rathore A, Sood I, et al. Long-term outcome of combined trabeculotomy–trabeculectomy by a single surgeon in patients with primary congenital glaucoma. *Eye*. 2018;32:426–432. doi:10.1038/eye.2017.207
11. Mullaney PB, Selleck C, Al-Awad A, Al-Mesfer S, Zwaan J. Combined trabeculotomy and trabeculectomy as an initial procedure in uncomplicated congenital glaucoma. *Arch Ophthalmol*. 1999;117(4):457–460. doi:10.1001/archophth.117.4.457
12. Al-Hazmi A, Zwaan J, Awad A, al-Mesfer S, Mullaney PB, Wheeler DT. Effectiveness and complications of mitomycin C use during pediatric glaucoma surgery. *Ophthalmology*. 1998;105(10):1915–1920. doi:10.1016/S0161-6420(98)91041-7
13. Essuman VA, Braimah IZ, Ndanu TA, Ntim-Amponsah CT. Combined trabeculotomy and trabeculectomy: outcome for primary congenital glaucoma in a West African population. *Eye (Lond)*. 2011;25:77–83. doi:10.1038/eye.2010.156
14. Khaw PT. What is the best primary surgical treatment for the infantile glaucomas? *Br J Ophthalmol*. 1996;80(6):495–496. doi:10.1136/bjo.80.6.495
15. Dureau P, Dollfus H, Cassegrain C, Dufer JL. Long-term results of trabeculectomy for congenital glaucoma. *J Pediatr Ophthalmol Strabismus*. 1998;35(4):198–202. doi:10.3928/0191-3913-19980701-05

Clinical Ophthalmology

Dovepress

Publish your work in this journal

Clinical Ophthalmology is an international, peer-reviewed journal covering all subspecialties within ophthalmology. Key topics include: Optometry; Visual science; Pharmacology and drug therapy in eye diseases; Basic Sciences; Primary and Secondary eye care; Patient Safety and Quality of Care Improvements. This journal is indexed on PubMed

Central and CAS, and is the official journal of The Society of Clinical Ophthalmology (SCO). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-ophthalmology-journal>