

Pain Management Providers in the Era of COVID-19: Who is Taking Care of Those Who Provide Care?

Gabriela Toutin Dias¹
Michael E Schatman²⁻⁴

¹Psychology Assessment Center, Massachusetts General Hospital, Boston, MA, USA; ²Department of Anesthesiology, Perioperative Care and Pain Medicine, NYU School of Medicine, New York, NY, USA; ³Division of Medical Ethics, NYU School of Medicine, New York, NY, USA; ⁴School of Social Work, North Carolina State University, Raleigh, NC, USA

The overall quality of chronic pain management in the United States is woefully poor, and unfortunately getting worse. Schatman and Fortino recently elucidated many of the causes of this deterioration,¹ although revisiting these deficiencies is beyond the scope of this analysis. As a result of this state, neither patients nor healthcare providers who treat those with chronic pain are particularly satisfied. For example, patients are likely to see themselves as being treated in a system in which their voices are progressively less likely to be heard, where there is a lack of communication and responsiveness, a lack of comprehensive pain care, and rushed visits.² On the other hand, providers are dissatisfied by issues including a lack of resources for providing adequate treatment(s), unrealistic patient expectations for pain relief, and their own inadequate educations regarding pain management.³

COVID-19 has certainly not helped improve the quality of chronic pain management. Much has been written over the past year and a half regarding the effect of the pandemic on treatment that patients have received. The early literature addressed the use of telehealth to provide behavioral services to chronic pain patients,⁴ along with best practices consensus statements, which were focused on interventional pain medicine.^{5,6} Progressively, other issues were addressed, including the impact of the pandemic on fellowship training,⁷ the safety of various medications during COVID-19,⁸⁻¹⁰ the risks of developing chronic pain as a result of COVID-19 infection and treatment,¹¹ manifestations of characterological disturbances among chronic pain sufferers triggered by the pandemic,¹² dealing with disruptions of research on pain management due to the pandemic,¹³ the implications of postponed joint replacement surgeries for pain management,¹⁴ the impact of social distancing and other COVID-19-related phenomena on psychosocial aspects of chronic pain,^{15,16} the impact of the redeployment of clinical staff from pain management departments due to the pandemic,¹⁷ and the effect of the pandemic on the opioid epidemic,^{18,19} among several other areas of investigation. The focus of the extant medical literature on pain patient well-being and threats to it is certainly appropriate. Glaringly absent, however, in a thorough review of the literature are articles addressing the impact of COVID-19 on providers of chronic pain management services.

Amidst myriad adjustments observed in pain management practices across the US as a result of the COVID-19 pandemic, this analysis aims to underscore the

Correspondence: Michael E Schatman
Department of Anesthesiology,
Perioperative Care and Pain Medicine,
NYU School of Medicine, 550 First
Avenue, New York, NY, 10016, USA
Tel +1 425-647-4880
Email Michael.Schatman@NYULangone.org

critical issue of emotional distress amongst clinicians in pain medicine. These unprecedented times have introduced novel stressors into a panorama already marked by grave challenges. A cross-sectional study conducted prior to the pandemic found that 60.4% of pain medicine physicians in the US reported high emotional exhaustion and 35.7% reported high depersonalization, a critical symptom in the diagnosis of burnout and a potential symptom observed in PTSD.²⁰ With the COVID-19 pandemic, pain management providers have experienced the additional burden of staffing shortages,²¹ limited hospital resources,²¹ extended workloads,²² and risk of infection,²³ all whilst bearing witness to ongoing death, suffering, and noncompliance to scientifically proven measures aimed at helping us battle COVID-19 and its variants. Specifically, amongst anesthesiologists and interventional physiatrists caring for chronic pain patients, the challenges exerted by COVID-19 have been especially burdensome, as many became frontline healthcare workers within Emergency Departments and Intensive Care Units across the country. With this call to action, interventionalists faced longer and irregular hours, insufficient personal protective equipment, increased risk of contamination, and exposure to challenging decision-making processes stemming from the need to allocate limited resources.²⁴

Additionally, health services across the country have reported significant increases in workplace violence. According to The Joint Commission on Accreditation of Healthcare Organizations, healthcare workers are 20% more likely to become victims of workplace violence compared to workers from other industries.²⁵ In 2015 alone, approximately 75% of reported workplace assaults transpired within healthcare and social service settings.²⁵ This violence has undoubtedly escalated due to the COVID-19 pandemic, as a result of significant staffing shortages, long wait times, and inadequate security and mental health support.²⁵ Workplace violence in the setting of chronic pain management is consistent with that seen in other fields of medicine. Studies have demonstrated that clinicians in this specialty experience workplace violence and threats of violence on a frequent basis, particularly within the context of complex medical situations, opioid management, and Workers' Compensation cases.²⁶

Consequently, the overwhelming impact of COVID-19 on healthcare systems and personnel has compounded a specialty, sufficiently challenging at its baseline, with substantial stressors, leading to critical emotional distress amongst its providers. While rates of burnout, insomnia,

and depression have increased amongst healthcare workers during this pandemic,²⁷ research depicting the exact impact of COVID-19 across pain medicine providers' mental health is needed. Previous data highlighting higher levels of burnout within this population compel us to hypothesize that this critical issue has only become worse.

Emotional distress and its associated consequences, including burnout, exert a multilayered effect. On an individual level, providers who experience burnout have a drastically reduced sense of well-being and are at increased risk for developing depression, substance abuse disorder, heart disease, and somatic symptoms.²⁸ Systemically, the implications of elevated rates of physician distress impact patient care and healthcare institutions. Burnout has been previously linked to suboptimal patient care, longer recovery times, and decreased patient satisfaction.²¹ Within healthcare institutions, burnout can lead to early retirement, job dissatisfaction, career change, and elevated turnover rates, likely due to augmented stress and significantly reduced professional fulfillment.^{21,29,30} A recent study including 20,000 clinical and non-clinical healthcare workers across the US indicated that 24% of physicians and 40% of nurses reported a moderate, likely, or definite intent to leave their current practices by 2022.²² Given the shortage of trained pain physicians and lengthy wait times prior to accessing treatment that were identified well prior to the onset of the COVID-19 crisis,³¹ this phenomenon spells potential disaster for many suffering from chronic pain.

The central question becomes: how can we address and effectively improve emotional distress amongst clinicians in pain medicine? This is by no means a simple and straightforward question, nor are we assuming to know its full answer. However, there are concrete steps we can take now, as clinicians and institutions, to validate and care for the suffering of those who care for others. First and foremost, effective change emerges from awareness, which calls for a deep understanding of which stressors most saliently contribute to emotional distress and burnout in providers who care for chronically ill patients. Once we have an accurate depiction of the roots of emotional distress within our specialty, we will be able to tackle the issue similarly to how it presents itself, ie, in a multilayered manner. That said, the emotional distress that providers are currently experiencing needs to be addressed with individual, team, institutional and systemic strategies. For example, a randomized clinical trial conducted prior to the pandemic determined that offering

clinicians protected time to address specific targets of emotional distress, including self-awareness and coping strategies, was successful in decreasing rates of depersonalization and increasing engagement at work.³² This study's findings illustrate the importance of associating individual and institutional initiatives to mitigate the current crisis of physician burnout and dissatisfaction, given that there is a collective responsibility to foster healthcare workers' well-being.³²

In sum, we believe that examining emotional distress amongst pain medicine clinicians will not only equip our specialty to better care for its providers, but will also shed light to the issue of mental health stigma amongst healthcare workers. The truth is that clinicians' emotional distress was already a public health crisis long before COVID-19. As we are now bedeviled with a pandemic that has heightened the suffering amongst those who dedicate their lives to caring for others, we urge individuals, institutions, and governmental bodies to implement concrete and effective changes that will attend to and mitigate clinicians' emotional distress.

Disclosure

Dr Michael E Schatman is a research consultant for Modoscript, outside the submitted work. The authors report no other conflicts of interest in this work.

References

- Schatman ME, Fortino MG. The problem (and the answer?) to the limited availability of pain psychologists: can clinical social workers help? *J Pain Res.* 2020;13:3525–3529. doi:10.2147/JPR.S297312
- Gruß I, Firemark A, McMullen CK, Mayhew M, DeBar LL. Satisfaction with primary care providers and health care services among patients with chronic pain: a mixed-methods study. *J Gen Intern Med.* 2020;35(1):190–197. doi:10.1007/s11606-019-05339-2
- Kim K, Rendon I, Starkweather A. Patient and provider perspectives on patient-centered chronic pain management. *Pain Manag Nurs.* 2021;22(4):470–477. doi:10.1016/j.pmn.2021.02.003
- Eccleston C, Blyth FM, Dear BF, et al. Managing patients with chronic pain during the COVID-19 outbreak: considerations for the rapid introduction of remotely supported (eHealth) pain management services. *Pain.* 2020;161(5):889–893. doi:10.1097/j.pain.0000000000001885
- Cohen SP, Baber ZB, Buvanendran A, et al. Pain management best practices from multispecialty organizations during the COVID-19 pandemic and public health crises. *Pain Med.* 2020;21(7):1331–1346. doi:10.1093/pm/pnaa127
- Shanthanna H, Strand NH, Provenzano DA, et al. Caring for patients with pain during the COVID-19 pandemic: consensus recommendations from an international expert panel. *Anaesthesia.* 2020;75(7):935–944. doi:10.1111/anae.15076
- Hagedorn JM, Moeschler S, Furnish T, Sayed D, Durbhakula S. Impact of COVID-19 on pain medicine fellowship training. *Reg Anesth Pain Med.* 2021;46(2):188–189. doi:10.1136/rapm-2020-101534
- Herzberg DL, Sukumaran HP, Viscusi E. NSAIDs for analgesia in the era of COVID-19. *Reg Anesth Pain Med.* 2020;45(9):677–678. doi:10.1136/rapm-2020-101584
- Manchikanti L, Kosanovic R, Vanaparthi R, et al. Steroid distancing in interventional pain management during COVID-19 and beyond: safe, effective and practical approach. *Pain Physician.* 2020;23(4S):S319–S350. doi:10.36076/ppj.2020/23/S319
- Chakravarthy K, Strand N, Frosch AS, et al. Recommendations and guidance for steroid injection therapy and COVID-19 vaccine administration from the American Society of Pain and Neuroscience (ASPEN). *J Pain Res.* 2021;14:623–629. doi:10.2147/JPR.S302115
- Kemp HI, Corner E, Colvin LA. Chronic pain after COVID-19: implications for rehabilitation. *Br J Anaesth.* 2020;125(4):436–440. doi:10.1016/j.bja.2020.05.021
- Shapiro H, Kulich RJ, Schatman ME. Manifestation of Borderline Personality symptomatology in chronic pain patients under stress: an understated and exacerbated consequence of the COVID-19 crisis. *J Pain Res.* 2020;13:1431–1439. doi:10.2147/JPR.S264761
- Coleman BC, Kean J, Brandt CA, Peduzzi P, Kerns RD. Adapting to disruption of research during the COVID-19 pandemic while testing nonpharmacological approaches to pain management. *Transl Behav Med.* 2020;10(4):827–834. doi:10.1093/tbm/ibaa074
- Cisternas AF, Ramachandran R, Yaksh TL, Nahama A. Unintended consequences of COVID-19 safety measures on patients with chronic knee pain forced to defer joint replacement surgery. *Pain Rep.* 2020;5(6):e855. doi:10.1097/PR9.0000000000000855
- Hruschak V, Flowers KM, Azizoddin DR, Jamison RN, Edwards RR, Schreiber KL. Cross-sectional study of psychosocial and pain-related variables among patients with chronic pain during a time of social distancing imposed by the coronavirus disease 2019 pandemic. *Pain.* 2021;162(2):619–629. doi:10.1097/j.pain.0000000000002128
- Lang-Ilievich K, Rumpold-Seitlinger G, Szilagyí IS, et al. Biological, psychological, and social factors associated with worsening of chronic pain during the first wave of the COVID-19 pandemic: a cross-sectional survey. *Br J Anaesth.* 2021;127(1):e37–e39. doi:10.1016/j.bja.2021.04.010
- Mullins CF, Harmon D, O'Connor T. Quantifying the impact of COVID-19 on chronic pain services in the Republic of Ireland. *Ir J Med Sci.* 2021;5:1–5.
- Ayad AE. Opioid epidemics during the pandemic: further insights to the same story. *J Opioid Manag.* 2021;17(1):9–12. doi:10.5055/jom.2021.0609
- Hale M, Garofoli M, Raffa RB. Benefit-risk analysis of buprenorphine for pain management. *J Pain Res.* 2021;14:1359–1369. doi:10.2147/JPR.S305146
- Kroll HR. A preliminary survey examining predictors of burnout in pain medicine physicians in the United States. *Pain Physician.* 2016;19(5):E689–696. doi:10.36076/ppj/2019.19.E689
- Aron R, Pawlowski J, Shukry M, Shillcutt S. The Impact of COVID-19 on the status of the anesthesiologists' well-being. *Adv Anesth.* 2021;39:149–167. doi:10.1016/j.aan.2021.07.009
- Sinsky CA, Brown RL, Stillman MJ, Linzer M. COVID-related stress and work intentions in a sample of US health care workers. *Mayo Clin Proc Innov Qual Outcomes.* 2021;5(6):1165–1173.
- Bradley M, Chahar P. Burnout of healthcare providers during COVID-19. *Cleve Clin J Med.* 2020. doi:10.3949/ccjm.87a.ccc051
- Almeida M. Burnout and the mental health impact of COVID-19 in anesthesiologists: a call to action. *J Clin Anesth.* 2021;68:110084. PMID: 33038719; PMCID: PMC7524672. doi:10.1016/j.jclinane.2020.110084
- The Joint Commission. Physical and verbal violence against health care workers. *Sentinel Event Alert.* 2018;59:1–9.
- Moman RN, Maher DP, Hooten WM. Workplace violence in the setting of pain management. *Mayo Clin Proc Innov Qual Outcomes.* 2020;4(2):211–215.

27. Shanafelt T, Ripp J, Trockel M. Understanding and addressing sources of anxiety among health care professionals during the COVID-19 pandemic. *JAMA*. 2020;323(21):2133–2134. doi:10.1001/jama.2020.5893
28. Romito BT, Okoro EN, Ringqvist JRB, Goff KL. Burnout and wellness: the anesthesiologist's perspective. *Am J Lifestyle Med*. 2020;15(2):118–125. doi:10.1177/1559827620911645
29. O'Brien JM, Goncin U, Ngo R, Hedlin P, Chakravarti A. Professional fulfillment, burnout, and wellness of anesthesiologists during the COVID-19 pandemic. *Can J Anaesth*. 2021;68(5):734–736. doi:10.1007/s12630-021-01916-4
30. Schneider BJ, Ehsanian R, Schmidt A, et al. The effect of patient satisfaction scores on physician job satisfaction and burnout. *Future Sci OA*. 2020;7(1):FSO657. doi:10.2144/fsoa-2020-0136
31. Breuer B, Pappagallo M, Tai JY, Portenoy RK. U.S. board-certified pain physician practices: uniformity and census data of their locations. *J Pain*. 2007;8(3):244–250. doi:10.1016/j.jpain.2006.08.009
32. West CP, Dyrbye LN, Rabatin JT, et al. Intervention to promote physician well-being, job satisfaction, and professionalism: a randomized clinical trial. *JAMA Intern Med*. 2014;174(4):527–533. doi:10.1001/jamainternmed.2013.14387

Journal of Pain Research

Dovepress

Publish your work in this journal

The Journal of Pain Research is an international, peer reviewed, open access, online journal that welcomes laboratory and clinical findings in the fields of pain research and the prevention and management of pain. Original research, reviews, symposium reports, hypothesis formation and commentaries are all considered for publication. The manuscript

management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-pain-research-journal>