

# The Appropriateness of Using Fixed Assay Cut-offs for Estimating Seroprevalence [Letter]

Yuting Da<sup>1,\*</sup>, Yuxuan Wu<sup>2,\*</sup>, Peiqing Quan<sup>1</sup>

<sup>1</sup>Outpatient Surgery Center, Obstetrics and Gynecology Hospital of Fudan University, Shanghai, People's Republic of China; <sup>2</sup>Anhui Medical University, Hefei City, Anhui Province, People's Republic of China

Correspondence: Peiqing Quan, Outpatient Surgery Center, Obstetrics and Gynecology Hospital of Fudan University, No. 128 Shenyang Road, Yangpu District, Shanghai, 200011, People's Republic of China, Email [quanpeiqing@163.com](mailto:quanpeiqing@163.com)

\*These authors contributed equally to this work

## Dear editor

We read this study<sup>1</sup> with great interest. This study aimed to investigate the seroprevalence of West Nile Virus (WNV) antibodies in humans, horses, and pigeons in the Eastern Province of Saudi Arabia. The results showed that the percentage of anti-WNV antibodies in the experiment including humans was found to be 9.6% (3.1% in females and 6.5% in males). This is an interesting finding due to the high prevalence of antibodies detected in human blood samples.

However, we have several points we wanted to mention regarding this study.

1. The cut-off value of seropositive WNV antibodies might need to be discussed in the text. Since there are no definite rules of the choice of the cut-off value,<sup>2</sup> the researchers followed the manufacturer's guides. According to previous studies,<sup>3</sup> the change of cut-off value might also change the sensitivity and specificity of the ELISA test.
2. There is a possibility that detection method might influence the results. The authors have already mentioned the difference of ELISA and microneutralization assay. A study<sup>4</sup> carried out the comparison of different methods for detecting antibody titers in the same blood sample, which is also a referential experiment for WNV future studies.
3. The quantitative detection of WNV antibodies might be reasonable for the prediction model setting of population titer. Currently, we follow the rule of 95% range as a fixed value. If the population-based data can be acquired, the comparison and data retrieval might be possible.
4. Past research<sup>5</sup> has identified that WNV might be transmitted through blood donation. Several confirmed cases in America have identified bloodborne transmission. The researchers could collect the clinical data of blood transfusions to determine the possibility of this route.

Nevertheless, thanks to the authors, we have evidence of anti-WNV antibodies detected in humans and pigeons. In future practice, the high seroprevalence of animals, such as horses and pigeons, should be taken into consideration in terms of the possibility of animal-human transmission. Besides, the blood bank might also need to confirm the possibility of bloodborne transmission.

## Disclosure

The authors report no conflicts of interest in this communication.

## References

1. Alkharsah KR, Al-Afaleq AI. Serological evidence of West Nile virus infection among humans, horses, and pigeons in Saudi Arabia. *Infect Drug Resist.* 2021;14:5595–5601. doi:10.2147/IDR.S348648

2. Kafatos G, Andrews NJ, McConway KJ, Maple PAC, Brown K, Farrington CP. Is it appropriate to use fixed assay cut-offs for estimating seroprevalence? *Epidemiol Infect.* 2016;144(4):887–895. doi:10.1017/S0950268815001958
3. Kim C-M, Kim D-M, Yun NR. Follow-up investigation of antibody titers and diagnostic antibody cutoff values in patients with scrub typhus in South Korea. *BMC Infect Dis.* 2021;21(1):69. doi:10.1186/s12879-020-05735-8
4. Safaian M, Ghosh A, Porras C, et al. Direct comparison of HPV16 serological assays used to define HPV-naïve women in HPV vaccine trials. *Cancer Epidemiol Biomarkers Prev.* 2012;21(9):1547–1554. doi:10.1158/1055-9965.EPI-12-0558
5. Centers for Disease Control and Prevention. Investigations of West Nile virus infections in recipients of blood transfusions. *MMWR Morb Mortal Wkly Rep.* 2002;51(43):973–974.

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Infection and Drug Resistance 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Infection and Drug Resistance editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

## Infection and Drug Resistance

Dovepress

### Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>

<https://doi.org/10.2147/IDR.S355939>