Comparison of Current Methods with Neutrophil-to-Lymphocyte Ratio in Predicting Stroke-Associated Pneumonia [Letter]

Jingfang Zou, Guangting Qiu

Department of Neurosurgery, Shanghai Fourth People's Hospital Affiliated to Tongji University School of Medicine, Shanghai, People's Republic of China

Correspondence: Guangting Qiu, Department of Neurosurgery, Shanghai Fourth People's Hospital Affiliated to Tongji University School of Medicine, No. 1279 Sanmen Road, Hongkou District, Shanghai, 200434, People's Republic of China, Tel/Fax +86-21-63240090, Email 18939739562@163.com

Dear editor

We read this study¹ with great interest. The concept of stroke-associated pneumonia (SAP) was first proposed by Hilker in 2003.² It is one of the important risk factors for death after stroke, and it increases the length of hospital stay and medical expenses, which brings a heavy burden to the family and society. In this article, the authors select a common clinical parameter in comparison of C reactive protein level for the prediction of the incidence of SAP. This is a novel idea and could be promoted in clinical practice.

However, there might be three points need to be mentioned for the readers.

1. The SAP criteria is slightly different compared to previous guidelines.³ In this study, the authors cited Chinese domestic expert consensus of SAP diagnosis. This might lead to slight bias in future research. Better illustrated data regarding the usage of different criteria might be a more convincing article.
2. For the application of prediction of SAP, we need to be more careful in first comparing the non-superiority of this prediction model with current available scores.

Since 2012, a number of studies⁴ have used multivariate regression models to design different SAP prediction scores, based on stroke-related risk factors, including stroke-induced immunosuppression, dysphagia, age, gender, smoking, stroke severity, stroke type, stroke location, level of consciousness, feeding method, application of acid inhibitors, admission to the intensive care department, combined with hypertension, diabetes, history of chronic respiratory diseases, and history of atrial fibrillation, etc. Among these risk factors, stroke-induced immunosuppression and dysphagia are important independent risk factors for SAP.

In 2018, a multicenter observational study⁵ of external validation of the SAP predicting score ISAN, A2DS2, and AIS-APS was carried out, with the conclusion of recommending AIS-APS score for future clinical trials.

Future comparison of the clinical parameter of Neutrophil-to-Lymphocyte Ratio with AIS-APS score will be more convincing for the neurologists in clinical practice.

1. A study by Cheng et al⁶ showed that higher monocyte-to-lymphocyte ratio (MLR) was significantly associated with SAP in AIS patients. Another study by Cao et al⁷ shows that a high MLR is an independent risk factor for SAP and has a predictive value for severe pneumonia in patients with SAP. Compared to this article, the authors use neutrophils rather than monocytes in the ratio. A comparison between these two might also be helpful in determining the clinical significance.

Nevertheless, thanks to the authors, we have a better understanding in interpreting the clinical values of the auxiliary examination. SAP-related research fields have accumulated more clinical evidence, especially the evidence from the Chinese population, and our understanding of SAP has been further deepened.

Acknowledgment

We would like to thank Professor Jiang Zhao for his contribution of this article.

Disclosure

The authors report no conflicts of interest for this communication.

References

1. Wu T, Zhang H, Tian X, Cao Y, Wei D, Wu X. Neutrophil-to-lymphocyte ratio better than high-sensitivity C-reactive protein in predicting stroke-associated pneumonia in afebrile patients. *Neuropsychiatr Dis Treat*. 2021;17:3589–3595. doi:10.2147/NDT.S340189
2. Hilker R, Poetter C, Findeisen N, et al. Nosocomial pneumonia after acute stroke: implications for neurological intensive care medicine. *Stroke*. 2003;34(4):975–981. doi:10.1161/01.STR.0000063373.70993.CD
3. Smith CJ, Kishore AK, Vail A, et al. Diagnosis of stroke-associated pneumonia: recommendations from the pneumonia in Stroke Consensus Group. *Stroke*. 2015;46(8):2335–2340. doi:10.1161/STROKEAHA.115.009617
4. Ji R, Shen H, Pan Y, et al. Novel risk score to predict pneumonia after acute ischemic stroke. *Stroke*. 2013;44(5):1303–1309. doi:10.1161/STROKEAHA.111.000598
5. Zapata-Arriaza E, Moniche F, Blanca P-G, et al. External validation of the ISAN, A2DS2, and AIS-APS scores for predicting stroke-associated pneumonia. *J Stroke Cerebrovasc Dis*. 2018;27(3):673–676. doi:10.1016/j.jstrokecerebrovasdis.2017.09.059
6. Cheng H-R, Song J-Y, Zhang Y-N, et al. High monocyte-to-lymphocyte ratio is associated with stroke-associated pneumonia. *Front Neurol*. 2020;11:575809. doi:10.3389/fneur.2020.575809
7. Cao F, Wan Y, Lei C, et al. Monocyte-to-lymphocyte ratio as a predictor of stroke-associated pneumonia: a retrospective study-based investigation. *Brain Behav*. 2021;11(6):e02141. doi:10.1002/brb3.2141

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Neuropsychiatric Disease and Treatment 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Neuropsychiatric Disease and Treatment editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Neuropsychiatric Disease and Treatment

Dovepress

Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS, and is the official journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>

<https://doi.org/10.2147/NDT.S355854>