eQTL Highlights the Potential Role of Negative Control of Innate Immunity in Kawasaki Disease

Sirui Song¹, Liqin Chen¹, Qianqian Ning², Danying Zhu¹, Feng Qiu², Guang Li², Hong Zhang³, Tingting Xiao¹, Guohui Ding⁴, Min Huang¹

¹Department of Cardiology, Shanghai Children's Hospital, Shanghai Jiao Tong University, Shanghai, People's Republic of China; ²Shanghai Center for Bioinformation Technology, Shanghai, People's Republic of China; ³Department of Laboratory, Shanghai Children's Hospital, Shanghai Jiao Tong University, Shanghai, People's Republic of China; ⁴International Human Phenome Institutes, Shanghai, 200235, People's Republic of China

Correspondence: Min Huang, Department of Cardiology, Shanghai Children's Hospital, Shanghai Jiao Tong University, Shanghai, People's Republic of China, Email huangmin@sjtu.edu.cn

Guohui Ding, International Human Phenome Institutes, Shanghai, 200235, People's Republic of China, Email dingguohui@ihup.org.cn

Purpose: Kawasaki disease (KD) is an acute systemic vasculitis mainly found in the medium-sized arteries, especially the coronary arteries. Immune system is involved in the pathogenesis of acute KD in children, but the functional differences in the immune system between healthy children and KD patients remain unclear.

Patients and Methods: A total of 190 KD patients and 119 healthy controls were recruited for the next-generation sequencing of 512 targeted genes from 4 immune-related pathways. Subsequently, the peripheral blood mononuclear cells (PBMCs) were isolated. RNA sequencing of the LPS treated PBMCs from additional 20 KD patients and 20 healthy controls was used to examine the differentially expressed genes (DEGs). Then, an expression quantitative trait locus (eQTL) analysis combined with previously analyzed RNA data were used to examine the DEGs. Finally, the serum levels of 13 cytokines were detected before and after LPS treatment in 40 samples to confirm the findings from eQTL analysis.

Results: A total of 319 significant eQTL were found, and both eQTL analysis and RNA sequencing showed some DEGs were involved in the connective tissue disorders and inflammatory diseases. DEGs that function to negatively regulate immunity were closely related to the pathogenesis of KD. In addition, the serum levels of IL-10 (an inflammatory and immunosuppressive factor) and SCD25 (an important immunosuppressant) reduced significantly in the KD patients.

Conclusion: Our study shows the expression of factors responsible for the negative control of innate immunity is altered, which plays an important role in the etiology of KD.

Keywords: Kawasaki disease, innate immunity, inflammation, next generation sequencing, cytokines

Introduction

Kawasaki disease (KD) as an acute, self-limited disease predominately affects children under 5 years.¹ KD is featured by the development of autoimmune-like polyangiitis and can cause damage to the coronary arteries (including coronary artery aneurysms and long-term cardiovascular sequences).² The incidence of reported KD is increasing over year, and the Asian population has a higher incidence. Moreover, the higher incidence of KD in siblings and twins suggests a genetic predisposition to KD.³ Some studies have investigated the KD-related genes from available databases and revealed some KD-related genes are mainly related to the immune response.⁴

Genome wide association study (GWAS) provides a tool for understanding the genetic changes in the KD. The single nucleotide polymorphisms (SNP) of inositol 1,4,5-trisphosphate 3-kinase C (ITPKC) have been found to be associated with the KD susceptibility and coronary risk in American and Japanese children. ITPKC can negatively regulate T-cell activation through a calcium signaling pathway, and the functional polymorphism of ITPKC may increase T-cell activation and cytokine expression.⁵ The coatomer protein complex beta-2 subunit (COPB2), a factor associated with T cell activity, is also related to the KD susceptibility.⁶ In humoral immunity, the polymorphism of IgG receptor gene

FCGR2A has also been found to be associated with the KD susceptibility in European and Asian populations. Immune cells (such as dendritic cells [DCs], neutrophils, monocytes/macrophages) express Fc fragment of IgG receptor IIa (FCGR2A) on the surface, which can transmit activation signals into cells when the receptor binds to the immune complex, leading to the immunosuppression in some cases.⁷ The self-limitation of KD, the therapeutic effectiveness of intravenous immunoglobulin (IVIG), and the functions of genes identified by GWAS suggest the important role of immunoregulation in the pathogenesis of KD.

GWAS mainly focuses on genome-wide SNPs information. However, few studies have focused on the roles of immunoregulation-related genes in the pathogenesis of KD. In the target capture sequencing, a specific probe is designed to capture the genome region of interest and then used to hybridize with the whole genome DNA; subsequently, the DNA fragments of target genome region are enriched and processed for sequencing. It is cost-effective and capable of finding disease-related genes as well. Meanwhile, the expression quantitative trait loci (eQTLs) is a key to map SNPs that affect gene expression.⁸

The present study was to investigate whether eQTL highlights the potential role of negative control of innate immunity in the KD. First, the genes related to four immune regulation pathways were investigated, and their expressions were compared between KD patients and healthy children. Then, the inflammatory process was simulated to further confirm the differentially expressed genes between the above two groups. Thereafter, the eQTLs analysis was employed to obtain the SNPs affecting gene expression. After stimulation, the secretion of cytokines was also evaluated. Our results showed some of genes and cytokines associated with immunoregulation were involved in the pathogenesis of KD.

Patients and Methods

Patients and Healthy Controls

The patients in this study were enrolled from Shanghai Children's Hospital between December 2015 and October 2017. All patients were recruited after informed consent was signed by their parents or guardians; this study was conducted in accordance with the Declaration of Helsinki and approved by the Ethics Committee of Shanghai Children's Hospital. The KD was diagnosed according to the diagnostic criteria for KD proposed by the American Heart Association in 2017.⁹ Children who had not received IVIG therapy were excluded. In the control group, healthy children who received routine physical examination were recruited from the Department of Outpatient. The healthy controls had no prior immunological disease and heart disease.

The enrolled 309 Han Chinese children in this study were unrelated. All of the 190 patients with KD were included in the next-generation sequencing (NGS) phase and 20 in the peripheral blood mononuclear cells (PBMCs) stimulation phase. Of the 119 healthy controls, 99 were included in the NGS phase, and 20 in the PBMC stimulation phase.

Target Capture Sequencing and Data Analysis

Blood samples (2–4 mL) were obtained from KD patients before IVIG treatment. Genomic DNA was extracted from 300 µL of whole peripheral blood using Genomic DNA Extraction Kit (Cat. No. DP329, TIANGEN Bioscience Beijing) according to the manufacturer's instructions. Two sources of information (one involved in 4 pathways and the other involved SNP previously published) were used to design hybridization probes in this study. First, 512 candidate genes among 4 pathways including Toll-like receptor signaling pathway, cytokine receptor interaction, TGF-beta signaling pathway and T cell receptor signaling pathway were selected. The capture region included all exons and the adjacent 15-bp intron of these genes. Then, 472 GWAS hotspots related to the KD according to GeneReviews (NCBI) were extracted. Sequences ±50 bp flanking the risk loci were included in this analysis. The final custom capture array was synthesized by NimbleGen, Roche. Target enrichment was performed using the custom capture array, and the enriched target fragments were subsequently sequenced on the Illumina HiSeq X10 platform.

The raw reads were mapped to the human reference genome (hg19) using the Burrows-Wheeler alignment (BWA v0.7.15).¹⁰ Then, Picard (v1.135) was used to process for PCR duplicates (<http://broadinstitute.github.io/picard/>). Variants in the target regions were detected using GATK (v3.7) HaplotypeCaller algorithm¹¹ and SNPs with “QD <

2.0 || FS > 60.0 || MQ < 40.0 || MQRankSum < -12.5 || ReadPosRankSum < -8.0" were excluded. Subsequently, functional annotation of filtered variants was performed using the ANNOVAR pipeline.¹²

Collection of Peripheral Blood Mononucleated Cells (PBMCs)

3 mL of blood was collected from 20 KD patients and 20 healthy children and then mixed with Ficoll-Paque PLUS (Cat. No.17544202, General Electric Co., USA), followed by centrifugation at 1000 g for 30 min at 4°C. The interphase PBMCs were collected and washed twice with RPMI 1640 medium (Cat. No.11875093, Life Technology Co., USA) including 10% fetal bovine serum (FBS) (Cat. No.10270106, Life Technology Co., USA). PBMCs were resuspended in plasma from the same person with 10% RPMI (No.11875093 Life Technology Co., USA) and then stored in the liquid nitrogen. The PBMCs were processed for further detection within a month.

LPS Stimulation and Immunoassays

Aliquots were thawed and centrifuged at 1000 rpm for 10 min at 24°C to precipitate cells. The cells were collected and incubated in RPMI 1640 supplemented with 10% FBS for 4 h at 37°C. The cells were collected after filtration and then re-suspended with RPMI at 1×10^6 /mL. The PBMCs were cultured at 1×10^6 cells/mL in an environment with 5% CO₂ at 37°C for 24 h.

To induce inflammation, lipopolysaccharides (LPS) from *Escherichia coli* 0111:B4 (Cat. No. L4391, Sigma Aldrich, USA) was added to the cells at 1 µg/µL, followed by incubation for 22 h. Then, the supernatant and cells were collected independently. The expression of 13 cytokines (IL-8, IL-1β, IL-2, IL-6, IL-10, IL-4, IL-5, IL-12P70, IL-17A, IL-18, TNF-α, INF-γ and SCD25) was detected with the Luminex 200 ProcartaPlex Mix&Match Human-13-plex kit (Affymetrix eBioscience, UK). These 13 cytokines have been previously reported to be associated with KD pathogenesis and play important roles in four immunoregulation pathways.

RNA Sequencing and Data Analysis

Total RNA was extracted from the cells using Trizol reagent and then reversely transcribed into cDNA using the PrimeScriptTM Master mix (Cat. No. RR036a TaKaRa). The RNA library was constructed using NEBNext[®] UltraTM II RNA Library Prep Kit for Illumina[®] kit and submitted to Illumina HiSeq X10 platform.

The data analysis of RNA sequencing was conducted according to previously reported.¹³ Briefly, raw reads from each sample were quality controlled by Q30. After 3' adaptor-trimming and removing low quality reads, the high-quality trimmed reads were mapped to the human reference genome hg19 (<ftp://ftp.ccb.jhu.edu/pub/infphilo/hisat2/data/grch37.tar.gz>) with HISAT v2.1.0. Then, StringTie (version 1.3.3b) was used for transcript assembly and estimating the expression level of each gene with a gene annotation file (ftp://ftp.ensembl.org/pub/release-75/gtf/homo_sapiens/Homo_sapiens.GRCh37.75.gtf.gz). The Fragments Per Kilobase per Million mapped reads (FPKM) was employed to determine gene expression levels. Finally, differentially expressed genes (DEGs) and transcripts were detected by R packages: Ballgown and DESeq. A value of P<0.05 was considered statistically significant.

Variant Filtering and eQTL Analysis

In order to evaluate the impact of filtered SNPs on the KD, association analysis was performed by using the R package SNPAssoc. SNPAssoc estimates the numbers, percentages, odds ratios and P-value for the likelihood ratio test of association. A value of P < 0.05 was considered statistically significant.

To further confirm the SNPs associated with KD, an eQTL analysis was conducted to identify the *cis* eQTLs using FastQTL (v2.184) in 20 KD samples. If the SNP location was within 1 Mb upstream or downstream of a gene's transcription starting site, this *cis* eQTL would be computed. FastQTL proposes an efficient permutation procedure to control for multiple testing and an adjusted P-value can be estimated. A significance threshold was set at 0.05, and therefore eQTLs with P-value<0.05 were considered statistically significant. The eQTL with the lowest FDR was selected for each gene to represent the best gene/eSNP pair.

Statistical Analysis

The Ingenuity Pathway Analysis (IPA) was carried out in order to analyze the molecular function, tox function and disease disorder of DEGs. The cytokine expression levels were analyzed using SPSS version 23.0 (IBM) and a value of two-sided $P < 0.05$ was considered statistically significant.

Results

Target Capture Sequencing of KD Patients

This study focused on the exon regions of 512 genes in the 4 KEGG pathways, including T cell receptor signaling pathway, Toll-like receptor signaling pathway, cytokine–cytokine interaction pathway and TGF- β signal pathway, and 472 risk loci of KD reported in previous GWAS studies ([Supplementary Table S1](#)). A total of 190 KD patients were recruited, including 34 patients with coronary artery lesions and 26 patients with IVIG resistance. The targeted regions were sequenced with an average depth of 440X ([Supplementary Table S2](#)).

A total of 12,470 SNPs ([Supplementary File S1](#)) were found in all samples. The annotation of SNPs showed that they consisted of 2247 nonsynonymous SNPs, 41 stopgain, 1 stoploss and others that do not affect the gene function ([Supplementary Tables S3](#) and [S4](#)). To confirm the risk SNPs, SNPpassco was employed to investigate the association of selected SNPs with KD.

A total of 2247 SNPs affected gene expression. Burden test showed there were 7 genes SNV sets that were closely associated with KD ($P < 0.05$), including 216 SNVs mutations ([Table 1](#)).

RNA Sequencing of PBMCs After LPS Stimulation

In the present study, LPS was used to stimulate PBMCs to simulate the process of superantigen activation. After the stimulation, the PBMCs were separated from 20 KD patients (including 8 patients with coronary artery lesions and 1 patient with IVIG resistance) and 20 controls. Total RNA was extracted, transcriptome data were compared to the reference genome, and 10 KD patients and 10 controls were selected for RNAseq data with a comparison ratio of more than 90%. The ballgown was used to screen out the genes with significant differences in the transcriptome data between KD patients and controls ([Table 2](#)). IPA software was used to analyze the physiological functions of DEGs and their roles in clinical disease processes.

IPA annotation showed these DEGs were related to the immune cell trafficking and cardiovascular system development and function ([Figure 1](#)). Then, the functions of genes with the most significant differences were further investigated. RUNX3 gene is related to the differentiation of regulatory T cells (Tregs), which is involved in the immune regulation.¹⁴ Results showed DEGs were mainly related to the connective tissue disorders and inflammatory diseases, which were consistent with the clinical manifestations of KD ([Figure 2](#)).

Table 1 The Number of SNVs Included in Differentially Expressed Genes Between KD Patients and Controls

Gene Name	SNVs
CD84	62
ELF5	2
IL-10	12
PIK3CG	38
PRKCQ	26
PTPN6	19
SMAD2	57

Table 2 Significant Differentially Expressed Genes Between KD Patients and Controls After RNAseq

Gene_Name	Gene_Id	fc	pval	qval
GIT1	ENSG00000108262	1.931794085	1.41E-06	0.016026
MIR500B	ENSG00000239057	5.054199512	3.43E-06	0.023294
RUNX3	ENSG00000020633	2.573926017	5.80E-06	0.028173
PLEKHG1	ENSG00000120278	1.703894538	8.29E-06	0.034766
SRCAP	ENSG00000080603	3.24041503	9.20E-06	0.034766
PLXND1	ENSG00000004399	3.520021248	1.22E-05	0.04161

eQTL Associated DNA and RNA Analysis

SNV sites in 20 KD patients that identified by the above-mentioned targeted sequencing showed correlation with the DEGs in the transcriptome sequencing data. A total of 356 SNVs were found to be associated with 313 DEGs (Table 3).

Then, 313 DEGs were subjected to IPA analysis. In terms of physiological functions, they were involved in the connective tissue development, cardiovascular system development and cell-mediated immune response (Figure 3). In terms of effects on the diseases, they were related to the connective tissue diseases, dermatological diseases and

Figure 1 Enrichment analysis of DEGs related to the physiological development and function in KD patients and controls. **Figure 2** Enrichment analysis of DEGs in disease disorder of KD patients and controls.

Table 3 eQTL Analysis for the Examination of DEGs

GeneSymbol	GeneID	Foldchange	Pvalue	Qvalue	SNVChromosome	SNVSite	SNVAccession	RefAllele	AltAllele	eQTLpvalue	eQTLslope
PIK3CD	ENSG00000171608	3.245557	<0.01	0.022091	chr1	9,784,423	rs11121484	C	T	0.04	-15.5955
PIK3CD	ENSG00000171608	3.245557	<0.01	0.022091	chr1	9,788,153	rs1141402	G	A	<0.01	-15.2027
ITPR3	ENSG00000096433	2.995176	<0.01	0.041427	chr6	33,638,180	rs2229634	C	T	0.01	2.50358
PRKCZ	ENSG00000067606	2.206851	<0.01	0.067363	chr1	1,148,100	rs34108055	G	C	0.02	3.85754
MAP2K1	ENSG00000169032	2.828066	<0.01	0.067363	chr15	67,361,308	rs7162912	T	G	0.05	-3.33218
MAP2K1	ENSG00000169032	2.828066	<0.01	0.067363	chr15	67,407,899	rs1438386	G	A	0.01	4.2629
ITGB2	ENSG00000160255	2.612719	<0.01	0.07445	chr21	46,330,628	rs2280965	C	T	0.03	13.775
MAVS	ENSG00000088888	2.341154	<0.01	0.115362	chr20	3,684,022	rs611847	A	G	0.02	-3.73908
MAVS	ENSG00000088888	2.341154	<0.01	0.115362	chr20	3,684,729	rs625372	T	C	0.02	-3.94639
OXT	ENSG00000101405	2.732799	<0.01	0.116585	chr20	3,682,242	rs45472900	C	T	<0.01	8.47991
CDC25B	ENSG00000101224	2.521221	<0.01	0.123854	chr20	3,668,082	rs656635	G	T	0.04	-8.13729
CDC25B	ENSG00000101224	2.521221	<0.01	0.123854	chr20	3,675,333	rs3746638	G	A	0.02	-5.0302
CDC25B	ENSG00000101224	2.521221	<0.01	0.123854	chr20	3,675,498	rs709012	T	G	0.01	-5.95212
CDC25B	ENSG00000101224	2.521221	<0.01	0.123854	chr20	3,677,736	rs673114	T	G	0.02	-5.0302
MAP4K1	ENSG00000104814	2.088039	<0.01	0.138198	chr19	39,398,631	rs3136644	A	G	<0.01	11.3568
MAP4K1	ENSG00000104814	2.088039	<0.01	0.138198	chr19	39,398,649	rs3136645	T	C	<0.01	8.36483
MAP4K1	ENSG00000104814	2.088039	<0.01	0.138198	chr19	39,399,195	rs1044284	A	G	<0.01	9.63068
MYBBP1A	ENSG00000132382	2.02601	<0.01	0.169123	chr17	5,412,361	rs8079034	C	T	0.03	7.0365
MYBBP1A	ENSG00000132382	2.02601	<0.01	0.169123	chr17	5,412,473	rs8079727	A	C	0.02	7.36876
MYBBP1A	ENSG00000132382	2.02601	<0.01	0.169123	chr17	5,412,822	rs16954813	G	A	0.02	7.46041
MYBBP1A	ENSG00000132382	2.02601	<0.01	0.169123	chr17	5,425,077	rs11651270	T	C	0.02	7.58837
DYRK1B	ENSG00000105204	2.146684	<0.01	0.183556	chr19	39,399,195	rs1044284	A	G	0.05	4.1065
MIR146B	ENSG00000202569	2.258973	<0.01	0.19091	chr10	1.05E+08	rs943037	C	T	0.04	12.6514
MIR146B	ENSG00000202569	2.258973	<0.01	0.19091	chr10	1.05E+08	rs10430665	C	T	0.04	12.6514
MIR146B	ENSG00000202569	2.258973	<0.01	0.19091	chr10	1.05E+08	rs10786736	G	C	0.04	12.6514
MIR146B	ENSG00000202569	2.258973	<0.01	0.19091	chr10	1.05E+08	rs17094683	G	T	0.04	12.6514

Figure 3 DEGs associated with SNVs involved in physiological development and function.

inflammatory diseases, which are consistent with the clinical phenotype of mucocutaneous damage in the KD at acute phase (Figure 4). The genes with the most significant changes in their expression (such as *PIK3CD*, *ITPR3* and *PRKCZ*) also play an important role in the immune regulation.^{15–17}

Expression of Immunoregulation Related DEGs Among 40 Samples

After expression analysis and eQTL associated analysis, four immunoregulation-related DEGs (*RUNX3*, *PIK3CD*, *ITPR3* and *PRKCZ*) were identified. Further examination showed their expression significantly decreased in the KD patients as compared to controls (Figure 5).

Cytokine Analysis After LPS Stimulation of PBMCs

The expression of 13 cytokines, including proinflammatory cytokines, anti-inflammatory cytokines, and cytokines previously reported to be associated with KD, was measured in the supernatant of cells after stimulation. Although the expression of immunoregulation-related cytokines IL-4 ($Z=-2.39$, $P=0.02$), SCD25 ($Z=-2.79$, $P=0.02$), and IL-10 ($Z=-3.10$, $P<0.01$) significantly increased after LPS stimulation, their expression was still lower in the KD group than in the control group (Figure 6C,G and K). The expression of other proinflammatory cytokines (such as TNF- α , INF- γ and IL-6) was comparable between KD patients and controls (Figure 6E,L and M).

Figure 4 DEGs associated with SNVs function in disease order.

Figure 5 Expression of immunoregulation related DEGs decreased in the KD patients.

Discussion

KD is a complex disease. Some studies have shown that the pathogenesis of KD is related to the immunological dysfunction. The genetic susceptibility is activated by antigens, causing immune activation and inflammatory reaction, which then produce corresponding clinical symptoms and vascular injury.¹⁸ However, the self-limitation in KD suggests the activation of anti-inflammatory pathways to regulate the immune system.

The mechanism underlying the IVIG-regulated inflammatory response in KD is still poorly understood. Studies have shown that, in vitro IVIG can negatively regulate T cell receptor signaling pathway, thereby inhibiting the proliferation and function of T lymphocytes.¹⁹ Thus, the present study aimed to investigate whether these genes associated with anti-inflammatory pathways were also involved in the pathogenesis of KD. Similarly, in Toll-like receptor pathway, stimulation of TLR2/TLR6 expression on mouse DCs induces their differentiation into resistant DCs which then secrete IL-10 to promote T-cell differentiation into Tregs.²⁰ Tregs production in mice is also induced by synergistic activation of T cells and smad3 nuclear factors in the TGF- β signaling pathway.^{21,22} These may help reduce the inflammatory

Figure 6 Expression of cytokines in the PBMCs from KD patients and controls before and after LPS stimulation (A–M).

response. There is evidence showing that the cytokine storm is present in the acute phase of KD, and cytokine interactions also influence the inflammatory response.^{23,24} In the present study, the chip covering genes from those four pathways was used. After sequencing, results showed the expression of some genes related to immune regulation was significantly different between KD patients and healthy controls. In the inflammatory response secondary to LPS stimulation, the DEGs between two groups mainly concentrated on the immune system and cardiovascular system. After eQTL association analysis, the genes that had mutation sites affecting their expression concentrated on those two systems. All of them were associated with vasculitis and the inflammatory response in the coronary arteries of KD.

RUNX3 was differentially expressed between KD group and control group. Although no evidence has indicated that its expression is affected by the mutation site, but, as a member of the *RUNX* gene, Rudra et al²⁵ found *RUNX* expression was up-regulated in the Treg cells, while that of *Foxp3* (a gene regulating Treg) decreased when *RUNX* was silenced. Thus, *RUNX* may affect the number and function of T cell subsets. In tumor-induced CD8⁺ Treg cells, *RUNX3* enhances the promoter activity of *Foxp3*.²⁶ In this study, our results showed that *RUNX3* expression decreased in the KD group, which may affect the immunomodulatory function of KD and be related to the KD pathogenesis. The targeted capture sequencing showed the expression of *PIK3CD*, *ITPR3* and *PRKCZ* in the KD group decreased and was affected by the mutant loci. Studies have shown that the *PIK3CD* family is involved in the regulation of cell proliferation, differentiation, apoptosis and glucose transport.^{27–29} In the primary immunodeficiency disease, there is *PIK3CD* gene mutation, which leads to the T lymphocyte injury and aging, and compromises the immune regulation.¹⁷ *ITPR3* receives the signals from second messenger *ITP*. *ITP* can be phosphorylated by *ITPKC* and negatively regulate the T cell signaling pathway by regulating Ca²⁺/nuclear factors. Huang et al reported that SNP of *ITPR3* gene (rs2229634) was related to KD in Taiwan children with concomitant coronary artery lesions (CAL).³⁰ This study showed that this SNV site was related to KD. In addition, *ITPR3* gene is involved in the regulation of T lymphocyte apoptosis, can activate T lymphocytes and plays an important role in the progression of KD.¹⁵ It is also related to a variety of autoimmune diseases, such as systemic lupus erythematosus and rheumatoid arthritis.³¹ *PRKCZ* affects *PKCζ* expression. As an isozyme of *PKC* family, *PKCζ* can regulate T cell division, and affect subsequent T cell function.¹⁶ Pro-mitogen-activated protein kinases (MAPKs) also play a key role in the cytokine secretion in the Tregs via a PKC-dependent manner. In neonates with immature immunity or patients with immune deficiency, the defective *PKCζ* expression and IL-12 production may be two main events related to the T cell function.³² *PKCζ* may affect the secretion of cytokines in T cells, which also affects the susceptibility to autoimmune disease and is involved in the disease progression. The reduced expression of these genes may be associated with the hyperactive immune response and compromised immunoregulation in the KD.

The expression of these cytokines was further detected. Results showed, among 13 cytokines, the expression of IL-10 and SCD25 (IL-2R) related to immunoregulation was significantly lower in the KD group after LPS stimulation than in the control group. On the contrary, some pro-inflammatory cytokines have been previously reported to be related to the KD, but there was no significant difference in the expression of these cytokines. This may be ascribed to the difference between in vitro and in vivo experiments. Many studies have suggested that IL-10 is an important cytokine in the pathogenesis of KD.³³ The IL-10 gene not only showed different SNV sites between KD group and control group, but also had decreased protein expression. IL-10 plays an important role in the immunoregulation and immunotherapy through Treg cells.³⁴ SCD25 (IL-2R) is an important cytokine related to the Treg cells and effector T cells.³⁵ As a part of the IL-2 receptor, SCD25 (IL-2R) can recognize T cell activity and has become a biomarker of some immune diseases and tumors.³⁶ Treg cells can secrete IL-2 which together with SCD25 (IL-2R) plays an immunomodulatory role.³⁷ The decreased SCD25 (IL-2R) expression after LPS stimulation indicates that the decreased activity of this pathway compromises the regulatory effect on the KD. There were no significant differences in other proinflammatory cytokines (INF-γ, TNF-α and IL-6) after LPS stimulation. Although previous studies have reported that these cytokines are significantly elevated in a variety of inflammatory diseases including KD, no studies have shown the dynamic changes of these cytokines in the KD at acute stage, and these cytokines have not yet become specific biological markers for KD diagnosis. Therefore, we speculate that the immunomodulatory related pathways and cytokines may become a breakthrough for exploring KD pathogenesis and biological markers for the diagnosis of KD.

There were still limitations in the present study. Although some immunoregulatory molecules were found to be involved in the pathogenesis of KD, they mainly concentrated on the immune system and cardiovascular system.

However, the DEGs affected by the SNVs have never been reported to be associated with KD, and the specific relationship between the hypothesized genes and cytokines has not been revealed. The genes related to anti-inflammatory pathways have no relationships with clinical phenotypes, disease severity, coronary artery lesions and alterations after IVIG therapy. Therefore, the mechanism by which these genes act in the pathogenesis of KD should be further investigated. In addition, the sample size was small, which limits the expansion of results.

Conclusions

Our study shows some genes and cytokines associated with immunoregulation are related to the pathogenesis of KD, which provides evidence for the development of therapies for KD.

Acknowledgments

This study was supported by the Hospital Development Center Research Funding (No SHDC12016119), Shanghai Science and Technology Committee Research Funding (No 17411954300), Shanghai Municipality Health Commission Research Funding (No 2019SY025, 20214Y0477), Shanghai Children's Hospital Clinical Research and Cultivation Funding (No 2019YLYZ01), and Open Project Program of Key Laboratory of Maternal and Child Diseases and Birth Defects, Sichuan University, Ministry of Education (No 2019KF02).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Dimitriadis VR, Brown AG, Gedalia A. Kawasaki disease: pathophysiology, clinical manifestations, and management. *Curr Rheumatol Rep*. 2014;16(6):423. doi:10.1007/s11926-014-0423-x
2. Cheung YF. Vascular health late after Kawasaki disease: implications for accelerated atherosclerosis. *Korean J Pediatr*. 2014;57(11):472–478. doi:10.3345/kjp.2014.57.11.472
3. Singh S, Vignesh P, Burgner D. The epidemiology of Kawasaki disease: a global update. *Arch Dis Child*. 2015;100(11):1084–1088. doi:10.1136/archdischild-2014-307536
4. Lv YW, Wang J, Sun L, et al. Understanding the pathogenesis of Kawasaki disease by network and pathway analysis. *Comput Math Methods Med*. 2013;2013:989307. doi:10.1155/2013/989307
5. Onouchi Y, Ozaki K, Burns JC, et al. A genome-wide association study identifies three new risk loci for Kawasaki disease. *Nat Genet*. 2012;44(5):517–521. doi:10.1038/ng.2220
6. Tsai FJ, Lee YC, Chang JS, et al. Identification of novel susceptibility loci for Kawasaki disease in a Han Chinese population by a genome-wide association study. *PLoS One*. 2011;6(2):e16853. doi:10.1371/journal.pone.0016853
7. Khor CC, Davila S, Breunis WB, et al. Genome-wide association study identifies FCGR2A as a susceptibility locus for Kawasaki disease. *Nat Genet*. 2011;43(12):1241–1246. doi:10.1038/ng.981
8. Westra HJ, Franke L. From genome to function by studying eQTLs. *Biochim Biophys Acta*. 2014;1842(10):1896–1902. doi:10.1016/j.bbdis.2014.04.024
9. McCrindle BW, Rowley AH, Newburger JW, et al. Diagnosis, treatment, and long-term management of Kawasaki disease: a scientific statement for health professionals from the American heart association. *Circulation*. 2017;135(17):e927–e999. doi:10.1161/CIR.0000000000000484
10. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*. 2009;25(14):1754–1760. doi:10.1093/bioinformatics/btp324
11. McKenna A, Hanna M, Banks E, et al. The genome analysis toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res*. 2010;20(9):1297–1303. doi:10.1101/gr.107524.110
12. Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res*. 2010;38(16):e164. doi:10.1093/nar/gkq603
13. Pertea M, Kim D, Pertea GM, Leek JT, Salzberg SL. Transcript-level expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. *Nat Protoc*. 2016;11(9):1650–1667. doi:10.1038/nprot.2016.095
14. Chakraborty S, Panda AK, Bose S, et al. Transcriptional regulation of FOXP3 requires integrated activation of both promoter and CNS regions in tumor-induced CD8(+) Treg cells. *Sci Rep*. 2017;7(1):1628. doi:10.1038/s41598-017-01788-z
15. Blackshaw S, Sawa A, Sharp AH, Ross CA, Snyder SH, Khan AA. Type 3 inositol 1,4,5-trisphosphate receptor modulates cell death. *FASEB J*. 2000;14(10):1375–1379. doi:10.1096/fj.14.10.1375
16. Metz PJ, Arsenio J, Kakaradov B, et al. Regulation of asymmetric division and CD8+ T lymphocyte fate specification by protein kinase C ζ and protein kinase C λ /t. *J Immunol*. 2015;194(5):2249–2259. doi:10.4049/jimmunol.1401652
17. Rae W, Ward D, Mattocks CJ, et al. Autoimmunity/inflammation in a monogenic primary immunodeficiency cohort. *Clin Transl Immunol*. 2017;6(9):e155. doi:10.1038/cti.2017.38
18. Yoon KL. Update of genetic susceptibility in patients with Kawasaki disease. *Korean J Pediatr*. 2015;58(3):84–88. doi:10.3345/kjp.2015.58.3.84
19. Kessel A, Ammuri H, Peri R, et al. Intravenous immunoglobulin therapy affects T regulatory cells by increasing their suppressive function. *J Immunol*. 2007;179(8):5571–5575. doi:10.4049/jimmunol.179.8.5571

20. Depaolo RW, Tang F, Kim I, et al. Toll-like receptor 6 drives differentiation of tolerogenic dendritic cells and contributes to LcrV-mediated plague pathogenesis. *Cell Host Microbe*. 2008;4(4):350–361. doi:10.1016/j.chom.2008.09.004
21. Maitra U, Davis S, Reilly CM, Li L. Differential regulation of Foxp3 and IL-17 expression in CD4 T helper cells by IRAK-1. *J Immunol*. 2009;182(9):5763–5769. doi:10.4049/jimmunol.0900124
22. Veldhoen M, Hocking RJ, Atkins CJ, Locksley RM, Stockinger B. TGFbeta in the context of an inflammatory cytokine milieu supports de novo differentiation of IL-17-producing T cells. *Immunity*. 2006;24(2):179–189. doi:10.1016/j.immuni.2006.01.001
23. Li Y, Zheng Q, Zou L, et al. Kawasaki disease shock syndrome: clinical characteristics and possible use of IL-6, IL-10 and IFN-γ as biomarkers for early recognition. *Pediatr Rheumatol Online J*. 2019;17(1):1. doi:10.1186/s12969-018-0303-4
24. Yamaji N, Da Silva Lopes K, Shoda T, et al. TNF-α blockers for the treatment of Kawasaki disease in children. *Cochrane Database Syst Rev*. 2019;8(8):Cd012448. doi:10.1002/14651858.CD012448.pub2
25. Rudra D, Egawa T, Chong MM, Treuting P, Littman DR, Rudensky AY. Runx-CBFBeta complexes control expression of the transcription factor Foxp3 in regulatory T cells. *Nat Immunol*. 2009;10(11):1170–1177. doi:10.1038/ni.1795
26. Klunker S, Chong MM, Mantel PY, et al. Transcription factors RUNX1 and RUNX3 in the induction and suppressive function of Foxp3+ inducible regulatory T cells. *J Exp Med*. 2009;206(12):2701–2715. doi:10.1084/jem.20090596
27. Cao XG, Kou CZ, Zhao YP, et al. Overexpression of LYRM1 induces mitochondrial impairment in 3T3-L1 adipocytes. *Mol Genet Metab*. 2010;101(4):395–399. doi:10.1016/j.ymgme.2010.08.010
28. Kim EH, Suresh M. Role of PI3K/Akt signaling in memory CD8 T cell differentiation. *Front Immunol*. 2013;4:20. doi:10.3389/fimmu.2013.00020
29. Wang D, Chen J, Chen H, et al. Leptin regulates proliferation and apoptosis of colorectal carcinoma through PI3K/Akt/mTOR signalling pathway. *J Biosci*. 2012;37(1):91–101. doi:10.1007/s12038-011-9172-4
30. Huang YC, Lin YJ, Chang JS, et al. Single nucleotide polymorphism rs2229634 in the ITPR3 gene is associated with the risk of developing coronary artery aneurysm in children with Kawasaki disease. *Int J Immunogenet*. 2010;37(6):439–443. doi:10.1111/j.1744-313X.2010.00943.x
31. Oishi T, Iida A, Otsubo S, et al. A functional SNP in the NKX2.5-binding site of ITPR3 promoter is associated with susceptibility to systemic lupus erythematosus in Japanese population. *J Hum Genet*. 2008;53(2):151–162. doi:10.1007/s10038-007-0233-3
32. Harb H, Irvine J, Amarasekera M, et al. The role of PKCζ in cord blood T-cell maturation towards Th1 cytokine profile and its epigenetic regulation by fish oil. *Biosci Rep*. 2017;37(2). doi:10.1042/BSR20160485
33. Ferdosian F, Dastgheib SA, Morovati-Sharifabad M, et al. Cumulative evidence for association between IL-10 polymorphisms and Kawasaki disease susceptibility: a systematic review and meta-analysis. *Fetal Pediatr Pathol*. 2021;40(2):153–165. doi:10.1080/15513815.2019.1686789
34. Hsu P, Santner-Nanan B, Hu M, et al. IL-10 potentiates differentiation of human induced regulatory T cells via STAT3 and Foxo1. *J Immunol*. 2015;195(8):3665–3674. doi:10.4049/jimmunol.1402898
35. Brusko TM, Wasserfall CH, Hulme MA, Cabrera R, Schatz D, Atkinson MA. Influence of membrane CD25 stability on T lymphocyte activity: implications for immunoregulation. *PLoS One*. 2009;4(11):e7980. doi:10.1371/journal.pone.0007980
36. Gupta R, Yadav A, Misra R, Aggarwal A. Urinary sCD25 as a biomarker of lupus nephritis disease activity. *Lupus*. 2015;24(3):273–279. doi:10.1177/0961203314555174
37. Meng X, Yang J, Dong M, et al. Regulatory T cells in cardiovascular diseases. *Nat Rev Cardiol*. 2016;13(3):167–179. doi:10.1038/nrcardio.2015.169

Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-general-medicine-journal>