Gene Expression Network Analysis Identifies Potential Targets for Prevention of Preeclampsia

Yu Xia¹⁻³, Yu-Dong Zhao⁴, Gui-Xiang Sun^{1,2}, Shuai-Shuai Xia¹, Zheng-Wang Yang³

¹Provincial Key Laboratory of TCM Diagnostics, Hunan University of Chinese Medicine, Changsha, Hunan Province, 410208, People's Republic of China; ²Institute of Chinese Medicine Diagnosis, Hunan University of Chinese Medicine, Changsha, Hunan Province, 410208, People's Republic of China; ³Department of Obstetrics and Gynecology, The First Hospital of Hunan University of Chinese Medicine, Changsha, Hunan Province, 410007, People's Republic of China; ⁴School of Clinical Medicine, Chengdu University of Traditional Chinese Medicine, Chengdu, 610075, People's Republic of China

Correspondence: Gui-Xiang Sun, Provincial Key Laboratory of TCM Diagnostics, Hunan University of Chinese Medicine, No. 300, Xueshi Road, Yuelu District, Changsha, Hunan Province, 410208, People's Republic of China, Tel +86-13787272837, Email 84663423@qq.com

Objective: Preeclampsia (PE) is a pregnancy-specific multisystem disease as well as an important cause of maternal and perinatal death. This study aimed to analyze the placental transcriptional data and clinical information of PE patients available in the published database and predict the target genes for prevention of PE.

Methods: The clinical information and corresponding RNA data of PE patients were downloaded from the GEO database. Cluster analysis was performed to examine the correlation between different genotyping genes and clinical manifestations. Then, bioinformatic approaches including GO, KEGG, WGCNA, and GSEA were employed to functionally characterize candidate target genes involved in pathogenesis of PE.

Results: Two PE datasets GSE60438 and GSE75010 were obtained and combined, thereby providing the data of 205 samples in total (100 non-PE and 105 PE samples). After eliminating the batch effect, we grouped and analyzed the integrated data, and further performed GSEA analysis. It was found that the genes in group 1 and group 2 were different from those in normal samples. Moreover, WGCNA analysis revealed that genes in group 1 were up-regulated in turquoise module, including *SASH1*, *PIK3CB* and *FLT-1*, while genes in group 2 were up-regulated in the blue and brown modules. We further conducted GO and KEGG pathway enrichment analyses and found that the differential genes in turquoise module were mainly involved in biological processes such as small molecular catabolic process, while being highly enriched in pathways, including MAPK signaling pathway and Rap1 signaling pathway.

Conclusion: FLT-1 was conventionally used to predict PE risk, and sFLT-1 could also be used as an indicator to evaluate PE treatment effect. As a candidate biomarker for predicting PE, SASH1 may participate in proliferation, migration, invasion and epithelial mesenchymal transformation of human trophoblast cells by regulating MAPK pathway and Rap1 signaling pathway, thus affecting the progression of PE. The mechanism allowing PIK3CB to regulate PE development was not clear, while the gene could be another candidate biomarker for PE risk prediction. This is an exploratory study and our findings were still required verification in further studies.

Keywords: preeclampsia, SASH1, PIK3CB, FLT-1, MAPK signaling pathway, Rap1 signaling pathway

Introduction

Preeclampsia (PE) is a common hypertensive disorder complicating pregnancy, with an incidence rate of approximately 2–5% worldwide.^{1,2} While 14% of all maternal deaths are caused by PE, it becomes the main reason for the increasing maternal and perinatal mortality.³ PE leads to maternal heart impairment. Compared with pregnant women with normal blood pressure, those with a history of gestational hypertension have a significantly increased risk of kidney and cardiovascular diseases and even impaired cognitive function in the later stage.^{4,5} In addition, PE may aggravate fetal intrauterine hypoxia, resulting in premature delivery,⁶ fetal growth retardation, or defective fetal heart development leading to fetal cardiac dysfunction.⁷ The etiology of PE is multifactorial and polygenic, which remains elusive despite

intensive research. According to the clinical features, several subtypes of PE can be sorted. In general, the development of PE can be divided into two stages: asymptomatic and symptomatic stages.⁸ Abnormal placental development in early pregnancy causes placental dysfunction and later leads to the release of factors into maternal blood, resulting in hypertension and organ impairment. The occurrence of PE is closely related to placental dysfunction. The researchers conducted in-depth research into the placenta as the root cause of PE, and as the research progressed, a more refined classification of PE stages was made.⁹

However, the molecular mechanisms underlying poor placentation remain largely unknown, although increasing research studies have applied RNA sequencing and other analytical approaches to explore placenta development. Some studies are sorting out the relationship between PE subtypes and specific genes.^{10,11} By analyzing the RNA profiles of 302 human placenta samples, Gong et al¹² demonstrated that elevated serum levels of follistatin-like 3 (FSTL3) in pregnant women were predictive of subsequent PE and fetal growth restriction (FGR). Moreover, reported by recent studies, plasma cell-free RNA (cfRNA) could exhibit specific patterns to indicate normal pregnancy progression and serve as the biomarker to detect the risk of PE months before clinical manifestations. While the blood sample collection only requires venipuncture for once, cfRNA signatures can track pregnancy progression at the placental level, on the maternal or fetal sides, and can effectively predict the occurrence of PE, with a sensitivity of 75% and a positive predictive value of 32.3%.¹³ However, a comprehensive analysis on the clinical data and genomic profiling in PE is still lacking. To protect the maternal and fetal health against PE, early detection of PE and timely treatment for pregnant women with PE are crucial, thereby avoiding adverse pregnancy outcomes. Early detection and timely treatment of pregnant women with PE are crucial for avoiding adverse pregnancy outcomes. Accordingly, it is necessary to identify early diagnostic markers and develop effective treatment methods. In this study, based on comparing the transcriptional profile of placental samples and the clinical information, the purpose was to evaluate the relationship between specific genotypes and clinical features of PE, which might contribute to the identification of candidate biomarkers for the prediction and prevention of PE.

Data and Methods

Data Download and Processing

Clinical information and transcriptional data were downloaded from GEO database (<https://www.ncbi.nlm.nih.gov/gds>). Inclusion criteria were as follows: 1) Pregnancy-induced hypertension, including gestational hypertension, preeclampsia, eclampsia, chronic hypertension with preeclampsia, and pregnancy complicated with chronic hypertension; 2) Samples with complete clinical information and human transcriptional data. Perl (V 5.18.4) was used to convert the probe name into gene name in downloaded data. R (R 4.1.1) language in combination with “limma” and “SVA” packages were employed to merge the downloaded data, take the intersection of the sorted genes, compute the mean value of the same gene, take log2 for the data with large gene expression, and carry out batch correction to eliminate the impact of batch effect. R language “ggplot2” package for PCA analysis was run to verify the effect of batch correction.

Classification, Correlation Analysis of Continuous Variables, and Classification Difference Analysis

The integrated data were classified by using “consumusclusterplus” package. The clinical data were sorted, and the correlation between age and gestational week of delivery was analyzed. According to the typing results, taking the mean difference >0.2 and $P < 0.05$ as the screening criteria, R language “limma” package was used to take the genes of different typing samples and were compared with those of normal samples, and potential biomarkers in different groups were screened.

GSEA and WGCNA

The differential genes were obtained, enriched and analyzed by gene set enrichment analysis (GSEA) (V4.1.0).¹⁴ Thereafter, the differences in specific differential genes between each typing and normal samples were compared. In

the meantime, R “WGCNA” package was used to calculate the threshold (optimal weight) of the co-expressive gene correlation network. Then, gene function modules were identified by dynamic segmentation algorithm (cutree dynamic), and genes with a similar expression pattern were assigned into one module. Finally, the dissimilarity between modules was calculated based on the module feature vector. The similar modules were combined, and the correlation between WGCNA gene function module and clinical features of samples was analyzed.

GO and KEGG Pathway Enrichment Analyses

GO and KEGG pathway enrichment analyses were performed by using R language software, Bioconductor plug-in, and clusterprofiler. Data were processed, and genes without ID were deleted. $P < 0.05$ was set as the standard for screening. The enrichment of different modules and genes was visually displayed, and R was run to obtain the bubble diagram of GO and KEGG enrichment analysis.

Results

Data Inclusion and Processing

According to the inclusion criteria, two PE chip datasets GSE60438 and GSE75010 were obtained from the GEO database and included in this study. The basic information of the two gene matrices were listed in Table 1, and the clinical data were listed in Tables 2 and Tables 3.

PCA Verification and Sample Typing

We conducted PCA analysis on the corrected data. As depicted in Figure 1, samples from GSE60438 and GSE75010 prior to correction were distributed on both sides of the figure (A), while those after correction were randomly distributed (B). This observation indicated that the influence of batch effect has been eliminated. We, therefore, typed the data by using R language. According to the consistent scoring results, the genes were divided into two groups. The analysis identified a weak correlation between different genotypes (C).

Table 1 Gene Expression Dataset of PE in GEO Database

GEO ID	Platform	Non-PE	PE
GSE60438	GPL10558	23	25
GSE75010	GPL6244	77	80

Table 2 Clinical Data of the GSE60438 Dataset

Grouping	Quantity	Tissue Type	Maternal Age (yrs)	Gestational Age (wks)	Infant Sex		Infant Weight (g)
					M	F	
PE	25	Decidua basalis	30.16*	32*	12	13	1662.32*
Non-PE	23	Decidua basalis	31.39*	39.04*	12	11	3290.57*

Note: *Represents the average.

Table 3 Clinical Data of the GSE75010 Dataset

Grouping	Quantity	Tissue Type	Maternal age (yrs)	Maternal BMI (kg/m ²)	Maternal Blood Type					Mean Umbilical PI	Max Systolic BP (mmHg)	Max Diastolic BP (mmHg)	Mode Protein-uria
					A	B	O	AB	NA				
PE	80	Placenta	33.19*	26.50*	30	12	33	4	1	1.39*	169.79*	107.28*	2.59*
Non-PE	77	Placenta	33.18*	24.49*	18	25	30	4	0	1.30*	136.12*	85.27*	0.47*

Note: *Represents the average.

Figure 1 PCA validation and sample typing. **(A)** PCA diagram before correction; **(B)** PCA diagram after correction; **(C)** The cluster of PE subtypes.

Correlation Analysis of Continuous Variables

We next employed R language to analyze the correlation between the age of pregnant women, gestational week of delivery and the classification. As shown in Figure 2, there were differences in the age of pregnant women and gestational week of delivery among different groups. According to the grouping results, we organize and analyze the genes of different groups. In group 1, the following genes had smaller p-values: *SASH1*, *PIK3CB*, *FLT-1*, *FLNB*, *FAM120A*, *ENG*, *MICAL3*, *SH3PXD2A*, *MYO7B*. While in group 2 is *SMARCA1*, *TSC22D1*, *NTRK2*, *HSD17B1*, *GSN*, *EPHB6*, *APLN*, *SULF2*, *CETN2*.

GSEA and WGCNA

We performed GSEA to examine specific differential genes in each group. As illustrated in Figure 3, the differential genes were different between genotyping samples and normal samples (A, B). We then constructed a gene co-expression module network by WGCNA, based on the differential genes. A total of 1356 differential genes in 205 samples were included in this study. TOM cluster analysis revealed an association of the gene module with the known phenotype, and the correlation coefficient between the feature vector of the module and the phenotype was calculated. Notably, we found that ME turquoise module was positively correlated with maternal age, maximum systolic blood pressure, maximum diastolic blood pressure, and the number of urinary proteins (C). Moreover, a heatmap of differential genes in the module

Figure 2 Correlation Analysis between maternal age, delivery gestational week, and classification. (A) Correlation analysis between maternal age and classification; (B) Correlation between gestational age and classification.

Note: * $P < 0.05$.

showed that while the group 1 genes were up-regulated in turquoise module, the group 2 genes were up-regulated in blue and brown modules (D). These data indicated that genes in turquoise module and group 1 were closely related to PE. Among those genes, according to the low p-values, three candidate genes were selected for subsequent analysis: *SASH1*, *PIK3CB* and *FLT-1*.

GO and KEGG Pathway Enrichment Analyses

We further performed GO and KEGG pathway enrichment analyses by using R language (Figure 4). GO enrichment analysis revealed that differential genes in the blue module and brown module were mainly involved in biological processes such as T cell activation and those including extracellular matrix organization, respectively, while the genes in the turquoise module are mainly associated with biological processes such as small molecular catalytic process (A). Meanwhile, KEGG pathway enrichment analysis showed that while differential genes in the blue module and brown module were highly enriched in pathways such as those involving cell adhesion molecules and pathways including those involving apoptosis, respectively, pathways of differential genes in turquoise module mainly included MAPK signaling pathway, Rap1 signaling pathway, and pathways involving valine, leucine and isoleucine degradation (B).

Discussion

It has been shown that under the condition of PE, great changes such as oxidative stress and inflammatory response occur in the intrauterine environment of pregnant women, leading to alterations in various signaling pathways. PE has a great impact on growth and development of the fetus, impairs the structure and function of fetal cardiovascular system or kidney, and causes hypertension.¹⁵ Current studies suggest that the pathogenesis of PE may be related to decidual macrophage dysfunction,¹⁶ placental angiogenesis disorder,¹⁷ placental aging,^{18,19} protein aggregation, and defective placental aggregation phagocytosis.²⁰ At the molecular level, the microRNA spectrum of blood exosomes has been used to evaluate the pathophysiology of preeclampsia and predict the disease,²¹ and there are more and more studies at gene level.^{22–25}

Zhang and Horvath²⁶ conducted the first WGCNA analysis in 2005 and then published WGCNA R software package in R based on theory of the research method in 2008. As an efficient and accurate bioinformatics approach, WGCNA has been widely used for the gene network analysis.²⁷

Herein, we intensively analyzed the published sequencing data in selected two datasets, including 205 samples (100 non-PE and 105 PE cases) with complete clinical phenotype information and transcriptome sequencing data. In the study, we constructed the co-expression network modules and performed clinical phenotype correlation analysis on the mRNAs

Figure 3 GSEA and WGCNA. **(A)** GSEA analysis on specific differential genes between group I and normal samples; **(B)** GSEA analysis on specific differential genes between group 2 and normal samples; **(C)** Module-trait relationships. **(D)** A heatmap of module genes.

obtained by placental tissue sequencing. The analysis revealed that MEturquoise module was positively correlated with maternal age, maximum systolic blood pressure, maximum diastolic blood pressure, and the number of urinary proteins. Further studies showed that the patients with group 1 are consistent with this law. The present study provided evidence that the up-regulated differential genes *SASH1*, *PIK3CB* and *FLT-1* may affect the progression of PE through MAPK signaling pathway and Rap1 signaling pathway.

Figure 4 Bubble Diagram of GO and KEGG enrichment analysis. (A) GO enrichment analysis bubble chart; (B) KEGG pathway enrichment analysis bubble chart.

Serebrov et al²⁸ reported that rs34845949 of the *SASH1* gene was significantly correlated with PE. The expression levels of *SASH1* in placental tissues were increased in patients with PE compared with healthy pregnancy,^{10,28,29} suggesting that *SASH1* gene might play a role in placental dysplasia in patients with preeclampsia. Furthermore, reported by published studies, the overexpression of *SASH1* could inhibit the proliferation of HTR-8/SVneo cells, possibly due to the inhibition on *CCND1* and *CCND3*, which arrested the cells in G1 phase. And the expression levels of *CCND1* and *CCND3* were associated with MAPK signaling.^{29,30} Ras/Raf/MEK/ERK pathway in MAPK can be activated by a variety of stimuli, and the activated Ras/Raf/MEK/ERK pathway regulate numerous physiological processes such as cell growth, development, division, and apoptosis. It has been demonstrated that *SASH1* interacts with MAP2K2 and MAP4K4. While MAP4K4 indirectly activates MAP2K2 via MEKK1, it may affect receptor signaling through Rac1-MEKK1-MAP4K4-JNK pathway and play an important role in cell migration and invasion.³¹ Human placental pluripotent mesenchymal stromal cells regulate trophoblast migration via Rap1 activation.³² Previous studies have shown that inhibition of MAPK signaling pathway suppresses trophoblast proliferation and migration, probably promoting the progression of PE.^{33,34} Based on the above observations, we speculate that *SASH1* may be involved in the progression of PE by affecting the proliferation and migration of trophoblasts via MAPK signaling pathway and Rap1 signaling pathway. At present, the molecular mechanism enabling *SASH1* to participated in the development of PE is still unclear. It is necessary to further study the specific role of *SASH1* in PE, especially by analyzing the differential protein expression levels. Based on the relationship between the differential expression patterns of *SASH1* and the specific clinical features in PE, the clinical significances of *SASH1* can be evaluated, including the potential as a biomarker for early diagnosis and as the target for novel PE therapy.

While PIK3 protein family plays a role in the regulation of various cell functions including cell growth, proliferation and survival, PIK3 is involved in controlling the migration and penetration of human cancer cells and trophoblasts.³⁵ Among the protein family, *PIK3CBBCL-2* is considered the core regulatory gene of decidua in eclampsia. Aberrant expression of *PIK3CBBCL-2* causes abnormal decidual apoptosis related to the pathogenesis of eclampsia.³⁶ Studies have confirmed that the expression of *PIK3CA* gene is up-regulated in PE patients as compared to the normal group ($P < 0.05$).³⁷

Among all potential target genes identified in the present study, *FLT-1* gene has been widely used in the animal modeling of PE and clinical trials. In this case, sFLT-1 overexpression is commonly used for generating the animal model of PE, whereas inhibiting FLT-1 can prevent PE.^{38,39} It has been shown that sFLT-1/PIGF ratio is valuable in predicting the occurrence of PE.^{40,41} Particularly, determination of sFLT-1/PIGF ratio becomes a valuable auxiliary means for PE diagnosis in the medical insurance system of France; a low ratio is regarded as a highly negative predictive value of PE, and unnecessary hospitalization and premature delivery can be avoided.⁴² Numerous studies analyzed the different critical values of the sFLT-1/PIGF ratio, thereby suggesting the ratio as an effective predictor for PE and PE-related complications.⁴³⁻⁴⁷ Moreover, some preceding studies suggested that the pathogenesis of PE

might be related to inhibition of MAPK pathway, thereby affecting the invasion of trophoblast cells and promoting the secretion of sFLT-1/sEng.⁴⁸ FLT-1 was found to be closely correlated with the occurrence and development of PE, and FLT-1 was significantly elevated in PE patients.⁴⁹ At present, the effective treatment of preeclampsia mainly relied on the termination of pregnancy. In recent years, some researchers explored the mechanism to reduce the level of sFLT-1 in pregnant women with preeclampsia, so as to restore the balance of angiogenesis. According to previous studies, blood sFLT-1 apheresis, a proton pump inhibitor, might interfere with placental renin-AGT uptake/transport, thereby reducing angiotensin-induced sFLT-1 synthesis;⁵⁰ aspirin nourishes HTR-8/SVneo, by inactivating NF- κ B signaling in lamellar cells, attenuated the role of sFLT-1 in oxidative stress and endothelial dysfunction, and reduced trophoblast apoptosis,⁵¹ suggesting that reducing sFLT-1 in PE patients could be applicable and effective. Concentration-related therapeutic approaches is expected to be important applications in the future clinical treatment for the prevention of preeclampsia.

In conclusion, WGCNA and the correlation analysis on published placental mRNA data and clinical information of PE patients revealed that genes in METurquoise module of group 1 patients were positively correlated with maternal age, maximum systolic blood pressure, maximum diastolic blood pressure, and the number of urinary proteins. Further studies showed that genes *SASH1*, *PIK3CB* and *FLT-1* were closely related to PE. Moreover, we found that MAPK signaling pathway and Rap1 signaling pathway were highly enriched in KEGG. FLT-1 was conventionally used to predict PE risk, and sFLT-1 could also be used as indicator to evaluate PE treatment effect. As a candidate biomarker for predicting PE, SASH1 could participate in the proliferation, migration, invasion and epithelial mesenchymal transformation of human trophoblast cells, by regulating MAPK pathway and Rap1 signaling pathway, thus affecting the progression of PE. The mechanism allowing PIK3CB to regulate PE development was not clear, although the gene could be another candidate biomarker for PE prediction. This is an exploratory study, and our findings still required verification by further studies.

Furthermore, this study was subject to limitations in the design. Firstly, due to the limitations of the GEO database, the current WGCNA analysis showed that group 1 had a higher correlation with clinical data in genetic data. Therefore, this study only discussed the objects related to group 1, while the genes of group 2 could be further studied in the future; secondly, this study was based on the previous research, which also required further investigation at the molecular level, so the mechanism underlying the regulatory roles of SASH1 and PIK3CB in PE could be further proved in the future.

Ethics Approval

GEO belongs to public database. The patients involved in the database have obtained ethical approval. Our study is based on open-source data, so there are no ethical issues and other conflicts of interest. After being reviewed by the Ethics Committee of First Hospital of Hunan University of Chinese Medicine, it can be exempted from ethical approval.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

This work was supported by the Special funds for traditional Chinese medicine subsidized by the central government of the State Administration of traditional Chinese Medicine (No. [2015]78).

Disclosure

The authors report no conflicts of interest in this work.

References

- Di Renzo GC, Fonseca E, Gratacos E; FIGO Working Group on Good Clinical Practice in Maternal-Fetal Medicine. Good clinical practice advice: first trimester screening and prevention of pre-eclampsia in singleton pregnancy. *Int J Gynaecol Obstet*. 2019;144(3):325–329. doi:10.1002/ijgo.12741
- Umesawa M, Kobashi G. Epidemiology of hypertensive disorders in pregnancy: prevalence, risk factors, predictors and prognosis. *Hypertens Res*. 2017;40(3):213–220. doi:10.1038/hr.2016.126
- World Health Organization. WHO recommendations: policy of interventionist versus expectant management of severe pre-eclampsia before term. Geneva; 2018.
- Garovic VD, White WM, Vaughan L, et al. Incidence and long-term outcomes of hypertensive disorders of pregnancy. *J Am Coll Cardiol*. 2020;75(18):2323–2334. doi:10.1016/j.jacc.2020.03.028
- Adank MC, Hussainali RF, Oosterveer LC, et al. Hypertensive disorders of pregnancy and cognitive impairment: a prospective cohort study. *Neurology*. 2021;96(5):e709–e18. doi:10.1212/WNL.00000000000011363
- Davies EL, Bell JS, Bhattacharya S. Preeclampsia and preterm delivery: a population-based case-control study. *Hypertens Pregnancy*. 2016;35(4):510–519. doi:10.1080/10641955.2016.1190846
- Thornburg KL, Drake R, Valent AM. Maternal hypertension affects heart growth in offspring. *J Am Heart Assoc*. 2020;9(9):e016538. doi:10.1161/JAHA.120.016538
- Roberts JM, Hubel CA. The two stage model of preeclampsia: variations on the theme. *Placenta*. 2009;30 Suppl A:S32–7. doi:10.1016/j.placenta.2008.11.009
- Redman C. The six stages of pre-eclampsia. *Pregnancy Hypertens*. 2014;4(3):246. doi:10.1016/j.preghy.2014.04.020
- Kaartokallio T, Cervera A, Kyllonen A, et al. Gene expression profiling of pre-eclamptic placentae by RNA sequencing. *Sci Rep*. 2015;5:14107. doi:10.1038/srep14107
- Ren Z, Gao Y, Gao Y, et al. Distinct placental molecular processes associated with early-onset and late-onset preeclampsia. *Theranostics*. 2021;11(10):5028–5044. doi:10.7150/thno.56141
- Gong S, Gaccioli F, Dopierala J, et al. The RNA landscape of the human placenta in health and disease. *Nat Commun*. 2021;12(1):2639. doi:10.1038/s41467-021-22695-y
- Rasmussen M, Reddy M, Nolan R, et al. RNA profiles reveal signatures of future health and disease in pregnancy. *Nature*. 2022;601(7893):422–427. doi:10.1038/s41586-021-04249-w
- Xu H, Xie Y, Sun Y, et al. Integrated analysis of multiple microarray studies to identify potential pathogenic gene modules in preeclampsia. *Exp Mol Pathol*. 2021;120:104631. doi:10.1016/j.yexmp.2021.104631
- Tenorio MB, Ferreira RC, Moura FA, et al. Cross-talk between oxidative stress and inflammation in preeclampsia. *Oxid Med Cell Longev*. 2019;2019:8238727. doi:10.1155/2019/8238727
- Rong M, Yan X, Zhang H, et al. Dysfunction of decidual macrophages is a potential risk factor in the occurrence of preeclampsia. *Front Immunol*. 2021;12:655655. doi:10.3389/fimmu.2021.655655
- Gebara N, Correia Y, Wang K, et al. Angiogenic properties of placenta-derived extracellular vesicles in normal pregnancy and in preeclampsia. *Int J Mol Sci*. 2021;22(10):5402. doi:10.3390/ijms22105402
- Scaife PJ, Simpson A, Kurlak LO, et al. Increased placental cell senescence and oxidative stress in women with pre-eclampsia and normotensive post-term pregnancies. *Int J Mol Sci*. 2021;22(14):7295. doi:10.3390/ijms22147295
- Hu X, Zhang L. Uteroplacental circulation in normal pregnancy and preeclampsia: functional adaptation and maladaptation. *Int J Mol Sci*. 2021;22(16):8622. doi:10.3390/ijms22168622
- Nakashima A, Shima T, Tsuda S, et al. Aggrephagy deficiency in the placenta: a new pathogenesis of preeclampsia. *Int J Mol Sci*. 2021;22(5):2432. doi:10.3390/ijms22052432
- Matsubara K, Matsubara Y, Uchikura Y, et al. Pathophysiology of preeclampsia: the role of exosomes. *Int J Mol Sci*. 2021;22(5):2572. doi:10.3390/ijms22052572
- Qin L, Yang Q, Fei Z, et al. Expression of lncRNA TINCR in the placenta of patients with pre-eclampsia and its effect on the biological behaviours of trophoblasts. *Zygote*. 2021;1–9. doi:10.1017/S0967199421000290
- Li XL, Zhang L, Hou B, et al. Expression of lncRNA MIR210HG in preeclampsia placental tissue and its functional analysis. *Zhonghua Fu Chan Ke Za Zhi*. 2021;56(6):425–433. doi:10.3760/cma.j.cn112141-20210118-00029
- Lekva T, Roland MCP, Estensen ME, et al. Dysregulated non-coding telomerase RNA component and associated exonuclease XRN1 in leucocytes from women developing preeclampsia-possible link to enhanced senescence. *Sci Rep*. 2021;11(1):19735. doi:10.1038/s41598-021-99140-z
- He J, Chen M, Xu J, et al. Identification and characterization of Piwi-interacting RNAs in human placentas of preeclampsia. *Sci Rep*. 2021;11(1):15766. doi:10.1038/s41598-021-95307-w
- Zhang B, Horvath S. A general framework for weighted gene co-expression network analysis. *Stat Appl Genet Mol Biol*. 2005;4:Article17. doi:10.2202/1544-6115.1128
- Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinform*. 2008;9:559. doi:10.1186/1471-2105-9-559
- Serebrova VN, Trifonova EA, Gabidulina TV, et al. [Detection of novel genetic markers of susceptibility to preeclampsia based on an analysis of the regulatory genes in the placental tissue]. *Mol Biol (Mosk)*. 2016;50(5):870–879. Russian. doi:10.1134/S0026893316050162
- Liu S, Jiang S, Huang L, et al. Expression of SASH1 in preeclampsia and its effects on human trophoblast. *Biomed Res Int*. 2020;2020:5058260. doi:10.1155/2020/5058260
- Radulovich N, Pham NA, Strumpf D, et al. Differential roles of cyclin D1 and D3 in pancreatic ductal adenocarcinoma. *Mol Cancer*. 2010;9:24. doi:10.1186/1476-4598-9-24
- Wang J, He Y, Zhang J-W, et al. SASH1 may interact with REK signaling pathways through MAP2K2 and MAP4K4. *Basic Clin Med*. 2014;34(11):1530–1536.
- Chen CP, Huang JP, Chu TY, et al. Human placental multipotent mesenchymal stromal cells modulate trophoblast migration via Rap1 activation. *Placenta*. 2013;34(10):913–923. doi:10.1016/j.placenta.2013.06.311

33. Zhou G, Li Z, Sun S, et al. TGF-beta1 alleviates HgCl2 induced apoptosis via P38 MAPK signaling pathway in human trophoblast cells. *Toxicol in Vitro*. 2019;61:104626. doi:10.1016/j.tiv.2019.104626
34. Li X, Wang Z, Liu G, et al. EIF3D promotes the progression of preeclampsia by inhibiting of MAPK/ERK1/2 pathway. *Reprod Toxicol*. 2021;105:166–174. doi:10.1016/j.reprotox.2021.09.006
35. Li MQ, Hou XF, Shao J, et al. The DSCs-expressed CD82 controls the invasiveness of trophoblast cells via integrinbeta1/MAPK/MAPK3/1 signaling pathway in human first-trimester pregnancy. *Biol Reprod*. 2010;82(5):968–979. doi:10.1095/biolreprod.109.080739
36. Tong J, Niu Y, Chen ZJ, et al. Comparison of the transcriptional profile in the decidua of early-onset and late-onset pre-eclampsia. *J Obstet Gynaecol Res*. 2020;46(7):1055–1066. doi:10.1111/jog.14257
37. Tsai K, Tullis B, Jensen T, et al. Differential expression of mTOR related molecules in the placenta from gestational diabetes mellitus (GDM), intrauterine growth restriction (IUGR) and preeclampsia patients. *Reprod Biol*. 2021;21(2):100503. doi:10.1016/j.repbio.2021.100503
38. Saad AF, Diken ZM, Kechichian TB, et al. Pravastatin effects on placental prosurvival molecular pathways in a mouse model of preeclampsia. *Reprod Sci*. 2016;23(11):1593–1599. doi:10.1177/1933719116648218
39. Saif J, Ahmad S, Rezai H, et al. Hydrogen sulfide releasing molecule MZe786 inhibits soluble Flt-1 and prevents preeclampsia in a refined RUPP mouse model. *Redox Biol*. 2021;38:101814. doi:10.1016/j.redox.2020.101814
40. Gao J, Shen J, Jiang Y, et al. Value of second trimester maternal serum sFlt-1, PlGF and their ratio in the prediction of preeclampsia. *Zhonghua Fu Chan Ke Za Zhi*. 2014;49(1):22–25.
41. Perales A, Delgado JL, De La Calle M, et al. sFlt-1/PlGF for prediction of early-onset pre-eclampsia: STEPS (Study of Early Pre-eclampsia in Spain). *Ultrasound Obstet Gynecol*. 2017;50(3):373–382. doi:10.1002/uog.17373
42. Mccarey C, Baert J, Mathey MP, et al. [Clinical value of angiogenic and anti-angiogenic marker assay in preeclampsia]. *Rev Med Suisse*. 2020;16(712):2031–2036. French
43. Zeisler H, Llurba E, Chantaine FJ, et al. Soluble fms-like tyrosine kinase-1 to placental growth factor ratio: ruling out pre-eclampsia for up to 4 weeks and value of retesting. *Ultrasound Obstet Gynecol*. 2019;53(3):367–375. doi:10.1002/uog.19178
44. Saleh L, Verdonk K, Jan Danser AH, et al. The sFlt-1/PlGF ratio associates with prolongation and adverse outcome of pregnancy in women with (suspected) preeclampsia: analysis of a high-risk cohort. *Eur J Obstet Gynecol Reprod Biol*. 2016;199:121–126. doi:10.1016/j.ejogrb.2016.02.013
45. Zeisler H, Llurba E, Chantaine F, et al. Predictive value of the sFlt-1: PlGFRatio in women with suspected preeclampsia. *N Engl J Med*. 2016;374(1):13–22. doi:10.1056/NEJMoa1414838
46. Verlohren S, Herraiz I, Lapaire O, et al. New gestational phase-specific cutoff values for the use of the soluble fms-like tyrosine kinase-1/placental growth factor ratio as a diagnostic test for preeclampsia. *Hypertension*. 2014;63(2):346–352. doi:10.1161/HYPERTENSIONAHA.113.01787
47. Rana S, Hacker MR, Modest AM, et al. Circulating angiogenic factors and risk of adverse maternal and perinatal outcomes in twin pregnancies with suspected preeclampsia. *Hypertension*. 2012;60(2):451–458. doi:10.1161/HYPERTENSIONAHA.112.195065
48. Liu X, Deng Q, Luo X, et al. Oxidative stress-induced Gadd45alpha inhibits trophoblast invasion and increases sFlt1/sEng secretions via p38 MAPK involving in the pathology of pre-eclampsia. *J Matern Fetal Neonatal Med*. 2016;29(23):3776–3785. doi:10.3109/14767058.2016.1144744
49. Shibuya M, Matsui H, Sasagawa T, et al. A simple detection method for the serum sFLT1 protein in preeclampsia. *Sci Rep*. 2021;11(1):20613. doi:10.1038/s41598-021-00152-6
50. Sun Y, Tan L, Neuman RI, et al. Megalin, proton pump inhibitors and the renin-angiotensin system in healthy and pre-eclamptic placentas. *Int J Mol Sci*. 2021;22(14):7407.
51. Zuo Q, Zou Y, Huang S, et al. Aspirin reduces sFlt-1-mediated apoptosis of trophoblast cells in preeclampsia. *Mol Hum Reprod*. 2021;27(1). doi:10.1093/molehr/gaaa089