Cancer Management and Research

ORIGINAL RESEARCH

LncRNA ASB16-AS1 Promotes Growth And Invasion Of Hepatocellular Carcinoma Through Regulating miR-1827/FZD4 Axis And Activating Wnt/β-Catenin Pathway

This article was published in the following Dove Press journal: Cancer Management and Research



Xiaoxiao Yao^{1,*} Guangqiang You^{1,*} Chen Zhou² Dan Zhang¹

¹Department of Hepatobiliary and Pancreatic Surgery, The Second Affiliated Hospital of Jilin University, Changchun 130041, People's Republic of China; ²Personnel Department, The First Affiliated Hospital of Jilin University, Changchun 130000, People's Republic of China

*These authors contributed equally to this work



Correspondence: Dan Zhang Department of Hepatobiliary and Pancreatic Surgery, The Second Affiliated Hospital of Jilin University, No. 218 Ziqiang Street, Nanguan District, Changchun 130041, People's Republic of China Email dan_zhang_1@163.com



Background: To date, although several long none ong RNAr ancRNAs) are reported to regulate hepatocellular carcinoma (HC) plevelopment, heir clationship still remains elusive. ASB16-AS1 is a poorly researced nov IncRNA. We aimed to investigate its function in HCC progression.

Methods: qRT-PCR and in situ hybridization (1.41) were used to analyze ASB16-AS1 expression in HCC tissues CCK8, Educatorporation and colony formation were used to determine cell proliferation Transwell as by was used to examine migration and invasion. Luciferase reporter assay was used to enalyze the interactions among ASB16-AS1, miR-1827 and FZD4.

Results: Bioinformatics may be identified ASB16-AS1 was overexpressed in HCC tissues, which a definite variated by qRT-PCR and in situ hybridization (ISH). Besides, ASB16-AS1 was defined as a potential indicator for HCC prognosis. Functional studies so wed AS 46 AS1 knockdown attenuated proliferation, migration and invasion of HCC cells of chanistically, ASB16-AS1 directly interacted with miR-1827 and promoted FZD4 expression by sponging miR-1827. Overexpressed FZD4 eventually activated Wnt/ β -catenin pathway and contributed to HCC progression.

Example Section First to identify ASB16-AS1 as an oncogene that enhances HC progression by modulating miR-1827/FZD4/Wnt/β-catenin pathways. **Keywords:** lncRNA, HCC, ASB16-AS1, miR-1827, FZD4

Introduction

Recent study has shown hepatocellular carcinoma (HCC) becomes one of the biggest problems for human health worldwide.¹ Every year, about 800,000 cases are diagnosed as HCC and it causes over 850,000 deaths per year.² As a major type of liver cancer, its pathogenesis still remains unclear.¹ Although several therapeutic methods developed, including surgery and radiotherapy, in past years, the outcomes of HCC are rather poor.³ The five-year survival rate in HCC patients is only about 30%.⁴ Hence, it is urgently required to investigate the in-depth mechanisms of HCC development.

Along with the advance of RNA-sequencing and bioinformatics methods, increasing long noncoding RNAs (lncRNAs) have been discovered in human transcriptome.⁵ LncRNA could be defined by over 200 nucleotides in length and

© 2019 Yao et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress.com/terms.php you hereby accept the and incorporate the Creative Commons Attribution – Non Commercial (unported, v3.0) License (http://treativecommons.org/licenses/by-no/3.0/). By accessing the work you hereby accept the firms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial uses of this work, lease see paragraphs 4.2 and 5 of our Terms (http://www.dovepress.com/terms.php). no potential for coding protein.⁶ Various studies show lncRNA is involved in many pathological processes, including tumorigenesis.⁷ Dysregulated lncRNA expression is associated with aberrant proliferation, metastasis or differentiation of tumor cells.⁸ For example, upregulation of lncRNA EPB41L4A-AS2 suppresses HCC progression via increasing FOXL1 expression.⁹ LncRNA MCM3AP-AS1 enhances HCC cell proliferation and metastasis through modulating miR-194-5p/FOXA1 signaling.¹⁰ LincRNA-p21 aberrant downregulation in squamous cell cancer leads to tumor aggravation via STAT3 pathway.¹¹ Obviously, lncRNAs play critical roles in cancer. And it is important to determine how lncRNA correlates with tumorigenesis.

ASB16-AS1 is a poorly characterized lncRNA. A recent study indicates that ASB16-AS1 promotes glioma development.¹² Its role in other cancers remains uncovered. Here, we discovered ASB16-AS1 was aberrantly upregulated in HCC using bioinformatics screening. We validated it and further revealed that ASB16-AS1 upregulation suggested a poor prognosis. Knockdown of ASB16-AS1 led to decreased proliferation and metastasis. Mechanistically, ASB16-AS1 was demonstrated to sponge miR-1827, leading to upregulation of FZD4 and activation of Wnt/β-catenin signaling. In conclusion, we uncovered novel mechanism of lncRNA ASB16-AS1/priP-1827/FZD4/Wnt/β-catenin signaling in HCC prognosion.

Materials And Method Human Samples

Fifty-one HCC samples and their adjacen normal controls were from The Second Affinated Hospital of Jup University. All tissues were store at liquit nitrogen. Our study was approved by the Ethic committee of The Second Affiliated Hospitar of Lilin Cliversity. Our study was conducted in accordance with the Declaration of Helsinki. Written inforced consent we obtained from patients.

Cell Culture

HCC cell lines and LO2 cells were obtained from the Type Culture Collection of the Chinese Academy of Sciences. Cell lines were cultured using DMEM (HyClone, Logan, UT, USA) containing 10% FBS (Hyclone) at 37° C in humidified incubator with 5% CO₂.

To obtain stable knockdown cell lines, shRNAs against ASB16-AS1 (5'-GGTTCTGAATCATTCAGTT-3') were

cloned into GV112 vector and transfected into cells using Lipofectamine 2000 (Invitrogen) at the concentration of 2 μ g per well, followed by selection using puromycin for 2 weeks.

qRT-PCR

RNA isolation along with qRT-PCR was completed using TRIzol reagent (Invitrogen) based on a previous work.¹³ GAPDH and U6 were the normalized controls, respectively. Relative expression was determined according to the $2^{-\Delta\Delta Ct}$ method.

CCK8 Assay

HCC cells were plated in a 96-well plate part cultured for described time. Then, CK8 strution was added and cell viability was determined by measuring absorbance at 450 nm.

Ethynyideoxyuridae (EdU) Analysis

Eduassay was completed following a previous work.¹⁴

Trans. I Assay

In cell assay was used to measure migration and invaon according to a previous work.¹⁴

Dual-Luciferase Reporter Assay

The binding site in ASB16-AS1 with miR-1827 was predicted using miRDB. And the binding site in FZD4 with miR-1827 was analyzed using TargetScan7.1. Then, the sequences of ASB16-AS1 and FZD4 3'-UTR containing the wild-type (WT) or mutant (MUT) binding site with miR-1827 were inserted into pGL3 vector (Promega). Then, reporters and miR-1827 mimic or control were transfected into Huh7 cells. Forty-eight hours later, the luciferase activity was measured by using the Dual-Luciferase Reporter Assay System (Promega Corporation).

Statistical Analysis

Statistical analysis was conducted by GraphPad Prism 6 software (GraphPad Software, Inc., La Jolla, CA, USA). Results were presented as mean \pm SD. Student's *t*-test or one-way ANOVA was used for statistical analysis. p<0.05 was considered as statistically significant.

Upregulated ASB16-AS1 Correlates With Poor Prognosis In HCC

To determine the correlation of lncRNAs with HCC, we performed bioinformatics analysis using Cancer RNA-seq Nexus tool (http://syslab4.nchu.edu.tw/). We identified that ASB16-AS1 was an upregulated lncRNA in HCC tissues (Figure 1A and B) according to this database. Another data in GEPIA database also validated this phenomenon (Figure 1C). Besides, qRT-PCR showed ASB16-AS1 was also upregulated in 51 HCC tissues (Figure 1D), which was further confirmed by in situ hybridization (Figure 1E). Moreover, ASB16-AS1 level was increased in advanced stages of tumor samples (Figure 1F). Then, HCC tissues were grouped into two subgroups based on ASB16-AS1 median expression value. Kaplan-Meier analysis indicated

ASB16-AS1 overexpression correlated with low survival rate (Figure 1G). Notably, the data in GEPIA database also showed ASB16-AS1 upregulation predicted low disease-free survival rate (Figure 1H). Thus, ASB16-AS1 is upregulated in HCC and may be a prognostic marker.

Effects Of ASB16-AS1 Knockdown On HCC Cells

To investigate the role of ASB16-AS1, we measured its expression in HCC cell lines. ASB16-AS1 was upregulated in tumor cell lines (Figure 2A). We then Jse Hun and Hep3B cells for further experiments. Though shRNA ASB16-AS1 expression was significatly declased in ese cell lines (Figure 2B). CCK8 2 ay showed A. NC S1 downregulation impaired the lifer on of Huh7 and Hep3B cells (Figure 2C). ation ar colony formation assays J incol



Figure I Upregulated ASB16-ASI correlates with poor prognosis in HCC. (A) Analysis of differentially expressed IncRNAs between HCC and control tissues using Cancer RNA-seq Nexus tool (http://syslab4.nchu.edu.tw/). (B) ASB16-ASI transcripts were indicated according to Cancer RNA-seq Nexus tool. (C) ASB16-ASI expression was analyzed using GEPIA tool (http://gepia.cancer-pku.cn/). (D) qRT-PCR analysis of ASB16-ASI expression in 51 pairs of HCC and adjacent normal tissues. (E) In situ hybridization (ISH) for ASB16-ASI expression in HCC and paired normal tissues. (F) Expression of ASB16-ASI in different stages of HCC tissues. (G) Overall survival rate was analyzed based on ASB16-ASI expression. (H) Disease-free survival rate was analyzed in HCC patients using GEPIA tool according to ASB16-ASI expression. *p<0.05.



Figure 2 Effects of ASB16-AS1 knockdown of ACC (A) Expression atterns of ASB16-AS1 in HCC cell lines. (B) shRNA-caused knockdown of ASB16-AS1 in Huh7 and Hep3B cells was validated by qRT-PCB, (C-E) Cell program and a sevaluated by CCK8, EdU incorporation, and colony formation assays. (F and G) Transwell assay for analysis of cell migration and invasion (C-E) Cell program and the sevaluated by CCK8, EdU incorporation, and colony formation assays. (F and G) Transwell assay for analysis of cell migration and invasion (C-E) Cell program and the sevaluated by CCK8, EdU incorporation, and colony formation assays. (F and G) Transwell assay for analysis of cell migration and invasion (C-E) Cell program and the sevaluated by CCK8, EdU incorporation, and colony formation assays. (F and G) Transwell assay for analysis of cell migration and invasion (C-E) Cell program and the sevaluated by CCK8, EdU incorporation, and colony formation assays. (F and G) Transwell assay for analysis of cell migration and invasion (C-E) Cell program and the sevaluated by CCK8, EdU incorporation, and colony formation assays. (F and G) Transwell assay for analysis of cell migration and invasion (C-E) Cell program and (C-E) Cell prog

also achieved similar und (Finare 2D and E). Next, cell migration and invesion we assessed by Transwell assay. Results showe that ACB16-As also ackdown reduced migration and investor of E e7 and Hep3B cells (Figure 2F and G). Hence, ASB10 that is an oncogene in HCC.

ASB16-ASI Regulates miR-1827/FZD4/ Wnt/β-Catenin Pathway

The molecular mechanism was then investigated. Through bioinformatics analyses, we identified ASB16-AS1 might sponge miR-1827 and miR-1827 may target FZD4. To demonstrate it, luciferase reporter vectors were constructed (Figure 3A). Luciferase reporter assay indicated that luciferase activity of ASB16-AS1-WT or FZD4-WT reporter was reduced by miR-1827 mimic in Huh7 cells (Figure 3B and C). Furthermore, biotin-labeled miR-1827 precipitated ASB16-AS1 in HCC cells (Figure 3D) while RNA immunoprecipitation (RIP) assay suggests miR-1827 associated with FZD4 mRNA (Figure 3E). Thus, miR-1827 directly interacted with ASB16-AS1 and targeted FZD4. Furthermore, ASB16-AS1 knockdown promoted miR-1827 expression (Figure 3F). Besides, miR-1827 mimic or ASB16-AS1 knockdown led to decreased expression of FZD4 (Figure 3G). Notably, miR-1827 inhibitor could rescue the effect of ASB16-AS1 knockdown (Figure 3G), suggesting ASB16-AS1 sponges miR-1827 to upregulate FZD4. FZD4 is an activator of Wnt/ β -catenin pathway in colorectal cancer.¹⁵ No evidence suggests a correlation between FZD4 and Wnt/ β -catenin pathway in HCC. Thus, we speculated





whether FZD4 also activat Wnt/ β -catenin pathway in ocl own of ASB16-AS1 suppressed HCC. Interestingly, D1, CAD2 and SOX4 (the clasexpression Ċ enin pathway) in Huh7 cells sical targ f Wnt/p . genes H). EZD4 overexpression rescued it (Figure (Figure 3H, dicating ASB16-AS1 activates Wnt/β-catenin pathway depending on FZD4. Moreover, bioinformatics analysis suggested FZD4 was upregulated in HCC tissues and qRT-PCR further validated it (Figure 3I and J), implying FZD4 as an oncogene in HCC.

FZD4 Rescues The Effects Of ASB16-AS1 Knockdown

Finally, to further determine whether FZD4 is critical for ASB16-AS1 function, rescue assays were conducted. As

presented by CCK8, EdU incorporation and colony formation assays, FZD4 overexpression rescued the effects of ASB16-AS1 knockdown on proliferation (Figure 4A–C). Similarly, FZD4 restoration also promoted migration and invasion in HCC cells with ASB16-AS1 knockdown (Figure 4D and E). Summarily, ASB16-AS1 promotes HCC progression by regulating miR-1827/FZD4 and activating Wnt/ β -catenin pathway.

Discussion

As a most common tumor, HCC causes large amounts of deaths.¹⁶ Finding effective biomarkers and therapeutic targets for HCC is a pivot task. In this study, we identified ASB16-AS1 is upregulated in HCC samples. Upregulation of ASB16-AS1 indicated low survival rate. Moreover,



Figure 4 FZD4 rescues the effects of ASB16-AS1 knockdown. (A-C) Cell proliferation was neasured by usin CCK8, EdU incorporation and colony formation assays. (D and E) Transwell assay was performed to determine migration and invasion. *p<0.05.

ASB16-AS1 knockdown suppressed proliferation, migration and invasion of HCC cells.

Recently, the function of lncRNAs in tu or ha peen widely explored. Growing evidences show ncRNA evert important roles in the development d p. ssion of ./-19 various cancers, including HC For ample, IncRNA LINC00461 is upregulated in CC and premotes cancer cell proliferation and avasion through miR-149-5p/ LRIG2 pathway.²⁰ Lncl A PVT1 regulates HCC progression via promotine autor agy.²¹ Overexpression of IncRNA LASP1-AS sign antly prinotes HCC growth and invasion oy in reasing e ression of LASP1.²² ASB16-AS is a ray winvestigated novel lncRNA. Only ved that ASB16-AS1 promotes progresa recent study Here, we validated ASB16-AS1 was sion of glioma. markedly upregulater in HCC tissues by bioinformatics approach and qRT-PCR detection. Furthermore, we found ASB16-AS1 expression was correlated with HCC prognosis. Then CCK8, EdU and colony formation assays demonstrated ASB16-AS1 knockdown suppressed HCC proliferation, migration and invasion. Thus, our data support ASB16-AS1 as an oncogene in HCC.

Mechanistically, lncRNAs have been reported to regulate gene expression by interacting with proteins or

s 14,23 Large amounts of studies support that m cRNAs are competing endogenous RNAs (ceRNAs) for tumor.^{9,20,21} niRNAs in For example, lncRNA CNQ1OT1 promotes colon cancer chemoresistance through sponging miR-34a and upregulating ATG4B.²⁴ LncRNA MALAT1 regulates stemness of colorectal cancer cells by interacting with miR-20b-5p to promote Oct4 expression.²⁵ In addition, lncRNA DGCR5 inhibits HCC cell proliferation and invasion via sponging miR-346 and activating KLF14.²⁶ However, how ASB16-AS1 exerts roles remains unclear. In our study, after bioinformatics prediction, we identified ASB16-AS1 may interact with miR-1827. Through luciferase reporter assay and pulldown, their direct interaction was confirmed. Moreover, miR-1827 expression was upregulated after ASB16-AS1 knockdown. Thus, ASB16-AS1 inhibited the activity of miR-1827.

Next, FZD4 was validated to be targeted by miR-1827. And results showed that FZD4 expression was suppressed by miR-1827 or knockdown of ASB16-AS1. Thus, we found that ASB16-AS1 inhibited miR-1827 to upregulate FZD4 expression. FZD4 is an important activator of Wnt/ β -catenin pathway while activation of Wnt/ β -catenin pathway promotes HCC progression.^{15,27} However, whether FZD4 activates Wnt/ β -catenin pathway is unknown. Our results showed that ASB16-AS1 increased activation of Wnt/ β -catenin pathway depending on FZD4 expression in HCC. Up to date, the function of FZD4 in HCC is uncovered. We found FZD4 expression is elevated in HCC tissues, implying an oncogenic role. Thus, rescue assays were performed. We showed that FZD4 restoration effectively rescued the effects of ASB16-AS1 knockdown. Thus, our data revealed ASB16-AS1 regulates miR-1827/FZD4/Wnt/ β -catenin signaling to initiate HCC tumorigenesis.

Collectively, ASB16-AS1 is upregulated in HCC and predicts poor prognosis. ASB16-AS1 overexpression promotes HCC development through regulating miR-1827/FZD4 axis and activating Wnt/ β -catenin pathway. Our research provides a new mechanism regulating HCC tumorigenesis. In our study, some limitations existed. For example, in vivo animal assays should be performed to test the function of ASB16-AS1/miR-1827/FZD4 axis.

Disclosure

The authors report no conflicts of interest in this work.

References

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jenal A. Global cancer statistics 2018: GLOBOCAN estimates of inclusion and mortality worldwide for 36 cancers in 185 courties. *CA Can J Clin.* 2018;68(6):394–424.
- Chen W, Zheng R, Baade PD, et al. Cancer pushics in phina, 201 CA Cancer J Clin. 2016;66(2):115–132.
- Altekruse SF, Henley SJ, Cucinelli J, Monor KA. Changing hepatocellular carcinoma incidence and liver carcinomorphility rates in the United States. *Am J Gastron*, vol. 2014;109(1542–553).
- Si T, Chen Y, Ma D, et al. The sarterial themoembolization prior to liver transplantation for relients with her occellular carcinoma: A meta-analysis. J Gastren levol Hepatol. 201, 2(7):1286–1294.
- Yan X, Hu Z, Feng Y et al. Conferences genomic characterization of long non-codin RNAs cross human cancers. *Cancer Cell*. 2015;28(4):529–540.
- 6. Wang KC HY. Decular techanisms of long noncoding RNAs. *11 Cell*. 11;43(0, 01–14.
- 7. Merey TR, Dieler ME, Mudick JS. Long non-coding RNAs: insight into frequence. *Rev Genet*. 2009;10(3):155–159.
- Leucci E, Ledramin R, Spinazzi M, et al. Melanoma addiction to the long non-comp RNA SAMMSON. *Nature*. 2016;531(7595):518–+. doi:10.1038/nat. 17161
- Wang YG, Wang T, Shi M, Zhai B. Long noncoding RNA EPB41L4A-AS2 inhibits hepatocellular carcinoma development by sponging miR-301a-5p and targeting FOXL1. *J Exp Clin Cancer Res.* 2019;38.
- Wang YF, Yang L, Chen TX, et al. A novel lncRNA MCM3AP-AS1 promotes the growth of hepatocellular carcinoma by targeting miR-194-5p/FOXA1 axis. *Mol Cancer*. 2019;18.

- 11. Jin SF, Yang X, Li JY, Yang WY, Ma HL, Zhang ZY. p53-targeted lincRNA-p21 acts as a tumor suppressor by inhibiting JAK2/STAT3 signaling pathways in head and neck squamous cell carcinoma. *Mol Cancer*. 2019;18.
- Zhang DL, Zhou HG, Liu J, Mao J. Long Noncoding RNA ASB16-AS1 Promotes Proliferation, Migration, and Invasion in Glioma Cells. *Biomed Res Int.* 2019;2019:10.
- Wang Y, Cheng Q, Liu J, Dong M. Leukemia stem cell-released microvesicles promote the survival and migration of myeloid leukemia cells and these effects can be inhibited by MicroRNA34a overexpression. *Stem Cells Int.* 2016;2016:9313425.
- 14. Guo D, Li Y, Chen Y, et al. DANCR promotes HCC progression and regulates EMT by sponging miR-27a-3p via ROCK1/LIMK1/ COFILIN1 pathway. *Cell Prolif.* 2019;52(4):e12628.
- Fang G, Ye BL, Hu BR, Ruan XV, VX. CircRNA_100290 promotes colorectal cancer program on throug miR-516b-induced downregulation of FZD4 expression and Wnt/bet materin signaling. *Biochem Biophys Res Commun.* 24:504(1):184–19.
- 16. Raoul JL, Forner A, Boltani L, Chenn TT, Klenkner R, de Baere T. Updated use of TACL for hepatocellum carry of treatment: how and when to use is based on anical evicence. *Cancer Treat Rev.* 2019;72:28–36.
- Liu YC, Li Y, Li F, Li Shao Y, Ja LP. SNHG15 functions as a tumor symposor in thyn Corper. J Cell Biochem. 2019;120 (4):612–612
- 18. Dong H, Hu J, Eu K, et al. Activation of LncRNA TINCR by acetylation comotes trastuzumab resistance and epithelialmesenchymal transition by targeting MicroRNA-125b in breast cancer. *Mol Center*. 2019;18(1):3.
- 20. Ji D, Wang Y, Li H, Sun B, Luo X. Long non-coding RNA INC00461/miR-149-5p/LRIG2 axis regulates hepatocellular carcinoma progression. *Biochem Biophys Res Commun.* 2019;512(2):176– 181.
- Yang L, Peng X, Jin H, Liu J. Long non-coding RNA PVT1 promotes autophagy as ceRNA to target ATG3 by sponging microRNA-365 in hepatocellular carcinoma. *Gene.* 2019;697:94–102.
- Yin L, Chen Y, Zhou Y, et al. Increased long noncoding RNA LASP1-AS is critical for hepatocellular carcinoma tumorigenesis via upregulating LASP1. J Cell Physiol. 2019;234(8):13493–13509.
- Zhu P, Wang Y, Huang G, et al. Inc-beta-Catm elicits EZH2-dependent beta-catenin stabilization and sustains liver CSC self-renewal. *Nat Struct Mol Biol.* 2016;23(7):631–639.
- 24. Li Y, Li C, Li D, Yang L, Jin J, Zhang B. IncRNA KCNQ10T1 enhances the chemoresistance of oxaliplatin in colon cancer by targeting the miR-34a/ATG4B pathway. *Onco Targets Ther.* 2019; 12:2649–2660.
- 25. Tang D, Yang Z, Long F, et al. Long noncoding RNA MALAT1 mediates stem cell-like properties in human colorectal cancer cells by regulating miR-20b-5p/Oct4 axis. *J Cell Physiol.* 2019;234(11): 20816–20828.
- Wang YG, Liu J, Shi M, Chen FX. LncRNA DGCR5 represses the development of hepatocellular carcinoma by targeting the miR-346/ KLF14 axis. J Cell Physiol. 2018;234(1):572–580.
- Yan X, Zhang D, Wu W, et al. Mesenchymal stem cells promote hepatocarcinogenesis via lncRNA-MUF interaction with ANXA2 and miR-34a. *Cancer Res.* 2017;77(23):6704–6716.

Cancer Management and Research

Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient.

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors. Submit your manuscript here: https://www.dovepress.com/cancer-management-and-research-journal

Dovepress

submit your manuscript | www.dovepress.com 9378 **Dove**Press