Relationship Between Sleep Duration and Stroke History in Middle-Aged and Elderly in Guiyang: A Cross-Sectional Survey

Li Liu¹, Jingyuan Yang¹, Junhua Wang¹, Meng Nie², Ziyun Wang¹, Han Guan¹, Jin Hu¹, Feng Hong¹

¹School of Public Health, The Key Laboratory of Environmental Pollution Monitoring and Disease Control, Ministry of Education, Guizhou Medical University, Guiyang, Guizhou, People's Republic of China; ²School of Public Health, Guizhou Medical University, Guiyang, Guizhou, People's Republic of China

Correspondence: Feng Hong, School of Public Health, The Key Laboratory of Environmental Pollution Monitoring and Disease Control, Ministry of Education, Guizhou Medical University, Guiyang, Guizhou, People's Republic of China, Tel +86-13984180981, Email 64048857@qq.com

Background: With over 2 million new cases annually, stroke is associated with the higher amount of disability-adjusted life-years lost than any other disease in China; however, the relationship between sleep time and stroke has not been concluded yet. Aim of this study was to analyze the relationship between sleep duration and stroke history in middle-aged and elderly people in Guiyang, China.

Methods: This study was a cross-sectional survey carried out in 40–99-years-old permanent residents of Guiyang. Yunyan, Wudang, and Baiyun districts and Xiuwen County were selected by stratified multilevel sampling for a face-to-face survey. Demographics, history of stroke, and self-reported sleep behavior data were collected, and multivariable logistic regression models were used to gradually adjust possible confounding factors.

Results: A total of 5065 participants were included, of them 126 (2.5%) had a history of stroke. Short sleep (<7 h) was observed in 11.0%, sufficient sleep (7–9 h) in 69.4%, and long sleep (>9 h) in 19.6%. Sleep duration and stroke prevalence showed a U-shaped distribution. When taking the sleep duration of 7–9 h as a reference, sleep duration >9 h was associated with stroke (all $P < 0.05$) in the univariable model (OR = 2.68, 95% CI: 1.83–3.93) and in the multivariable models 1 (OR = 2.35, 95% CI: 1.59–3.47), 2 (OR = 2.27, 95% CI: 1.53–3.37), 3 (OR = 2.25, 95% CI: 1.51–3.33), and 4 (OR = 2.11, 95% CI: 1.39–3.19). There were no significant differences between the <7 and 7–9 h groups ($P > 0.05$).

Conclusion: Thus, long sleep duration (>9 h) is independently associated with history of stroke in middle-aged and elderly people in Guiyang.

Keywords: stroke, sleep duration, prevalence, middle-aged, elderly, China

Introduction

Stroke is a cerebrovascular disease that rapidly leads to localized or diffuse brain dysfunction due to acute cerebral circulation disorder.¹ Stroke features a high mortality rate (about 5.5 million each year) and high morbidity rate, as well as long-term disability in up to 50% of survivors.^{2,3} The incidence of stroke in China is 298 per 100,000 person-years in rural areas and 204 per 100,000 person-years in urban areas,⁴ with geographic variations across the country⁵ and the incidence is increasing over time.⁶ Stroke is, therefore, a significant public health burden. The risk factors for stroke include among others a history of transient ischemic attacks, smoking, metabolic syndrome, hypertension, heavy alcohol use, diabetes mellitus type 2, high cholesterol levels, carotid artery stenosis, obesity, and poor physical status.^{7,8}

Sleep duration and quality are able to affect not only the quality of life but also physical health.^{9,10} Previously, sleep duration was reported to be associated with hypertension, diabetes, and other diseases,⁹ and frequent snoring during sleep has been associated with the risk of stroke in various populations.^{10–12} Furthermore, recent studies showed that both short and long duration of sleep are predictors of cardiovascular outcomes and relationship between sleep duration and the risk of stroke follows a J-shaped curve, as stroke may follow a shift in sleep patterns, and sleep patterns in turn might be

affected by a history of stroke and related changes in everyday life.^{13,14} When considering a normal sleep duration of 7 h as a reference (9), sleep duration <7 h will increase the risk of stroke by 7% for every 1 h of shorter sleep, and >7 h will increase the risk of stroke by 17% for every 1 h of sleep duration >7 h.¹⁴

Nevertheless, considering the wide variation in the incidence of stroke among different regions in China, it is possible that the relationship between sleep duration and the risk of stroke also varies considerably.^{15,16} Because of the low prevalence of stroke in the community, it was hypothesized that in Guiyang, a city in southwest China, currently in the process of gradual urbanization with relatively high level of medical services, sleep time may differ along with the relationship to stroke caused by it. Epidemiological studies with sufficiently large population samples can provide reliable evidence to answer the necessary research questions discussed above.

Therefore, the purpose of this study was to investigate the relationship between sleep duration and history of stroke in the middle-aged and elderly people in Guiyang through a survey of more than 10,000 residents in Guiyang. The research results can provide a reference for the prevention and control of stroke.

Materials and Methods

Study Design and Participants

The survey adopted stratified and multilevel sampling methods and was held from July 2017 to August 2017. According to the geographical location, Guiyang city was divided into three types of regions, and in each region, a district was randomly selected according to the random number table method, and then 1 township (town or community) was randomly selected from each district. In the final selected area, the survey was conducted with the family as the unit, and all middle-aged and elderly members were included in this study as the subjects of the survey. The inclusion criteria were 1) voluntary participation; 2) >40 years of age, and 3) During investigation, local residents who have left Guiyang for <12 months or migrant residents who have lived for >6 months in Guiyang in the past 12 months. Those potential participants who were unable to finish the survey after three attempts including contacting relatives were excluded. The study was conducted in accordance with the Declaration of Helsinki and approved by the ethics committee of Drug Clinical Trial of Affiliated Hospital of Guizhou Medical University, and all participants signed the informed consent form.

Survey Method

The questionnaire of the 2017 Guiyang Residents Disease Spectrum Survey was adopted, with the household as a unit, and a face-to-face questionnaire survey was conducted by all residents who met the eligibility criteria in the household. In principle, the questionnaire had to be answered face to face. For those who could not answer the questionnaire due to physical, mental, intellectual, and age factors, their families answered the questionnaire on their behalf. The survey contents included general situation (sex, age, nationality, education, marital status, and annual family income), stroke history, sleep behavior (including sleep duration and sleep quality), and lifestyle behaviors (smoking, drinking, and exercise frequency).

Definition of Relevant Indicators and Grouping

For participants who acknowledged the history of stroke, diagnosis was confirmed by contacting a local hospital. Sleep duration was calculated from “sleep duration and getting up time in the past month” questionnaire section, and the participants were divided into <7, 7–9, and >9 h. Referring to the sleep duration standard,¹⁷ <7 h was regarded as short sleep duration, 7–9 h was considered as average sleep duration, and >9 h was considered as long sleep duration. Sleep quality was obtained from the “self-evaluation of night sleep quality in the past month” questionnaire section. In this study, sleep quality was self-reported by participants using a five-point Likert scale as “very good, good, fair, poor, and very poor”. However, the number of “very good” and “very poor” cases was small, so we combined the data for the statistical assessment (very good was considered as good, and very poor was considered as poor). Thus, sleep quality, as well as general health status, were divided into “good” (including “very good” and “good”), average, and poor (including “very poor” and “poor”). Hypertension, diabetes, and coronary heart disease were reported by the participants, according

to a previously known diagnosis. Overweight and obesity were evaluated based on the body mass index (BMI, kg/m²). In this study, the Chinese Adult Weight Determination Standard was adopted to classify overweight and obesity.¹⁸ BMI < 18.5 was lean, 18.5 ≤ BMI < 24.0 was normal, 24.0 ≤ BMI < 28.0 was overweight, and BMI ≥ 28.0 was obesity. Exercising was defined as physical activity specifically aimed at improving health or fitness (excluding walking to/from work or other everyday activities) and according to frequency exercising was divided into 0–3, 4–5, and 6–7 days/week.

Calculation of Sample Size

The sample size was calculated using the following equation¹⁹ $n = \frac{Z_{1-\alpha/2}^2 \times pq}{d^2}$

The adopted prevalence of stroke was 5%,^{20,21} and the allowable error was $d = 0.2p$, α Take 0.05 and the calculated sample size is 1900. Considering the design effect $deff = 2$,²² the total sample size $n = n \times Deff = 1900 \times 2 = 3800$; Considering the 20% loss of follow-up rate, at least 4750 cases should be included in this study.

Statistical Method

EpiData 3.1 (Centers for Disease Control, Atlanta, GA, USA) was used to establish the database. All data were double-entered and verified by comparison of the independent two entries. SPSS 25.0 (IBM Corp, Armonk, NY, USA) was used for statistical analysis. The data are described as means ± standard deviations and n (%). The distribution of sleep duration and the prevalence of stroke according to different characteristics were tested using the chi-square test. The relationship between sleep duration and stroke was analyzed by a two-classification logistic regression model. First, univariable logistic regression analysis was carried out with sleep duration as the independent variable and stroke prevalence as the dependent variable. Then, the possible confounding factors were gradually added using a multivariable logistic regression analysis ([Supplementary Table S1](#)): Model 1: adjusted for sex and age; Model 2: On the basis of Model 1, adjusted for education, occupation, marital status and migrant workers; Model 3: on the basis of Model 2, adjusted for smoking, drinking, and sitting or leaning time every day; and Model 4: on the basis of Model 3, further adjusted for health status, sleep quality, hypertension, diabetes, and coronary heart disease. GraphPad Prism 8.4.2 (GraphPad Software Inc., San Diego, CA, USA) was used to draw the forest map of the results of the multivariate analysis. Two-sided P-values <0.05 were considered statistically significant.

Results

Characteristics of the Participants

A total of 5065 participants were included in the analysis, aged between 40 and 99, with a Median age of 57.5 (48.8, 67.9) years. Among them, 2327 (45.9%) were men, and 2738 (54.1%) were women. By profession, participants were mainly farmers (26.7%) and unemployed/out of work (24.5%), 53.5% of them had a primary school education or below, 81.7% were married, 46.9% ($n = 2373$) felt that their health status was “good” and 17.8% (902) felt that their health status was “poor”, and 2816 (55.6%) suffered from chronic diseases ([Table 1](#)).

Distribution of Self-Reported Sleep Duration with Different Characteristics

Among the 5065 subjects, 555 (11.0%) had slept <7 h, 3516 (69.4%) slept 7–9 h, and 994 (19.6%) slept >9 h. People of different ages, education, occupation, marital status, health status, sleep quality, regular three meals, dietary pattern, smoking, and drinking had different sleep durations (all $P < 0.05$) ([Table 1](#)). With increasing age, the proportion of people who slept for 7–9 h gradually decreased ($P_{trend} < 0.001$) ([Table 1](#)).

Prevalence of Stroke

A total of 126 middle-aged and elderly people suffered from a stroke in this survey, with a prevalence of 2.5%, including 55 men (prevalence of 2.4%) and 71 women (prevalence of 2.6%). The prevalence of stroke was the lowest (1.8%) among the subjects with 7–9 h of sleep duration, while the prevalence of stroke among the subjects with <7 and >9 h of sleep duration increased continuously and showed a U shape distribution. The prevalence of stroke among the subjects with long sleep duration (>9 h) was higher than that of the subjects with short sleep duration (<7 h). With increasing age,

Table 1 Characteristics of 5065 Middle-Aged and Elderly People in Guiyang City in 2017 According to Sleep Duration

Item	Grouping	n	%	Sleep Duration			χ^2	P
				<7 h	7–9 h	>9 h		
Gender	Male	2327	45.9	277 (11.9)	1600 (68.8)	450 (19.3)	3.967	0.138
	Female	2738	54.1	278 (10.2)	1916 (70.0)	544 (19.3)		
Age	40~	1453	28.7	155 (10.7)	1037 (71.4)	261 (18.0)	70.380	<0.001
	50~	1382	27.3	179 (13.0)	975 (70.5)	228 (16.5)		
	60~	1154	22.8	137 (11.9)	796 (69.0)	221 (19.2)		
	70~	768	15.2	68 (8.9)	521 (67.8)	179 (23.3)		
	≥80	308	6.1	16 (5.2)	187 (60.7)	105 (34.1)		
Marital Status	Married	4140	81.7	450 (10.9)	2918 (70.5)	772 (18.6)	14.906	0.001
	Others	925	18.3	105 (11.4)	598 (64.6)	222 (24.0)		
Body mass index	Lean	2716	53.6	29 (10.8)	176 (65.4)	64 (23.8)	6.449	0.375
	Normal	269	5.3	286 (10.5)	1883 (69.3)	547 (20.1)		
	Overweight	1538	30.4	174 (11.3)	1080 (70.2)	284 (18.5)		
	Obese	542	10.7	66 (12.2)	377 (69.9)	99 (18.3)		
Chronic diseases		2816	55.6	324 (11.5)	1930 (68.5)	562 (20.0)	2.805	0.246
Hypertension		1092	21.6	110 (10.1)	742 (67.9)	240 (22.0)	5.354	0.069
Diabetes		268	5.3	29 (10.8)	179 (66.8)	60 (22.4)	1.387	0.500
CHD		119	2.3	13 (10.9)	79 (66.4)	27 (22.7)	0.744	0.689
Stroke		126	2.5	15 (11.9)	64 (50.8)	47 (37.3)	27.136	<0.001
Smoking	Never	2996	59.2	292 (9.7)	2114 (70.6)	590 (19.7)	14.412	0.006
	Current	1702	33.6	226 (13.3)	1147 (67.4)	329 (19.3)		
	Previous	367	7.2	37 (10.1)	255 (69.5)	75 (20.4)		
Drinking alcohol	Never	2996	59.2	301 (10.0)	2080 (69.4)	615 (20.5)	16.091	0.003
	Current	1608	31.7	201 (12.5)	1133 (70.5)	274 (17.0)		
	Previous	461	9.1	53 (11.5)	303 (65.7)	105 (22.8)		
Sleep quality	Good	3038	60.0	286 (9.4)	2154 (70.9)	598 (19.7)	43.090	<0.001
	General	1030	20.3	103 (10.0)	728 (70.7)	199 (19.3)		
	Poor	997	19.7	166 (16.6)	634 (63.6)	197 (19.8)		
Exercise frequency (day/week)	0–3	3704	73.1	410 (11.1)	2509 (67.7)	785 (21.2)	23.059	<0.001
	4–5	229	4.5	25 (10.9)	169 (73.8)	35 (15.3)		
	6–7	1132	22.3	120 (10.6)	838 (74.0)	174 (15.4)		

Abbreviation: CHD, coronary heart disease.

the prevalence of stroke gradually increased ($P_{\text{trend}} < 0.001$). The prevalence of stroke among middle-aged and elderly people with different occupations, education, marital status, health status, chronic disease, sleep quality, smoking, sitting or leaning time and drinking status was significantly different (all $P < 0.05$) (Table 2).

Relationship Between Self-Reported Sleep Duration and Stroke History

The results of the univariable and multivariable logistic regression analysis showed that when taking the sleep duration of 7–9 h as reference, sleep duration >9 h was associated with reported history of stroke (all $P < 0.05$) in the univariable model (OR = 2.68, 95% CI: 1.83–3.93) and in the multivariable model 1 (OR = 2.35, 95% CI: 1.59–3.47), model 2 (OR = 2.27, 95% CI: 1.53–3.37), model 3 (OR = 2.25, 95% CI: 1.51–3.33), and model 4 (OR = 2.11, 95% CI: 1.39–3.19). There were no significant differences between the <7 and 7–9 h groups ($P > 0.05$) (Table 3 and Figures 1 and Figures 2).

Discussion

Sleep duration is associated with the risk of stroke,^{13,14} but the incidence of stroke varies considerably in different countries and often even within one country.^{2–5} Therefore, the relationship between sleep duration and stroke could also vary. The relationship between sleep time and stroke has not been concluded yet in China. This study aimed to study the relationship between self-reported sleep duration and history of stroke in middle-aged and elderly people in Guiyang, China.

Table 2 Prevalence of Stroke in 5065 Middle-Aged and Elderly People in Guiyang City in 2017

Item	Grouping	With Stroke	Without Stroke	χ^2/Z	P
Gender	Male	55 (2.4)	2272 (97.6)	0.273	0.601
	Female	71 (2.6)	2667 (97.4)		
Age	40~	6 (0.4)	1447 (99.6)	74.885	<0.001
	50~	22 (1.6)	1360 (98.4)		
	60~	41 (3.6)	1113 (96.4)		
	70~	36 (4.7)	732 (95.3)		
	≥80	21 (6.8)	287 (93.2)		
Marital Status	Married	94 (2.3)	4046 (97.7)	4.406	0.036
	Others	32 (3.5)	893 (96.5)		
Health status	Good	24 (1.0)	2349 (99.0)	154.659	<0.001
	General	27 (1.5)	1763 (98.5)		
	Poor	75 (8.3)	827 (91.7)		
Chronic diseases		112 (4.0)	2704 (96.0)	58.012	<0.001
Sleep duration	<7 h	15 (2.7)	540 (97.6)	27.136	<0.001
	7–9 h	64 (1.8)	3452 (98.2)		
	>9 h	47 (4.7)	947 (95.3)		
Sleep quality	Good	50 (1.6)	2988 (98.4)	26.901	<0.001
	General	31 (3.0)	999 (97.0)		
	Poor	45 (4.5)	952 (95.5)		
Smoking	Never	78 (2.6)	2918 (97.4)	6.149	0.046
	Current	33 (1.9)	1669 (98.1)		
	Previous	15 (4.1)	352 (95.9)		
Drinking alcohol	Never	88 (2.9)	2908 (97.1)	18.877	<0.001
	Current	19 (1.2)	1589 (98.8)		
	Previous	19 (4.1)	442 (95.9)		
Exercise frequency (day/week)	0–3	86 (2.3)	3618 (97.7)	1.678	0.432
	4–5	6 (2.6)	223 (97.4)		
	6–7	34 (3.0)	1098 (96.0)		
BMI	Lean	64 (2.4)	2652 (97.6)	2.348	0.503
	Normal	7 (2.6)	262 (97.6)		
	Overweight	45 (2.9)	1493 (97.1)		
	Obese	10 (1.8)	532 (98.2)		
Hypertension		88 (8.1)	1004 (91.9)	178.111	<0.001
Diabetes		20 (7.5)	248 (92.5)	28.873	<0.001
CHD		12 (10.1)	107 (89.9)	28.989	<0.001
Sitting or leaning time		5.00 (3.00, 7.00)	4.50 (3.00, 6.00)	–2.202	0.028

Abbreviations: BMI, body mass index (kg/m^2); CHD, coronary heart disease.

Based on the conducted population-based survey and research, with a large sample size and strong statistical power, the relationship between sleeping habits and stroke was explored for the community populations in Guiyang City. Due to the relatively high level of medical services in Guiyang City compared to cities around, the overall incidence of stroke was low, but after controlling the possible confounding risk factors, the independent association between long sleep duration (>9 h) and history of stroke was still identified, which provides a scientific basis for future in-depth research on the impact of sleep on stroke and intervention studies.

In this study, the prevalence of stroke was 2.5% among middle-aged and elderly people aged ≥40 years old in Guiyang,^{5,20,21,23} 2.4% for men and 2.6% for women, without significant difference, gradually increased with age. Found prevalence values are slightly lower than reported in neighboring provinces, which contributes to the epidemiological study of stroke distribution over China.^{5,20,21,23} Differences might be partly explained by a higher level of medical services in Guiyang City compared to cities around; in addition, the prevalence of stroke in individuals with

Table 3 Correlation Between Sleep Duration and Stroke in Middle-Aged and Elderly People in Guiyang City, OR (95% CI)

Item		Sleep Duration		
		<7 h	7–9 h	>9 h
Univariable model	–	1.498 (0.848–2.648)	1	2.677 (1.825–3.927)*
Multivariable model	1	1.600 (0.900–2.843)	1	2.346 (1.589–3.465)*
	2	1.718 (0.963–3.065)	1	2.272 (1.533–3.367)*
	3	1.700 (0.952–3.036)	1	2.245 (1.514–3.331)*
	4	1.673 (0.911–3.071)	1	2.107 (1.391–3.190)*

Notes: The univariable factor model includes sleep duration only. Model 1: sleep duration, gender, and age. Model 2: Model 1, educational level, occupation, marital status, and migrant workers. Model 3: Model 2, smoking, drinking, and sitting or leaning time every day. Model 4: Model 3, health status, sleep quality, hypertension, diabetes, and coronary heart disease. *P<0.05.

hypertension, diabetes, and coronary heart disease was 8.1%, 7.5%, and 10.1%, respectively, which was significantly higher than in individuals without these diseases, also in line with previous studies.^{5,7,8,20,21,23}

Of the 5065 middle-aged and elderly participants in Guiyang, most (69.4%) had reported normal sleep duration (7–9 h), 11% had short sleep duration (<7 h), while 19.6% reported long sleep duration (>9 h). Age, education, occupation, marital status, health status, sleep quality, smoking status, and drinking status were all associated with sleep duration. Moreover, the worse the sleep quality, the greater the proportion of people who have short sleep duration, as observed by previous studies,¹⁶ confirming that quality²⁴ of sleep appears to depend on its quantity among other factors.²⁴

This study showed that the prevalence of stroke in middle-aged and elderly people with different sleep duration was different. The prevalence of stroke in people with normal sleep duration (7–9 h) was the lowest, at 1.8%, while in people with long sleep duration (>9 h) it was equal to 4.7%, still higher than that in people with short sleep duration (<7 h) (2.7%): so called a “U” distribution between sleep duration and stroke. This relationship was still observed after adjusting for the other risk factors of stroke, being in line with previous studies.^{11,13,14,16,25,26}

Various reasons might explain the relationship between long sleep and stroke. As our study was retrospective by the nature, it is possible that in some of the participants sleep patterns have shifted to the longer sleep duration even after the stroke. However, previous studies have reported that long sleep is associated with elevated inflammatory markers,^{27,28} and inflammation is involved in the pathogenesis of stroke.^{7,8,29} Long sleep is also associated with high blood lipid levels,^{30,31} which directly contribute to atherogenesis. In addition, the elevated prevalence of metabolic syndrome,³² carotid artery stenosis,³³ and white matter hyperintensities^{34,35} is observed in people with long sleep duration. Indirect associations could be found between long sleep and a more sedentary life,^{36,37} depression, sleep apnea, and lower socioeconomic status,^{12,38,39} which are all risk factors for stroke.^{7,8} The research hypothesis implied that in Guiyang, a city in southwest China, the process of gradual urbanization led to the significant change of lifestyle in elderly, and thus sleep time may differ along with the relationship to stroke caused by it. With the continuous development of China's economy, many cities are in the process of urbanization right now, and the change of lifestyle will be relevant to other developing regions as a point of comparison. In addition, the correlation degree is relatively strong; thus, it could be applied to the situation of younger people, in which sleep time shortens for other reasons, so in the future, it might lead to

Figure 1 Univariable model for the association between sleep duration and the risk of stroke. Single factor model.

Figure 2 4 multivariate models for association between sleep duration and the risk of stroke. **(A)** Model 1: sleep duration, gender, and age. **(B)** Model 2: Model 1, educational level, occupation, marital status and migrant workers. **(C)** Model 3: Model 2, smoking, drinking, and sitting or leaning time every day. **(D)** Model 4: Model 3, health status, sleep quality, hypertension, diabetes, and coronary heart disease.

an earlier age of stroke, an increased risk of stroke, and thus an increased healthcare burden. Based on that, future studies should focus on sleep duration, stroke pathogenesis, and its molecular epidemiology.

This study has some limitations. It was a cross-sectional survey, preventing any determination of a cause-to-effect relationship. As it was not possible to assess sleep objectively in a sample of this magnitude, sleep duration, stroke history, and other indicators were reported by the participants, resulting in inevitable subjectivity and biases. No blood parameters or intensity, method and type of exercises were analyzed in this study. Furthermore, stroke history and sleep behavior change before baseline were not collected

in questionnaire. Due to the retrospective nature of the stroke, it is possible that in some of the participants sleep patterns have shifted to the longer sleep duration after the stroke (as in some of them it might have shifted to shorter sleep pattern). Thus, due to the lack of prospective observation, the causal relationship between sleep duration and stroke disease cannot be obtained, and further epidemiological verification is needed.

Conclusions

In conclusion, a long sleep duration (>9 h) is independently associated with history of stroke in middle-aged and elderly people in Guiyang. Short sleep duration was not associated with a history of stroke in this population.

Abbreviations

BMI, body mass index.

Data Sharing Statement

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Ethics Approval and Informed Consent

The study was approved by the ethics committee of Drug Clinical Trial of Affiliated Hospital of Guizhou Medical University and all participants signed the informed consent form.

Acknowledgments

Thanks to Guiyang Health and Family Planning Commission, Guiyang Center for Disease Control and Prevention, the health and family planning bureau of each investigation district and county, the township health center, the community health service center, the subdistrict office for the organization and coordination of this investigation. Thanks to the investigation site health and family planning professionals, Guizhou Medical University School of Public Health disease spectrum project team to participate in the investigation of teachers and students for the smooth implementation of the investigation paid hard work. Thanks to all the respondents for their support and cooperation.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

This work was supported by the Project on Investigation of Disease Spectrum of Residents in Guiyang City (BCXZFCG2016-023A). This work was supported by the National Key R&D Program “Precision Medicine Initiative” of China (Grant no: 2017YFC0907301).

Disclosure

The authors declare no conflicts of interest in this work.

References

1. Sacco RL, Kasner SE, Broderick JP, et al. An updated definition of stroke for the 21st century: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2013;44(7):2064–2089. doi:10.1161/STR.0b013e318296aeca
2. Lunardi Baccetto S, Lehmann C. Microcirculatory changes in experimental models of stroke and CNS-injury induced immunodepression. *Int J Mol Sci*. 2019;20:5184.
3. Donkor ES. Stroke in the 21(st) century: a snapshot of the burden, epidemiology, and quality of life. *Stroke Res Treat*. 2018;2018:3238165. doi:10.1155/2018/3238165

4. Wu S, Wu B, Liu M, et al. Stroke in China: advances and challenges in epidemiology, prevention, and management. *Lancet Neurol.* 2019;18(4):394–405. doi:10.1016/S1474-4422(18)30500-3
5. Jiang B, Wang WZ, Chen H, et al. Incidence and trends of stroke and its subtypes in China: results from three large cities. *Stroke.* 2006;37:63–68. doi:10.1161/01.STR.0000194955.34820.78
6. Hu S, Cui B, Mlynash M, et al. Stroke epidemiology and stroke policies in China from 1980 to 2017: a systematic review and meta-analysis. *Int J Stroke.* 2020;15:18–28. doi:10.1177/1747493019873562
7. Meschia JF, Bushnell C, Boden-Albala B, et al. Guidelines for the primary prevention of stroke: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke.* 2014;45:3754–3832. doi:10.1161/STR.0000000000000046
8. Allen CL, Bayraktutan U. Risk factors for ischaemic stroke. *Int J Stroke.* 2008;3(2):105–116. doi:10.1111/j.1747-4949.2008.00187.x
9. Cabeza de Baca T, Chayama KL, Redline S, et al. Sleep debt: the impact of weekday sleep deprivation on cardiovascular health in older women. *Sleep.* 2019;42:zsz149.
10. Hermann DM, Bassetti CL. Role of sleep-disordered breathing and sleep-wake disturbances for stroke and stroke recovery. *Neurology.* 2016;87(13):1407–1416. doi:10.1212/WNL.00000000000003037
11. Li J, McEvoy RD, Zheng D, et al. Self-reported snoring patterns predict stroke events in high-risk patients with OSA: post hoc analyses of the SAVE Study. *Chest.* 2020;158(5):2146–2154. doi:10.1016/j.chest.2020.05.615
12. Yaggi HK, Concato J, Kernan WN, et al. Obstructive sleep apnea as a risk factor for stroke and death. *N Engl J Med.* 2005;353(19):2034–2041. doi:10.1056/NEJMoa043104
13. Cappuccio FP, Cooper D, D'Elia L, et al. Sleep duration predicts cardiovascular outcomes: a systematic review and meta-analysis of prospective studies. *Eur Heart J.* 2011;32(12):1484–1492. doi:10.1093/eurheartj/ehr007
14. Li W, Wang D, Cao S, et al. Sleep duration and risk of stroke events and stroke mortality: a systematic review and meta-analysis of prospective cohort studies. *Int J Cardiol.* 2016;223:870–876. doi:10.1016/j.ijcard.2016.08.302
15. Li LJ, Yang Y, Guan BY, et al. Insomnia is associated with increased mortality in patients with first-ever stroke: a 6-year follow-up in a Chinese cohort study. *Stroke Vasc Neurol.* 2018;3:197–202. doi:10.1136/svn-2017-000136
16. Ji A, Lou H, Lou P, et al. Interactive effect of sleep duration and sleep quality on risk of stroke: an 8-year follow-up study in China. *Sci Rep.* 2020;10(1):8690. doi:10.1038/s41598-020-65611-y
17. Hirshkowitz M, Whiton K, Albert SM, et al. National Sleep Foundation's sleep time duration recommendations: methodology and results summary. *Sleep Health.* 2015;1(1):40–43. doi:10.1016/j.sleh.2014.12.010
18. Consultation WE. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet.* 2004;363(9403):157–163. doi:10.1016/S0140-6736(03)15268-3
19. Zhan S. *Epidemiology (Seventh Edition, National University Course Book, for Preventive Medical Specialty)*. Chinese ed. Beijing: People's Medical Publishing House; 2017.
20. Yi X, Luo H, Zhou J, et al. Prevalence of stroke and stroke related risk factors: a population based cross sectional survey in southwestern China. *BMC Neurol.* 2020;20(1):5. doi:10.1186/s12883-019-1592-z
21. Wang W, Jiang B, Sun H, et al. Prevalence, incidence, and mortality of stroke in China: results from a nationwide population-based survey of 480 687 adults. *Circulation.* 2017;135(8):759–771. doi:10.1161/CIRCULATIONAHA.116.025250
22. Li XS, HaoYT, LiuMN. *Health Statistics*. 8th ed. Beijing: People's Medical Publishing House; 2017.
23. Yang JJ, Zhou C, Huang Y, et al. Icotinib versus whole-brain irradiation in patients with EGFR-mutant non-small-cell lung cancer and multiple brain metastases (BRAIN): a multicentre, Phase 3, open-label, parallel, randomised controlled trial. *Lancet Respir Med.* 2017;5:707–716. doi:10.1016/S2213-2600(17)30262-X
24. Bin YS. Is sleep quality more important than sleep duration for public health? *Sleep.* 2016;39:1629–1630. doi:10.5665/sleep.6078
25. Patyar S, Patyar RR. Correlation between sleep duration and risk of stroke. *J Stroke Cerebrovasc Dis.* 2015;24(5):905–911. doi:10.1016/j.jstrokecerebrovasdis.2014.12.038
26. Zhou L, Yu K, Yang L, et al. Sleep duration, midday napping, and sleep quality and incident stroke: the Dongfeng-Tongji cohort. *Neurology.* 2020;94(4):e345–e356. doi:10.1212/WNL.00000000000008739
27. Patel SR, Zhu X, Storfer-Isser A, et al. Sleep duration and biomarkers of inflammation. *Sleep.* 2009;32(2):200–204. doi:10.1093/sleep/32.2.200
28. Dowd JB, Goldman N, Weinstein M. Sleep duration, sleep quality, and biomarkers of inflammation in a Taiwanese population. *Ann Epidemiol.* 2011;21(11):799–806. doi:10.1016/j.annepidem.2011.07.004
29. Su H. Inflammation and genetic factors in stroke pathogenesis. *Neuroimmunol Neuroinflamm.* 2017;4:260–262. doi:10.20517/2347-8659.2017.22
30. Yang L, Yang H, He M, et al. Longer sleep duration and midday napping are associated with a higher risk of CHD incidence in middle-aged and older Chinese: the Dongfeng-Tongji Cohort Study. *Sleep.* 2016;39(3):645–652. doi:10.5665/sleep.5544
31. Zhan Y, Chen R, Yu J. Sleep duration and abnormal serum lipids: the China health and nutrition survey. *Sleep Med.* 2014;15(7):833–839. doi:10.1016/j.sleep.2014.02.006
32. Yang L, Xu Z, He M, et al. Sleep duration and midday napping with 5-year incidence and reversion of metabolic syndrome in middle-aged and older Chinese. *Sleep.* 2016;39(11):1911–1918. doi:10.5665/sleep.6214
33. Abe T, Aoki T, Yata S, et al. Sleep duration is significantly associated with carotid artery atherosclerosis incidence in a Japanese population. *Atherosclerosis.* 2011;217(2):509–513. doi:10.1016/j.atherosclerosis.2011.02.029
34. Ramos AR, Dong C, Rundek T, et al. Sleep duration is associated with white matter hyperintensity volume in older adults: the Northern Manhattan Study. *J Sleep Res.* 2014;23(5):524–530. doi:10.1111/jsr.12177
35. DeBette S, Beiser A, DeCarli C, et al. Association of MRI markers of vascular brain injury with incident stroke, mild cognitive impairment, dementia, and mortality: the Framingham Offspring Study. *Stroke.* 2010;41(4):600–606. doi:10.1161/STROKEAHA.109.570044
36. Lakerveld J, Mackenbach JD, Horvath E, et al. The relation between sleep duration and sedentary behaviours in European adults. *Obes Rev.* 2016;17(Suppl S1):62–67. doi:10.1111/obr.12381
37. Ford ES, Caspersen CJ. Sedentary behaviour and cardiovascular disease: a review of prospective studies. *Int J Epidemiol.* 2012;41(5):1338–1353. doi:10.1093/ije/dys078
38. Patel SR, Malhotra A, Gottlieb DJ, et al. Correlates of long sleep duration. *Sleep.* 2006;29(7):881–889. doi:10.1093/sleep/29.7.881

39. Pan A, Sun Q, Okereke OI, et al. Depression and risk of stroke morbidity and mortality: a meta-analysis and systematic review. *JAMA*. 2011;306(11):1241–1249. doi:10.1001/jama.2011.1282

Neuropsychiatric Disease and Treatment

Dovepress

Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS, and is the official journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>