

# Comparative Study of Perioperative and Oncological Outcomes Between Elderly Patients and Younger Patients Who Received Radical Cystectomy and Pelvic Lymph Node Dissection: A Single-Center Retrospective Study

Haixin Wang<sup>1-4,\*</sup>, Haiwen Huang<sup>1-3,\*</sup>, Meixia Shang<sup>5</sup>, Han Hao<sup>1-3</sup>, Zhijun Xi<sup>1-3</sup>

<sup>1</sup>Department of Urology, Peking University First Hospital, Beijing, People's Republic of China; <sup>2</sup>Institute of Urology, Peking University, National Urological Cancer Center, Beijing, People's Republic of China; <sup>3</sup>National Research Center for Genitourinary Oncology, Beijing, People's Republic of China; <sup>4</sup>Department of Urology, Yankuang New Journey General Hospital, Zoucheng, Shandong, People's Republic of China; <sup>5</sup>Department of Medical Statistics, Peking University First Hospital, Beijing, People's Republic of China

\*These authors contributed equally to this work

Correspondence: Zhijun Xi; Han Hao, Department of Urology, Peking University First Hospital, 8 Xishiku Street, Xicheng District, Beijing, 100034, People's Republic of China, Tel +86-10-83572481, Fax +86-10-66175710, Email xizhijun@hsc.pku.edu.cn; haohan1122@vip.sina.com

**Purpose:** To compare the perioperative and survival outcomes of patients over 75 years and younger patients who received radical cystectomy.

**Patients and methods:** A total of 119 patients aged  $\geq 75$  years and 488 patients aged  $< 75$  years were enrolled. All patients underwent radical cystectomy with pelvic lymph node dissection. Clinical characteristics and perioperative outcomes were compared between the two groups. Overall survival and progression-free survival were analyzed by using the Kaplan–Meier method. Cox regression analysis and logistic regression analysis were used to identify the risk factors affecting the outcomes observed.

**Results:** There was no significant difference in perioperative complications between the elderly patient group and the younger patient group ( $p = 0.349$ ). The 5-year overall survival of elderly patients was lower than that of young patients ( $p < 0.001$ ). Age  $\geq 75$  years was a risk factor for overall survival (HR = 1.69 [95% CI: 1.22–2.35];  $p = 0.002$ ) and progression-free survival (HR = 1.69 [95% CI: 1.14–2.50];  $p = 0.008$ ) for patients who received radical cystectomy but was not a poor risk factor for major complications (HR = 1.25 [95% CI: 0.47–3.31];  $p = 0.658$ ) after radical cystectomy. In addition, preoperative renal insufficiency was associated with a higher risk of major complications.

**Conclusion:** In our cohort, compared with younger patients, elderly patients aged  $\geq 75$  years had worse survival outcomes, but age  $\geq 75$  years was not a risk factor for major complications after radical cystectomy with pelvic lymph node dissection. Radical surgery should be encouraged for elderly patients who can tolerate aggressive treatments.

**Keywords:** radical cystectomy, elderly, survival outcomes, complications

## Introduction

Bladder cancer is the 10th most common and 13th most deadly cancer worldwide<sup>1</sup> and accounts for 3% of global cancer diagnoses. Age is considered to be one of the risk factors for the development of bladder cancer.<sup>2</sup> In the United States, approximately 80% of cases of bladder cancer are diagnosed in adults aged 65 or older,<sup>1</sup> and approximately 32% of diagnoses are made in those between 75 and 84.<sup>3</sup> According to data published in 2019, the incidence rate of bladder cancer in China was 5.80 per 100,000 in 2015, and the incidence increased with age.<sup>4</sup> Radical cystectomy with pelvic lymph node dissection (PLND) is the gold standard for muscle-invasive bladder cancer (MIBC) and recurrent high-grade

non-muscle-invasive bladder cancer (NMIBC).<sup>5</sup> However, its morbidity of complications was as high as 54–58% because of great surgical trauma.<sup>6–8</sup> For elderly patients, who have more comorbidities and poorer physical conditions, the morbidity of complications increases to 54–72%.<sup>3,8–10</sup>

With the improvement of surgical techniques and perioperative management, radical cystectomy could be an acceptable option for elderly patients. Although some studies have illustrated the decrement of overall survival for elderly patients receiving radical cystectomy, its survival outcomes remain controversial because of poor comparability.<sup>11–13</sup>

This study aims to compare the perioperative and survival outcomes of patients over 75 years old and younger patients receiving radical cystectomy, and the risk factors affecting the outcomes observed were identified.

## Methods

### Patient Population

In the present study, retrospective analysis of data from chart review was performed. The inclusion criteria were as follows: (1) patients underwent radical cystectomy from January 2006 to April 2017 at Peking University First Hospital; and (2) postoperative pathology indicated urothelial carcinoma of the bladder. The exclusion criteria were (1) distant metastasis ( $n=23$ ); (2) PLND not performed ( $n=164$ ); and (3) loss to follow-up ( $n=227$ ) and lack of complete survival data ( $n=5$ ). Finally, we enrolled a total of 607 consecutive patients with bladder cancer who underwent radical cystectomy plus PLND. Imaging and pathological examination were performed for all patients before the operation. The indications for radical cystectomy were T2–4aN0–xM0 tumor, high-risk and recurrent NMIBC, and BCG-resistant Tis, as well as extensive papillary disease that could not be controlled with transurethral resection of bladder tumor and intravesical therapy alone. En bloc resection of the bladder, removal of internal iliac lymph nodes, external iliac lymph nodes, obturator lymph nodes, and presacral lymph nodes were carried out for all patients, and extracorporeal urinary diversion was performed after radical cystectomy, which included ureterocutaneostomy, ileal conduit or orthotopic neobladder. This study was approved by the clinical research ethics committee of Peking University First Hospital, Beijing, China.

## Outcomes Measures

### Clinical and Pathological Data

Clinical and pathological data were obtained from chart review. Patients were categorized into two groups: young patients aged  $< 75$  years and elderly patients aged  $\geq 75$  years. Clinical characteristics, including the American Society of Anesthesiologists (ASA) score, time of operation, estimated blood loss (EBL), postoperative stay, and complication rate, were compared between the two groups. Renal function was analyzed on the basis of the last recorded preoperative serum creatinine (sCr) value, and  $sCr \geq 1.5$  mg/dL was classified as renal insufficiency. Complications within 30 days after radical cystectomy were extracted from the review of electronic patient records, and each complication was graded according to the Clavien–Dindo classification system.<sup>14,15</sup> The highest Clavien–Dindo class was summarized for each patient. Major complication was characterized as complications of Clavien–Dindo class III–V. In addition, the perioperative comorbidity assessment for elderly patients was performed using the Charlson Comorbidity Index.<sup>16</sup> Histological type, surgical margin status, lymph node yield, and the number of positive lymph nodes were collected in pathological data. The TNM staging system of bladder cancer of the American Joint Committee on Cancer Staging Manual 8th edition was used in this study.

### Oncologic Outcomes

Postoperative surveillance for all patients was performed through outpatient visits. All patients received examinations, including physical examination, laboratory measurements of blood and urine, and computer tomography scans every six months until the third year and annually thereafter. Overall survival (OS) was characterized as the period from the operation to death from any cause, cancer-specific survival (CSS) referred to the period from the operation to cancer-related death, and progression-free survival (PFS) was defined as the period from the operation to the time of recurrence of cancer.

## Statistical Analysis

For the continuous variables, the Kolmogorov-Smirnov test was used first to identify normality. The independent *t* test was used to compare the variables following a normal distribution, and variables following a nonnormal distribution were analyzed by the Mann–Whitney *U*-test. The  $\chi^2$ -test or Fisher's exact test was used to compare the disordered categorical variables, while the ordered categorical variables were analyzed by using the Mann–Whitney *U*-test. The OS and PFS of all patients were determined by using Kaplan–Meier curves. Cox regression analysis of all patients was used to identify the risk factors affecting the prognosis outcomes, adjusting for age (<75, ≥75), ASA score, pathologic stage (Ta/Tis/T1, T2, T3, T4), pathologic nodal stage (N0, N+), major complications (no, yes), chemotherapy (no, yes), and surgery margin (no, yes). The proportional hazards (PH) assumption was examined using log-minus-log plots of the hazard functions. There were no significant violations of the proportional hazards assumption. Logistic regression analysis of all patients was used to identify the risk factors associated with the major complications. Variables that showed a univariate relationship with major complications or that were considered clinically relevant were entered into the logistic regression model. In addition, to identify the factors affecting the OS of elderly patients, Cox regression analysis of elderly patients was also performed, adjusting for sex (female, male), Charlson comorbidity index (≤2, ≥3), type of urinary diversion (ileal conduit, ureterocutaneostomy), pathologic stage (Ta/Tis/T1, T2, T3, T4), pathologic nodal stage (N0, N+), major complications (no, yes), and chemotherapy (no, yes). All analyses in this study were performed using IBM SPSS version 25 (IBM Corporation, Armonk, NY, USA), and all *p* values were two-sided. *p* < 0.05 was considered to be statistically significant.

## Results

### Patient Characteristics

There were 607 patients with bladder cancer enrolled in the present study. Among them, 119 patients were aged ≥ 75 years, whose median age was 77 (interquartile range 76–80) years, and 488 patients were aged < 75 years, with a median of 62 (interquartile range 56–68) years. The median follow-up time was 52 months (interquartile range 29–78).

The perioperative and pathological outcomes are presented in Table 1. Compared with younger patients, elderly patients had higher ASA scores (*p* < 0.001) and higher clinical T stages (*p* = 0.002). However, there was no significant difference in pathological T stage or pathological nodal stage between the two groups. In addition, more elderly patients received ureterocutaneostomy (52.9% vs 14.5%, *p* < 0.001), and fewer elderly patients were treated with neoadjuvant or adjuvant chemotherapy (7.6% vs 23.2%, *p* < 0.001). The time of operation was shorter in the group of elderly patients (259 vs 313 mins, *p* < 0.001), but no significant difference was found in the postoperative complications and postoperative stay.

### Oncologic Outcomes

To identify oncologic outcomes of radical cystectomy between the two groups, Kaplan–Meier curves (see Figure 1) were used to analyze OS and PFS. The 5-year OS of elderly patients was lower than that of young patients (*p* < 0.001), which was 0.456 versus 0.628. Furthermore, the 5-year PFS was 0.620 versus 0.670, and the 5-year CSS was 0.638 versus 0.725 in the elderly patient group and young patient group, respectively, but there was no significant difference between the two groups. In addition, Cox regression analysis was performed to further verify the factors affecting the OS and PFS of all patients. As shown in Table 2, age ≥ 75 years (HR = 1.69 [95% CI: 1.22–2.35]; *p* = 0.002), higher pathological T stage (*p* < 0.001), higher pathological nodal stage (HR = 1.97 [95% CI: 1.41–2.75]; *p* < 0.001), and major complications (HR = 2.71 [95% CI: 1.56–4.70]; *p* < 0.001) were risk factors for OS in patients who underwent radical cystectomy. Age ≥ 75 years (HR = 1.69 [95% CI: 1.14–2.50]; *p* = 0.008), higher pathological T stage (*p* < 0.001), and higher pathological nodal stage (HR = 1.67 [95% CI: 1.17–2.39]; *p* = 0.005) were poor prognostic factors for PFS.

Moreover, Cox regression analysis was used to identify the factors affecting the OS of elderly patients. We found that male sex (HR = 4.01 [95% CI: 1.39–11.51]; *p* = 0.010), Charlson comorbidity index ≥ 3 (HR = 2.29 [95% CI: 1.06–4.98]; *p* = 0.036), ureterocutaneostomy (HR = 1.84 [95% CI: 1.02–3.34]; *p* = 0.043), T4 (HR = 2.63 [95% CI: 1.04–

**Table I** The Clinicopathologic Characteristics of All Patients

	<75 Years (n =488)	≥75 Years (n =119)	p
Male	428 (87.7%)	99 (83.2%)	0.192
Age	62 (56–68)	77 (76–80)	<0.001
ORC	324 (66.4%)	88 (73.9%)	0.113
LRC	164 (33.6%)	31 (26.1%)	
BMI (kg/m <sup>2</sup> )	24.0 (22.0–26.5)	23.8 (21.5–26.1)	0.176
Preoperative renal insufficiency	35 (7.2%)	10 (8.4%)	0.646
ASA score			<0.001
1	46 (9.4%)	2 (1.7%)	
2	400 (82.0%)	84 (70.6%)	
3	41 (8.4%)	32 (26.9%)	
4	1 (0.2%)	1 (0.8%)	
Charlson comorbidity index			0.107
≤2	459 (94.1%)	107 (89.9%)	
≥3	29 (5.9%)	12 (10.1%)	
Type of urinary diversion			<0.001
Ureterocutaneostomy	71 (14.5%)	63 (52.9%)	
Ileal conduit	384 (78.7%)	56 (47.1%)	
Orthotopic neobladder	33 (6.8%)	0 (0%)	
Time of operation (mins)	312.5 (245–383)	259 (216–335)	<0.001
EBL (L)	500 (200–1000)	500 (200–800)	0.626
Clavien-Dindo class			0.349
0	155 (31.8%)	33 (27.7%)	
1	6 (1.2%)	2 (1.7%)	
2	306 (62.7%)	77 (64.7%)	
3	14 (2.9%)	4 (3.4%)	
4	6 (1.2%)	3 (2.5%)	
5	1 (0.2%)	0 (0%)	
Major complication	21 (4.3%)	7 (5.9%)	0.462
Postoperative stay (days)	10 (8–14)	10 (8–17)	0.714
Clinical T stage			0.002
Ta and Tis and T1	94 (19.3%)	15 (12.6%)	
T2	209 (42.8%)	40 (33.6%)	
T3	134 (27.5%)	46 (38.7%)	
T4	51 (10.5%)	18 (15.1%)	
Pathological T stage			0.587
Ta and Tis and T1	139 (28.5%)	37 (31.1%)	
T2	176 (36.1%)	42 (35.3%)	
T3	107 (21.9%)	25 (21.0%)	
T4	66 (13.5%)	15 (12.6%)	
Pathological nodal stage			0.199
N0	395 (80.9%)	103 (86.6%)	
N1	38 (7.8%)	4 (3.4%)	
N2	43 (8.8%)	8 (6.7%)	
N3	12 (2.5%)	4 (3.4%)	

(Continued)

**Table 1** (Continued).

	<75 Years (n =488)	≥75 Years (n =119)	p
Pathologic grade			0.132
Low grade	50 (10.3%)	18 (15.1%)	
High grade	437 (89.7%)	101 (84.9%)	
Lymph node yield	11 (7–15)	8 (5–12)	0.225
Negative margin	483 (99.0%)	117 (98.3%)	0.628
Positive margin	5 (1.0%)	2 (1.7%)	
No neoadjuvant/adjuvant chemotherapy	375 (76.8%)	110 (92.4%)	<0.001
Neoadjuvant/adjuvant chemotherapy	113 (23.2%)	9 (7.6%)	

**Note:** Major complication: Clavien-Dindo class III–V.

**Abbreviations:** ORC, open radical cystectomy; LRC, laparoscopic radical cystectomy; BMI, Body Mass Index; EBL, estimated blood loss; ASA, American Society of Anesthesiologists.

6.65];  $p = 0.042$ ), and major complications (HR = 5.05 [95% CI: 1.87–13.64];  $p = 0.001$ ) were associated with poor OS outcomes for elderly patients (see in [Table 3](#)).


## Perioperative Complications

There was no significant difference in terms of perioperative complications ( $p = 0.349$ ), major complications ( $p = 0.462$ ), or estimated blood loss ( $p = 0.626$ ) between the group of elderly patients and the group of younger patients. Then, the risk factors for major complications were identified in this study. As shown in [Table 4](#), age  $\geq 75$  years was not a risk factor for major complications (OR = 1.25 [95% CI: 0.47–3.31];  $p = 0.658$ ), while preoperative renal insufficiency was associated with a higher risk of major complications (OR = 4.33 [95% CI: 1.64–11.38];  $p = 0.003$ ).

## Discussion

In this study, we focused on the perioperative and survival outcomes of patients over 75 years old and younger patients receiving radical cystectomy. The OS of elderly patients was lower than that of younger patients, while there was no significant difference in PFS between the two groups. Furthermore, age  $\geq 75$  years was one of the poor prognostic factors of OS and PFS for patients who received radical cystectomy. In addition, we found that there was no significant difference in perioperative complications between the elderly patient group and the younger patient group. Age  $\geq 75$  years was not a risk factor for major complications, while preoperative renal insufficiency was associated with a higher risk of major complications.

Radical cystectomy with PLND remains the standard treatment for nonmetastatic muscle-invasive bladder cancer.<sup>5</sup> However, as a challenging operation, its morbidity of complications is high, with a risk of 40–65%.<sup>17</sup> High therapy-induced risk of severe complications and shorter estimated life expectancy caused many elderly patients to refuse more aggressive treatment.<sup>18</sup> Past studies have suggested that only some elderly patients receive radical surgeries, and cystectomy rates decrease with advanced age.<sup>19</sup> However, aggressive surgical management of bladder cancer in elderly patients would significantly improve oncological outcomes;<sup>20</sup> in patients with T3 MIBC left untreated, the OS was less than 1 year.<sup>21</sup> In addition, with the improvement of surgical instruments and perioperative care and with the spread of minimally invasive surgery, the surgical safety of elderly patients has been greatly improved. We also compared the perioperative and oncological outcomes of patients in the first 5 years and last 5 years ([Supplementary Figure 1](#) and [Table 1](#)). For the 2012–2017 group, the time of operation, EBL, and complication rate decreased, and the postoperative stay increased, with the improvement of overall survival. A recent study from China also illustrated that elderly patients who received robot-assisted radical cystectomy showed no increased perioperative risks compared with younger patients.<sup>22</sup>



**Figure 1** Kaplan-Meier curves of OS probability ( $p < 0.001$ ), PFS ( $p = 0.131$ ), and CSS probability ( $p = 0.065$ ) in elderly patients and younger patients who received radical cystectomy and pelvic lymph node dissection. The 5-year OS of elderly patients and young patients was 0.456 versus 0.628, respectively. The 5-year PFS was 0.620 versus 0.670, and the 5-year CSS was 0.638 versus 0.725 in the elderly patient group and the young patient group, respectively.

**Abbreviations:** OS, overall survival; PFS, progression-free survival; CSS, cancer-specific survival.

**Table 2** Multivariable Cox Regression Analysis of Variables Associated with Overall Survival and Progression-Free Survival for All Patients Received Radical Cystectomy

Variable	Overall Survival			Progression-Free Survival		
	Hazard Ratio	95% CI	p	Hazard Ratio	95% CI	p
Age (years)						
<75	-	referent	-	-	referent	-
≥75	1.69	1.22–2.35	0.002	1.69	1.14–2.50	0.008
ASA			0.023			0.637
I	-	referent	-	-	referent	-
2	2.10	0.98–4.50	0.057	1.43	0.72–2.83	0.308
3	3.15	1.38–7.21	0.006	1.70	0.77–3.76	0.194
4	5.93	0.70–50.01	0.102	1.51	0.19–12.28	0.702
Pathologic stage			<0.001			<0.001
Ta and Tis and T1	-	referent	-	-	referent	-
T2	1.38	0.90–2.09	0.137	1.43	0.88–2.32	0.147
T3	2.87	1.86–4.45	<0.001	2.86	1.74–4.68	<0.001
T4	3.27	2.05–5.23	<0.001	2.60	1.51–4.46	0.001
Pathologic nodal stage						
N0	-	referent	-	-	referent	-
N+	1.97	1.41–2.75	<0.001	1.67	1.17–2.39	0.005
Major complications						
No	-	referent	-	-	referent	-
Yes	2.71	1.56–4.70	<0.001	1.18	0.48–2.91	0.713
No chemotherapy	-	referent	-	-	referent	-
Chemotherapy	1.20	0.86–1.69	0.286	2.91	2.06–4.10	<0.001
Negative margin	-	referent	-	-	referent	-
Positive margin	0.40	0.06–2.89	0.366	0.38	0.05–2.72	0.332

**Note:** Major complication: Clavien-Dindo class III–V.

**Abbreviation:** ASA, American Society of Anesthesiologists.

In some previous studies performed in the United States, France, Canada, and Australia, worse overall survival was found in elderly patients receiving radical cystectomy, but most studies defined elderly patients as patients aged >65 or 70 years.<sup>23–25</sup> Considering that the median age at diagnosis of MIBC was above 70 years and an aging population, an age cutoff of 75 years for defined elderly patients was applied in our study. We found that overall survival was worse in patients older than 75 years old, and age ≥ 75 years was associated with poor oncological outcomes. More comorbidities and poor health status could account for the worse overall survival of elderly patients. In our study, a high Charlson Comorbidity Index score was identified as a risk factor for the overall survival of elderly patients. However, not only overall survival but also progression-free survival was worse for elderly patients, although there was no significant difference in PFS, which means that there might be some other reasons responsible for the worse outcomes. Neoadjuvant chemotherapy and adjuvant chemotherapy have been proven to be significantly beneficial to survival.<sup>26,27</sup> However, the use of neoadjuvant chemotherapy and adjuvant chemotherapy decreased for elderly patients because of impaired renal function and poor performance status.<sup>28,29</sup>

Moreover, perioperative complication was found to be a risk factor for overall survival in this study, although there was no difference in terms of any perioperative complications or major complications between elderly patients and younger patients, which was the same as some previous studies.<sup>30,31</sup> The improvement of surgical instruments and advances in perioperative care may cover up the difference in complications caused by increasing age. Patel et al showed



**Table 3** Univariable/Multivariable Cox Regression Analysis of Variables Associated with Overall Survival for Elderly Patients

Variable	Univariable Cox Regression Analysis			Multivariable Cox Regression Analysis		
	Hazard Ratio	95% CI	p	Hazard Ratio	95% CI	p
Sex						
Female	–	referent	–	–	referent	–
Male	4.099	1.473–11.403	0.007	4.005	1.394–11.505	0.010
Charlson comorbidity index						
≤2	–	referent	–	–	referent	–
≥3	3.166	1.516–6.614	0.002	2.291	1.055–4.978	0.036
Type of urinary diversion						
Ileal conduit	–	referent	–	–	referent	–
ureterocutaneostomy	1.666	0.949–2.925	0.075	1.842	1.018–3.333	0.043
Pathologic stage			0.011			0.185
Ta and Tis and T1	–	referent	–	–	referent	–
T2	1.116	0.540–2.306	0.766	1.598	0.725–3.519	0.245
T3	1.705	0.792–3.669	0.172	2.150	0.922–5.013	0.077
T4	3.445	1.530–7.756	0.003	2.626	1.038–6.646	0.042
Pathologic nodal stage						
N0	–	referent	–	–	referent	–
N+	2.323	1.184–4.557	0.014	1.766	0.814–3.832	0.150
Major complications						
No	–	referent	–	–	referent	–
Yes	4.459	1.865–10.660	0.001	5.047	1.867–13.644	0.001
Chemotherapy						
Yes	–	referent	–	–	referent	–
No	2.012	0.856–4.729	0.109	1.941	0.797–4.726	0.144

**Note:** Major complication: Clavien-Dindo class III–V.

an improving trend of perioperative mortality at high volume centers.<sup>25</sup> It is necessary to refer elderly patients who need aggressive surgery to high-volume hospitals.

In our study, preoperative renal insufficiency was found to be a risk factor for major perioperative complications in elderly patients, which means that the physical status of elderly patients affects the perioperative outcomes after radical cystectomy. Hence, the potential benefits and harms from radical cystectomy should be weighed before surgery. It is important to recognize elderly patients suitable for curative treatment. With increasing age, physical function and functional reserve decreased, resulting in the decrement of tolerance for aggressive treatment. Frailty is the phenotype of declining physiologic function and loss of functional reserve across organ systems, leading to vulnerability against disease and death.<sup>32,33</sup> Frailty rates were proven to be associated with higher complications after radical cystectomy.<sup>34</sup> The incidence of frailty increased with advanced age. It is necessary to screen frailty for elderly patients before surgery in routine clinical practice. The use of some form of geriatric assessments was beneficial to select the appropriate candidates for aggressive treatments for urologists.<sup>35</sup> Geriatric assessments for elderly patients are a multidomain assessment of functional, medical, and psychosocial aspects of health. However, there was no recommendation of optimal assessment tools for clinical practice. In our center, a multidisciplinary assessment was performed for all elderly patients before radical cystectomy. Further studies of the best tools of geriatric assessments should be encouraged.

For elderly patients with nonmetastatic bladder cancer, radical cystectomy improves the survival outcomes, although it is inferior to younger patients, and the morbidity of perioperative complications is similar to that of younger patients.



**Table 4** Logistic Regression Analysis of Variables Associated with Major Complication for All Patients

	Odds Ratio	95% CI	p
Age (years)			
<75	–	referent	–
≥75	1.25	0.47–3.31	0.658
Sex			
Female	–	referent	–
Male	1.92	0.44–8.44	0.388
ASA			0.232
I	–	referent	–
2	1.68	0.22–13.21	0.620
3	4.35	0.48–39.51	0.191
4	–	–	–
Preoperative renal insufficiency			
No	–	referent	–
Yes	4.33	1.64–11.38	0.003
Type of operation			
ORC	–	referent	–
LRC	0.56	0.22–1.47	0.24
Type of urinary diversion			0.794
ureterocutaneostomy	–	referent	–
Ileal conduit	0.91	0.31–2.66	0.869
Orthotopic neobladder	1.58	0.21–11.99	0.659
Time of operation (continuous)	1.00	0.99–1.01	0.175

**Abbreviations:** ORC, open radical cystectomy; LRC, laparoscopic radical cystectomy; ASA, American Society of Anesthesiologists.

Hence, radical surgeries should be encouraged for elderly patients who could tolerate aggressive treatments. However, more studies are needed to explore better approaches to select the appropriate candidates.

There were certain limitations in our study. First, there were some inherent selection biases because of the retrospective design. Second, the elderly patients enrolled were highly selective, and patients who received radical cystectomy but not lymphonodectomy were excluded from this study, which was too idealistic. In clinical practice, to decrease operative time and reduce the risk of perioperative complications, pelvic lymph node dissection is often omitted.<sup>36</sup> Furthermore, there were no data on 30-, 60-, and 90-day mortality because of the retrospective nature of the study, which was associated with inpatient treatment.

## Conclusion

In our cohort, compared with younger patients, elderly patients aged  $\geq 75$  years had worse survival outcomes, but age  $\geq 75$  years was not a risk factor for major complications after radical cystectomy with pelvic lymph node dissection. Radical surgery should be encouraged for elderly patients who could tolerate aggressive treatments.

## Data Sharing Statement

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

## Ethics Approval

The present study was approved by the clinical research ethics committee of Peking University First Hospital (Protocol number: 2015[977]). The informed consent requirement was exempted by the ethics committee because of the retrospective design. The data used in this study were anonymized and maintained with confidentiality, and the study complied with the Declaration of Helsinki.

## Acknowledgments

This study was supported by the National Natural Science Foundation of China (grant number: 81272829).

## Funding

This study was supported by the National Natural Science Foundation of China (grant number: 81272829).

## Disclosure

The authors declare that they have no conflicts of interest.

## References

1. Saginala K, Barsouk A, Aluru JS, Rawla P, Padala SA, Barsouk A. Epidemiology of bladder cancer. *Med Sci*. 2020;8(1). doi:10.3390/medsci8010015
2. Burger M, Catto JW, Dalbagni G, et al. Epidemiology and risk factors of urothelial bladder cancer. *Eur Urol*. 2013;63(2):234–241. doi:10.1016/j.eururo.2012.07.033
3. Taylor JA 3rd, Kuchel GA. Bladder cancer in the elderly: clinical outcomes, basic mechanisms, and future research direction. *Nat Clin Pract Urol*. 2009;6(3):135–144. doi:10.1038/ncpuro1315
4. Chen W, Zheng R, Baade PD, et al. Cancer Statistics in China, 2015. *CA: a Cancer Journal for Clinicians*. 2016;66(2):115–132. doi:10.3322/caac.21338
5. Witjes JA, Bruins HM, Cathomas R, et al. EAU guidelines on muscle invasive and metastatic bladder cancer. *EAU Guidelines*; 2021.
6. Novara G, De Marco V, Aragona M, et al. Complications and mortality after radical cystectomy for bladder transitional cell cancer. *J Urol*. 2009;182(3):914–921. doi:10.1016/j.juro.2009.05.032
7. Hautmann RE, de Petriconi RC, Volkmer BG. Lessons learned from 1000 neobladders: the 90-day complication rate. *J Urol*. 2010;184(3):990–994; quiz 1235. doi:10.1016/j.juro.2010.05.037
8. Wuethrich PY, Vidal A, Burkhard FC. There is a place for radical cystectomy and urinary diversion, including orthotopic bladder substitution, in patients aged 75 and older: results of a retrospective observational analysis from a high-volume center. *Urol Oncol*. 2016;34(2):58e19–27. doi:10.1016/j.urolonc.2015.08.011
9. Farnham SB, Cookson MS, Alberts G, Smith JA Jr, Chang SS. Benefit of radical cystectomy in the elderly patient with significant co-morbidities. *Urol Oncol*. 2004;22(3):178–181. doi:10.1016/j.urolonc.2003.12.005
10. Donat SM, Siegrist T, Cronin A, Savage C, Milowsky MI, Herr HW. Radical cystectomy in octogenarians—does morbidity outweigh the potential survival benefits?. *J Urol*. 2010;183(6):2171–2177. doi:10.1016/j.juro.2010.02.015
11. Konety BR, Joslyn SA. Factors influencing aggressive therapy for bladder cancer: an analysis of data from the SEER program. *J Urol*. 2003;170(5):1765–1771. doi:10.1097/01.ju.0000091620.86778.2e
12. Prout GR, Wesley MN, Yancik R, Ries LAG, Havlik RJ, Edwards BK. Age and comorbidity impact surgical therapy in older bladder carcinoma patients. *Cancer*. 2005;104(8):1638–1647. doi:10.1002/cncr.21354
13. Koppie TM, Serio AM, Vickers AJ, et al. Age-adjusted Charlson comorbidity score is associated with treatment decisions and clinical outcomes for patients undergoing radical cystectomy for bladder cancer. *Cancer*. 2008;112(11):2384–2392. doi:10.1002/cncr.23462
14. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg*. 2004;240(2):205–213. doi:10.1097/01.sla.0000133083.54934.ae
15. Clavien PA, Barkun J, de Oliveira ML, et al. The Clavien-Dindo classification of surgical complications: five-year experience. *Ann Surg*. 2009;250(2):187–196. doi:10.1097/SLA.0b013e3181b13ca2
16. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40(5):373–383. doi:10.1016/0021-9681(87)90171-8
17. Challacombe BJ, Bochner BH, Dasgupta P, et al. The role of laparoscopic and robotic cystectomy in the management of muscle-invasive bladder cancer with special emphasis on cancer control and complications. *Eur Urol*. 2011;60(4):767–775. doi:10.1016/j.eururo.2011.05.012
18. Fonteyne V, Ost P, Bellmunt J, et al. Curative treatment for muscle invasive bladder cancer in elderly patients: a systematic review. *Eur Urol*. 2018;73(1):40–50. doi:10.1016/j.eururo.2017.03.019
19. Fedeli U, Fedewa SA, Ward EM. Treatment of muscle invasive bladder cancer: evidence from the National Cancer Database, 2003 to 2007. *J Urol*. 2011;185(1):72–78. doi:10.1016/j.juro.2010.09.015
20. Hollenbeck BK, Miller DC, Taub D, et al. Aggressive treatment for bladder cancer is associated with improved overall survival among patients 80 years old or older. *Urology*. 2004;64(2):292–297. doi:10.1016/j.urol.2004.03.034
21. Marshall VF, Whitmore WF Jr. The surgical treatment of cancers of the urinary bladder. *Cancer*. 1956;9(3):609–619. doi:10.1002/1097-0142(195605/06)9:3<609::aid-cncr2820090327>3.0.co;2-m

22. Xie S, Zhao Z, Feng B, et al. A comparative study of perioperative and survival outcomes of robot-assisted radical cystectomy in patients over 80 and under 80 years old. *World J Surg Oncol*. 2021;19(1):202. doi:10.1186/s12957-021-02312-4
23. Guillotreau J, Miocinovic R, Gamé X, et al. Outcomes of laparoscopic and robotic radical cystectomy in the elderly patients. *Urology*. 2012;79(3):585–590. doi:10.1016/j.urolgy.2011.11.042
24. Fairey AS, Kassouf W, Aprikian AG, et al. Age  $\geq$  80 years is independently associated with survival outcomes after radical cystectomy: results from the Canadian Bladder Cancer Network Database. *Urol Oncol*. 2012;30(6):825–832. doi:10.1016/j.urolonc.2011.07.014
25. Patel MI, Bang A, Gillatt D, Smith DP. Contemporary radical cystectomy outcomes in patients with invasive bladder cancer: a population-based study. *BJU Int*. 2015;116(Suppl 3):18–25. doi:10.1111/bju.13152
26. Yin M, Joshi M, Meijer RP, et al. Neoadjuvant chemotherapy for muscle-invasive bladder cancer: a systematic review and two-step meta-analysis. *oncologist*. 2016;21(6):708–715. doi:10.1634/theoncologist.2015-0440
27. Leow JJ, Martin-Doyle W, Rajagopal PS, et al. Adjuvant chemotherapy for invasive bladder cancer: a 2013 updated systematic review and meta-analysis of randomized trials. *Eur Urol*. 2014;66(1):42–54. doi:10.1016/j.eururo.2013.08.033
28. Leveridge MJ, Siemens DR, Mackillop WJ, et al. Radical cystectomy and adjuvant chemotherapy for bladder cancer in the elderly: a population-based study. *Urology*. 2015;85(4):791–798. doi:10.1016/j.urolgy.2014.12.027
29. Reardon ZD, Patel SG, Zaid HB, et al. Trends in the use of perioperative chemotherapy for localized and locally advanced muscle-invasive bladder cancer: a sign of changing tides. *Eur Urol*. 2015;67(1):165–170. doi:10.1016/j.eururo.2014.01.009
30. Fairey A, Chetner M, Metcalfe J, et al. Associations among age, comorbidity and clinical outcomes after radical cystectomy: results from the Alberta Urology Institute radical cystectomy database. *J Urol*. 2008;180(1):128–34; discussion 134. doi:10.1016/j.juro.2008.03.057
31. Schmid M, Rink M, Traumann M, et al. Evidence from the “PROspective MulticEnTer RadIcal Cystectomy Series 2011 (PROMETRICS 2011)” study: how are preoperative patient characteristics associated with urinary diversion type after radical cystectomy for bladder cancer?. *Ann Surg Oncol*. 2015;22(3):1032–1042. doi:10.1245/s10434-014-4029-3
32. Clegg A, Young J, Iliffe S, Rikkert MO, Rockwood K. Frailty in elderly people. *Lancet*. 2013;381(9868):752–762. doi:10.1016/S0140-6736(12)62167-9
33. Grimberg DC, Shah A, Molinger J, et al. Assessments of frailty in bladder cancer. *Urol Oncol*. 2020;38(9):698–705. doi:10.1016/j.urolonc.2020.04.036
34. Chappidi MR, Kates M, Patel HD, et al. Frailty as a marker of adverse outcomes in patients with bladder cancer undergoing radical cystectomy. *Urol Oncol*. 2016;34(6):256.e1–6. doi:10.1016/j.urolonc.2015.12.010
35. Gadzinski AJ, Psutka SP. Risk stratification metrics for bladder cancer: comprehensive geriatric assessments. *Urol Oncol*. 2020;38(9):725–733. doi:10.1016/j.urolonc.2020.01.003
36. Abdollah F, Sun M, Shariat SF, et al. The importance of pelvic lymph node dissection in the elderly population: implications for interpreting the 2010 National Comprehensive Cancer Network practice guidelines for bladder cancer treatment. *J Urol*. 2011;185(6):2078–2084. doi:10.1016/j.juro.2011.02.022

## Cancer Management and Research

Dovepress

### Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>