

Efficacy of Moxibustion Smoke for Stage I Post-Stroke Shoulder-Hand Syndrome: Protocol for a Multi-Center, Single-Blind Randomized Sham-Controlled Trial [Response to Letter]

Xiaonan Meng¹, Liping Wang¹, Chunying Li¹, Sen Gao², Haikuo Yu³, Lufen Zhang⁴, Jie Sun⁵

¹Acupuncture Department, Beijing Huguosi TCM Hospital, affiliated with Beijing University of Chinese Medicine, Beijing, People's Republic of China; ²Rehabilitation Department, Beijing Huguosi TCM Hospital, affiliated with Beijing University of Chinese Medicine, Beijing, People's Republic of China; ³Rehabilitation Department, Xuanwu Hospital Capital Medical University, Beijing, People's Republic of China; ⁴School of Acupuncture-Moxibustion and Tuina, Beijing University of Chinese Medicine, Beijing, People's Republic of China; ⁵Rehabilitation Department of Integrated Chinese and Western Medicine, Beijing Xiaotangshan Hospital, Beijing, People's Republic of China

Correspondence: Jie Sun, Rehabilitation Department of Integrated Chinese and Western Medicine, Beijing Xiaotangshan Hospital, No. 390 Spring Road, Changping District, Beijing 102211, People's Republic of China, Email jiesun1109@gmail.com

Dear editor

In a letter to editor titled “Efficacy of Moxibustion Smoke for Stage 1 Post-Stroke Shoulder-Hand Syndrome: Protocol for a Multi-Center, Single-Blind Randomized Sham-Controlled Trial [Letter]”, Chen et al kindly mentioned some issues to discuss and offered some suggestions for our study protocol. As a result of their constructive recommendations, this study will be carried out in an even more accurate and scientific manner. Therefore, we'd be delighted to answer their questions.

To begin with, the visual analogue scale (VAS) for pain was designated as the primary outcome because pain contributes to the further exacerbation of regional edema and motor dysfunction in patients with shoulder-hand syndrome.¹⁻³ Given that a VAS for pain is a subjective rating, this measure could be affected by psychological factors. In order to mitigate such effects and maintain adequate blinding, we have referenced some additional prior publications.^{4,5} Before treatments, the patients will be informed that they will receive either traditional moxibustion with a variety of levels of moxibustion smoke concentration (MSC) and heat sensation, or minimal moxibustion with relatively lower levels of MSC and heat sensation, both of which are associated with positive outcomes in clinical practice. A blinding assessment will be conducted upon the completion of all interventions. Our intention is that priming participants with this information in advance will help to counteract the effects of any type of bias, as well as the Hawthorne effect, that might otherwise occur.

The moxibustion apparatus applied in this study, the Happy-all Moxibustion Device (manufactured by the Chongqing Happy-all Medical Device Co., Ltd), has already been thoughtfully designed with considerations to prevent burns to the regional skin by either adjusting the distance between the moxa and skin, or the number of open ventilation holes in the wall of the apparatus. In our study, since indirect moxibustion will be utilized with the goal of producing a comfortable sensation of radiant heat,⁶ the dermal temperature in the region receiving moxibustion will also be recorded using a non-contact thermometer (BOSCH GIS500, Germany) after the moxa has been lit. According to our pre-test assessment, the regional dermal temperature ranges from approximately 38°C to 45°C, a spectrum that is not expected to pose a significant risk of burns to the skin. The application of an ointment barrier could potentially influence clinical effects through reducing the amount of radiant heat that is locally absorbed, as well as permeation of the local acupoints by moxibustion smoke, so we have decided against utilizing such a barrier. However, we feel that the above measures are more than adequate to ensure a safe and clinically effective study.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Cacchio A, De Blasis E, Necozone S, Di Orio F, Santilli V. Mirror therapy for chronic complex regional pain syndrome type 1 and stroke. *N Engl J Med*. 2009;361(6):634–636. doi:10.1056/NEJMc0902799
2. Treede RD, Rief W, Barke A, et al. Chronic pain as a symptom or a disease: the IASP Classification of Chronic Pain for the International Classification of Diseases (ICD-11). *Pain*. 2019;160(1):19–27. doi:10.1097/j.pain.0000000000001384
3. Peng L, Zhang C, Zhou L, Zuo HX, He XK, Niu YM. Traditional manual acupuncture combined with rehabilitation therapy for shoulder hand syndrome after stroke within the Chinese healthcare system: a systematic review and meta-analysis. *Clin Rehabil*. 2018;32(4):429–439. doi:10.1177/0269215517729528
4. Sun Y, Liu Y, Liu B, et al. Efficacy of Acupuncture for Chronic Prostatitis/Chronic Pelvic Pain Syndrome: a Randomized Trial. *Ann Intern Med*. 2021;174(10):1357–1366. doi:10.7326/M21-1814
5. Witt C, Brinkhaus B, Jena S, et al. Acupuncture in patients with osteoarthritis of the knee: a randomised trial. *Lancet*. 2005;366(9480):136–143. doi:10.1016/S0140-6736(05)66871-7
6. Sun C, Li Y, Kuang J, Liang X, Wu J, Ji C. The thermal performance of biological tissue under moxibustion therapy. *J Therm Biol*. 2019;83:103–111. doi:10.1016/j.jtherbio.2019.05.018

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Journal of Pain Research 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Journal of Pain Research editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Journal of Pain Research

Dovepress

Publish your work in this journal

The Journal of Pain Research is an international, peer reviewed, open access, online journal that welcomes laboratory and clinical findings in the fields of pain research and the prevention and management of pain. Original research, reviews, symposium reports, hypothesis formation and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-pain-research-journal>

<https://doi.org/10.2147/JPR.S367017>