

# Impact of the *ABCB1* Drug Resistance Gene on the Risk Factors of Patients with COVID-19 and Its Relationship with the Drugs Used

Irma Olarte Carrillo<sup>1</sup>, Christian Omar Ramos Peñafiel<sup>2</sup>, Iveth Mendoza Salas<sup>2</sup>, Rafael Cerón Maldonado<sup>1</sup>, Anel Irais García Laguna<sup>1</sup>, Adrián De la Cruz Rosas<sup>1</sup>, Carlos Martínez Murrillo<sup>2</sup>, Adolfo Martínez Tovar<sup>1</sup>

<sup>1</sup>Laboratorio de Biología Molecular, Servicio de Hematología, Hospital General de México, “Dr. Eduardo Liceaga”, Ciudad de México, Mexico;

<sup>2</sup>Servicio de Hematología, Hospital General de México, “Dr. Eduardo Liceaga”, Ciudad de México, Mexico

Correspondence: Adolfo Martínez Tovar, Laboratorio de Biología Molecular, Servicio de Hematología, Hospital General de México “Dr. Eduardo Liceaga”, Ciudad de México, Mexico, Email [mtadolfo73@hotmail.com](mailto:mtadolfo73@hotmail.com)

**Objective:** In the last two years progress was made in molecular, physio pathological understanding and the form of transmission of COVID-19, and different therapeutic strategies have been explored to deal with the situation of the pandemic. However, the evaluation of certain genes that participate in the metabolism and transport of these drugs has not been fully explored. A lack of response to treatment and a lower survival have been observed that may be due to the presence of the *ABCB1* drug resistance gene. Our research group analyzed whether the expression levels of the *ABCB1* gene are associated with comorbidities, treatments, overall survival and risk of death in patients with severe COVID-19.

**Methods:** The expression levels of the *ABCB1* gene were analyzed by RT-qPCR in 61 patients diagnosed with COVID-19. The association between the levels of expression, the risk variables and different treatments were determined by the Chi-Square test and the Fisher’s exact test. Global Survival (GS) was determined by the Kaplan–Meier method. The impact of high levels of expression and the risk of death was performed by odds ratio.

**Results:** The different risk variables showed that patients with either high or absent levels of *ABCB1* gene expression presented a greater risk of death (OR 3.08, 95%, CI 1.02–9.26) as well as need for ventilatory support (OR 2.8, 95%, CI 0.98 –8.5). Patients with diabetes and COVID-19, treated with metformin, were associated with a lower risk of death (OR 1.11, 95%, CI 0.38–3.22). OS with respect to high or absent levels of expression of the *ABCB1* gene was lower.

**Conclusion:** High levels or null expression of the *ABCB1* gene are associated with a higher risk of death or progression of the disease, the use of metformin in patients with COVID-19 confers a lower risk of death.

**Keywords:** *ABCB1*, SARS-CoV2, drug transporter genes

## Background

The Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) is the virus that causes the disease by Coronavirus 2019 (COVID-19).<sup>1</sup> SARS-CoV-2 was reported for the first time in the Chinese city of Wuhan and has spread rapidly around the world through human interactions. WHO declared the COVID-19 outbreak as a pandemic on March 12, 2020, as a result of the increase in the infection rate outside of China.<sup>2</sup>

SARS-CoV-2 belongs taxonomically to the family of the Coronavirus, and the Sarbecovirus subgenus, which contains other species that cause human diseases from mild to severe.<sup>3</sup>

The mortality rate of COVID-19 is lower than that of severe acute respiratory syndrome (SARS) and the Middle East respiratory syndrome (MERS), both by coronaviruses.<sup>4</sup> However, COVID-19 is highly contagious and spreads rapidly, as it can be transmitted by respiratory droplets. The common symptoms observed in patients with COVID-19 are fever, cough, severe headache, myalgia and fatigue.<sup>5,6</sup>

The incubation period of a COVID-19 infection is usually 1 to 14 days and can be extended up to 24 days.<sup>7</sup> The health conditions of most patients with COVID-19 are mild, but they can become serious, especially among the elderly or with underlying diseases, such as chronic or cardiovascular pulmonary disease.<sup>8</sup> Various factors such as male gender, diabetes, hypertension, alterations in blood counts and hypoalbuminemia have been associated with a rapid progression towards respiratory failure.<sup>9,10</sup>

Infection by other viruses such as influenza A or B can cause symptoms similar to those of COVID-19, which can complicate the distinction between this and other diseases, especially during the flu season.<sup>11,12</sup>

The current gold standard for the diagnosis of COVID-19 is based on the detection of the virus RNA in respiratory samples of nasopharyngeal swabs or bronchial aspirates by means of a molecular test by quantitative reverse transcription polymerase chain reaction (RT-qPCR).<sup>13</sup> This is a sensitive and specific method for detecting SARS-CoV-2, with different diagnostic protocols, including target primer sequences available in the WHO public database.<sup>14</sup>

The test has its limitations since false negatives can be presented if the amount of viral genome is insufficient; also, it is estimated that the incubation period of COVID-19 is 5 days, and false negative results are common within 7 days afterwards the infection. In addition, the analyzes and studies of imaging and artificial intelligence (AI) have helped achieve a more accurate diagnosis, as well as the evaluation of patients during the monitoring of the therapy and treatment of COVID-19.<sup>15,16</sup>

Treatments in patients with COVID-19 are based on different combinations of medicines and treatment schemes (antiviral and/or immunotherapy and/or steroids) with a high potential for interactions,<sup>17,18</sup> and one of the least evaluated effects is either the expression or the blocking of different drug-resistance genes. Some examples are the families of *ABC* membrane transporters (ATP-Binding Cassette) that include 49 different types of transporters, including the P-glycoprotein (P-gp).<sup>19,20</sup> Their main function is cell detoxification, and are distributed along the hemato-encephalic barrier, placenta, kidney and liver.<sup>21–23</sup> The role of the expression of *ABC* genes in COVID-19 is still unknown, and various drugs used for COVID-19 may show some interaction with the *ABC* family, such as azithromycin (*ABCB1*), lopinavir/ritonavir (*ABCC2*) or antimalarial agents like chloroquine.<sup>24,25</sup> In Mexico, the mortality associated with COVID-19 in severely ill patients is higher than expected due to a large extent by related risk factors.<sup>26</sup> Our working group has evaluated the impact of *ABCB1* overexpression in patients with acute leukemia, which causes a decrease in survival.<sup>27</sup> The main objective of this study is to establish the impact of comorbidities, drugs used and the expression of *ABCB1* gene on the overall survival and risk of death in COVID-19 patients.

## Methods

### Study Design

This was an exploratory type study, performed during the second wave of COVID-19 in our country. All patients were in severe condition of intensive therapy, treated with different treatment schemes. Sixty-one patients with a diagnosis of COVID-19 admitted to the Intensive Care Unit of the General Hospital of Mexico “Dr. Eduardo Liceaga” were included, and their informed consent was obtained for the molecular analysis of the *ABCB1* gene. The study was conducted in accordance with the Declaration of Helsinki. The project was approved by the Research, Ethics and Biosecurity Committees of the General Hospital of Mexico “Dr. Eduardo Liceaga” D1/15/103/03/57.

### Isolation of Peripheral Blood Mononuclear Cells by Density Gradient

EDTA-Anticoagulated peripheral blood was obtained by venopunction. Mononuclear cells were isolated by density gradient centrifugation of an identical volume of peripheral blood and Lymphoprep™ (density  $1.077 \pm 0.001$  g/mL, Axis-Shield PoC AS, Oslo, Norway). The preparations were centrifuged in a Sorvall ST8 centrifuge (Thermo Fisher Scientific, Leicestershire, UK) at 300 xg for 15 minutes, the cell fractions were homogenized in TRIzol reagent and stored at  $-80^{\circ}\text{C}$  until RNA extraction.

### RNA Extraction and cDNA Synthesis

Total RNA isolation was performed according to the manufacturer’s protocol (Invitrogen corporation, Carlsband, CA). The isolated RNA was dissolved in nuclease free water (Promega, Madison, WI). Complementary DNA (cDNA) was synthesized from 2 µg of RNA using Oligo (dT) and with MML-V RT enzyme (Promega Biotech AB).

## Levels of Expression of *ABCB1* Gene

The expression levels of the *ABCB1* gene were determined by RT-qPCR using gene-specific TaqMan™ hydrolysis probe (Hs01069047, Applied Biosystems, Inc. Foster City, CA, USA). Normalization was made with the endogen *GUSB* gene (Hs00939627). Each sample was analyzed in triplicate.

Relative expression levels were calculated using method  $2^{-\Delta\Delta Ct}$ . The high and low expression cut-points were determined according to the average values observed in a previous study.<sup>27</sup>

## Criterion and Type of Treatment

The treatment of COVID-19 included the combination of high-flow oxygen with the best available therapy, according to the local recommendations and the time of recruitment. 8.2% (n = 5) initiated treatment with hydroxychloroquine, 44.3% (n = 27) with metformin hydrochloride, 36.1% (n = 22) with azithromycin, 55.7% (n = 34) with atorvastatin and 95.2% (n = 58) with low molecular weight heparin, all during the hospital period. 49.2% (n = 30) required ventilatory support, which modified the treatment according to each particular case, being mainly vasopressor drugs, sedation, analgesia, parenteral nutrition and anticoagulation therapy. No patient received Tocilizumab, convalescent plasma, nor plasma replacement. Only an individual in critical condition received Baricitinib for a short period of time. The average of hospital stay was 38.78 days, 57.4% died due to complications associated with COVID-19.

## Statistical Analysis

The association between the levels of expression, the risk variables and different treatments were determined by the Chi-Square test and the Fisher exact test. Global survival (GS) was determined by the Kaplan–Meier method. The impact of high levels of expression and the risk of death was performed through the odds ratio. The statistical software used was SPSS version 22.0.

## Results

A total of 61 patients diagnosed with COVID-19 were included, 59% corresponded to the male gender (n = 36), obesity was the most frequent comorbidity (45.9%, n = 28), followed by non-insulin-dependent diabetes (36.1%, n = 22) and hypertension (29.5%, n = 18). One-third of the patients had a history of smoking (32.8%) and about half of alcoholism (42.6%). The average of leukocytes when admitted was  $13.1 \times 10^3 /\mu\text{L}$  ( $3.1\text{--}55.4 \times 10^3 /\mu\text{L}$ ), 54.1% (n = 33) had leukocytosis at the time of diagnosis. Neutrophilia was reported by 77% (n = 47) and 65.6% (n = 40) counted with lymphopenia. Eosinopenia was constant, unlike thrombocytopenia (minor counts of  $100 \times 10^3 /\mu\text{L}$ ) which was infrequent (3.3%). 74.3% (n = 52) had a neutrophil-lymphocyte index greater than 7 with an average of 18.1 (2–123). The average of bilirubin was 0.42 mg/dL, 15.7% (n = 11) had bilirubin levels greater than 1.5 mg/dL. The average of albumin was 3.3 g/dL (1.7–4.1 g/dL), with 57.1% (n = 40) of individuals with hypoalbuminemia. No patient presented renal failure when admitted (Table 1). When we analyzed the expression levels of the *ABCB1* gene in patients with COVID-19, we found that 37.7% (n = 23) had high levels, 36% (n = 22) low levels and for 26.2% (n = 16) It was absent. When evaluating the different risk factors such as diabetes, hypertension, smoking, alcohol consumption and overweight with respect to the expression levels of the *ABCB1* gene, the results showed that alcohol consuming patients were associated with groups of patients expressing either high levels or absence of the *ABCB1* gene ( $p \leq 0.05$ , 95% CI). No association of risk factors was observed with respect to low levels of gene expression (Table 2).

When analyzing the effect of the treatments used in patients with metformin, atorvastatin, hydroxychloroquine, azithromycin and anticoagulants, the results showed that groups of patients with high and negative levels of expression of the *ABCB1* gene were associated with the use of atorvastatin and metformin ( $p \leq 0.05$ , 95% CI), unlike low levels of gene expression (Table 3).

The different risk variables showed that patients with high levels and with absence of expression of the *ABCB1* gene presented greater risk of death (Odds ratio [OR] 2.8, 95% CI 0.98–8.5) and need for ventilatory support (OR 3.08, 95% CI 1.02–9.26). Other risk factors such as male gender (OR 2.54, 95% CI 0.89–7.27) and age (OR 5.19, 95% CI 1.48–18.2) were associated with risk of death.

**Table 1** General Characteristics in COVID-19 Patients

	Mean	Range
Age, years	49	16–80
Sex; %		
Male, 59%	36	–
Female, 41%	25	–
Hemoglobin, g/dL	14.02	2.90–18.60
Hematocrit, %	38.89	8.90–55.30
WBC count, 10 <sup>3</sup> /μL	13.19	3.10–55.40
Neutrophils, 10 <sup>3</sup> /μL	19.90	1.80–98.00
Lymphocytes, 10 <sup>3</sup> /μL	4.14	0.10–31.12
Monocytes, 10 <sup>3</sup> /μL	0.51	0.10–6.20
Platelets, 10 <sup>3</sup> /μL	272	13–652
MPV, fL	7.93	5.10–10.80
DB, mg/dL	0.46	0–1.10
TB, mg/dL	0.43	0–3.80
PT, 10 <sup>3</sup> /μL	6.41	4.30–8.20
ALB, g/dL	3.32	1.75–4.10
ALT, u/L	75	9–265
AST, u/L	96	12–788
LDH, u/L	542	214–1518

**Abbreviations:** MPV, mean platelet volume; DB, direct bilirubin; TB, total bilirubin; PT, prothrombin; ALB, albumin; ALT, alanine transferase; AST, aspartate transferase; LDH, lactate dehydrogenase.

**Table 2** ABCB1 Levels Associated with Risk Factors in COVID-19 Patients

Risk Factors		ABCB1 Levels	
		Low n (%)	High and Absent n (%)
DM	No	15 (24.6)	24 (39.3)
	Yes	7 (11.5)	15 (24.6)
SAH	No	18 (29.5)	25 (41.0)
	Yes	4 (6.6)	14 (23.0)
Smoking	No	12 (19.7)	29 (47.5)
	Yes	10 (16.4)	10 (16.4)
Alcoholism	No	8 (13.1)	27 (44.3)
	Yes	14 (23.0)	12 (19.7)*
Overweight	No	11 (18.0)	22 (36.1)
	Yes	11 (18.0)	17 (27.9)

**Note:** Chi-square test, \*p<0.05.

**Abbreviations:** DM, diabetes mellitus; SAH, systemic arterial hypertension.

It is important to note that patients with diabetes which were treated with metformin due to their condition were associated with a lower risk of death (OR 1.11.95% CI 0.38–3.22), as well as patients who presented low levels of expression of the *ABCB1* gene (OR 0.34, 95% CI 0.11–1.08) [Table 4](#).

Finally, in the overall survival (OS) at 78 days of follow-up with respect to the expression levels of the *ABCB1* gene, it was found that the high and absent expression groups showed an average of 11.18 days (9–13.37) and an OS of 33.3%, while those who presented low levels had an average of 67.42 days (56.3–78.7) and an OS of 59.1% (p = 0.00, Log Rank Test). [Figure 1](#).

**Table 3** Treatment in COVID-19 Patients Associated with ABCBI Levels

Treatment		ABCBI Levels	
		Low n (%)	High and Absent n (%)
Metformin	No	8 (13.1)	26 (42.6)
	Yes	14 (23.0)	13 (21.3)*
Atorvastatin	No	5 (8.2)	22 (36.1)
	Yes	17 (27.9)	17 (27.9)*
Hydroxychloroquine	No	19 (31.1)	37 (60.7)
	Yes	3 (4.9)	2 (3.3)
Azithromycin	No	13 (21.3)	26 (42.6)
	Yes	9 (14.8)	13 (21.3)
Anticoagulant	No	2 (3.3)	1 (1.6)
	Yes	20 (32.8)	38 (62.3)

Note: Chi-square test, \*p<0.05.

**Table 4** Factors Associated with Mortality and Respiratory Support

	Mechanical Ventilation			Death		
	OR	(95% CI)	p	OR	(95% CI)	p
Male genre	3.33	(1.139, 9.782)	0.027*	2.54	(0.890, 7.278)	0.081
DM	2.51	(0.856, 7.391)	0.093	1.11	(0.386, 3.220)	0.838
Age over 55 years	1.62	(0.561, 4.729)	0.369	5.194	(1.481, 18.216)	0.010*
Hypertension	1.43	(0.475, 4.344)	0.520	1.11	(0.366, 03.370)	0.852
ABCBI levels						
High /Absent	3.08	(1.024, 9.262)	0.045	2.88	(0.981, 8.503)	0.054
Low	0.32	(0.108, 0.976)	0.045	0.34	(0.117, 1.018)	0.054


Note: Chi-square test, \*p<0.05.

Abbreviations: DM, diabetes mellitus; OR, odds ratio; CI, confidence interval.

## Discussion

The COVID-19 has currently become the leading cause of death worldwide, causing a great economic and social crisis. The WHO has reported more than 462 million confirmed cases of SARS-CoV2 infection around the world with more than ~6 million deaths.<sup>28</sup> Adverse drug reactions (ADR) associated with current treatments for COVID-19 are one of the main causes of death in patients with multiple comorbidities.<sup>29,30</sup> In a recent pharmacovigilance study conducted in Spain, an incidence 4.75 times higher to present ADR in severe patients with COVID-19 compared to patients with some other condition and treated with azithromycin (8.4%), dexamethasone (7.6%), lopinavir-ritonavir (7.4%) and chloroquine (CQ)/hydroxychloroquine (HCQ) (6.9%) was reported.<sup>31</sup> Another study conducted in Brazil with 402 patients with COVID-19 indicated that chloroquine (CQ) (OR = 5.4, 95% CI: 1.9–15.6) and hydroxychloroquine (HCQ) (OR = 2.1 95% CI: 1.2–3.6) were the only drugs associated with severe ADR.<sup>32</sup>

At present there are no specific therapeutic recommendations for the treatment of COVID-19.<sup>29,33</sup> The treatment based on chloroquine, a drug used as immunosuppressor in lupus and rheumatoid arthritis, has shown an in vitro antiviral effect against influenza and different types of coronavirus, including COVID-19.<sup>34,35</sup> When analyzing the available evidence, its use did not impact mortality, but it did increase the risk of complications, mainly cardiovascular.<sup>36</sup> Also, a drug widely used is azithromycin, which contributes to preventing the entry of the virus to the cell via CD147 with potential antiviral activity, and it is generally used in combination with hydroxychloroquine.<sup>37,38</sup> Other drugs with immunomodulatory effect are statins and metformin hydrochloride, its use became a daily practice in our country due to the high prevalence of diabetes and obesity.<sup>39,40</sup> It has been shown that a group of patients with COVID-19 suffering



**Figure 1** Overall survival at 78 days of follow up with respect to the expression levels of the *ABCB1* gene.

from chronic type 2 diabetes, being treated with metformin, increased their survival, which was associated with a lower risk of death.<sup>41–43</sup> In conjunction with anticoagulation and various anti-inflammatory drugs, the combination of these drugs was a fundamental part of the treatment for COVID-19 during the first months of pandemic in Mexico.

We evaluated the expression levels of *ABCB1* genes in COVID-19 patients and found high levels of expression, 37.7%. Several studies have shown that the expression levels of the mRNA are increased by the response to substrates of the *ABCB1* pump, and within the substrates described are azithromycin and atorvastatin, as well as self-medicated antibiotics.<sup>44</sup> The use of these drugs would explain the increase in levels, unlike the healthy population. It is important to highlight that patients who presented high levels of expression mostly had associated comorbidities such as diabetes and hypertension, and most of them were of the male gender.

It has been previously demonstrated that the expression of the *ABCB1* gene is altered in different metabolic diseases including diabetes.<sup>45</sup>

When analyzing the risk factors of patients with COVID-19 associated with high levels of expression and absence of expression of *ABCB1* gene, alcohol consumption was significant, this is consistent with the work of Haaman et al 2019, in which prolonged exposure to drug abuse, ethanol and cocaine alters the expression of *ABCB1* mRNA.<sup>46</sup> When the impact of the different treatments was evaluated, we observed increased expression levels in patients who did not consume metformin. Metformin inhibits the expression of the NF- $\kappa$ B nuclear factor, inducing the apoptosis of hepatocytes and inhibiting the expression of the *ABCB1* gene.<sup>47,48</sup>

Finally, the group of patients with either high levels or absence of expression of the *ABCB1* gene showed a lower survival rate, unlike the group of patients with low levels. These results are very similar to other studies that explain that the increased expression of these genes is related to the expulsion of the drugs used in the treatments, and, therefore, the failure thereof and the absence of expression would be causing an intoxication with the drugs by not protecting against xenobiotic agents in the different tissues, which leads to an organic failure, and this agrees with the reported in other ailments.<sup>49,50</sup>

## Conclusion

The alteration in the expression of the *ABCB1* gene is associated with an increased risk of death or progression of the disease; the failure of previous treatments can be explained by the interaction of the drugs with the flow pump, modifying its distribution in different tissues.

## Abbreviations

SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; COVID-19, coronavirus disease 2019; ABCB1, ATP-binding cassette B1; P-gp, P-glycoprotein; GAPDH, glyceraldehyde 3-phosphate dehydrogenase; *NF-kB*, nuclear factor *kappa* B; OS, overall survival.

## Data Sharing Statement

The datasets used and/or analysed during the current study available from the corresponding author on reasonable request.

## Ethics Approval and Consent to Participate

Written informed consent was obtained from all subjects and the study was approved by the Ethics Committee of the Hospital General Mexico “Dr Eduardo Liceaga”.

## Acknowledgments

The authors thank Direccion de Investigación Hospital General de Mexico.

## Author contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

## Funding

This work was supported by the Direccion de Investigacion Hospital General Mexico “Dr Eduardo Liceaga” (DI/20/204/04/31, DI/20/204/04/41).

## Disclosure

The authors declare that they have no competing interests.

## References

1. World Health Organization. WHO director-general's opening remarks at the media briefing on COVID-19 - 11 March 2020. World Health Organization; 2020. Available from: <https://www.who.int/director-general/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19—11-march-2020>. Accessed April 26, 2020.
2. Khailany RA, Safdar M, Ozaslan M. Genomic characterization of a novel SARS-CoV-2. *Gene Report*. 2020;19:100682. doi:10.1016/j.genrep.2020.100682
3. Walls AC, Park YJ, Tortorici MA, Wall A, McGuire AT, Veesler D. Structure, function, and antigenicity of the SARS-CoV-2 spike glycoprotein. *Cell*. 2020;181(2):281–292.e6. doi:10.1016/j.cell.2020.02.058



4. Yüce M, Filiztekin E, Özkaya KG. COVID-19 diagnosis -a review of current methods. *Biosens Bioelectron.* 2021;172:112752. doi:10.1016/j.bios.2020.112752
5. Zhu N, Zhang D, Wang W, et al.; China Novel Coronavirus Investigating and Research Team. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med.* 2020;382(8):727–733. doi:10.1056/NEJMoa2001017
6. Ye Z, Zhang Y, Wang Y, Huang Z, Song B. Chest CT manifestations of new coronavirus disease 2019 (COVID-19): a pictorial review. *Eur Radiol.* 2020;30(8):4381–4389. doi:10.1007/s00330-020-06801-0
7. Hui DS, Azhar EI, Madani TA, et al. The continuing 2019-nCoV epidemic threat of novel coronaviruses to global health - The latest 2019 novel coronavirus outbreak in Wuhan, China. *Int J Infect Dis.* 2020;91:264–266. doi:10.1016/j.ijid.2020.01.009
8. Gardner W, States D, Bagley N. The coronavirus and the risks to the elderly in long-term care. *J Aging Soc Policy.* 2020;32(4–5):310–315. doi:10.1080/08959420.2020.1750543
9. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet.* 2020;395(10229):1054–1062. doi:10.1016/S0140-6736(20)30566-3
10. Guan WJ, Liang WH, Zhao Y, et al.; China Medical Treatment Expert Group for COVID-19. Comorbidity and its impact on 1590 patients with COVID-19 in China: a nationwide analysis. *Eur Respir J.* 2020;55(5):2000547. doi:10.1183/13993003.00547-2020
11. Zhang J, Wang X, Jia X, et al. Risk factors for disease severity, unimprovement, and mortality in COVID-19 patients in Wuhan, China. *Clin Microbiol Infect.* 2020;26(6):767–772. doi:10.1016/j.cmi.2020.04.012
12. Zhao W, Zhong Z, Xie X, Yu Q, Liu J. Relation between chest CT findings and clinical conditions of coronavirus disease (COVID-19) pneumonia: a multicenter study. *AJR Am J Roentgenol.* 2020;214(5):1072–1077. doi:10.2214/AJR.20.22976
13. Böger B, Fachi MM, Vilhena RO, Cobre AF, Tonin FS, Pontarolo R. Systematic review with meta-analysis of the accuracy of diagnostic tests for COVID-19. *Am J Infect Control.* 2021;49(1):21–29. doi:10.1016/j.ajic.2020.07.011
14. Theagarajan LN. Group testing for COVID-19: how to stop worrying and test more. Cornell University; 2020.
15. Lauer SA, Grantz KH, Bi Q, et al. The incubation period of coronavirus disease 2019 (COVID-19) from publicly reported confirmed cases: estimation and application. *Ann Intern Med.* 2020;172(9):577–582. doi:10.7326/M20-0504
16. Naseem M, Akhund R, Arshad H, Ibrahim MT. Exploring the potential of artificial intelligence and machine learning to combat COVID-19 and existing opportunities for LMIC: a scoping review. *J Prim Care Community Health.* 2020;11:2150132720963634. doi:10.1177/2150132720963634
17. Arora G, Joshi J, Mandal RS, Shrivastava N, Virmani R, Sethi T. Artificial intelligence in surveillance, diagnosis, drug discovery and vaccine development against COVID-19. *Pathogens.* 2021;10(8):1048. doi:10.3390/pathogens10081048
18. Gautret P, Lagier JC, Parola P, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *Int J Antimicrob Agents.* 2020;56(1):105949. doi:10.1016/j.ijantimicag.2020.105949
19. Liu X. Transporter-mediated drug-drug interactions and their significance. *Adv Exp Med Biol.* 2019;1141:241–291. doi:10.1007/978-981-13-7647-4\_5
20. Choi YH, Yu AM. ABC transporters in multidrug resistance and pharmacokinetics, and strategies for drug development. *Curr Pharm Des.* 2014;20(5):793–807. doi:10.2174/138161282005140214165212
21. Pohl PC, Klafke GM, Júnior JR, Martins JR, da Silva Vaz I Jr, Masuda A. ABC transporters as a multidrug detoxification mechanism in *Rhipicephalus (Boophilus) microplus*. *Parasitol Res.* 2012;111(6):2345–2351. doi:10.1007/s00436-012-3089-1
22. Sun M, Kingdom J, Baczyk D, Lye SJ, Matthews SG, Gibb W. Expression of the multidrug resistance P-glycoprotein, (ABCB1 glycoprotein) in the human placenta decreases with advancing gestation. *Placenta.* 2006;27(6–7):602–609. doi:10.1016/j.placenta.2005.05.007
23. Shen S, Zhang W. ABC transporters and drug efflux at the blood-brain barrier. *Rev Neurosci.* 2010;21(1):29–53. doi:10.1515/revneuro.2010.21.1.29
24. Senarathna SM, Page-Sharp M, Crowe A. The interactions of p-glycoprotein with antimalarial drugs, including substrate affinity, inhibition and regulation. *PLoS One.* 2016;11(4):e0152677. doi:10.1371/journal.pone.0152677
25. Takahashi T, Luzum JA, Nicol MR, Jacobson PA. Pharmacogenomics of COVID-19 therapies. *NPJ Genomic Med.* 2020;5(1):35. doi:10.1038/s41525-020-00143-y
26. Olivas-Martínez A, Cárdenas-Fragoso JL, Jiménez JV, et al. In-hospital mortality from severe COVID-19 in a tertiary care center in Mexico city: causes of death, risk factors and the impact of hospital saturation. *PLoS One.* 2021;16(2):e0245772. doi:10.1371/journal.pone.0245772
27. Olarte Carrillo I, Ramos Peñafiel C, Miranda Peralta E, et al. Clinical significance of the ABCB1 and ABCG2 gene expression levels in acute lymphoblastic leukemia. *Hematology.* 2017;22(5):286–291. doi:10.1080/10245332.2016.1265780
28. World Health Organization. WHO coronavirus (COVID-19) dashboard. (s. f.). With vaccination data. Available from: <https://covid19.who.int/>. Accessed March 18, 2022.
29. Lemaitre F, Solas C, Grégoire M, et al.; French Society of Pharmacology, Therapeutics (SFPT), the International Association of Therapeutic Drug Monitoring, Clinical Toxicology (IATDMCT). Potential drug-drug interactions associated with drugs currently proposed for COVID-19 treatment in patients receiving other treatments. *Fundam Clin Pharmacol.* 2020;34(5):530–547. doi:10.1111/fcp.12586
30. Sun J, Deng X, Chen X, et al. Incidence of adverse drug reactions in COVID-19 patients in China: an active monitoring study by hospital pharmacovigilance system. *Clin Pharmacol Ther.* 2020;108(4):791–797. doi:10.1002/cpt.1866
31. Ramírez E, Urroz M, Rodríguez A, et al. Incidence of suspected serious adverse drug reactions in corona virus disease-19 patients detected by a pharmacovigilance program by laboratory signals in a tertiary hospital in Spain: cautionary data. *Front Pharmacol.* 2020;11:602841. doi:10.3389/fphar.2020.602841
32. Melo J, Duarte EC, Moraes MV, Fleck K, Silva A, Arrais P. Adverse drug reactions in patients with COVID-19 in Brazil: analysis of spontaneous notifications of the Brazilian pharmacovigilance system. Reações adversas a medicamentos em pacientes com COVID-19 no Brasil: análise das notificações espontâneas do sistema de farmacovigilância brasileiro. *Cad Saude Publica.* 2021;37(1):e00245820. doi:10.1590/0102-311X00245820
33. Biswas M, Roy DN. Potential clinically significant drug-drug interactions of hydroxychloroquine used in the treatment of COVID-19. *Int J Clin Pract.* 2021;75(11):e14710. doi:10.1111/ijcp.14710
34. Ibáñez S, Martínez O, Valenzuela F, Silva F, Valenzuela O. Hydroxychloroquine and chloroquine in COVID-19: should they be used as standard therapy? *Clin Rheumatol.* 2020;39(8):2461–2465. doi:10.1007/s10067-020-05202-4
35. Ebina-Shibuya R, Namkoong H, Horita N, et al. Hydroxychloroquine and chloroquine for treatment of coronavirus disease 19 (COVID-19): a systematic review and meta-analysis of randomized and non-randomized controlled trials. *J Thorac Dis.* 2021;13(1):202–212. doi:10.21037/jtd-20-2022



36. Celotto S, Veronese N, Barbagallo M, et al. An umbrella review of systematic reviews with meta-analyses evaluating positive and negative outcomes of hydroxychloroquine and chloroquine therapy. *Int J Infect Dis.* 2021;103:599–606. doi:10.1016/j.ijid.2020.12.018
37. Damle B, Vourvahis M, Wang E, Leaney J, Corrigan B. Clinical pharmacology perspectives on the antiviral activity of azithromycin and use in COVID-19. *Clin Pharmacol Ther.* 2020;108(2):201–211. doi:10.1002/cpt.1857
38. Fantini J, Chahinian H, Yahi N. Synergistic antiviral effect of hydroxychloroquine and azithromycin in combination against SARS-CoV-2: what molecular dynamics studies of virus-host interactions reveal. *Int J Antimicrob Agents.* 2020;56(2):106020. doi:10.1016/j.ijantimicag.2020.106020
39. Singh AK, Singh R. Is metformin ahead in the race as a repurposed host-directed therapy for patients with diabetes and COVID-19? *Diabetes Res Clin Pract.* 2020;165:108268. doi:10.1016/j.diabres.2020.108268
40. El-Arabey AA, Abdalla M. Metformin and COVID-19: a novel deal of an old drug. *J Med Virol.* 2020;92(11):2293–2294. doi:10.1002/jmv.25958
41. Gao Y, Liu T, Zhong W, et al. Risk of metformin in patients with type 2 diabetes with COVID-19: a preliminary retrospective report. *Clin Transl Sci.* 2020;13(6):1055–1059. doi:10.1111/cts.12897
42. Scheen AJ, Marre M, Thivolet C. Prognostic factors in patients with diabetes hospitalized for COVID-19: findings from the CORONADO study and other recent reports. *Diabetes Metab.* 2020;46(4):265–271. doi:10.1016/j.diabet.2020.05.008
43. Holman N, Knighton P, Kar P, et al. Risk factors for COVID-19-related mortality in people with type 1 and type 2 diabetes in England: a population-based cohort study. *Lancet Diabetes Endocrinol.* 2020;8(10):823–833. doi:10.1016/S2213-8587(20)30271-0
44. Liu L, Liu XD. Alterations in function and expression of ABC transporters at blood-brain barrier under diabetes and the clinical significances. *Front Pharmacol.* 2014;5:273. doi:10.3389/fphar.2014.00273
45. Yan R, Luo J, He X, Li S. Association between ABC family variants rs1800977, rs4149313, and rs1128503 and susceptibility to type 2 diabetes in a Chinese Han population. *J Int Med Res.* 2020;48(8):300060520941347. doi:10.1177/0300060520941347
46. Hammad AM, Alasmari F, Sari Y, Scott Hall F, Tiwari AK. Alcohol and cocaine exposure modulates ABCB1 and ABCG2 transporters in male alcohol-preferring rats. *Mol Neurobiol.* 2019;56(3):1921–1932. doi:10.1007/s12035-018-1153-2
47. Parissenti AM, Gannon BR, Villeneuve DJ, Kirwan-rhude AF, Chadderton A, Glück S. Lack of modulation of MDR1 gene expression by dominant inhibition of cAMP-dependent protein kinase in doxorubicin-resistant mcf-7 breast cancer cells. *Int J Cancer.* 1999;82(6):893–900.
48. Ramos-Peñafiel C, Olarte-Carrillo I, Cerón-Maldonado R, et al. Effect of metformin on the survival of patients with ALL who express high levels of the ABCB1 drug resistance gene. *J Transl Med.* 2018;16(1):245. doi:10.1186/s12967-018-1620-6
49. Delou J, Vignal GM, Índio-do-Brasil V, et al. Loss of constitutive ABCB1 expression in breast cancer associated with worse prognosis. *Breast Cancer.* 2017;9:415–428. doi:10.2147/BCTT.S131284
50. Litviakov NV, Cherdyntseva NV, Tsyganov MM, et al. Deletions of multidrug resistance gene loci in breast cancer leads to the down-regulation of its expression and predict tumor response to neoadjuvant chemotherapy. *Oncotarget.* 2016;7(7):7829–7841. doi:10.18632/oncotarget.6953

## Infection and Drug Resistance

Dovepress

### Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>