Neurocognitive impairment after off-pump and on-pump coronary artery bypass graft surgery – an Iranian experience

Mehdi Farhoudi¹
Kaveh Mehrvar²
Abbas Afrasiabi³
Rezayat Parvizi³
Ahmad Ali Khalili³
Babak Nasiri³
Khosrow Hashemzadeh³
Kamyar Ghabili⁴

¹Neurosciences Research Center, Imam Reza Hospital, ²Razi Hospital, ³Cardiovascular Research Center, ⁴Tuberculosis and Lung Disease Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Coronary artery bypass graft (CABG) surgery is one of the most commonly performed surgical procedures worldwide, and it may be accompanied by postoperative neurocognitive impairment. Although this complication has been attributed to the use of cardiopulmonary bypass, it is still a matter of debate whether the switch from on-pump to off-pump technique affects the cognitive function.

Objective: The aim of this study was to compare the impact of the on-pump and off-pump techniques on neurocognitive impairment in low-risk CABG surgery groups.

Methods: In a descriptive and analytic study, 201 CABG patients with left-ventricular ejection fraction >30%, and without cardiac arrhythmia were enrolled. Before the elective operation, all patients underwent neurological examination and neurocognitive test, Mini-Mental State Examination (MMSE). Two months following the operation, both on- and off-pump, the patients were re-examined by MMSE to detect any neurocognitive impairment.

Results: Out of 154 patients included in the study, 95 (61.6%) and 59 (38.3%) patients were in off-pump and on-pump groups, respectively. Mean age of the patients was 57.17 ± 9.82 years. A 2-month postoperative neurocognitive impairment was detected among 17 patients of on-pump group (28.8%) and in 28 cases of off-pump group (29.4%) ($P = 0.54$). The mean postoperative MMSE scores were not comparable between groups (25.01 ± 4.49 in off-pump group versus 23.73 ± 4.88 in on-pump group, $P = 0.09$).

Conclusion: The present study revealed that in low-risk patients undergoing CABG surgery, either the techniques of on-pump or off-pump did not differ regarding the neurocognitive outcome 2 months after the procedure.

Keywords: neurocognitive impairment, off-pump, on-pump, coronary artery bypass graft surgery

Introduction

Coronary artery bypass graft (CABG) surgery is one of the most commonly performed surgical procedures worldwide, intended to treat ischemic heart disease and alleviate angina pectoris.^{1–3} Despite advances in the surgical procedure, neurological injury is still an important complication after CABG surgery and is found in two types.⁴ The first type of neurological injury occurs in 3%–6% of the cases with manifestations such as stroke, transient ischemic attack, and coma. On the other hand, impairment of cognitive function including defects of attention, concentration, short-term memory, fine motor function, and speed of mental and motor responses is found in the milder type second.^{4,5} With an incidence of 20%–80%, neurocognitive impairment after cardiac surgery has been largely attributed to the use of cardiopulmonary bypass (CPB).^{6–9}

Correspondence: Kamyar Ghabili
Tuberculosis and Lung Disease Research Center, Tabriz University of Medical Sciences, Tabriz, Iran
Tel +989144106136
Fax +984113361120
Email kghabili@gmail.com

CPB increases the permeability of the blood–brain barrier and generates microemboli, which may affect cognitive function.^{10,11} Recently, cardiac stabilizer devices have been developed to facilitate CABG without using CPB (off-pump CABG surgery). Although diminished cerebral embolism in off-pump CABG has been highlighted,¹² it is still controversial whether the switch from on-pump to off-pump technique fully alleviates the cognitive dysfunction. Minor improvements in cognitive function shortly after operation,^{4,12,13} better cognitive outcome,^{14,15} and no improvement in neurocognitive status after off-pump CABG have been hitherto reported in off-pump CABG in comparison to the on-pump method.¹⁶ Due to the aforementioned controversy, the present study was designed to compare the impact of the on-pump and off-pump techniques on neurocognitive impairment in low-risk CABG groups.

Methods

In a descriptive, analytic, and prospective follow-up study, 201 candidates for CABG surgery were enrolled at Madani Heart Hospital in Tabriz, Iran. Inclusion criteria were elective CABG, lack of concomitant cardiac arrhythmia, and left-ventricular ejection fraction (LVEF) >30%. Patients undergoing emergency CABG or CABG combined with other surgical procedures such as valve replacement or carotid endarterectomy and those with history of neurological and psychiatric diseases were excluded from the study. Before operation, all patients underwent neurological and neurocognitive (Mini-Mental State Examination, MMSE) examination.

Patients were assigned to on- or off-pump groups. CABG was carried out through median sternotomy. The anesthetic technique was standardized and all patients received the similar protocol. Follow-up study was performed up to

2 months after operation by neurological and neurocognitive examination. A decline of ≥ 1 score in MMSE after surgery was defined as cognitive dysfunction. All data are presented as mean values $\pm$ SD. Statistical analysis was performed with SPSS for Windows (version 13.0; SPSS Inc., Chicago, IL) using independent samples *t*-test, paired samples *t*-test, and chi-squared test, whenever appropriate. A *P* value of less than 0.05 was considered to represent significant difference. The Research Vice-Chancellor of Tabriz Medical Faculty approved this study.

Results

Out of 201 patients enrolled in the study, 47 patients refused or were unable to return for follow-up. Data from 154 patients were therefore analyzed: 95 (61.6%) in off-pump group and 59 (38.3%) in on-pump group. Out of 154 patients, 123 (79.8%) were male and 31 (20.1%) were female. The mean age of patients was 57.17 ± 9.82 years ranging from 31–82. Mean LVEF was $50.65 \pm 9.41\%$. The patients' baseline characteristics are shown in Table 1.

There was no intraoperative accident. Two patients in on-pump group (2.5%) and nine patients in off-pump group (7.4%) had postoperative myocardial infarction ($P > 0.05$). In each group, there were two postoperative strokes. There was only one death in off-pump group; the cause was severe cardiac disease and postoperative cardiac arrest, and no neurological complication was found. In on-pump group, there was no significant difference between the mean preoperative and postoperative MMSE scores (23.24 ± 5.03 versus 23.73 ± 4.88 , $P = 0.16$). However, in off-pump group, the mean 2-month postoperative MMSE score was significantly higher than the mean preoperative MMSE score (25.01 ± 4.49 versus 24.18 ± 4.51 , $P = 0.007$). On the basis of the MMSE results, cognitive impairment was detected in

Table 1 Baseline participants' characteristics and pre- and postoperative MMSE scores (n = 201)

	Off-pump group (n = 95)	On-pump group (n = 59)	P value
Gender (male:female)	77:18	46:13	0.39 ¹
Age (years)	56.34 ± 9.00	58.51 ± 10.97	0.18 ²
Coronary vessel disease	–	–	0.09 ¹
1-vessel disease	4 (4.2%)	0 (0)	–
2-vessel disease	34 (35.7%)	15 (25.4%)	–
≥ 3 -vessel disease	57 (60%)	43 (72.8%)	–
Diabetes mellitus	24 (25.2%)	7 (11.8%)	0.03 ^{1*}
History of hypertension	40 (42.1%)	30 (50.8%)	0.21 ¹
Preoperative MMSE score	24.18 ± 4.51	23.24 ± 5.03	0.23 ²
Postoperative MMSE score	25.01 ± 4.49	23.73 ± 4.88	0.09 ²
Postoperative neurocognitive impairment	28 (29.4%)	17 (28.8%)	0.54 ¹

Notes: ¹Chi-square test; ²Independent samples *t*-test. *Statistically significant ($P < 0.05$).

Abbreviation: MMSE, Mini-Mental State Examination.

21.2% of the cases in on-pump group and 23.1% in off-pump ($P = 0.54$, Table 1).

Discussion

The results of this study showed no significant difference in postoperative neurocognitive impairment between on-pump and off-pump low-risk CABG surgery groups. This is in agreement with the findings of some previous studies that failed to detect any difference in neurocognitive impairment between on- and off-pump techniques.^{16–20} Jensen et al recently found similar cognitive outcome 3 months after off-pump and on-pump surgery in a group of elderly with high risk for CABG surgery.¹⁸ Interestingly, Ernest et al revealed no difference in neurocognitive state at 2 and 6 months following CABG surgery between both groups, with the exception that fewer off-pump patients showed impairment on verbal fluency at 6 months.²⁰

In contrast to the findings of the present study, some investigations have demonstrated favorable results for patients who underwent off-pump CABG surgery. Murkin et al²¹ and Zamvar et al¹⁵ reported less neurocognitive impairment among off-pump patients at first week postoperatively, while Motallebzadeh et al found better neurocognitive function at discharge from hospital but no difference at 6 months postoperatively.⁴ Stroobant et al concluded contrarily, ie, no difference at discharge but better function at 6 months postoperatively for off-pump group.²² However, van Dijk et al showed that low-risk patients avoiding the use of CPB had no effect on 5-year cognitive outcomes.²³ Altogether, such a matter of controversy in neurocognitive impairment between on- and off-pump CABG surgery has been attributed to the variations in the tests used, the time points of assessment, the definition of impairment, and the statistical methods used for comparing groups.^{4,24} On the other hand, factors other than CPB may be responsible for cognitive decline, such as anesthesia and the generalized inflammatory response that is associated with major surgical procedures.^{4,25}

The present study has several limitations. First, assessment of neurocognitive function was done only by one neuropsychological test (MMSE) and ≥ 1 score decline in MMSE was defined as cognitive impairment. Second, MMSE is a simple screening instrument which might not be reliable in detection of subtle cognitive impairment. Therefore, a more rigorous set of neuropsychological tests is required in further investigations. Third, MMSE scores were not corrected for educational attainment and age. This might explain quite low preoperative MMSE scores in the present study.²⁶ Fourth, the results of this investigation cannot be extrapolated to

older patients with more advanced coronary artery disease or higher preoperative risks. On the other hand, large sample size along with the follow-up rate of 76% which is comparable to the previous studies is among the advantages of the current study.^{4,7,27}

On the basis of these findings, we conclude that in low-risk patients undergoing CABG surgery, avoiding the use of CPB had no effect on cognitive outcome 2 months after the procedure.

Acknowledgment

This study was financially supported by a grant from the Research Vice-Chancellor office of Tabriz University of Medical Sciences.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Yoon BW, Bae HJ, Kang DW, et al. Intracranial cerebral artery disease as a risk factor for central nervous system complications of coronary artery bypass graft surgery. *Stroke*. 2001;32:94–99.
2. Farhoudi M, Afrasiabi A, Tarzamni MK, Khoshnan M, Arami MA. Transcranial and carotid Doppler study in coronary artery bypass graft patient. *Neurosciences*. 2004;9:186–189.
3. Farhoudi M, Parvizi R, Bilehjani E, Tarzamni MK, Mehrvar K, Safaiyan A. Preoperative transcranial and carotid Doppler study in coronary artery bypass graft patients. *Neurosciences*. 2007;12:42–45.
4. Motallebzadeh R, Bland JM, Markus HS, Kaski JC, Jahangiri M. Neurocognitive function and cerebral emboli: randomized study of on-pump versus off-pump coronary artery bypass surgery. *Ann Thorac Surg*. 2007;83:475–482.
5. Roach GW, Kanchuger M, Mangano CM, et al. Adverse cerebral outcomes after coronary bypass surgery. Multicenter Study of Perioperative Ischemia Research Group and the Ischemia Research and Education Foundation Investigators. *N Engl J Med*. 1996;335:1857–1863.
6. Vingerhoets G, van Nooten G, Vermassen F, de Soete G, Jannes C. Short-term and long-term neuropsychological consequences of cardiac surgery with extracorporeal circulation. *Eur J Cardiothorac Surg*. 1997;11:424–431.
7. van Dijk D, Jansen EW, Hijman R, et al. Cognitive outcome after off-pump and on-pump coronary artery bypass graft surgery: a randomized trial. *JAMA*. 2002;287:1405–1412.
8. Newman MF, Mathew JP, Grocott HP, et al. Central nervous system injury associated with cardiac surgery. *Lancet*. 2006;368:694–703.
9. van Dijk D, Keizer AM, Diephuis JC, Durand C, Vos LJ, Hijman R. Neurocognitive dysfunction after coronary artery bypass surgery: a systematic review. *J Thorac Cardiovasc Surg*. 2000;120:632–639.
10. Harris DN, Bailey SM, Smith PL, Taylor KM, Oatridge A, Bydder GM. Brain swelling in first hour after coronary artery bypass surgery. *Lancet*. 1993;342:586–587.
11. Lund C, Hol PK, Lundblad R, et al. Comparison of cerebral embolization during off-pump and on-pump coronary artery bypass surgery. *Ann Thorac Surg*. 2003;76:765–770.
12. Bowles BJ, Lee JD, Dang CR, et al. Coronary artery bypass performed without the use of cardiopulmonary bypass is associated with reduced cerebral microemboli and improved clinical results. *Chest*. 2001;119:25–30.

13. Lee JD, Lee SJ, Tsushima WT, et al. Benefits of off-pump bypass on neurologic and clinical morbidity: a prospective randomized trial. *Ann Thorac Surg.* 2003;76:18–25.
14. Diegeler A, Hirsch R, Schneider F, et al. Neuromonitoring and neurocognitive outcome in off-pump versus conventional coronary bypass operation. *Ann Thorac Surg.* 2000;69:1162–1166.
15. Zamvar V, Williams D, Hall J, et al. Assessment of neurocognitive impairment after off-pump and on-pump techniques for coronary artery bypass graft surgery: prospective randomised controlled trial. *BMJ.* 2002;325:1268.
16. Lloyd CT, Ascione R, Underwood MJ, Gardner F, Black A, Angelini GD. Serum S-100 protein release and neuropsychologic outcome during coronary revascularization on the beating heart: a prospective randomized study. *J Thorac Cardiovasc Surg.* 2000;119:148–154.
17. Lund C, Sundet K, Tennøe B, et al. Cerebral ischemic injury and cognitive impairment after off-pump and on-pump coronary artery bypass grafting surgery. *Ann Thorac Surg.* 2005;80:2126–2131.
18. Jensen BO, Hughes P, Rasmussen LS, Pedersen PU, Steinbrüchel DA. Cognitive outcomes in elderly high-risk patients after off-pump versus conventional coronary artery bypass grafting: a randomized trial. *Circulation.* 2006;113:2790–2795.
19. Vedin J, Nyman H, Ericsson A, Hylander S, Vaage J. Cognitive function after on or off pump coronary artery bypass grafting. *Eur J Cardiothorac Surg.* 2006;30:305–310.
20. Ernest CS, Worcester MU, Tatoulis J, et al. Neurocognitive outcomes in off-pump versus on-pump bypass surgery: a randomized controlled trial. *Ann Thorac Surg.* 2006;81:2105–2114.
21. Murkin JM, Newman SP, Stump DA, Blumenthal JA. Statement of consensus on assessment of neurobehavioral outcomes after cardiac surgery. *Ann Thorac Surg.* 1995;59:1289–1295.
22. Stroobant N, van Nooten G, Bellegheem Y, Vingerhoets G. Short-term and long-term neurocognitive outcome in on-pump versus off-pump CABG. *Eur J Cardiothorac Surg.* 2002;22:559–564.
23. van Dijk D, Spoor M, Hijman R, et al. Cognitive and cardiac outcomes 5 years after off-pump vs on-pump coronary artery bypass graft surgery. *JAMA.* 2007;297:701–708.
24. Mahanna EP, Blumenthal JA, White WD, et al. Defining neuropsychological dysfunction after coronary artery bypass grafting. *Ann Thorac Surg.* 1996;61:1342–1347.
25. Moller JT, Cluitmans P, Rasmussen LS, et al. Long-term postoperative cognitive dysfunction in the elderly ISPOCD1 study. ISPOCD investigators. International Study of Post-Operative Cognitive Dysfunction. *Lancet.* 1998;351:857–861.
26. Crum RM, Anthony JC, Bassett SS, Folstein MF. Population-based norms for the Mini-Mental State Examination by age and educational level. *JAMA.* 1993;269:2386–2391.
27. Newman MF, Kirchner JL, Phillips-Bute B, et al. Longitudinal assessment of neurocognitive function after coronary-artery bypass surgery. *N Engl J Med.* 2001;344:395–402.

Neuropsychiatric Disease and Treatment

Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS, and is the official

Submit your manuscript here: <http://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>

Dovepress

journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.